



American Society of Hematology
2021 L Street NW, Suite 900,
Washington, DC 20036
Phone: 202-776-0544 | Fax 202-776-0545
editorial@hematology.org

Recombinant *Erwinia* Asparaginase (JZP458) in Acute Lymphoblastic Leukemia: Results from the Phase 2/3 AALL1931 Study

Tracking no: BLD-2022-016923R1

Luke Maese (Huntsman Cancer Institute, University of Utah, Primary Children's Hospital, United States) Mignon Loh (Seattle Children's Hospital, the Ben Town Center for Childhood Cancer Research, University of Washington, Seattle, WA., United States) Mi Rim Choi (Jazz Pharmaceuticals, United States) Tong Lin (Jazz Pharmaceuticals, United States) Etsuko Aoki (Jazz Pharmaceuticals, United States) Michelle Zanette (Jazz Pharmaceuticals, United States) Shirali Agarwal (Jazz Pharmaceuticals, United States) Robert Iannone (Jazz Pharmaceuticals, United States) Jeffrey Silverman (Jazz Pharmaceuticals, United States) Lewis Silverman (Dana-Farber Cancer Institute, United States) Elizabeth Raetz (New York University, United States) Rachel Rau (Texas Children's Cancer and Hematology Center, Baylor College of Medicine, United States)

Abstract:

AALL1931 (NCT04145531), a phase 2/3 study conducted in collaboration with the Children's Oncology Group, investigated the efficacy and safety of JZP458 (asparaginase *erwinia chrysanthemi* (recombinant)-rywn), a recombinant *Erwinia* asparaginase derived from a novel expression platform, in patients with acute lymphoblastic leukemia/lymphoblastic lymphoma who developed hypersensitivity/silent inactivation to *E. coli*-derived asparaginases. Each dose of a pegylated *E. coli*-derived asparaginase remaining in patients' treatment plan was substituted by six doses of intramuscular (IM) JZP458 on Monday/Wednesday/Friday (MWF). Three dosing regimens were evaluated: Cohort 1a, 25 mg/m² MWF; Cohort 1b, 37.5 mg/m² MWF; Cohort 1c, 25/25/50 mg/m² MWF. The primary and key secondary efficacy outcomes were the proportion of patients maintaining adequate nadir serum asparaginase activity (NSAA ≥ 0.1 IU/mL) at 72 hours as well as at 48 hours during the first treatment course, respectively. 167 patients were enrolled: Cohort 1a (n=33), 1b (n=83), and 1c (n=51). Mean SAA levels (IU/mL) at 72-hr were 0.16 for Cohort 1a, 0.33 for Cohort 1b, and 0.47 for Cohort 1c; and 0.45, 0.88, and 0.66 at 48-hr, respectively. The proportion of patients achieving NSAA ≥ 0.1 IU/mL at 72-hr and 48-hr in Cohort 1c was 90% (44/49) and 96% (47/49), respectively. Simulated data from a population pharmacokinetic model matched the observed data well. Grade 3/4 treatment-related adverse events (TRAEs) occurred in 86/167 (51%) patients; TRAEs leading to discontinuation included pancreatitis (6%), allergic reactions (5%), increased alanine aminotransferase (1%), and hyperammonemia (1%). Study results demonstrate that IM JZP458 at 25/25/50 mg/m² MWF is efficacious and has a safety profile consistent with those of other asparaginases.

Conflict of interest: COI declared - see note

COI notes: Luke Maese: Served on an advisory board and speakers bureau for Jazz Pharmaceuticals; Mignon L. Loh: Nothing to disclose; Mi Rim Choi: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Tong Lin: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Etsuko Aoki: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Michelle Zanette: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Shirali Agarwal: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Robert Iannone: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Jeffrey A. Silverman: was an employee of Jazz Pharmaceuticals at the time of the study and holds stock ownership and/or stock options in Jazz Pharmaceuticals; Lewis B. Silverman: Served on scientific advisory boards for Jazz Pharmaceuticals and Servier Pharmaceuticals; Elizabeth A. Raetz: Received institutional research funding from Pfizer and serves on a Data and Safety Monitoring Board for BMS; Rachel E. Rau: Served on advisory boards for Jazz Pharmaceuticals and Servier Pharmaceuticals

Preprint server: No;

Author contributions and disclosures: Conception and design: Luke Maese, Mignon L. Loh, Mi Rim Choi, Robert Iannone, Jeffrey A. Silverman, Tong Lin, Elizabeth A. Raetz, and Rachel E. Rau; collection and assembly of data: Luke Maese, Mignon L. Loh, Mi Rim Choi, Tong Lin, Etsuko Aoki, Michelle Zanette, Shirali Agarwal, Jeffrey A. Silverman, Lewis B. Silverman, Elizabeth A. Raetz, and Rachel E. Rau; data analysis and interpretation: all authors; manuscript writing: all authors; final approval of manuscript: all authors; accountable for all aspects of the work: all authors.

Non-author contributions and disclosures: Yes; Medical writing and editorial support were provided by Austin Horton, PhD, of Cello Health Communications/SciFluent Communications, Inc., and were financially supported by Jazz Pharmaceuticals.

Agreement to Share Publication-Related Data and Data Sharing Statement: All relevant data, including the study protocol, are provided within the manuscript and supporting files.

Clinical trial registration information (if any): ClinicalTrials.gov Identifier: NCT04145531

Recombinant *Erwinia* Asparaginase (JZP458) in Acute Lymphoblastic Leukemia: Results from the Phase 2/3 AALL1931 Study

Luke Maese,¹ Mignon L. Loh,² Mi Rim Choi,³ Tong Lin,³ Etsuko Aoki,³ Michelle Zanette,⁴ Shirali Agarwal,³ Robert Iannone,⁴ Jeffrey A. Silverman,³ Lewis B. Silverman,⁵ Elizabeth A. Raetz,⁶ Rachel E. Rau⁷

¹Huntsman Cancer Institute, University of Utah, Primary Children's Hospital, Salt Lake City, UT, USA; ²Ben Towne Center for Childhood Cancer Research, Seattle Children's Research Institute and Department of Pediatrics, Seattle Children's Hospital, University of Washington, Seattle, WA, USA; ³Jazz Pharmaceuticals, Palo Alto, CA, USA; ⁴Jazz Pharmaceuticals, Philadelphia, PA, USA; ⁵Dana-Farber Cancer Institute, Boston Children's Hospital, Boston, MA, USA; ⁶New York University Langone Health, New York, NY, USA; ⁷Texas Children's Cancer and Hematology Center, Baylor College of Medicine, Houston, TX, USA.

Corresponding Author:

Luke Maese
Huntsman Cancer Institute
University of Utah
Primary Children's Hospital
100 N. Mario Capecchi Drive
Salt Lake City, UT 84113
Email: Luke.Maese@hsc.utah.edu
Phone: 801-662-4700

Running Title:

Efficacy and safety of JZP458 in ALL/LBL

Word counts

Abstract: 249/250 words

Text: 3266/4000 words

Figure/tables: 6/7 items

References: 26/100

Key Points:

- Intramuscular JZP458 at 25/25/50 mg/m² M/W/F was effective in ALL/LBL patients with a safety profile similar to other asparaginases.
- JZP458 overcomes one of the most significant drug shortage issues facing patients with ALL/LBL.

Explanation of Novelty

Asparaginases are part of the cornerstone of ALL therapy, but global shortages of *Erwinia*-derived asparaginases have resulted in reduced availability. To overcome this, JZP458, a recombinant form of *Erwinia chrysanthemi* asparaginase derived from a novel *Pseudomonas fluorescens* expression platform was developed. This study showed that intramuscular JZP458 is efficacious and safe and provides a reliable treatment option for patients who develop hypersensitivity to *E. coli*-derived asparaginases.

Abstract

AALL1931 (NCT04145531), a phase 2/3 study conducted in collaboration with the Children's Oncology Group, investigated the efficacy and safety of JZP458 (asparaginase *erwinia chrysanthemi* (recombinant)-rywn), a recombinant *Erwinia* asparaginase derived from a novel expression platform, in patients with acute lymphoblastic leukemia/lymphoblastic lymphoma who developed hypersensitivity/silent inactivation to *E. coli*-derived asparaginases. Each dose of a pegylated *E. coli*-derived asparaginase remaining in patients' treatment plan was substituted by six doses of intramuscular (IM) JZP458 on Monday/Wednesday/Friday (MWF). Three dosing regimens were evaluated: Cohort 1a, 25 mg/m² MWF; Cohort 1b, 37.5 mg/m² MWF; Cohort 1c, 25/25/50 mg/m² MWF. The primary and key secondary efficacy outcomes were the proportion of patients maintaining adequate nadir serum asparaginase activity (NSAA ≥ 0.1 IU/mL) at 72 hours as well as at 48 hours during the first treatment course, respectively. 167 patients were enrolled: Cohort 1a (n=33), 1b (n=83), and 1c (n=51). Mean SAA levels (IU/mL) at 72-hr were 0.16 for Cohort 1a, 0.33 for Cohort 1b, and 0.47 for Cohort 1c; and 0.45, 0.88, and 0.66 at 48-hr, respectively. The proportion of patients achieving NSAA ≥ 0.1 IU/mL at 72-hr and 48-hr in Cohort 1c was 90% (44/49) and 96% (47/49), respectively. Simulated data from a population pharmacokinetic model matched the observed data well. Grade 3/4 treatment-related adverse events (TRAEs) occurred in 86/167 (51%) patients; TRAEs leading to discontinuation included pancreatitis (6%), allergic reactions (5%), increased alanine aminotransferase (1%), and hyperammonemia (1%). Study results demonstrate that IM JZP458 at 25/25/50 mg/m² MWF is efficacious and has a safety profile consistent with those of other asparaginases.

Introduction

Asparaginases are proteins originating from bacteria employed in the treatment of acute lymphoblastic leukemia (ALL) and lymphoblastic lymphoma (LBL), and come with a high allergenic and immunogenic potential.¹ Hypersensitivity reactions ranging from localized erythema to systemic anaphylaxis are the most common dose-limiting adverse reactions (AEs), occurring in up to 30% of patients receiving *Escherichia coli*-derived L-asparaginases.²

Additionally, antibody development can lead to inactivation of L-asparaginase activity in the absence of clinical symptoms, termed silent inactivation,² which is monitored via measurement of serum asparaginase activity (SAA). An SAA level of ≥ 0.1 IU/mL correlates with complete asparagine depletion.³ In the setting of a hypersensitivity reaction or silent inactivation to an *E. coli*-derived asparaginase, it is recommended to switch patients to an asparaginase preparation with a distinct immunogenic profile, such as an *Erwinia chrysanthemi*-derived asparaginase.³

JZP458 is a recombinant *Erwinia* asparaginase with an identical amino acid sequence to native *Erwinia* asparaginase (Erwinaze[®]/Erwinase[®]) but is derived from a novel *Pseudomonas fluorescens* expression platform. JZP458 is expected to have minimal immunologic cross-reactivity to *E. coli*-derived asparaginase preparations and has comparable *in vitro* activity to native *Erwinia*-derived asparaginase.⁴⁻⁷

In a phase 1 study in adult healthy volunteers, JZP458 was evaluated at a starting dose of 25 mg/m², chosen to provide similar asparaginase activity to native *Erwinia* asparaginase at 25,000 IU/m² based on *in vitro* analytical comparability. Results from that phase 1 study demonstrated that a single dose of 25 mg/m² intramuscular (IM) JZP458 resulted in similar SAA to 25,000 IU/m² of native *Erwinia* asparaginase.⁴

The AALL1931 study was initiated in patients with ALL or lymphoblastic lymphoma (LBL) and was conducted in collaboration with the Children's Oncology Group (COG). Based on interim data from this trial, JZP458 (Rylaze[®]; asparaginase erwinia chrysanthemi (recombinant)-rywn) was approved by the US Food and Drug Administration (FDA) in June 2021 for the treatment of ALL/LBL in patients ≥ 1 month of age who have developed hypersensitivity to *E. coli*-derived asparaginase, following review under the Real-Time Oncology Review (RTOR) program.⁸

Here, we report the efficacy and safety results from Part A of AALL1931, which investigated the IM route of administration in patients with ALL or LBL who developed hypersensitivity or silent inactivation to *E. coli*-derived asparaginases.

Methods

Study Design and Treatment

AALL1931 (ClinicalTrials.gov Identifier: NCT04145531) is a pivotal phase 2/3, open-label, multicenter, dose-confirmation and pharmacokinetic (PK) study to assess the efficacy and safety of JZP458, designed to allow additional cohorts to be enrolled at a new dose following review of safety and efficacy results of the current cohort by the Study Data Review Committee (SDRC) during the dose-confirmation phase. Part A investigated the IM route of administration of JZP458, while Part B (ongoing) investigated the intravenous route of administration. Each remaining pegylated *E. coli*-derived asparaginase dose on a patient's treatment plan was replaced by one course of IM JZP458 (six doses administered on a Monday/Wednesday/Friday [MWF] schedule over two weeks), with the initial dose starting on either M, W, or F, to align with the patient's planned chemotherapy schedule. All other chemotherapy continued according to the original treatment protocol for the patient's ALL/LBL.

The starting dose for IM JZP458 was 25 mg/m² (Cohort 1a) on MWF, based on data from the phase 1 study.⁴ Subsequent to a review of Cohort 1a nadir SAA (NSAA) and safety data by the SDRC, Cohort 1b opened at 37.5 mg/m² on MWF. Population pharmacokinetic (PopPK) modeling and simulations were used throughout the conduct of the AALL1931 study to inform dosing decisions. After a preliminary PopPK model based on data from Cohorts 1a and 1b predicted that a regimen of 25/25/50 mg/m² MWF would be required to achieve 72-hour NSAA levels ≥ 0.1 IU/mL to support MWF dosing, Cohort 1c was initiated to evaluate this regimen.

Patients

Patients were enrolled in Part A from December 2019 to March 2021 at approximately 70 sites across North America. The study was conducted in accordance with Good Clinical Practice guidelines and the Declaration of Helsinki. All patients or their guardians (for patients <18 years old) provided written informed consent and assent when indicated. The protocol and all amendments were approved by the institutional review board or ethics committee at each participating institution.

Pediatric or adult patients with newly diagnosed ALL/LBL were eligible if they developed a grade ≥ 3 allergic reaction (National Cancer Institute Common Terminology Criteria for Adverse Events version 5.0) to a pegylated *E. coli*-derived asparaginase or silent inactivation (definition in **Supporting Information**). Patients were considered ineligible if they had previously received native *Erwinia*-derived asparaginase or JZP458, had relapsed ALL/LBL, or a history of grade ≥ 3 pancreatitis. Full eligibility criteria are presented in the **Supporting Information**.

Study Endpoints and Assessments

The primary efficacy endpoint of AALL1931 was the proportion of patients with the last 72-hour NSAA level ≥ 0.1 IU/mL (to evaluate SAA coverage from Friday to Monday), and the key

secondary endpoint was the proportion of patients with the last 48-hour NSAA level ≥ 0.1 IU/mL (to evaluate SAA coverage from Monday to Wednesday or Wednesday to Friday, depending on the patient's scheduled starting day) during the first course of IM JZP458 administration. Additional secondary endpoints included the proportion of patients with the last 48-hour and 72-hour NSAA levels ≥ 0.4 IU/mL during the first course, characterization of the PK of IM JZP458 using a PopPK approach based on SAA, and assessment of immunogenicity following repeat administration. Plasma asparagine concentration was included as an additional assessment. Blood samples for SAA and asparagine were collected at pre-specified timepoints (**Table S1**) and transferred on dry ice to bioanalytical labs. SAA samples were analyzed at Charles River Laboratories (Skokie, IL) using a validated enzyme activity assay in human serum over the range of 0.0349 to 0.2096 IU/mL. Asparagine samples were assayed for plasma asparagine concentration by Syneos Health (Princeton, NJ), using a validated liquid chromatography tandem mass spectrometry method over the range of 0.025 to 10.0 $\mu\text{g/mL}$. Safety and tolerability were assessed by the occurrence of treatment-emergent AEs and treatment-related AEs. AEs were considered to be related to JZP458 if the event followed a reasonable temporal sequence from administration and satisfied ≥ 1 instances of clinical evidence: 1) the event followed a known or suspected response pattern to the study drug; 2) the event disappeared after stopping the study drug; or 3) the event reappeared after the study drug was restarted.

PopPK Modeling and Simulations

A PopPK model was developed for JZP458 with SAA data from AALL1931 using nonlinear mixed effects modeling (NONMEM; version 7.3) to describe the PK of JZP458 after IM administration. SAA was the basis of the PK assessment in this study.

The PopPK model included 2,687 SAA data points from 166 patients who received IM JZP458 in this study: 32 patients from Cohort 1a, 83 patients from Cohort 1b, and 51 patients from Cohort 1c. The effects of potential covariates were evaluated to identify the covariates likely to contribute to the variability of JZP458 pharmacokinetics.

The model was used to simulate the SAA profiles for a virtual population with 2,000 subjects (1,000 pediatric; 1,000 adult) to explore the likelihood of achieving a therapeutic target NSAA level of ≥ 0.1 IU/mL based on different doses, schedules, and routes of administration. The virtual population ranged from 1 month to 80 years old to represent real world scenarios. Dosing regimens tested in the AALL1931 study were simulated, and one dosing regimen not tested in the study was also simulated.

Statistical Analysis

The primary and key secondary efficacy endpoints were assessed by calculating the 95% Wald confidence interval (CI) around the proportion of patients with NSAA levels ≥ 0.1 IU/mL. The efficacy analysis population included patients who received ≥ 1 dose of JZP458 and had at least one 48-hour or 72-hour NSAA assessment within the protocol defined sample collection window (± 2 hours) in the first course. The safety analysis population included patients who received ≥ 1 dose of JZP458. The PK analysis set included patients who received ≥ 1 dose of JZP458 and had ≥ 1 postdose evaluable SAA value.

Cohort Determination

An interim analysis including data from Cohort 1a (25 mg/m² MWF) and Cohort 1b (37.5 mg/m² MWF) was conducted by the SDRC after 51 patients in Cohort 1b completed Course 1. The SDRC recommended initiating Cohort 1c based on the results of preliminary PopPK modeling

and simulation while continuing enrollment in Cohort 1b. Analysis of all patients enrolled in Part A was conducted after 51 patients in Cohort 1c (25/25/50 mg/m² MWF) completed Course 1.

This report uses data current as of July 19, 2021. Additional details are presented in the

Supporting Information.

Data Sharing Statement

All relevant data are provided with the manuscript and supporting files. Jazz has established a process to review requests from qualified external researchers for data from Jazz-sponsored clinical trials in a responsible manner that includes protecting patient privacy, assurance of data security and integrity, and furthering scientific and medical innovation. Additional details on Jazz Pharmaceuticals data sharing criteria and process for requesting access can be found at:

<https://www.jazzpharma.com/science/clinical-trial-data-sharing/>.

Results

Patient Characteristics and Disposition

AALL1931 enrolled 167 patients to the three IM cohorts (Cohort 1a, n = 33; Cohort 1b, n = 83; Cohort 1c, n = 51). As of the cut-off date for this analysis, 28 (17%) patients were still receiving treatment, 104 (62%) patients had completed all planned doses of JZP458 treatment, and 35 (21%) patients had discontinued JZP458 treatment (**Figure 1**). The most common reason for discontinuation was AEs (13% [n = 21]). The median (range) number of planned courses of JZP458 was 5 (1, 14) for Cohort 1a, 5 (1, 15) for Cohort 1b, and 5 (1, 15) for Cohort 1c. At the data cut-off date, the median (range) number of JZP458 courses received was 5 (1, 14) for Cohort 1a, 5 (1, 15) for Cohort 1b, and 4 (1, 11) for Cohort 1c (some patients were still receiving treatment).

Baseline demographic and disease characteristics are presented in **Table 1**. Median (range) age at enrollment was 10 (1, 25) years, with 87% of patients aged <18 years.

Efficacy

Following the first treatment course of JZP458, the proportion (95% CI) of patients who achieved NSAA levels ≥ 0.1 IU/mL at 72 hours was 64% (47%, 82%) in Cohort 1a, 91% (84%, 97%) in Cohort 1b, and 90% (81%, 98%) in Cohort 1c. At 48 hours, over 95% of patients in each cohort (Cohort 1a, 97% [91%, 100%]; Cohort 1b, 99% [96%, 100%]; Cohort 1c, 96% [90%, 100%]) achieved NSAA levels ≥ 0.1 IU/mL (**Figure 2A**). The proportions of patients who achieved NSAA levels ≥ 0.4 IU/mL following the first treatment course are presented in **Table S2**. The mean (interquartile range^{25,75}) NSAA levels at 72 hours were 0.16 (0.09, 0.23) in Cohort 1a, 0.33 (0.17, 0.41) in Cohort 1b, and 0.47 (0.17, 0.59) in Cohort 1c. At 48 hours, mean NSAA levels were 0.45 (0.27, 0.65) in Cohort 1a, 0.88 (0.47, 1.09) in Cohort 1b, and 0.66 (0.33, 0.89) in Cohort 1c. **Figure 2B** shows that the mean (95% CI) NSAA levels (IU/mL) during the first treatment course at the last 72 and the last 48 hours were ≥ 0.1 IU/mL in all cohorts.

PopPK Modeling and Simulations

The clinical efficacy of JZP458 was demonstrated by a combination of observed data and modeled results, which included observed NSAA levels at the protocol-specified time points and PopPK modeling and simulation results. The PopPK model that best described the PK (based on SAA) of JZP458 following IM administration was a 1-compartment IM model with mixed order absorption and linear elimination, with BSA included as an allometric covariate on JZP458 SAA clearance (CL) and volume of distribution (V), accompanied with self-reported race (ie, Black/African American) and disease subtype (ie, T-cell ALL) as categorical covariates on

JZP458 SAA CL. This model indicated that the simulated proportion of patients achieving NSAA ≥ 0.1 IU/mL exceeded 90% in both Black/African American and non-Black/African American patients, suggesting that no clinically significant difference would be expected in patients following the proposed BSA-based dosing. Therefore, no dose modification is recommended based on ethnicity, including African American patients.

Using this PopPK model, simulations suggested that when JZP458 is administered IM at 25/25/50 mg/m² MWF, 92.1% (95% CI: 90.9%, 93.3%) of patients were expected to achieve the last 72-hour NSAA levels ≥ 0.1 IU/mL and 93.8% (95% CI: 92.7%, 94.9%) of patients were expected to achieve the last 48-hour NSAA levels ≥ 0.1 IU/mL.

Asparagine Depletion

Depletion of plasma asparagine was observed after IM JZP458 administration for all patients in all dosing cohorts. In Course 1, mean predose (baseline) L-asparagine concentrations ranged from 7.59 to 11.32 $\mu\text{g/mL}$ for all dose levels and schedules, consistent with literature-reported values.⁹ For all cohorts following IM JZP458 administration, mean plasma asparagine levels rapidly declined from predose levels to levels below or near the assay lower limit of quantitation (0.025 $\mu\text{g/mL}$); reduced plasma levels lasted throughout the treatment duration of Course 1 up to predose 6, when the last sample was collected (**Figure 3**). Plasma asparagine levels over time were similar across all 3 dosing schedules (ie, MWF vs WFM vs FMW). Four patients had transient low-level increases in plasma asparagine (Cohort 1a, 3 patients; Cohort 1b, 1 patient).

Safety

One hundred and twenty-four (74.3%) patients experienced treatment-related AEs and 86 (51.5%) patients experienced grade 3/4 treatment-related AEs (**Table 2**). The most common non-hematologic grade 3/4 treatment-related AEs were febrile neutropenia (9.0%), increased ALT

(7.8%), and nausea (5.4%; **Table 3**). In total, 21 (12.6%) patients discontinued JZP458 due to treatment-related AEs: pancreatitis (6.0%), allergic reactions (5.4%) including anaphylaxis (1.8%), increased ALT (0.6%), and hyperammonemia (0.6%). Treatment-emergent AEs that led to death were sepsis (Cohort 1a, n = 1), aspiration pneumonia (Cohort 1b, n = 1), and multi-organ failure (Cohort 1b, n = 1), none of which were deemed to be related to JZP458 treatment.

Treatment-related AEs of special interest included allergic reactions, pancreatitis, and thrombosis; incidence by preferred term is presented in **Table S4**. Allergic reactions related to JZP458 were observed in 16 (9.6%) patients, including 3 (1.8%) with anaphylactic reaction (**Table 3**). Incidence of grade 3/4 treatment-related AEs of special interest among all cohorts were as follows: allergic reactions (5.4%), pancreatitis (6.0%), and thrombosis (1.2%).

Treatment-related AEs of hepatotoxicity, predominately elevation of liver enzymes, increased bilirubin and/or ammonia of any grade, occurred in 33 (19.8%) patients (**Table S5**). Treatment-related hypertriglyceridemia occurred in 12 (7.2%) patients. Grade 4 hypertriglyceridemia, which was deemed related to JZP458, was reported in 1 (3.0%) patient in Cohort 1a and 1 (2.0%) patient in Cohort 1c.

Discussion

Incorporation of L-asparaginase into multi-agent regimens has been of significant importance in improving outcomes of children and adolescents with ALL, and omission of asparaginase (or low/non-detectable serum levels after its administration) may lead to higher risk of relapse and, in patients with recurrent disease, a poorer response to reinduction therapy.¹⁰⁻¹² *E. coli*-derived asparaginases (particularly pegylated formations) are typically used in frontline ALL protocols. Asparaginases derived from *Erwinia chrysanthemi* have a distinct immunologic profile and have

long been the only option available to patients unable to receive *E. coli*-derived formulations due to hypersensitivity or silent inactivation. In recent years, switching from an *E. coli*-derived asparaginase to an *Erwinia*-derived preparation has proved challenging due to manufacturing issues with native asparaginase *Erwinia chrysanthemi* (Erwinaze[®]/Erwinase[®]), leading to global shortages.¹³ Due to this critical unmet medical need, JZP458 was developed as a novel asparaginase product to provide a reliable treatment option for patients who develop hypersensitivity to *E. coli*-derived asparaginases. The JZP458 clinical program, including the pivotal phase 2/3 AALL1931 trial, was conducted in close collaboration with the COG and the FDA under the accelerated RTOR program.

The clinical effectiveness of asparaginase is thought to be based on adequate depletion of asparagine^{3,14,15}; however, accurate measurement of asparagine levels is technically challenging and not practical in the clinical setting. Thus, serum asparaginase activity serves as a surrogate marker for asparagine depletion in clinical trials and for patient care.^{3,4} Although one study of pegylated *E. coli*-derived asparaginase has suggested that a higher level of asparaginase activity (≥ 0.4 IU/mL) may be needed for complete asparagine depletion, particularly in the cerebrospinal fluid,¹⁶ the preponderance of the literature suggests the NSAA threshold of ≥ 0.1 IU/mL as the target to ensure therapeutic benefit from asparaginase therapy.^{3,14,15} Therefore, an NSAA threshold of ≥ 0.1 IU/mL was used to define the primary and key secondary efficacy endpoints in the JZP458 AALL1931 study while an NSAA threshold of ≥ 0.4 IU/mL was explored as a secondary endpoint.

An interim analysis of data from the first 102 patients enrolled in AALL1931 (primarily in Cohorts 1a and 1b), in conjunction with PopPK modeling and simulations, suggested that 94% (95% CI: 93%, 95%) of patients would be expected to maintain NSAA levels ≥ 0.1 IU/mL at 48

hours at 25 mg/m² with 7 doses per 2-week course.¹⁷ These data along with interim safety data were considered sufficient by the FDA to support the approval of JZP458 at 25 mg/m² every 48 hours under the RTOR program, providing timely access to an *Erwinia*-derived asparaginase to patients in need. The FDA accepted the observed and simulated data to meet the required efficacy target as the basis for its decision. Data from all 167 IM patients enrolled in Part A of AALL1931 were provided to the FDA to pursue a label update to include the IM dosing regimen with 25/25/50 mg/m² MWF, which may be the preferred option at some institutions that cannot easily administer asparaginase in the outpatient setting over the weekend.

Here, we report the results for all 167 patients enrolled in Part A of AALL1931, including 51 patients in Cohort 1c. Results from this follow-up analysis demonstrate that the IM dosing regimen of 25/25/50 mg/m² MWF (Cohort 1c), which intentionally used a higher Friday dose to improve the 72-hour SAA coverage, is efficacious and tolerable, based on observed data and model-based simulations as well as the observed safety profile. The observed data indicated similar proportions of patients achieved 72-hr NSAA levels ≥ 0.1 IU/mL in Cohort 1c (90%) and Cohort 1b (91%) although the mean SAA levels at the last 72-hour was higher in Cohort 1c than in Cohort 1b (0.47 [95% CI: 0.35, 0.59] IU/mL vs 0.33 [95% CI: 0.27, 0.39] IU/mL). However, the study was not powered to compare results among cohorts. When the observed data were used to refine the PopPK model, simulations indicate that 92.1% (95% CI: 90.9%, 93.3%) of patients are expected to achieve the last 72-hour NSAA levels ≥ 0.1 IU/mL with JZP458 administered IM at 25/25/50 mg/m² MWF. PopPK modeling and simulation has several important advantages in that it uses pooled data from all patients, all dose levels, all time points, and across all courses and it uses continuous rather than categorical data, resulting in more precise estimation of NSAA levels in a large population as manifested by the tight 95% CI.

The study was not designed to test for statistically significant differences between safety profiles among cohorts. However, the totality of observed safety data from Cohort 1c appeared similar to the safety data from Cohorts 1a (25 mg/m² MWF) and 1b (37.5 mg/m² MWF) and were consistent with what has been described for other asparaginases.^{1,2,18-20} Limitations of this study include the single-arm design. In addition, at the data cut-off date for the results presented in this manuscript, some patients had not completed treatment; therefore, the complete safety follow-up data will need to be reported in a future manuscript.

Conclusions

In conclusion, IM JZP458 using the dosing regimen of 25/25/50 mg/m² MWF is efficacious, tolerable, and achieves NSAA levels ≥ 0.1 IU/mL at both 48 and 72 hours in the vast majority of patients. The safety profile of JZP458 is consistent with what has been described with other asparaginase products.^{1,2,18-20} Data from all 167 patients treated with IM JZP458 in Part A of AALL1931 were provided to the FDA to support a label update to include the IM dosing regimen of 25/25/50 mg/m² MWF, in supplement of the approved dosing of 25 mg/m² every 48 hours. Future studies are needed to determine the impact of JZP458 on patient experience and outcomes. These dosing regimens optimize SAA coverage and provide flexibility and optimize alignment with many pediatric oncology protocols throughout the world. Importantly, with its reliable manufacturing process, and the efficacy and safety demonstrated in AALL1931, JZP458 addresses one of the most significant drug shortages impacting the care of patients with ALL/LBL.

Acknowledgments

This study was conducted in collaboration with the Children's Oncology Group (COG). The authors thank all investigators who contributed to this project, the patients who participated in this study, the staff members at the study sites, the study data review committee, and the staff members involved in data collection and analyses. Medical writing and editorial support were provided by Austin Horton, PhD, of Cello Health Communications/SciFluent Communications, Inc., and were financially supported by Jazz Pharmaceuticals.

Author Contributions:

Conception and design: Luke Maese, Mignon L. Loh, Mi Rim Choi, Robert Iannone, Jeffrey A. Silverman, Tong Lin, Elizabeth A. Raetz, and Rachel E. Rau

Collection and assembly of data: Luke Maese, Mignon L. Loh, Mi Rim Choi, Tong Lin, Etsuko Aoki, Michelle Zanette, Shirali Agarwal, Jeffrey A. Silverman, Lewis B. Silverman, Elizabeth A. Raetz, and Rachel E. Rau

Data analysis and interpretation: All authors

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

Disclosures of Conflicts of Interest:

Luke Maese: Served on an advisory board and speakers bureau for Jazz Pharmaceuticals

Mignon L. Loh: Nothing to disclose

Mi Rim Choi: Employee of and holds stock ownership and/or stock options in Jazz

Pharmaceuticals

Tong Lin: Employee of and holds stock ownership and/or stock options in Jazz Pharmaceuticals

Etsuko Aoki: Employee of and holds stock ownership and/or stock options in Jazz

Pharmaceuticals

Michelle Zanette: Employee of and holds stock ownership and/or stock options in Jazz

Pharmaceuticals

Shirali Agarwal: Employee of and holds stock ownership and/or stock options in Jazz

Pharmaceuticals

Robert Iannone: Employee of and holds stock ownership and/or stock options in Jazz

Pharmaceuticals

Jeffrey A. Silverman: was an employee of Jazz Pharmaceuticals at the time of the study and holds stock ownership and/or stock options in Jazz Pharmaceuticals

Lewis B. Silverman: Served on scientific advisory boards for Jazz Pharmaceuticals and Servier Pharmaceuticals

Elizabeth A. Raetz: Received institutional research funding from Pfizer and serves on a Data and Safety Monitoring Board for BMS

Rachel E. Rau: Served on advisory boards for Jazz Pharmaceuticals and Servier Pharmaceuticals

Funding

This study was funded by Jazz Pharmaceuticals.

List of Previous Presentations of This Study

Initial results of this study were presented at the 63rd American Society of Hematology (ASH) Annual Meeting and Exposition; December 11-14, 2021; Atlanta, Georgia; Poster No #2307; results of this study were also presented at the American Society of Clinical Oncology Annual Meeting; June 3-7, 2022; Chicago, Illinois; Abstract No #7001.

References

1. Hijiya N, van der Sluis IM. Asparaginase-associated toxicity in children with acute lymphoblastic leukemia. *Leuk Lymphoma*. 2016;57(4):748-757.
2. Raetz EA, Salzer WL. Tolerability and efficacy of L-asparaginase therapy in pediatric patients with acute lymphoblastic leukemia. *J Pediatr Hematol Oncol*. 2010;32(7):554-563.
3. van der Sluis IM, Vrooman LM, Pieters R, et al. Consensus expert recommendations for identification and management of asparaginase hypersensitivity and silent inactivation. *Haematologica*. 2016;101(3):279-285.
4. Lin T, Hernandez-Illas M, Rey A, et al. A randomized phase I study to evaluate the safety, tolerability, and pharmacokinetics of recombinant *Erwinia* asparaginase (JZP-458) in healthy adult volunteers. *Clin Transl Sci*. 2021;14(3):870-879.
5. Willer A, Gerss J, König T, et al. Anti-*Escherichia coli* asparaginase antibody levels determine the activity of second-line treatment with pegylated *E coli* asparaginase: a retrospective analysis within the ALL-BFM trials. *Blood*. 2011;118(22):5774-5782.
6. Wang B, Relling MV, Storm MC, et al. Evaluation of immunologic crossreaction of antiasparaginase antibodies in acute lymphoblastic leukemia (ALL) and lymphoma patients. *Leukemia*. 2003;17(8):1583-1588.
7. Zalewska-Szewczyk B, Gach A, Wyka K, Bodalski J, Młynarski W. The cross-reactivity of anti-asparaginase antibodies against different L-asparaginase preparations. *Clin Exp Med*. 2009;9(2):113-116.
8. RYLAZE™ (asparaginase erwinia chrysanthemi (recombinant)-rywn) [package insert]. Leinster, Ireland: Jazz Pharmaceuticals Ireland Limited; June 2021.

9. Tong WH, Pieters R, Kaspers GJ, et al. A prospective study on drug monitoring of PEGasparaginase and *Erwinia* asparaginase and asparaginase antibodies in pediatric acute lymphoblastic leukemia. *Blood*. 2014;123(13):2026-2033.
10. Bender C, Maese L, Carter-Febres M, Verma A. Clinical utility of pegaspargase in children, adolescents and young adult patients with acute lymphoblastic leukemia: a review. *Blood Lymphat Cancer*. 2021;11:25-40.
11. Jarrar M, Gaynon PS, Periclou AP, et al. Asparagine depletion after pegylated *E. coli* asparaginase treatment and induction outcome in children with acute lymphoblastic leukemia in first bone marrow relapse: a Children's Oncology Group study (CCG-1941). *Pediatr Blood Cancer*. 2006;47(2):141-146.
12. Gupta S, Wang C, Raetz EA, et al. Impact of asparaginase discontinuation on outcome in childhood acute lymphoblastic leukemia: a report from the Children's Oncology Group. *J Clin Oncol*. 2020;38(17):1897-1905.
13. Maese L, Rizzari C, Coleman R, Power A, van der Sluis I, Rau RE. Can recombinant technology address asparaginase *Erwinia chrysanthemi* shortages? *Pediatr Blood Cancer*. 2021;68(10):e29169.
14. Asselin B, Rizzari C. Asparaginase pharmacokinetics and implications of therapeutic drug monitoring. *Leuk Lymphoma*. 2015;56(8):2273-2280.
15. Pieters R, Hunger SP, Boos J, et al. L-asparaginase treatment in acute lymphoblastic leukemia: a focus on *Erwinia* asparaginase. *Cancer*. 2011;117(2):238-249.
16. Angiolillo AL, Schore RJ, Devidas M, et al. Pharmacokinetic and pharmacodynamic properties of calaspargase pegol *Escherichia coli* L-asparaginase in the treatment of patients with

acute lymphoblastic leukemia: results from Children's Oncology Group Study AALL07P4. *J Clin Oncol*. 2014;32(34):3874-3882.

17. US Food and Drug Administration. Center for Drug Evaluation and Research.

Application number 761179Orig1s000. Multi-discipline review.

https://www.accessdata.fda.gov/drugsatfda_docs/nda/2021/761179Orig1s000MultidisciplineR.pdf. Accessed 20 October 2021.

18. Vrooman LM, Kirov, II, Dreyer ZE, et al. Activity and toxicity of intravenous *Erwinia* asparaginase following allergy to *E. coli*-derived asparaginase in children and adolescents with acute lymphoblastic leukemia. *Pediatr Blood Cancer*. 2016;63(2):228-233.

19. Avramis VI, Sencer S, Periclou AP, et al. A randomized comparison of native *Escherichia coli* asparaginase and polyethylene glycol conjugated asparaginase for treatment of children with newly diagnosed standard-risk acute lymphoblastic leukemia: a Children's Cancer Group study. *Blood*. 2002;99(6):1986-1994.

20. van der Sluis IM, de Groot-Kruseman H, Te Loo M, et al. Efficacy and safety of recombinant *E. coli* asparaginase in children with previously untreated acute lymphoblastic leukemia: a randomized multicenter study of the Dutch Childhood Oncology Group. *Pediatr Blood Cancer*. 2018;65(8):e27083.

21. Lin T, Dumas T, Kaullen J, et al. Population pharmacokinetic model development and simulation for recombinant *Erwinia* asparaginase produced in *Pseudomonas fluorescens* (JZP-458). *Clin Pharmacol Drug Dev*. 2021;10(12):1503-1513.

Table 1. Demographic and Baseline Disease Characteristics (Safety Analysis Set)

Characteristic	Cohort 1a: 25 mg/m² MWF (n = 33)	Cohort 1b: 37.5 mg/m² MWF (n = 83)	Cohort 1c: 25/25/50 mg/m² MWF (n = 51)	Total (N = 167)
Age, years				
Median (range)	11 (1, 24)	8 (1, 20)	12 (3, 25)	10 (1, 25)
<6, n (%)	9 (27)	24 (29)	11 (22)	44 (26)
6 to <12, n (%)	9 (27)	34 (41)	14 (27)	57 (34)
12 to <18, n (%)	7 (21)	20 (24)	18 (35)	45 (27)
≥18, n (%)	8 (24)	5 (6)	8 (16)	21 (13)
Sex, n (%)				
Male	17 (52)	55 (66)	31 (61)	103 (62)
Race, n (%)				
American Indian or Alaska native	0	0	3 (6)	3 (2)
Asian	1 (3)	5 (6)	1 (2)	7 (4)
Black or African American	3 (9)	11 (13)	8 (16)	22 (13)
White	24 (73)	58 (70)	33 (65)	115 (69)
Multiple	1 (3)	0	0	1 (1)
Not reported	4 (12)	9 (11)	6 (12)	19 (11)
BMI, median (range), kg/m ²	19.9 (13.4, 42.6)	17.9 (13.7, 30.7)	18.4 (13.8, 42.0)	18.4 (13.4, 42.6)
BSA, median (range), m ²	1.28 (0.44, 2.53)	1.01 (0.56, 2.26)	1.29 (0.54, 2.43)	1.19 (0.44, 2.53)
Primary disease, n (%)				
ALL				
B-ALL	27 (82)	60 (72)	37 (73)	124 (74)
T-ALL	4 (12)	13 (16)	9 (18)	26 (16)
LBL				
B-LBL	0	0	1 (2)	1 (1)
T-LBL	2 (6)	10 (12)	4 (8)	16 (10)
Eligibility criteria met, n (%)				
Grade ≥3 allergic reaction to an <i>E. coli</i> -derived asparaginase ^a	27 (82)	75 (90)	44 (86)	146 (87)
Silent inactivation	3 (9)	3 (4)	1 (2)	7 (4)
Allergic reaction with inactivation	3 (9)	5 (6)	6 (12)	14 (8)

MWF, Monday, Wednesday, Friday; BMI, body mass index; BSA, body surface area; ALL, acute lymphoblastic leukemia; B-ALL, B-cell acute lymphoblastic leukemia; T-ALL, T-cell acute lymphoblastic leukemia; B-LBL, LBL; lymphoblastic lymphoma; B-cell lymphoblastic lymphoma; T-LBL, T-cell lymphoblastic lymphoma; COG, Children's Oncology Group.

Patients received various ALL/LBL treatment protocols including (but not limited to) COG AALL0434, AALL0932, AALL1131, AALL1231, AALL1731, and AALL1732, as well as DFC16-001 and TOT17.

^aAll patients in Part A received pegaspargase prior to entering the study.

Table 2. Summary of AEs

	Cohort 1a: 25 mg/m ² MWF (n = 33)	Cohort 1b: 37.5 mg/m ² MWF (n = 83)	Cohort 1c: 25/25/50 mg/m ² MWF (n = 51)	Total (N = 167)
Patients, n (%)				
Any treatment-related AE	21 (64)	66 (80)	37 (73)	124 (74)
Any treatment-related grade 3/4 AE	14 (42)	46 (55)	26 (51)	86 (51)
Any treatment-related serious AE	5 (15)	28 (34)	12 (24)	45 (27)
Any treatment-related AE leading to study drug discontinuation	2 (6)	14 (17)	5 (10)	21 (13)
Any treatment-related AE leading to death	0	0	0	0

AE, adverse event; MWF, Monday/Wednesday/Friday.

Table 3. Most Commonly Reported ($\geq 10\%$ in Any Cohort) Non-Hematologic Treatment-related AEs^a

Patients, n (%)	Cohort 1a: 25 mg/m ² MWF (n = 33)		Cohort 1b: 37.5 mg/m ² MWF (n = 83)		Cohort 1c: 25/25/50 mg/m ² MWF (n = 51)	
	All grades	Grade 3/4	All grades	Grade 3/4	All grades	Grade 3/4
Nausea	7 (21)	1 (3)	20 (24)	6 (7)	13 (25)	2 (4)
Vomiting	7 (21)	0	22 (27)	3 (4)	10 (20)	1 (2)
ALT increased	2 (6)	2 (6)	14 (17)	8 (10)	8 (16)	3 (6)
Decreased appetite	3 (9)	1 (3)	12 (14)	3 (4)	8 (16)	1 (2)
Fatigue	3 (9)	1 (3)	17 (20)	0	2 (4)	0
AST increased	2 (6)	0	12 (14)	4 (5)	3 (6)	0
Abdominal pain	2 (6)	0	9 (11)	2 (2)	6 (12)	0
Febrile neutropenia	2 (6)	2 (6)	11 (13)	11 (13)	2 (4)	2 (4)
Hyperglycemia	3 (9)	1 (3)	5 (6)	2 (2)	6 (12)	2 (4)
Treatment-related AEs of special interest						
Allergic reaction ^b	2 (6)	2 (6)	11 (13)	6 (7)	3 (6)	1 (2)
Pancreatitis ^c	0	0	6 (7)	6 (7)	6 (12)	4 (8)
Thrombosis ^d	0	0	2 (2)	2 (2)	0	0

Treatment-related AE, adverse event; MWF, Monday/Wednesday/Friday; ALT, alanine aminotransferase; AST, aspartate aminotransferase; MedDRA, *Medical Dictionary for Regulatory Activities*.

^aTreatment-related AEs were reported as MedDRA (version 22.1) preferred terms.

^bIncludes drug hypersensitivity, anaphylactic reaction, hypersensitivity, infusion-related reaction, rash maculo-papular, rash, urticaria, and rash erythematous.

^cIncludes pancreatitis and acute pancreatitis. Reporting of pancreatitis by the preferred terms “pancreatitis” or “acute pancreatitis” was assigned by the investigator. However, there was no clear distinction between the two preferred terms.

^dIncludes superior sagittal sinus thrombosis, pulmonary embolism, and jugular vein thrombosis.

Figure Legends

Figure 1. CONSORT Diagram of Patient Disposition by Cohort.

Figure 2. A) Proportion of Patients Achieving NSAA levels ≥ 0.1 IU/mL in the First Treatment Course^a and B) Mean NSAA Levels for Evaluable Patients in the First Treatment Course.^b

NSAA, nadir serum asparaginase activity; CI, confidence interval; MWF, Monday/Wednesday/Friday.

^aError bars represent 95% CIs calculated by the Wald method.

^bError bars represent 95% CIs.

Figure 3. Mean SAA-Time Profiles and Corresponding Mean Plasma L-Asparagine Levels in the First Treatment Course by Cohort and Dosing Schedule.

SAA, serum asparaginase activity; MWF, Monday/Wednesday/Friday; W/F/M, Wednesday/Friday/Monday; F/M/W, Friday/Monday/Wednesday; LLOQ, lower limit of quantitation.

The LLOQ were as follows: SAA = 0.0350 IU/mL; L-asparagine = 0.0250 μ g/mL. Values below the LLOQ were graphed as 0 on the linear scale or omitted from the semi-logarithmic scale.

Only SAA is graphed on a semi-logarithmic scale.

^aFor Cohort 1a, the sample size for mean SAA levels and plasma asparagine levels ranged from 8 to 12 patients for all timepoints. For Cohort 1b, the sample size ranged from 15 to 22 patients for all timepoints. For Cohort 1c, the sample size ranged from 17 to 19 patients for all timepoints.

^bFor Cohort 1a, the sample size for mean SAA levels and plasma asparagine levels was 7 patients for all timepoints. For Cohort 1b, the sample size ranged from 27 to 32 patients for all timepoints. For Cohort 1c, the sample size ranged from 18 to 23 for all timepoints.

^cFor Cohort 1a, the sample size for mean SAA levels and plasma asparagine levels ranged from 6 to 8 patients for all timepoints. For Cohort 1b, the sample size ranged from 23 to 29 patients for all timepoints. For Cohort 1c, the sample size ranged from 6 to 8 patients for all timepoints.

Figure 1.




Figure 2.




Figure 3.

SAA

L-asparagine concentration

Monday/Wednesday/Friday^a

Wednesday/Friday/Monday^b

Friday/Monday/Wednesday^c

Mean SAA (IU/mL)

Plasma L-asparagine levels ($\mu\text{g/mL}$)

Time

predose 1

2.5 h postdose 1

48 h postdose 1

72 h postdose 2

2.5 h postdose 3

48 h postdose 4

72 h postdose 5

Downloaded from https://academic.oup.com/blood/article-pdf/dof/10.1182/blood.2022016923/1921795/blood.2022016923.pdf by guest on 16 September 2022

