□□ - 2026-03-12

□□□□: 2026-03-12 23:51

□□□□□□: 129

1 Immune-microbiome coordination defines interferon setpoints in healthy humans

- □□□□□□: 45 / 60
- **AML** □□□□: 23 / 30
- □□□□□□□: 7 / 10
- □□□□: 66 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□

□□: Coronary artery disease is the leading cause of death worldwide. Recently, hundreds of genomic loci have been shown to increase risk for the disease, however, the molecular mechanisms underlying signals from risk loci remain largely unclear. Here, we integrate the latest statistics of coronary artery disease genetics from over one million individuals with epigenetic data from 45 cell types to identify genes and transcription factors whose regulation is affected by variants. Applying two…

—

4 Midostaurin response in AML is shaped by a progenitor-like cell state selectively targeted by SMAC mimetics

- □□□□□□: 36 / 60
- **AML** □□□□: 21 / 30
- □□□□□□□: 5 / 10
- □□□□: 53 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□

- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□HAstV1-8□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

- [redacted]: 43 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]

[redacted]: A greater understanding of the biology of nevi will provide insights into the etiology of melanoma. Our large-scale meta-analysis of 14 nevus genome-wide association studies (GWAS) includes 85,965 individuals of European ancestry. We identify 29 nevus-associated loci ($p < 5 \times 10^{(-8)}$), of which 24 have not been previously reported in a GWAS conducted for nevus count alone. We further identify 255 candidate genes for nevus loci, including SIKE1 which is involved in immune response regulation.…

—

10 Qiangji Decoction regulates AMPKα/Drp1/Nrf2 pathway to ameliorate neural damage induced by D-galactose by suppressing oxidative stress and neuronal apoptosis

- [redacted]: 33 / 60
- **AML** [redacted]: 10 / 30
- [redacted]: 6 / 10
- [redacted]: 42 / 100
- [redacted]: 2026-03-12
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]AML [redacted]

[redacted]: This study aims to investigate the mechanism by which Qiangji Decoction(QJD) regulates the adenosine 5′-monophosphate(AMP)-activated protein kinase α(AMPKα)/dynamin-related protein 1(Drp1)/nuclear factor E2 related factor 2(Nrf2) pathway to ameliorate oxidative stress and neuronal apoptosis induced by D-galactose. Seventy-two C57BL/6 mice were randomized into the control, model, low-dose Qiangji Decoction(L-QJD), medium-dose Qiangji Decoction(M-QJD), high-dose Qiangji Decoction(H-QJD), and…

—

11 Identification of altered immune landscape at single-cell resolution in NSCLC brain metastasis and its association with poor immune checkpoint inhibitor responses

- [redacted]: 35 / 60
- **AML** [redacted]: 10 / 30
- [redacted]: 6 / 10
- [redacted]: 42 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] NSCLC [redacted] AML [redacted]

[redacted]: Brain metastasis, a common complication of advanced non-small cell lung cancer (NSCLC), leads to poor prognosis and reduces the efficacy of immune checkpoint inhibitors (ICIs). However, the mechanisms underlying this diminished ICI response remain unclear. Here we perform single-cell RNA sequencing on 101,959 tumor-infiltrating immune cells from primary tumors and brain metastases to delineate their distinct immune landscapes. Brain metastases display profound immunosuppressive reprogramming,…

—

12 Maternal microbiome-derived propionate regulates offspring myelination via histone lactylation

- □□□□□□: 34 / 60
- **AML** □□□□: 9 / 30
- □□□□□□□: 5 / 10
- □□□□: 41 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□-□□-□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

□□: The maternal gut microbiome plays a crucial role in regulating offspring neurodevelopment through microbial metabolite signaling, yet its influence on CNS myelinogenesis, a pivotal process for neural circuit maturation, remains poorly understood. Here, using antibiotic-induced maternal dysbiosis models, we identify propionate (PA), a short-chain fatty acid (SCFA) derived from the maternal microbiome, as a key epigenetic modulator of oligodendrocyte precursor cell (OPC) differentiation. Maternal…

—

13 Simulating the Photochemical Birth of the Hydrated Electron in Liquid Water

- □□□□□□: 40 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 40 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

15 NAD⁺ controls circadian rhythmicity during cardiac aging

- □□□□□□: 34 / 60
- **AML** □□□□: 9 / 30
- □□□□□□□: 5 / 10
- □□□□: 39 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□ NAD^+□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□/□□□□□□□□□□□□□□□□□□□□□□□□

□□: Disruption of the circadian clock as well as reduced NAD^(+) levels are both hallmarks of aging. While circadian rhythms and NAD^(+) metabolism have been linked in heart disease, their relationship during cardiac aging is less clear. Here, we show that aging leads to disruption of diurnal gene expression in the heart. Long-term supplementation with the NAD^(+) precursor nicotinamide riboside (NR) boosts NAD^(+) levels, reprograms the diurnal transcriptome, and reverses naturally occurring…

—

16 Biological valorization of methane and nitrogen gas-derived ammonia via methanotrophic bacteria for gut-beneficial nutrients

- □□□□□□: 33 / 60
- **AML** □□□□: 7 / 30
- □□□□□□□: 7 / 10
- □□□□: 39 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□

18 The Transposable Element-PARP Axis Underpins Synthetic Lethality and Immunogenic Vulnerability in Blood Cancer

- □□□□□□: 24 / 60
- **AML** □□□□: 20 / 30
- □□□□□□□: 5 / 10
- □□□□: 35 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□ Highlight □□□□□□□□□□“recent evidence”□□□□ TE-PARP □□

□□: Transposable elements (TEs) are emerging regulators of hematopoiesis and leukemia, creating vulnerabilities exploitable for therapy. Recent evidence shows that TE reactivation induces innate immune signalling, DNA damage responses and dependence on Poly(ADP-ribose) polymerase (PARP)-mediated protection, enabling synthetic lethality with PARP inhibition even in homologous-recombination-proficient leukemias with epigenetic gene mutations. In this article, we highlight the biology underpinning this…

—

19 Unbiased recording and identification of thymic cellular interactomes using synthetic Notch receptors

- □□□□□□: 27 / 60
- **AML** □□□□: 10 / 30
- □□□□□□□: 5 / 10
- □□□□: 35 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□ T □□□□□□□□ AML□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Cellular interactions between thymocytes and other immune and stromal thymic cells play a key role in T cell maturation and homeostasis. Previous efforts delineating the cellular interactomes that support T cell development have mostly relied on imaging techniques, genetic deletion of essential molecular factors and bone marrow chimeras. Here, using synthetic NOTCH receptors we took a direct and unbiased genetic approach to fluorescently label cells in physical contact with CD4^(+) and…

—

20 Resident tissue macrophages maintain intraocular pressure homeostasis

- □□□□□□: 35 / 60
- **AML** □□□□: 6 / 30
- □□□□□□□: 5 / 10
- □□□□: 34 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□

—

21 DPEP2 suppresses hyperinflammation via metabolic reprogramming of macrophages in sepsis

- □□□□□□: 29 / 60
- **AML** □□□□: 13 / 30
- □□□□□□□: 5 / 10
- □□□□: 33 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□+□□□□□□□□□□□□□□□□□□

□□: Sepsis-induced excessive inflammation contributes to mortality, but restricting hyperinflammation in sepsis remains challenging. Here, we identify dipeptidase 2 (DPEP2) as an immunotherapeutic target in sepsis by integrating single-cell and bulk RNA sequencing data from septic patients. In patients with sepsis, peripheral monocytes/macrophages have reduced DPEP2 expression, with DPEP2 levels negatively correlating with inflammation severity, disease progression, and clinical outcomes. In vitro,…

—

22 Blood phosphorylated tau elevation as a biomarker in immunoglobulin light chain and transthyretin amyloidosis

- □□□□□□: 30 / 60
- **AML** □□□□: 7 / 30
- □□□□□□□: 6 / 10
- □□□□: 32 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ p-tau □□□□□□ PNP □ AL □□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□□□□

structural dynamics over space and time. In this study, we introduce a photochromic molecule, spiropyran, to modify a specific residue of proteins, thereby achieving a reversible,…

—

24 Anti-PD-1 antibody combined with P-GEMOX chemotherapy versus P-GEMOX chemotherapy with or without autologous stem-cell transplantation for previously untreated advanced natural killer/T cell lymphoma: a retrospective cohort study

- □□□□□□: 31 / 60
- **AML** □□□□: 6 / 30
- □□□□□□□: 6 / 10
- □□□□: 32 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ NKTCL □□□□□□□□□□□□□□□ AML □□□□□□□□LSC□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: We evaluated the effectiveness of adding anti-PD-1 antibody to P-GEMOX regimen in previously untreated advanced-stage natural killer/T-cell lymphoma (NKTCL) compared to P-GEMOX alone or with autologous stem cell transplantation (ASCT). 418 patients (135 and 283 in the immunochemotherapy and chemotherapy groups, respectively) were analyzed from 15 Chinese centers (2014-2023). The median follow-up was 40.7 months. The immunochemotherapy group had a higher objective response rate (ORR) (89.6%…

—

25 Extracellular vesicle engineering using a small scaffold protein

- □□□□□□: 26 / 60
- **AML** □□□□: 12 / 30
- □□□□□□□: 5 / 10
- □□□□: 31 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□ EV □□□□ scaffold□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Extracellular vesicles (EVs) are promising drug-delivery vehicles owing to their biocompatibility and low immunogenicity. Genetic engineering of a membrane-bound EV-sorting scaffold protein empowers EVs by installing targeting moieties on the surface and enriching therapeutic cargo in the lumen. However, the choice of scaffold proteins with simple structures and short sequences is limited. Here, we conduct mass spectrometry-based proteomic studies and identify ENPP1 as a superior scaffold…

—

26 SHMT2 deficiency disrupts transcriptional regulation through homocysteine-mediated suppression of histone lactylation in Huntington's disease models

- □□□□□□: 26 / 60
- **AML** □□□□: 9 / 30
- □□□□□□□: 5 / 10
- □□□□: 31 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□ HD □□-□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□

□□: Huntington’s disease (HD) is a fatal neurodegenerative disorder characterized by progressive motor dysfunction, cognitive decline, and striatal neuron degeneration, primarily affecting medium spiny neurons (MSNs). Despite extensive research, the underlying metabolic vulnerabilities contributing to HD pathogenesis remain poorly understood. In this study, we employ RNA sequencing (RNA-seq) and metabolomics analyses to identify marked dysregulation of one-carbon metabolism in HD. We validate that…

—

27 The role of amygdala GABA neurons in controlling stress and reproduction in female mice

- □□□□□□: 29 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 5 / 10
- □□□□: 30 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Stress can disrupt menstrual cycles, impair fertility and cause reproductive disfunction. The posterodorsal medial amygdala (MePD) integrates stress signals and regulates the gonadotropin-releasing hormone (GnRH) pulse generator through a dense network of GABA and Urocortin-3 (UCN3) neurons, yet the mechanisms underlying the circuitry remain poorly understood. Here, we combine in vivo mini-endoscopic calcium imaging, optogenetics, clustering analysis, and computational modeling to investigate…

—

28 Intraoperative fluorescence-guided fresh frozen sectioning for margin control in head and neck cancer: phase 2 clinical trial

- □□□□□□: 29 / 60
- **AML** □□□□: 6 / 30
- □□□□□□□: 6 / 10
- □□□□: 30 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□/□□□

□□: Accurate and robust prediction of drug-target affinity (DTA) plays a critical role in drug discovery. While deep learning has advanced DTA prediction, existing methods struggle with limited training data and poor generalization. In this study, we propose AdaMBind, a novel DTA prediction model based on meta-learning framework with an adaptive task module designed for low-data scenarios. It employs a dynamic “easy-to-hard” task scheduling mechanism to enhance training efficiency and robustness.…

—

30 Learning turbulent flows with generative models for super resolution and sparse flow reconstruction

- □□□□□□□: 34 / 60
- **AML** □□□□□: 4 / 30
- □□□□□□□□: 4 / 10
- □□□□□: 30 / 100
- □□□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□□

□□□□□: □□□□□□□□□□□□□□□□□□□□□□□□ AI □□□□□□□□ AML□□□□□/□□□□□□□□□□□□□□□□□□□

- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□/□□□□□□□□□□□□□□□□□□□

35 No increased risk of acute myeloid leukemia in adults with primary immune thrombocytopenia treated with thrombopoietin receptor agonists: a French nationwide population-based study

- [redacted]: 24 / 60
- **AML** [redacted]: 12 / 30
- [redacted]: 3 / 10
- [redacted]: 28 / 100
- [redacted]: 2026-03-12
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted]“[redacted]”[redacted]

[redacted]: Not available.

—

36 Behavioral screening defines the molecular Parkinsonism-related subgroups in Drosophila

- [redacted]: 29 / 60
- **AML** [redacted]: 6 / 30
- [redacted]: 5 / 10
- [redacted]: 28 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted]PD[redacted] AML[redacted]

[redacted]: Parkinson’s disease (PD) and related familial Parkinsonism are defined by motor dysfunction, but the specific upstream molecular causes of these clinical symptoms can vary widely. We hypothesize that these causes converge onto a limited number of core cellular pathways. To investigate this, we created a collection of 24 genetically well-controlled Drosophila models of familial forms of PD and related mono-genic forms of Parkinsonism. Using unbiased behavioral screening and machine learning we…

—

37 A systematic review of machine and deep learning techniques for acute lymphoblastic leukemia diagnosis

- [redacted]: 21 / 60
- **AML** [redacted]: 12 / 30
- [redacted]: 6 / 10
- [redacted]: 27 / 100
- [redacted]: 2026-03-10
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] ALL [redacted] AI [redacted] AML [redacted]

[redacted]: Acute lymphoblastic leukemia (ALL) is a hematological malignancy characterized by the rapid proliferation of immature white blood cells in the bone marrow. Early and accurate diagnosis is essential for improving clinical outcomes; however, distinguishing between lymphocytes and lymphoblasts poses significant challenges owing to their subtle morphological similarities. Traditional manual diagnostic methods, which rely on expert evaluations, are inherently time-consuming and subject to human…

—

38 In situ synchrotron X-ray scattering reveals organic-mediated scaling mechanisms on desalination membranes

- □□□□□□: 33 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 26 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□/□□□□□□□□□□□□□□□□□□□

41 ULK4 and CDKN2A polymorphisms influence the risk of developing monoclonal gammopathy of undetermined significance

- □□□□□□: 30 / 60
- **AML** □□□□: 6 / 30
- □□□□□□□: 4 / 10
- □□□□: 25 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ MM □□ MGUS □□□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□

□□: Monoclonal gammopathy of undetermined significance (MGUS) is a necessary precursor condition to multiple myeloma (MM). Given the role of autophagy in modulating MM risk, we investigated whether genetic variation in autophagy-related genes influences susceptibility to MGUS. We analyzed the association of 34,042 common autophagy-related single nucleotide polymorphisms (SNPs) with MGUS across six independent cohorts, five from Europe and one from North America, comprising 2317 MGUS cases and…

—

42 Protection of telomeres 1b safeguards the Arabidopsis genome by regulating ROS homeostasis

- □□□□□□: 27 / 60
- **AML** □□□□: 5 / 30
- □□□□□□□: 5 / 10
- □□□□: 25 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□□□□

□□: Telomeres safeguard the genome, acting as sentinels of oxidative stress and preventing chromosome ends from eliciting a DNA damage response. PROTECTION OF TELOMERES 1 (POT1) is a highly conserved telomere protein, essential for chromosome integrity and telomeric DNA replication. Arabidopsis thaliana encodes two divergent POT1 paralogs: AtPOT1a stimulates telomerase activity, but AtPOT1b function is unknown. Here we show that AtPOT1b modulates reactive oxygen species (ROS) homeostasis. Oxidative…

—

43 Reconstructing fine-scale 3D wind fields with terrain-informed machine learning

- □□□□□□: 28 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 5 / 10
- □□□□: 25 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□/□□□□□□□□□ AI □□□□□□□□□□□AML □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Fine-scale near-surface wind field prediction is essential for a wide range of applications. However, most operational and AI-based weather models operate at kilometer-scale resolution, where terrain-induced wind features such as slope jets, flow deflection, and recirculation are systematically averaged out. Here we

introduce FuXi-CFD, a machine learning-based framework designed to generate detailed three-dimensional (3D) near-surface wind fields at 30-meter horizontal resolution, using only…

—

44 Predicting rice drought-responsive genes via distance-based prototypical graph neural network with path aggregation mechanism

- [illegible]: 24 / 60
- **AML** [illegible]: 7 / 30
- [illegible]: 5 / 10
- [illegible]: 24 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: Drought is a primary factor that adversely impacts rice yield and quality. Identifying drought-responsive genes is essential for developing drought-responsive cultivars. Recently, graph neural network methods based on embedding learning have shown considerable success in biological networks. However, challenges still remain in adequately capturing node attribute features, representing topological structures, and addressing class imbalance, which may constrain the model's predictive capability.…

—

45 Abrasion-resistant wearable skins based on bilayered solid/liquid stretchable conductors

- [illegible]: 24 / 60
- **AML** [illegible]: 4 / 30
- [illegible]: 5 / 10
- [illegible]: 24 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible] LSC[illegible]

[illegible]: Soft bioelectronic skins represent next-generation wearable technologies for continuous and unobtrusive biomonitoring, enabled by diverse active materials and topological designs. Despite encouraging progress, most wearable skins reported to date remain unable to match the abrasion-resistant sensing functions of human skin. Here, we present abrasion-resistant wearable skins based on a bilayer stretchable conductor architecture. The top layer, composed of silver particle (AgPs)-impregnated…

—

46 HK2-driven histone H3K18 lactylation promotes stromal cell senescence and decidualization deficiency in URSA via CUX1-mediated SASP factor transcription

- [illegible]: 23 / 60
- **AML** [illegible]: 9 / 30
- [illegible]: 5 / 10
- [illegible]: 24 / 100
- [illegible]: 2026-03-10
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]-[illegible]-SASP [illegible] URSA [illegible] AML [illegible]

□□: CONCLUSIONS: Our findings define a novel HK2-H3K18la-CUX1-SASP signaling axis that drives URSA pathogenesis by linking metabolic reprogramming with epigenetic regulation. This work highlights CUX1 as a potential therapeutic target for correcting decidualization deficiency in URSA.

—

47 Design of miniprotein inhibitors targeting complement C9 to block membrane attack complex assembly

- □□□□□□: 22 / 60
- **AML** □□□□: 8 / 30
- □□□□□□□: 5 / 10
- □□□□: 23 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

□□: The regionalized structure of the intestinal epithelium is critical for its function, and the risk for certain diseases has a regional bias. However, how regionalization is established and how it influences disease susceptibility remain poorly understood. Here, we investigated the role of the gut microbiome-the regionalized community of microorganisms in the intestinal lumen-in promoting regionalization of the colon. We found that the proximo-distal identity of colonocytes along the organ's…

—

50 Multi-modal characterization of nitrate reduction nano-catalysts with periodic strain distribution

- □□□□□□: 27 / 60
- **AML □□□□**: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 23 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□/□□□□□□□□□□□□□□□□□□□□□□□□□□□□AML □□□□□□ □□□□□□□□

□□: Strain engineering serves as a pivotal strategy to optimize catalytic activity in electrocatalysis. However, the catalyst sizes under industrial conditions are usually large and even beyond nanometer regime. The critical methodological limitations on strain imaging of such catalysts with both large field of view and high spatial resolution obscure the mechanistic understanding of strain-performance correlations. Here, we present an optimized four-dimensional scanning transmission electron…

—

51 Perovskite solar cells with enhanced thermal fatigue resistance under extreme temperature cycling

- □□□□□□: 30 / 60
- **AML □□□□**: 2 / 30
- □□□□□□□: 5 / 10
- □□□□: 23 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]/[illegible] AML [illegible] LSC[illegible]

[illegible]: Two-dimensional ferromagnetic materials hold great promise for advancing low-power, high-integrated spintronic devices due to their atomic flat surfaces and versatile interfacial modulation. However, achieving a combination of room-temperature, field-free spin-orbit torque switching with tunable polarity in wafer-scale vdW heterostructures remains a significant challenge. Here, we demonstrate polarity-tunable, field-free spin-orbit torque switching in an all-vdW Bi(2)Te(3)/Fe(4)GeTe(2)…

—

53 Longitudinal mental health data collected via the Corona Health smartphone app during COVID-19

- [illegible]: 24 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 5 / 10
- [illegible]: 22 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]LSC[illegible]

[illegible]: Mental health impacts during the COVID-19 pandemic underscored the importance of real-time assessment methods to capture population-level changes (e.g., longitudinal changes in quality of life). This dataset contains questionnaire responses collected with the Corona Health app, a multilingual mHealth app available on Android and iOS platforms. The dataset includes baseline from 2,704 participants (i.e., adults aged 18 years and older, living in Germany) and 11,541 repeated ecological momentary…

—

54 Amorphous/crystalline interwoven multipods with high Co/Ni activity for wide-temperature-range sodium-sulfur batteries

- [illegible]: 26 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 5 / 10
- [illegible]: 22 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: Sluggish kinetics caused by 16-electron transfer hinders development of wide-temperature-range sodium-sulfur batteries. Here we report Sn-doped CoNiS multipods with an amorphous-crystalline interwoven structure. Employed as a positive electrode catalyst, the resulting sodium-sulfur battery exhibits a discharge capacity of 1320.8 mAh g^(−1) at 3 A g^(−1) after 1200 cycles at room temperature, together with stable and high-capacity electrochemical performance ranged from −20 to 50 °C. It has been…

—

55 Structural dynamics and immunogenicity of the recombinant and outer membrane vesicle-embedded Meningococcal antigen NadA

- [illegible]: 26 / 60

- **AML □□□□**: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 22 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□

□□: Structural knowledge of antigens in their native state can drive the design of optimized vaccine antigens that mimic the native epitope exposure and conformation. Here, by hydrogen-deuterium exchange mass spectrometry, we assessed the structural features of Neisseria Adhesin A (NadA), a meningococcal trimeric outer membrane protein, included as soluble recombinant antigen in the 4CMenB vaccine. We propose a structural annotation of the recombinant NadA and compare its structural dynamics with…

—

56 Immune austerity: How the body balances its books when calories run low

- □□□□□□: 23 / 60
- **AML □□□□**: 6 / 30
- □□□□□□□: 5 / 10
- □□□□: 22 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ Menezes-Silva et al.□ Immunity □□□□□□□□□□□□□□□□□□□□□□

□□: Immune responses are energetically costly and must adapt to systemic nutrient availability. How immunity is reorganized to preserve host glucose during caloric restriction remains unclear. In this issue of Immunity, Menezes-Silva et al. show that dietary restriction, via glucocorticoids, rewires innate and adaptive immunity, and this leads to the conservation of glucose while maintaining protective host defense.

—

57 A hierarchical theranostic nanoagent for multimodal imaging and targeted foam cell intervention in atherosclerosis

- □□□□□□: 21 / 60
- **AML □□□□**: 6 / 30
- □□□□□□□: 5 / 10
- □□□□: 21 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□

□□: Atherosclerosis (AS) is a leading cause of cardiovascular and cerebrovascular disease and is driven by lipid accumulation and chronic inflammation within arterial plaques. Foam cells play a central role in plaque growth and destabilization, yet their precise diagnosis and targeted regulation remain challenging. Here we present a hierarchical nanoagent that integrates multimodal imaging with selective foam cell intervention for improved AS management. The molecular probe enables near-infrared II…

—

58 Dynamic stabilization of a mechanical oscillator in the absence of any stable feature

- □□□□□□: 27 / 60
- **AML □□□□**: 3 / 30

- □□□□□□□: 5 / 10
- □□□□: 21 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□□□□

61 Lichtman MA, Rowe JM. Hyperleukocytic leukemias: rheological, clinical, and therapeutic considerations. Blood. 1982;60(2):279-283

- □□□□□□: 16 / 60
- **AML** □□□□: 14 / 30
- □□□□□□□: 5 / 10
- □□□□: 20 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□ 1982 □□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□

□□: No abstract

—

62 Big-eaters choking? Macrophages can't stomach microplastics

- □□□□□□: 21 / 60
- **AML** □□□□: 5 / 30
- □□□□□□□: 5 / 10
- □□□□: 20 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□

- [redacted]: 20 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]/[redacted]

[redacted]: Gel-based soft materials are attractive for flexible electronics and biointerfaces but are often limited by insufficient mechanical robustness and constrained functional integration. Here, we introduce a geometry-programmed self-wrinkling strategy that enables the spontaneous formation of aligned wrinkle architectures during thermal-evaporative gelation of poly(vinyl alcohol)-based organo-hydrogels. Without external patterning or post-processing, this process produces materials with enhanced…

—

65 Antibody responses against bacterial glycans affinity mature and diversify in germinal centers

- [redacted]: 25 / 60
- **AML** [redacted]: 4 / 30
- [redacted]: 5 / 10
- [redacted]: 20 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]

[redacted]: Anti-carbohydrate antibodies (Abs) play crucial roles in pathogen control, but their generation remains poorly understood. Here, we studied how B cell immunity against the Streptococcus pyogenes surface polysaccharide is influenced by age and antigen (Ag) exposure across human blood, spleen, and tonsils. The glycan-targeted response shifted from immunoglobulin M (IgM) biased in children toward IgG and IgA biased in adults. Both natural colonization and controlled human infection with S. pyogenes…

—

66 A foundation model for multi-task cross-distribution restoration of fluorescence microscopy images

- [redacted]: 21 / 60
- **AML** [redacted]: 6 / 30
- [redacted]: 5 / 10
- [redacted]: 20 / 100
- [redacted]: 2026-03-11
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AI [redacted] AML [redacted]/[redacted]

[redacted]: Deep learning has demonstrated remarkable abilities in restoring fluorescence microscopy images degraded by noise, blur, or undersampling. However, most existing models are task-specific and trained on limited, homogeneously distributed data, which restricts their generalizability and practicality. Here, we present FluoResFM, a foundation model for multi-task and cross-distribution fluorescence microscopy image restoration in a unified model. FluoResFM leverages textual prior information to…

—

67 DDX5 orchestrates RNA homeostasis to ensure oocyte developmental competence

- [redacted]: 23 / 60

- **AML** □□□□: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 19 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ RNA □□□□ AML □□□□□/□□□□□□□□□□□□□□□□

□□: Oocyte development requires tight regulation of transcription and RNA metabolism, which is coordinated by RNA-binding proteins, whose roles in mammalian oogenesis remain incompletely understood. Here, we identify the DEAD-box RNA helicase DDX5 as a key regulator of RNA homeostasis in oocytes. Oocyte-specific deletion of DDX5 leads to female sterility, which is characterized by defective chromatin remodeling, meiotic arrest, increased aneuploidy, and fertilization failure. Mechanistically, DDX5…

—

68 Scalable and stretchable 1D multifunctional fibers for multimodal sensing and stimulation

- □□□□□□: 22 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 5 / 10
- □□□□: 19 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

70 Highly ordered mesoporous TiO₂ nanomeshes with tunable pore periodicity via self-limiting modular monolayer assembly of monomicelles

- □□□□□□□: 24 / 60
- **AML** □□□□□: 2 / 30
- □□□□□□□□: 5 / 10
- □□□□□: 19 / 100
- □□□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□□

□□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

- **AML** □□□□: 9 / 30
- □□□□□□□: 5 / 10
- □□□□: 19 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ NK □□□□□□□□□□□□□□□□□□□□

- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□LSC□□□□□□□□□□□□□□□□□□□

- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□

- □□: □□□□

□□□□: □□□□□□□□□□□□□□ TNBC □□ HORMAD1“□□”□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□+□□□□□□□□□□□□□□□

- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□

- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□/□□□□□□□□□□□□□□□□□□□□

□□: Spin defects in solids offer promising platforms for quantum sensing and memory due to their long coherence times and optical addressability. Here, we integrate a single nitrogen-vacancy (NV) center in diamond with scanning probe microscopy to detect, read out, and spatially map spin-based quantum sensors at the nanoscale. Using the boron vacancy (VB-) center in hexagonal boron nitride-an emerging two-dimensional spin system-as a model, we detect its electron spin resonance indirectly via…

—

89 CF2H: a cell-free two-hybrid platform for rapid protein binder screening

- □□□□□□: 19 / 60
- **AML** □□□□: 7 / 30
- □□□□□□□: 5 / 10
- □□□□: 16 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ de novo □□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

□□: Protein binders that detect, activate, inhibit, or otherwise modulate their targets are pivotal for biomedical applications. With the increasing accuracy and accessibility of de novo protein design, faster and cheaper experimental screening methods would democratize and accelerate the identification of high-affinity binders. Here we present Cell-Free Two-Hybrid (CF2H), a rapid and sensitive method for detecting high-affinity protein-protein interactions (PPI) that does not require cloning,…

—

90 Massive-scale spatial multiplexing of multimode VCSELs with a 3D-printed photonic lantern

- □□□□□□: 21 / 60
- **AML** □□□□: 2 / 30
- □□□□□□□: 5 / 10
- □□□□: 16 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□ AML□□□□□□□□□/□□□□□□□□□□□□□□□□

95 RejuAgro A as an antimicrobial for fire blight control of pome fruits and beyond

- □□□□□□: 22 / 60
- **AML** □□□□: 1 / 30
- □□□□□□□: 5 / 10
- □□□□: 14 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]

[illegible]: No abstract

—

99 Predictors of Recurrence after Curative Surgery for Stage I-III Rectal Adenocarcinoma: A Single-Center Retrospective Cohort Study

- [illegible]: 21 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 4 / 10
- [illegible]: 13 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]LSC[illegible]

[illegible]: CONCLUSION: Our study identifies tumor downstaging after neoadjuvant therapy as a prognostically relevant factor. This finding emphasizes the necessity of incorporating treatment response into postoperative risk stratification to tailor an individualized surveillance strategy.

—

100 Combined physical and pharmacological anabolic osteoporosis therapies increase bone response and mechanoregulation in female mice

- [illegible]: 18 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 5 / 10
- [illegible]: 13 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: Bone's ability to adapt to mechanical demands is governed by mechanoregulation, the process by which cells sense and respond to mechanical stimuli to maintain skeletal integrity. In osteoporosis, increased bone resorption activity leads to structural deterioration and elevated fracture risk. While existing pharmacological therapies aim to restore bone mass to reduce fracture risk, it is unclear how they modulate mechanoregulation, especially when combined with physical interventions. Here, we…

—

101 Transfusing HPA-mismatched platelets to mimic fetomaternal hemorrhage elicits fetal/neonatal alloimmune thrombocytopenia

- [illegible]: 18 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 4 / 10
- [illegible]: 13 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] FNAIT [illegible] AML [illegible]

□□: Fetal/neonatal alloimmune thrombocytopenia (FNAIT) is a bleeding disorder in which maternal antibodies target fetal and neonatal platelet alloantigens, most commonly human platelet alloantigen-1a (HPA-1a), resulting in fetal and neonatal thrombocytopenia severe enough to cause life-threatening organ bleeds, such as intracranial hemorrhage. Hemolytic disease of the fetus and newborn (HDFN) is an analogous disease caused by maternal exposure to fetal red blood cell (RBC) alloantigens, most…

—

102 Ultra-efficient physical field computing by complex-valued network quantization

- □□□□□□: 18 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 4 / 10
- □□□□: 13 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

□□: Background: During the SARS-CoV-2 pandemic, many countries restricted visits to nursing homes to reduce infections among residents and staff. This also affected the residents' relatives, as personal contact was no longer possible. Aim: As part of a research network, we investigated the experiences, challenges, strategies, and consequences of visiting restrictions in nursing homes during the early phase of the pandemic from the perspective of relatives and stakeholders advocating residents'…

—

105 Intrinsic capacity and stroke risk in a multiple cohort study

- □□□□□□: 20 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 4 / 10
- □□□□: 10 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□ LSC□□□□□□□□□□□□□□□□□□□

108 Textured piezoelectric ceramics with reduced grain size for high-frequency transducer applications

- □□□□□□□: 17 / 60
- **AML** □□□□□: 1 / 30
- □□□□□□□□: 4 / 10
- □□□□□: 8 / 100
- □□□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□□

□□□□□: □□□□□□□□

- □□□□□□□: 3 / 10
- □□□□: 7 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□/AML □□□□□□□□□□□□□□□□

□□: Unconventional superconductivity typically emerges out of a strongly correlated normal state, manifesting as a highly renormalised Fermi liquid or a strange metal with T-linear resistivity. In Ruddlesden-Popper bilayer nickelates, superconductivity with a critical temperature T(c) exceeding 80 and 40 K has been respectively realised in pressurised bulk crystals and epitaxially strained thin films. These advancements call for the characterisation of fundamental normal-state and superconducting…

—

112 Current treatment algorithm: diffuse large B cell lymphoma

- □□□□□□□: 9 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□□: 5 / 10
- □□□□: 7 / 100
- □□□□: 2026-03-10
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□ DLBCL □□□□□□□□□□□□□□□□□□□□

- [illegible]: 5 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]/[illegible]

[illegible]: No abstract

—

115 Pediatric acute erythroid leukemia with NUP98::KDM5A fusion

- [illegible]: 5 / 60
- **AML** [illegible]: 12 / 30
- [illegible]: 3 / 10
- [illegible]: 5 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]/[illegible] AML [illegible]

[illegible]: No abstract

—

116 When CAR T cells become the disease

- [illegible]: 9 / 60
- **AML** [illegible]: 9 / 30
- [illegible]: 2 / 10
- [illegible]: 5 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]

[illegible]: No abstract

—

117 Circulating T follicular helper cells as emerging biomarkers in pediatric Evans syndrome

- [illegible]: 8 / 60
- **AML** [illegible]: 7 / 30
- [illegible]: 4 / 10
- [illegible]: 5 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: Not available.

—

118 The rationale for targeting nectin-4 in pancreatic cancer

- [illegible]: 15 / 60
- **AML** [illegible]: 2 / 30
- [illegible]: 5 / 10

- □□□□: 5 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□ PDAC □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□□□□

□□: Nectin-4 is an immunoglobulinlike cell adhesion molecule that contributes to tumor aggressiveness and immune evasion in pancreatic ductal adenocarcinoma (PDAC). Its high and selective expression on pancreatic cancer cells, together with the clinical efficacy of nectin-4-targeted antibody-drug conjugates in other solid tumors, establishes nectin-4 as a compelling therapeutic target in PDAC.

—

119 Diversify and conquer: Antibody feedback in the germinal center

- □□□□□□: 12 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 4 / 10
- □□□□: 5 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

122 National statistics are in crisis around the world - and the impacts will be severe

- [illegible]: 11 / 60
- **AML** [illegible]: 2 / 30
- [illegible]: 4 / 10
- [illegible]: 4 / 100
- [illegible]: 2026-03-12
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: No abstract

—

123 Helicobacter pylori Screen-and-Treat Programs for Gastric Cancer Prevention - IARC Working Group Report

- [illegible]: 13 / 60
- **AML** [illegible]: 1 / 30
- [illegible]: 3 / 10
- [illegible]: 4 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: No abstract

—

124 Design of a configurable SoC for Alzheimer’s disease detection based on multimodal signals

- [illegible]: 10 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 3 / 10
- [illegible]: 4 / 100
- [illegible]: 2026-03-11
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible] LSC[illegible]

[illegible]: No abstract

—

125 AAPM task group 317 report: A joint AAPM and ESTRO report on brachytherapy catheter, needle, and applicator tracking technology

- [illegible]: 11 / 60
- **AML** [illegible]: 1 / 30
- [illegible]: 4 / 10
- [illegible]: 4 / 100
- [illegible]: 2026-03-10
- **DOI**: N/A
- [illegible]: [illegible]

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□ LSC□□□□□□□□□□□□□□□□□□□

—

129 Microglia Dld-K127 lactylation promotes Parkinson's disease via regulating the metabolism of lactate-pyruvate transformation

- □□□□□□: 12 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 3 / 10
- □□□□: 1 / 100
- □□□□: 2026-03-11
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□