追觅 - 2026-03-21

生成时间: 2026-03-21 07:54

分析文章数量: 60

1 Spatial heterogeneity of MDSCs mediated by ANXA1-FPRs signaling drives immune suppression in OSCC progression

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 20 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 46 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇利用单细胞与空间转录组解析 OSCC 免疫抑制微环境的观察性研究，聚焦 MDSCs 异质性与 ANXA1-FPRs，但与 AML 无直接关联，仅在肿瘤微环境层面存在有限可迁移性。；置信度：低

摘要: Limited efficacy of immunotherapy in oral squamous cell carcinoma (OSCC) is driven by an immunosuppressive tumor microenvironment, yet the role of intratumoral spatial heterogeneity in immune responses remains unclear. Here, we employ single-cell and spatial transcriptomics to dissect the cellular composition and spatial organization of OSCC. We find CD8^(+) T cells are spatially localized yet functionally suppressed in late-stage OSCC, while myeloid-derived suppressor cells (MDSCs) transition…

—

2 Chromatin reorganization drives overexpression of a Btaf1 variant underpinning hematopoietic aging

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 20 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 45 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；该文提出染色质重塑驱动特定转录本的 HSC 衰老机制，具有潜在原创性，但基于极有限摘要文本，无法评估证据强度与 AML 直接迁移价值。；置信度：低

摘要: Age-associated hematopoietic stem cell (HSC) dysfunction is accompanied by dramatic transcription changes, but it remains unclear whether specific transcripts could orchestrate these HSC aging phenotypes. Here, we perform epigenetic profiling in male mice to investigate the regulatory mechanisms underlying the HSC aging transcriptome and screen for potential aging driver genes. We identify a looping structure formed between part of the Btaf1 gene and the whole Ide gene in old HSCs which is…

—

3 Celcomen: spatial causal disentanglement for single-cell and tissue perturbation modeling

- **研究本身价值**: 36 / 60
- **AML 借鉴价值**: 13 / 30

- **表达与可复用性**: 6 / 10
- **最终总分**: 43 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯方法学创新论文，聚焦通用空间转录组因果建模，与 AML 基础研究无任何直接关联，仅在方法层面可能存在极有限的间接启发。；置信度：低

摘要: Celcomen leverages a mathematical causality framework to disentangle intra- and inter-cellular gene regulation programs in spatial transcriptomics data through a generative graph neural network. It is a first step towards perturbation models of Virtual Tissues and can generate post-perturbation counterfactual spatial transcriptomics, thereby offering access to experimentally inaccessible samples. We validated its disentanglement, identifiability of causal structure, and counterfactual prediction…

—

4 NANP targeting radiosensitizes glioblastoma through TNFR1 sialylation-driven mesenchymal shift

- **研究本身价值**: 34 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 42 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文在 GBM 放射抵抗领域提出 NANP 新靶点及糖基化机制，具有一定原创性，但属于非 AML 研究，AML 借鉴价值极低。；置信度：低

摘要: Glioblastoma (GBM) patients have dismal survival due to resistance to initial ionizing radiation therapy (RT). Clonal evolution analysis reveals no dominant RT-resistant clones, prompting a genome-wide CRISPR screen to identify radiosensitizing targets. The screening highlights DNA damage response genes, validating the effectiveness of our approach. N-acylneuraminate-9-phosphatase (NANP), a critical enzyme in the sialic acid synthetic pathway, is top-ranked in the screening and associated with…

—

5 Comparative effectiveness of HMA with venetoclax vs intensive chemotherapy in AML with very high-risk cytogenetics

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 15 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 39 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦临床治疗选择，但完全缺乏基础机制、生信或功能验证，对 AML 基础研究而言仅为背景参考而非核心贡献。；置信度：低

摘要: The optimal frontline therapy for acute myeloid leukemia (AML) with very high-risk cytogenetics (vHRC)-defined by complex karyotype (CK), monosomal karyotype (MK), or

inv(3)/t(3;3)-remains uncertain. We retrospectively analyzed 358 newly diagnosed AML-vHRC cases treated at five academic centers (2014-2024), stratified by intensive chemotherapy (IC) vs hypomethylating agent plus venetoclax (HMA+ven). Cytogenetic features included CK in 90.2%, MK in 64%, and inv(3)/t(3;3) in 9.8%; TP53 mutations…

—

6 Psoriasis modulates inflammatory bowel disease risk and intestinal epithelium lipid metabolism via IL-1β-producing macrophages

- **研究本身价值**: 35 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 37 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于临床队列与小鼠模型提出银屑病-IBD 共病的新机制，但与 AML 基础研究领域完全无关。；置信度：低

摘要: Patients with psoriasis have an increased risk of developing inflammatory bowel disease, yet the mechanisms underlying this comorbidity remain unclear. In a clinical cohort, we observed an inverse correlation between psoriasis severity and postprandial plasma apolipoprotein B48 levels, a finding recapitulated in experimental psoriasis mouse models, indicating impaired intestinal lipid handling. To directly interrogate this process, we developed a recombinant photoconvertible apolipoprotein B…

—

7 Inhibition of p300/CREBBP catalytic activity drives context-dependent transcriptional activation in AML

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 20 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 37 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文提出一个在 AML 中看似矛盾的转录激活现象，但本次评估仅基于标题与部分摘要，核心证据、实验细节及验证结果均未呈现，无法判断结论是否成立。；置信度：低

摘要: The lysine acetyltransferase (KAT) activity of p300/CREBBP has traditionally been linked to transcriptional activation. This has been attributed largely to acetylation of histone residues such as H3K27ac, a defining hallmark of active regulatory elements. Here we show that, in acute myeloid leukemia (AML), inhibition of p300/CREBBP catalysis can paradoxically increase transcription. We combined time-resolved dynamics of nascent and total transcription with chromatin binding dynamics of…

—

8 Lactate metabolism and protein lactylation in colorectal cancer: from metabolic reprogramming to epigenetic regulation

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 10 / 30

- **表达与可复用性**: 5 / 10
- **最终总分**: 36 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇以 CRC 为焦点的综述文章，整合乳酸代谢与 Kla 的现有文献，但无原创实验数据，与 AML 基础研究无直接关联。；置信度：低

摘要: Colorectal cancer (CRC) exhibits profound metabolic reprogramming, in which excessive lactate accumulation remodels the tumor microenvironment and promotes immune suppression, angiogenesis, and therapeutic resistance. Recent studies reveal that lactate also serves as a substrate for lysine lactylation (Kla), linking metabolic overflow to epigenetic regulation. This review focuses on CRC but also incorporates mechanistic data from other tumor models when CRC-specific evidence is limited,…

—

9 IVNS1ABP mutation drives cellular senescence in newly identified progeroid neuropathy

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 35 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇针对罕见遗传性早老综合征的初步机制探索，采用 isogenic iPSC 模型，但与 AML 基础研究（LSC、克隆演化、耐药、造血微环境等）无逻辑可迁移性，AML 借鉴价值极低。；置信度：低

摘要: We identified a new progeroid syndrome with severe neuropathy and intellectual deficits but its underlying cellular and molecular mechanism is unknown. Exome sequencing revealed a homozygous mutation in the IVNS1ABP gene, which encodes IVNS1ABP, an influenza virus non-structural protein-1 binding protein. To investigate disease mechanisms, we generated isogenic induced pluripotent stem cells (iPSCs) from patient fibroblasts and differentiated them into neural progenitor cells (NPCs). Mutant…

—

10 Enhancer-mediated Etv4 activation stimulates osteogenic differentiation

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 35 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦骨质疏松相关增强子异常与成骨分化机制，与 AML（急性髓系白血病）基础研究无直接关联。；置信度：低

摘要: Enhancers, as cis-regulatory elements, play pivotal roles in transcriptional homeostasis. The abnormality in enhancers is highly associated with various diseases, including osteoporosis. However, the landscape of active enhancers underlying bone diseases remains incomplete. By conducting an integrative analysis of transcriptome and ChIP-seq data, we identify enh11 as an active enhancer during osteoblastogenesis. CRISPR/Cas9-mediated deletion of enh11 inhibits cell differentiation of…

—

11 Targeting BMP and TAZ/TEAD mechanotransduction pathways impairs acute myeloid leukemia chemoresistance

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 19 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；文章基于摘要提出一个将骨髓力学微环境与 AML 耐药关联的新视角，但提供的文本仅限标题与部分摘要，无任何实验数据、方法、结果或验证信息，无法实质评估研究质量。；置信度：低

摘要: Despite extensive research and intensive use of chemotherapies in clinics, the 5-year overall survival of acute myeloid leukemia (AML) patients does not exceed 20%. The clonal expansion of leukemic blasts leads to modifications of the bone marrow physical properties, including increased extracellular matrix stiffening, upregulation of intramedullary pressure and reduction of the space available for cells. These biomechanical modifications are speculated to alter therapeutic response and cause…

—

12 Engineering programmable CAR and antigen pairing via drug-gated light activation

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 33 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦固体瘤 CAR-T 工程优化的方法学文章，与 AML 基础研究（LSC、克隆演化、耐药、微环境等）无直接关联，仅在 CAR-T 通用技术层面有极有限的间接启发。；置信度：低

摘要: Chimeric antigen receptor (CAR) T-cell immunotherapy has achieved clinical success, but remains limited in solid tumors by antigen escape and tumor heterogeneity. Here, we engineer a high-affinity R-phycoerythrin (PE)-binding monobody to create a PE-programmable CAR toward user-defined antigens sequentially or simultaneously. To reduce off-tumor toxicity, we implement a drug-gated light-activation approach (DGLA) strategy to spatially confine CAR function. We further couple DGLA-controlled tumor…

—

13 CD177^+ neutrophil-platelet aggregates contribute to thromboinflammation via NETs in necrotizing enterocolitis

- **研究本身价值**: 34 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 33 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦早产儿坏死性小肠结肠炎（NEC）的局部免疫血栓形成，与AML基础研究无直接关联，核心结论依赖临床样本与功能验证但本文仅提供摘要片段，无法充分评估。；置信度：低

摘要: Necrotizing enterocolitis (NEC) is a severe gastrointestinal disease affecting premature infants, yet its precise pathogenic mechanisms remain unclear. While immunothrombosis, an interplay between inflammation and coagulation, is well recognized in systemic inflammatory diseases, its role in NEC pathogenesis and local intestinal regulation has been incompletely characterized. Here, we combine multi-omics profiling, histopathological analyses, and functional studies using clinical NEC samples and…

—

14 Serial PNA-Transformer-Based Virtual Screening Identifies Nanomolar DYRK1A Inhibitors for Pancreatic Ductal Adenocarcinoma

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇计算化学驱动的药物筛选论文，聚焦PDAC靶点鉴定，证据链极短且完全与AML无关。；置信度：低

摘要: Dual-specificity tyrosine-regulated kinase 1A (DYRK1A) is a promising therapeutic target for pancreatic ductal adenocarcinoma (PDAC). Herein, we developed an integrated AI and structure-based pipeline featuring a Serial PNA-Transformer graph neural network, which achieved a test AUC of 0.8901. Multistage screening of 21,738 compounds prioritized 232 candidates across 10 chemical clusters. Enzymatic assays confirmed three hits with IC(50) values <500 nM; notably, CX-6258 (IC(50) = 473.7 nM)…

—

15 An orthogonal RNA aptamer for targeted RNA regulation in phase-separated subcellular compartments

- **研究本身价值**: 33 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用 RNA 生物学工具开发论文，基于有限摘要提出新平台，但无任何实验数据、验证结果或 AML 相关内容，学术可信度与 AML 借鉴价值均极低。；置信度：低

摘要: Subcellular compartments organize RNAs into phase-separated condensates, significantly influencing RNA metabolism. However, the study of how specific RNAs regulate interacting factors and their phenotypic outcomes is hindered by the lack of advanced imaging and regulation tools. To address this, we developed an orthogonal RNA aptamer system, Clivia-HT, which integrates a fluorescent imager with a ribonuclease-targeting chimera (RIBOTAC) degrader. This platform enables simultaneous RNA…

—

16 Transplantation of encapsulated mitochondria alleviates dysfunction in mitochondrial and Parkinson’s disease models

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦线粒体递送技术创新的治疗性研究，与 AML 基础研究无直接关联，迁移价值极低。；置信度：低

摘要: Mitochondrial transplantation holds significant potential for the treatment of mitochondrial diseases. However, how to efficiently deliver exogenous mitochondria to somatic cells or tissues remains unresolved. We present a mitochondrial transplantation approach to deliver mitochondria into the cells and tissues of mice and monkeys with high efficiency, based on encapsulating mitochondria with vesicles derived from the plasma membrane of erythrocytes. Treatment with encapsulated mitochondria…

—

17 TCB-Kla: a hybrid deep learning model for predicting human lysine lactylation sites by incorporating transformer encoder and multi-scale CNN

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用计算生物信息学工具开发论文，聚焦 PTM 位点预测，但与 AML 或造血系统无任何直接关联或实验映射。；置信度：低

摘要: Lysine lactylation as a newly discovered post-translational modification of proteins, plays a key role in various cellular processes. It can be stimulated by lactate and regulates gene expression and life activities. Mass spectrometry is currently the fundamental method for identifying post-translational modification sites, but this approach is often time-consuming and labour-intensive. In this study, we propose a hybrid deep learning model, termed TCB-Kla, for predicting human lysine…

—

18 Systematic identification of variant-specific RNA structure-small molecule interactions exemplified by RNA G-quadruplexes

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯方法学导向的论文，聚焦 RNA 靶向小分子与变异的交互鉴定，但与 AML 基础研究无任何直接关联，且本次评估仅基于标题与摘要片段。；置信度：低

摘要: Individual genetic variations, such as cancer-associated somatic mutations, alter RNA structures, thereby potentially enhancing or inhibiting the binding of RNA-targeting small molecules. However, to date, no approach has been available to identify these variant-specific RNA-small molecule interactions due to technical limitations. Here, we present Binding- and Vinyl-Quinazolinone-Induced Deletion-Based Mutational Profiling (BIVID-MaP), a high-throughput method for detecting RNA-small molecule…

—

19 Concise synthesis of bufogargarizin B by a conformation-controlled skeletal reorganization approach

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯有机化学总合成论文，聚焦 19-norbufadienolide 类天然产物的骨架构建，与 AML（急性髓系白血病）基础研究无任何生物学、机制或临床关联。；置信度：低

摘要: Bufogargarizins are a class of unique 19-norbufadienolides isolated from the venom of Bufo bufo gargarizans. Here we report a concise synthesis of bufogargarizin B from inexpensive, commercially available dehydroepiandrosterone by means of a conformation-controlled skeletal reorganization approach. The synthesis features the rapid construction of the 5/7/6/5 tetracyclic core framework by means of a SmI(2)-mediated ketone-allylic acetate transannular cyclization, and the installation of…

—

20 Permeable nanoreactor eye drop for enzymatic cascade-mediated treatment for acute retinal injury model mimicking geographic atrophy

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-18

- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为眼科 AMD 相关纳米递送治疗研究，与 AML 基础研究无直接或可迁移关联。；置信度：低

摘要: Geographic atrophy (GA), the most prevalent form of age-related macular degeneration (AMD), remains clinically challenging due to its complex pathogenesis. Using an acute retinal injury model that mimics GA pathology (oxidative stress-induced retinal apoptosis), we identify enzymatic cascade restoration as a promising therapeutic strategy. To overcome ocular delivery barriers, we engineer zwitterionic nanocages for co-delivering catalase (CAT) and superoxide dismutase (SOD) (Z(CAT&SOD)). These…

—

21 Predator-mediated local convergence fosters global microbial community divergence

- **研究本身价值**: 33 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇生态学领域跨尺度整合研究，设计思路较新颖，但与 AML 基础研究（包括 LSC、微环境、克隆演化等）完全无关。；置信度：低

摘要: Understanding how microbial communities assemble is central to predicting ecosystem function. Although predators strongly influence bacterial communities through predation, the role of microbial predators in modulating global microbial divergence and convergence patterns remains largely neglected. Here, we integrated global-scale amplicon sequencing data, controlled field experiments, and reconstructions of natural and synthetic communities to examine predator-mediated community assembly…

—

22 Treatment of Acute Myeloid Leukemias and Myelodisplastic Syndromes Relapsing After Allogeneic Stem Cell Transplantation: An In-Depth Analysis of the GITMO AML/MDS-Relapse Registry Study

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 15 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；文章仅呈现临床结局比较结论（HMA ORR 较低但 OS 相当，DLI 独立有益），无任何机制、功能或基础层面探讨，属于纯临床回顾性注册分析。；置信度：低

摘要: CONCLUSIONS: Despite the lower ORR observed with HMA-based therapy, the long-term OS was comparable to that observed with other therapies. The immune control of the disease relapse with DLI is of benefit, independently from the salvage therapy. .

—

23 Histone lactylation-driven feedback loop modulates pyrimidine metabolism to promote oral carcinogenesis

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；基于极有限的标题与摘要片段，该文仅初步提示乳酰化在 OLK→OSCC 进展中的水平变化及代谢-表观反馈，但证据链极不完整，且与 AML 完全无关。；置信度：低

摘要: Metabolic reprogramming and epigenetic alterations promote oral squamous cell carcinoma (OSCC). Lactate-dependent histone modification is a novel histone mark that connects the epigenetic process of lactylation to glycolytic metabolites. However, the role of histone lactylation in oral carcinogenesis remains poorly understood. In this study, the levels of histone lactylation in oral leukoplakia (OLK) and OSCC tissues were determined by immunohistochemistry. The involvement of histone lactylation…

—

24 Mitochondrial control of glycerolipid synthesis by a PEP shuttle

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦通用细胞代谢转运机制的文章，与 AML 基础研究无直接或可迁移关联。；置信度：低

摘要: Mitochondria provide a variety of metabolites, in addition to ATP, to meet cell-specific needs. One such metabolite is phosphoenolpyruvate (PEP), which contains a higher-energy phosphate bond than ATP and has diverse biological functions. However, how mitochondria-generated PEP is delivered to the cytosol and fulfills cell-specific requirements remains elusive. Here, we show that SLC25A35 regulates mitochondrial PEP efflux and glyceroneogenesis in lipogenic cells that utilize the pyruvate-to-PEP…

—

25 The B cell receptor-mTOR signaling axis restricts the accumulation of lung tissue-resident memory B cells after influenza infection

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 完全无关的肺部感染免疫机制研究，仅基于标题与摘要片段，无法判断其对 AML 任何核心问题的直接或可迁移价值。；置信度：低

摘要: Lung tissue-resident memory B (B(RM)) cells are important in establishing protective immunity against respiratory pathogens. However, the pathways regulating the accumulation of lung B(RM) cells are poorly understood. Here, we show that the transcription factors NFATC1 (nuclear factor of activated T cells 1) and EGR2 (early growth response 2) repress lung B(RM) cell accumulation after influenza infection, whereas IKZF1 (IKAROS family zinc finger 1) enhances it. IKZF1 restrains B cell receptor…

—

26 Prasugrel inhibits TLR7-driven autoimmunity in systemic lupus erythematosus by acetylating cGAS

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文在 SLE 领域提出 cGAS 作为新靶点及 prasugrel 再利用潜力，但与 AML 基础研究无直接关联，且本次评估仅基于标题+摘要片段，证据链极不完整。；置信度：低

摘要: Systemic lupus erythematosus (SLE) has a complex, multifactorial etiology, which contributes to a lack of definitive cure and limited treatment efficacy. Here, we report that cyclic GMP-AMP synthase (cGAS) is significantly activated in SLE patients. We further demonstrate that cGAS deletion protects mice from lupus-like symptoms induced by the TLR7 agonist imiquimod (IMQ). In a screen of 3,159 FDA-approved drugs, we identify the antiplatelet agent prasugrel as a potent cGAS inhibitor….

—

27 Bst2-targeted senotherapy restores visual function by eliminating senescent retinal cells

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦视网膜衰老与退行性疾病的非 AML 机制研究，仅基于摘要片段无法判断其证据链完整性，与 AML 基础研究关联极弱。；置信度：低

摘要: Senescent cells contribute to degenerative processes in multiple tissues, including the retina. In the retinal pigment epithelium (RPE), their accumulation is closely associated with retinal aging and disease progression. Eliminating senescent RPE cells has shown therapeutic potential, but conventional senolytics often lack the specificity required to spare non-senescent cells, raising safety concerns. To overcome this, we performed integrated transcriptomic analyses of male mouse-derived RPE…

—

28 Polycomb repressive-deubiquitinase complex safeguards oocyte epigenome and female fertility by restraining Polycomb activity

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇生殖生物学/表观遗传领域的机制研究，聚焦卵母细胞非经典染色质景观的调控，与 AML 基础研究（LSC、表观遗传异常、克隆演化等）无直接关联，仅在泛表观遗传层面存在极弱的可类比性。；置信度

摘要: Mouse oocytes exhibit a unique chromatin landscape characterized by broad H3K27ac and H3K27me3 domains, demarcating euchromatin and facultative heterochromatin, respectively. However, the mechanisms underlying this non-canonical landscape remain elusive. Here we report BAP1, a core component of the Polycomb Repressive-Deubiquitinase (PR-DUB) complex, as a key negative regulator of Polycomb activity during oogenesis. BAP1 restricts pervasive H2AK119ub1 accumulation and protects oocyte-specific…

—

29 Randomized trial of GvHD Prophylaxis in Haploidentical PBSC Transplantation: ATG, PTCy, and Low-Dose Combination Therapy

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯临床 III 期随机试验文章，并非 AML 基础机制、生信或单细胞研究，提供的仅为标题+部分摘要（未含结果、数据、图表、结论或方法细节），对 AML 基础研究的直接价值极低。；置信度：低

摘要: The optimal graft-versus-host disease (GvHD) prophylaxis strategy in haploidentical peripheral blood stem cell transplantation (haplo-PBSCT) remains controversial. In this open-label, phase Ⅲ study, patients aged 14-70 years with acute myeloid leukemia or myelodysplastic syndromes with excess blasts Ⅰ or Ⅱ were randomized (2:1:1) to receive low-dose ATG (5 mg/kg)/PTCy (50 mg/kg), standard-dose ATG (total dose 10 mg/kg), or a PTCy (total dose 100 mg/kg)-based regimen for GvHD prophylaxis. The…

—

30 Isomer design unlocks rainbow phosphorescence

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的有机材料领域实验研究，基于有限摘要无法判断其在自身领域证据强度，对 AML 无任何逻辑迁移价值。；置信度：低

摘要: Achieving predictable color-tunable organic room-temperature phosphorescence (RTP) remains challenging due to limited understanding of triplet-state regulation in heteroaromatic systems. Carbazole and benzindole isomers provide an ideal platform to clarify how nitrogen positional isomerism governs triplet exciton behavior and emission energetics. Here, we establish a unified comparative framework to systematically investigate carbazole together with Bd[f], Bd[e], and Bd[g]. Nitrogen-site…

—

31 Reversible DNA condensation drives natural transformation

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦细菌抗生素耐药传播的体外单分子机制研究，与 AML 或任何造血/白血病相关内容完全无关。；置信度：低

摘要: Natural transformation drives the spread of antibiotic resistance among bacteria. The DNA receptor ComEA is essential for transporting external transforming DNA into the periplasm by an unknown mechanism. Here, single-molecule optical tweezers and electron microscopy approaches show that Geobacillus stearothermophilus ComEA forms dynamic oligomers on DNA that can switch between two conformations depending on local concentration. When ComEA sparsely decorates DNA, it forms bridging oligomers that…

—

32 Real-world treatment patterns and clinical outcomes in patients with AML from 65 to 74 years unfit for first-line intensive chemotherapy in Japan

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 11 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦临床实践的回顾性观察性研究，而非 AML 基础机制、生信或单细胞研究，提供文本仅为标题+摘要片段，未包含任何结果、统计或图表，对基础研究借鉴价值极低。；置信度：低

摘要: Venetoclax (VEN), a BCL-2 inhibitor, was approved in Japan in March 2021, for acute myeloid leukemia (AML). We retrospectively analyzed the impact of VEN approval on treatment patterns and outcomes in older AML patients aged 65-74 years unfit for intensive chemotherapy in Japan. Using the Hokkaido Leukemia Net database, we categorized 101 patients into pre-VEN (n = 46) and post-VEN (n = 55) cohorts, excluding those who had acute promyelocytic leukemia or received intensive chemotherapy….

—

33 Lactylation Converts ABHD6 into a Mitochondrial Regulator that Drives Lenvatinib Resistance in Hepatocellular Carcinoma

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；基于极有限的标题与部分摘要，该文提出 ABHD6 乳酸化介导线粒体动态改变作为 HCC 耐药新机制，但无任何实验细节、验证数据或图表支持，无法判断结论是否成立。；置信度：低

摘要: Hepatocellular carcinoma (HCC) frequently develops resistance to lenvatinib, a frontline tyrosine kinase inhibitor. Resistance arises from heterogeneous mechanisms involving metabolic reprogramming and mitochondrial adaptation, implicating regulators of these processes as potential therapeutic targets. Here, we identified α/β hydrolase domain containing 6 (ABHD6) as a critical driver of lenvatinib resistance by perturbing mitochondrial dynamics. Ligand-binding at the S148 catalytic site…

—

34 Molecular mechanism of menthol-induced TRPV5 channel inhibition

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文仅以电生理和 cryo-EM 结构数据初步描述 menthol 对 TRPV5 的抑制作用，但内容极端有限且与 AML/造血系统完全无关。；置信度：低

摘要: TRPV5 channels play a critical role in calcium homeostasis and are implicated in various pathophysiological conditions. Here, we demonstrate that the monoterpene menthol, commonly used for pain and inflammation management, is an inhibitor of TRPV5. Electrophysiology experiments reveal that menthol blocks ion conduction through a slow blocker mechanism. Using single-particle cryo-EM, we determine the structure of menthol-bound TRPV5, which shows menthol interacting with W583, a residue previously…

—

35 Hk2 Promotes Postburn Intestinal Injury by Facilitating Pyroptosis in Intestinal Epithelial Cells Through Enhancing Gsdmc2 Lactylation

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-19

- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦烧伤后肠道代谢-表观遗传-细胞死亡机制的非血液学研究，与 AML 基础研究无直接关联。；置信度：低

摘要: CONCLUSION: Hk2 knockdown mitigated postburn intestinal injury by inhibiting pyroptosis through decreased Gsdmc2 lactylation, providing a theoretical basis for developing clinical treatment strategies for this condition.

—

36 Histone H3K9 lactylation activates the TXNIP/NLRP3 pathway to drive macrophage inflammation after spinal cord injury

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 SCI 后代谢-表观遗传-炎症轴的机制研究，核心发现限于患者血清乳酸升高与小鼠模型中的乳酸积累，但与 AML 基础研究无直接关联。；置信度：低

摘要: Spinal cord injury (SCI) induces metabolic and immune disruptions that impede tissue repair. However, the underlying mechanisms are poorly understood. In this study, we identified lactate accumulation as a critical driver of macrophage-mediated inflammation through histone H3K9 lactylation (H3K9la). Targeted metabolomics revealed elevated serum lactate levels in SCI patients, which were linked to increased glycolysis and lactate dehydrogenase activity. In mice, lactate accumulation after SCI was…

—

37 Ferric Carboxymaltose Increases Fracture Risk in Patients and Reduces Bone Formation in Mice with Iron Deficiency Anemia

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦 IDA 治疗的骨骼副作用（低磷血症相关），证据初步但完全非 AML 研究，AML 借鉴价值极低。；置信度：低

摘要: Modern intravenous (IV) iron formulations allow treatment of iron deficiency anemia (IDA) with one or two infusions. Ferric carboxymaltose (FCM) is a widely used IV iron, which causes hypophosphatemia in the majority of patients. Osteomalacia and fractures are increasingly recognized after repeated infusions of FCM. It is unknown why ferric derisomaltose (FDI) rarely causes hypophosphatemia. In this study, we compare the effects of FCM and FDI on fracture risk and investigate potential…

—

38 Low-intensity focused ultrasound-activated piezoelectric gel bandage for diabetic wound repair and neuropathic pain relief

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦 DFU 领域中神经病变与局部免疫紊乱的交互问题，创新性地引入物理激活材料，但与 AML 基础研究（LSC、克隆演化、微环境、表观遗传等）无任何逻辑关联。；置信度：低

摘要: Nociceptive hypersensitivity and prolonged wound healing due to the interaction of peripheral neuropathy and local immune disorders are key issues that need to be addressed in diabetic foot ulcer (DFU) regenerative repair. Here, we present a low-intensity focused ultrasound (LIFU)-activated piezoelectric gel bandage for DFU wound repair and neuropathic pain relief. Acting as an artificial “skin”, this self-powered, functionalized bandage not only temporarily shields tissue from external…

—

39 Fast-swimming biohybrid OstraBot with self-trained high-strength muscles

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇工程生物学领域的应用创新文章，与 AML（急性髓系白血病）基础研究无任何逻辑关联。；置信度：低

摘要: Limited muscle force generation remains a major bottleneck in developing stronger, faster, and more efficient biohybrid robots. We present a fully autonomous self-training platform that strengthens skeletal muscle tissues by harnessing their robust spontaneous contractions. This approach produced muscle actuators with a maximum force of 7.05 mN and a stress of 8.51 mN/mm^2, the highest reported for C2C12-derived muscle actuators. To demonstrate their capabilities, we developed a twin-tail…

—

40 A bioinspired microdevice unifying energy storage and actuation through hydration control

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；该文为典型的材料科学/工程学研究，与 AML（急性髓系白血病）基础研究在研究对象、机制或临床转化上无任何重叠或可迁移逻辑。；置信度：低

摘要: Biological systems seamlessly integrate energy storage and actuation within compact architectures, whereas synthetic approaches largely implement these functions as separate components. Conjugated polymers can couple both, yet their operation relies on ion insertion accompanied by hydration water within the polymer backbone, creating an intrinsic trade-off between performance and stability. Here we show that anion hydration governs this trade-off. In-operando Raman spectroscopy and time-resolved…

—

41 The double-edged sword role of tumor-associated macrophages: preventing or causing resistance to immunotherapy

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇泛癌领域的概述性讨论，未涉及任何 AML 特异性内容或原始实验数据。；置信度：低

摘要: Immunotherapy has transformed cancer treatment, yet only a subset of patients achieves durable responses. Although macrophages play a key role in modulating immune response and could be used to enhance immunotherapy response, accumulating evidence highlights tumor-associated macrophages (TAMs) as both preventors or mediators of cancer, and its related therapies. TAMs contribute to resistance through multiple mechanisms, such as promoting T-cell exclusion, suppressing antitumor immunity via…

—

42 Leveraging large language models to populate structured clinical case report forms from unstructured medical notes in radiation oncology

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇高度应用导向的临床数据处理短文，仅通过结论摘要展示 LLM 效率，但与 AML 基础研究（LSC、克隆演化、表观遗传、耐药等）完全无关。；置信度：低

摘要: CONCLUSION: This study demonstrated the effectiveness of LLMs in structuring clinical data from medical non-English notes, with high accuracy in extracting and categorizing information. While multi-institutional validation is needed, the results indicate a significant healthcare impact through efficient data management, processing notes in 16 s, and accurately populating CRFs with minimal staff involvement.

—

43 A snow-fire bridge mechanism for the 2025 Southern California winter wildfire

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇气候科学领域的关联性+模拟研究，与 AML 基础研究（包括 LSC、克隆演化、耐药、表观遗传、微环境等核心问题）完全无关。；置信度：低

摘要: In January 2025, a rare and highly destructive wildfire devastated Southern California, becoming the costliest wildfire event in recorded history. The unusual timing of this wildfire suggests the possibility of unique remote, large-scale climatic precursors that differ from those of previous wildfires. Here, through observational analysis and large-ensemble numerical simulations, we identify that western Eurasian snow cover reduction is associated with weather conditions favorable for wildfires…

—

44 Atom-economy upcycling of commodity thermoset polyurethane into photocuring 3D printing resins based on selective cleavage-crosslink strategy

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯材料科学领域的回收技术创新论文，与 AML、白血病、造血干细胞或任何肿瘤微环境/表观遗传/代谢机制完全无关。；置信度：低

摘要: Polyurethane waste poses significant environmental challenges due to its large bulk volume and thermoset nature. Current recycling methods suffer from high reagent consumption, urethane byproduct generation, and limited economic value. Here, we well address these issues through a selective cleavage-crosslink strategy using food-grade ethyl acetoacetate (EAA) to break biuret crosslinks while preserving urethane/urea bonds in non-crosslinked regions virtually intact. This strategy minimizes…

—

45 Assessing the effectiveness of riparian buffers in protecting biodiversity: a meta-analysis

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇生态学领域的元分析文章，与 AML（急性髓系白血病）基础研究、造血干/祖细胞、肿瘤微环境或任何血液学机制完全无关。；置信度：低

摘要: Riparian buffers - vegetation strips retained or established along waterways - are mandated in many countries for water and soil protection, yet their role in biodiversity conservation remains unclear. Our global meta-analysis compares forested riparian buffers with converted riparian zones in human-modified landscapes across forest biomes. We find that forested riparian buffers support higher animal biodiversity than converted riparian zones at both local and landscape scales, particularly in…

—

46 Explainable graph learning for multimodal single-cell data integration

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用单细胞数据整合方法学短摘要，无任何 AML 或造血相关内容，生物学机制与 AML 特异性完全缺失。；置信度：低

摘要: CONCLUSIONS: SCPRO-VI offers a robust and interpretable framework for integrating paired single-cell multi-omic data. Its ability to enhance cell type separation and uncover meaningful sub-clusters suggests its utility in advancing our understanding of cellular diversity and regulatory mechanisms in complex tissues.

—

47 Delactylation of viral proteins by SIRT1 suppresses influenza A virus replication

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文提出乳酰化修饰在 IAV 感染中的新角色及 SIRT1 调控机制，但仅基于标题与摘要片段，无法验证任何实验证据或机制链条。；置信度：低

摘要: Lactylation, a protein post-translational modification driven by lactate, plays crucial roles in various physiological and pathological processes. However, the function and underlying mechanisms of lactylation in infection and pathogenesis of influenza A virus (IAV) remain poorly understood. Here, we demonstrated that sirtuin 1 (SIRT1) suppressed IAV replication by mediating delactylation of viral proteins. We found that IAV infection significantly downregulated the expression of SIRT1….

—

48 FUT8 reprograms glycolytic metabolism to promote PKM2 lactylation and drive clear cell renal cell carcinoma progression

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于 ccRCC 特有的 VHL 丢失/HIF 激活背景，探索 FUT8 在代谢层面的新作用，但与 AML 基础研究无直接关联。；置信度：低

摘要: Clear cell renal cell carcinoma (ccRCC) is characterized by the loss of the von Hippel-Lindau (VHL) gene, leading to constitutive activation of hypoxia-inducible transcription factors (HIFs) and metabolic reprogramming toward aerobic glycolysis. Although core fucosylation catalysed by fucosyltransferase 8 (FUT8) is known to regulate receptor signaling and tumor malignancy, its role in metabolic regulation of ccRCC remains poorly defined. Here, we demonstrate that FUT8 knockdown significantly…

—

49 Efficient control of enterochromaffin versus islet differentiation from human pluripotent stem cell-derived pancreatic progenitors

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦糖尿病相关干细胞分化的增效协议研究，与 AML 造血系统、LSC 或白血病机制无任何直接或间接关联。；置信度：低

摘要: Knowledge of the molecular cues guiding pancreatic development is critical to developing beta cell replacement therapies for the treatment of diabetes. We compare different methods of pancreatic endocrine differentiation from human pluripotent stem cells (hPSCs) and establish sequences of patterning that can selectively increase the frequencies of islet-like or off-target enterochromaffin (EC)-like cells, thereby significantly increasing islet-like cellular yield and glucose-stimulated insulin…

—

50 Carbon, nitrogen, and sulfur cycling unveil deep-sea microbial niches in the Atacama Trench

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇描述性环境微生物生态学研究，仅基于摘要片段显示为初步功能注释，未涉及任何血液学、白血病或造血相关内容，与 AML 基础研究无逻辑关联。；置信度：低

摘要: Deep-sea sediments host microbial communities essential to nitrogen, sulfur, and carbon cycling, yet those in the Southern Hemisphere remain understudied. In this work, we analyze sediments from an oceanic trench off northern Chile using metatranscriptomics, aiming to relate community composition with functional activity. Most assayed sites form part of a broadly uniform deep-sea ecosystem fueled by the degradation of marine snow through diverse aerobic and anaerobic pathways. Nitrogen cycling…

—

51 Deep-learning-based de novo discovery and design of therapeutics that reverse disease-associated transcriptional phenotypes

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用计算方法学论文，未涉及任何具体疾病（包括 AML）的生物学验证或机制探索，属于“方法先行、应用待证”类型。；置信度：低

摘要: Identifying drugs that reverse disease-associated transcriptomic features has been widely explored for drug repurposing, but its potential for de novo drug discovery remains underexplored. Here, we present gene expression profile predictor on chemical structures (GPS), a deep-learning-based drug discovery platform, guided by transcriptomic features, that screens large compound libraries and optimizes lead molecules. We first develop a model that captures transcriptomic perturbation signatures…

—

52 D-lactate Drives Lysine D-lactylation to Regulate Metabolism in Escherichia coli

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文在细菌领域提出 D-乳酰化新 PTM 机制，但基于有限摘要文本，证据链完全缺失，且与 AML 基础研究无任何逻辑关联；置信度：低

摘要: L-lactate-drived lysine L-lactylation (K(L-la)) has emerged as a key regulator in diverse cellular processes and disease pathogenesis. While lactate predominantly exists as the L-isomer in eukaryotes, both L- and D-lactate are present in some bacteria. However, it remains unclear whether D-lactate can drive post-translational modification (PTM) to exert biological functions. Here, we reported that D-lactate-derived lysine D-lactylation (K(D-la)) serves as a PTM in E. coli. We demonstrated that…

—

53 Decoding the immune-tumor synapse for data-driven design of next-generation immunotherapies

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-18
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；文章仅基于概念概述提出技术整合路径，科学问题重要但缺乏任何具体数据、机制或验证支持，且与 AML 特异性基础机制无直接关联。；置信度：低

摘要: The immunological synapse (IS) formed between cytotoxic CD8^(+) T lymphocytes (CTLs) and tumor cells represents the critical interface where many immunotherapies act, including therapeutic antibodies and chimeric antigen receptor immunotherapy. However, detailed characterization of the CTL-tumor IS has remained limited, primarily due to the transient and dynamic nature of this structure. Recent advances in proteomics, transcriptomics, high-throughput imaging, and machine learning are now…

—

54 Engineering synthetic cells with intramembrane domains possessing distinct bilayer asymmetries

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇合成生物学领域的膜工程方法学短文，与 AML 或造血系统基础研究无任何直接或间接关联。；置信度：低

摘要: Our understanding of how membrane asymmetry governs biological function is limited by the lack of techniques to produce model membranes which can reliably and accurately mimic cellular membrane asymmetry. Not only in terms of asymmetric lipid distribution, but also how that asymmetry can be confined to specific lateral locations across the membrane. Here we present an inverted emulsion method that can be used to produce synthetic cells with symmetric and asymmetric bilayers, as well as phase…

—

55 FGF1 orchestrates circadian hepatic triglyceride secretion

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 MASLD 代谢与昼夜节律的非 AML 机制研究，与血液学/白血病基础领域无直接关联，仅摘要片段无法支撑任何可信结论。；置信度：低

摘要: Metabolic dysfunction-Associated Steatotic Liver Disease (MASLD) represents a global health crisis associated with dysregulated hepatic triglyceride (TG) synthesis, oxidation and secretion. Despite progress in targeting hepatic lipid synthesis/oxidation for MASLD treatment and a well-documented relationship between circadian rhythms and lipid metabolism, the adaptive mechanisms coordinating TG secretion with circadian timing remain incompletely understood. Here we identify an autocrine…

—

56 Cryo-EM structure of TRPM1 reveals a non-canonical architecture with an inverted transmembrane domain

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦视觉相关膜蛋白结构解析的生物物理学论文，与 AML（急性髓系白血病）基础研究完全无关。；置信度：低

摘要: Transient receptor potential melastatin 1 (TRPM1) is a membrane protein essential for vision in dim light, and mutations in TRPM1 cause complete congenital stationary night blindness. Although TRPM1 shares sequence similarity to other TRPM ion channels such as TRPM3, whether it independently functions as an ion channel remains controversial. This controversy is largely caused by TRPM1′s challenging biochemical behaviors that prevent detailed molecular characterization. In this work, we isolate…

—

57 Epigenetic regulation in a high-sugar environment (Review)

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇聚焦糖尿病领域的表观遗传综述，与 AML 基础研究无直接关联，仅在代谢-表观遗传交叉点存在极弱的潜在迁移可能性。；置信度：低

摘要: The prevalence of diabetes and its complications has become a major global health challenge, with its pathological process closely linked to the phenomenon of ‘metabolic memory’ induced by persistent hyperglycemia. Epigenetic regulation is recognized as the core molecular mechanism underpinning this process. The present review systematically elucidated how the hyperglycemic microenvironment profoundly regulates cellular functions and drives the onset and progression of diabetes and its vascular…

—

58 scZiva: imputation method for single-cell RNA-seq data with zero-inflated variational autoencoder

- **研究本身价值**: 15 / 60
- **AML 借鉴价值**: 16 / 30
- **表达与可复用性**: 1 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文仅提供标题，无摘要、正文或任何结果细节，无法进行实质学术评估。；置信度：低

摘要: No abstract

—

59 Lysosome-targeting live attenuated influenza vaccines elicit robust and broad immunity in mice

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦流感疫苗设计的工程学创新文章，但与 AML（急性髓系白血病）基础研究完全无关，仅能作为极边缘的蛋白降解思路参考。；置信度：低

摘要: The lysosome is a cell’s endogenous machinery responsible for degrading proteins. Here we describe two lysosome-targeting live attenuated vaccine approaches, LYTAR 1.0 and LYTAR 2.0, by harnessing the lysosome to conditionally degrade viral proteins of influenza virus. LYTAR 1.0 incorporates a conditionally removable lysosome-targeting motif at the N- or C-terminus of viral proteins. LYTAR 2.0 allows flexible placement of lysosome-targeting motifs at internal or terminal sites of viral proteins⋯.

—

60 Aptamers with magnetically tunable affinity for divalent cobalt ions

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-19
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇化学/生物物理领域的方法学探索论文，与 AML 基础研究完全无关，仅摘要显示创新点局限于体外适配体亲和力调控。；置信度：低

摘要: There is considerable interest in the ability to modulate biological processes with magnetic fields. Here we demonstrate a strategy for selecting aptamers that exhibit enhanced binding to paramagnetic metal ions under a strong magnetic field. Using a high-magnetic-field (HM)-SELEX method targeting Co^(2+), we identified two classes of aptamers

with magnetically-modulated binding behavior. One displayed a gradual 2-3-fold increase in affinity as magnetic field strength increased, while the other…

—