追觅 - 2026-03-17

生成时间: 2026-03-17 07:13

分析文章数量: 38

1 An international multi-centre study to develop and validate federated learning-based prognostic models for anal cancer

- **研究本身价值**: 38 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 41 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一项方法学导向的临床生信研究，在其自身领域（罕见癌数据隐私保护）具有一定价值，但与 AML 基础研究完全无关。；置信度：低

摘要: Precision oncology relies on access to high-quality data for increasingly smaller patient subgroups. The international atomCAT consortium investigates the potential of federated learning to support this, using anal cancer as a rare cancer exemplar. Here, we show that federated multivariable Cox models trained across 14 centres (1428 patients) and externally validated in two additional centres (277 patients) achieve consistent calibration and discrimination during leave-one-centre-out and…

—

2 AARS2 R199C mutation induces lactylation-driven premature ovarian insufficiency phenotypes partially reversible by SIRT3

- **研究本身价值**: 35 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 36 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文首次构建纯合 Aars2 R194C 敲入小鼠模型，验证了该突变在体内诱导 POI 相关表型并提出乳酸化机制，但与 AML 基础研究完全无关。；置信度：低

摘要: Premature ovarian insufficiency (POI) often arises from genetic causes, yet the pathogenic consequences of many variants remain undefined. The AARS2 R199C mutation has been repeatedly reported in patients, but its physiological effects were unknown. Here, we generated the first homozygous Aars2 R194C knock-in mouse to model this variant in vivo. Female knock-in mice showed irregular estrous cycles, reduced fecundity, altered endocrine profiles, and accelerated depletion of the primordial…

—

3 AARS1-mediated lactylation of STAT1 drives immune evasion

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 35 / 100
- **发布日期**: 2026-03-15

- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；摘要提出一个新兴乳酰化机制，但因缺乏任何实验证据支持，结论目前仅为“声称”而非“证实”。；置信度：低

摘要: Lactate accumulates in large amounts in tumor cells due to the Warburg effect. However, the role of lactate-mediated lactylation, a post-translational modification, in regulating tumor immunity remains unclear. Here, we report that lactate-driven lactylation of STAT1 K193 inhibits interferon (IFN)-γ signaling pathway-mediated tumor immunity. Mechanistically, AARS1 lactylates STAT1 K193 and inhibits its binding to JAK2 and phosphorylation, thereby disrupting tumor responsiveness to IFN-γ, which…

—

4 Global literature review and survey of implementation constraints on natural climate solutions

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇聚焦气候变化缓解策略落地障碍的描述性综述与调查文章，与 AML 血液学、白血病干细胞、机制或肿瘤微环境研究无任何关联或逻辑迁移性。；置信度：低

摘要: Natural Climate Solutions - protection, restoration, and improved management of lands and waters that reduce greenhouse gasses - have large climate change mitigation potential. However, lack of comprehensive information on implementation challenges hinders the adoption of Natural Climate Solutions and the delivery of near-term mitigation. Using a global survey of Natural Climate Solutions projects and a systematic review of recent studies, we map 46 constraints in eight categories, yielding…

—

5 Global coincident bursts of high frequency oscillations across the human cortex coordinate large-scale memory processing

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇神经生理学观察性研究，聚焦大脑记忆加工的振荡协调机制，与 AML 基础研究（包括 LSC、克隆演化、耐药、微环境等）完全无关。；置信度：低

摘要: Oscillations in the high gamma and ripple frequency ranges are known to coordinate local hippocampal and neocortical neuronal assemblies during memory encoding and recall. Here, we explored spatiotemporal dynamics and the role of global coordination of these fast oscillatory discharges across the sensory and associational cortical areas in

distinct phases of memory processing. Individual bursts of high frequency oscillations were detected in intracranial recordings from epilepsy patients…

—

6 Multitrait GWAS and functional validation reveal genetic loci for gastric cancer

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦胃癌遗传学的多性状 GWAS 文章，与 AML 基础研究在疾病类型、核心机制（LSC、克隆演化、分化阻滞、耐药等）上无任何逻辑交集。；置信度：低

摘要: Gastric cancer is a leading cause of cancer-related death, particularly in East Asia, yet its genetic basis remains incompletely understood. Using genome-wide association study data from BioBank Japan, we show that gastric cancer shares significant genetic associations with 25 other phenotypes, most notably cardiovascular conditions such as angina pectoris and myocardial infarction. Mendelian randomization supports a causal role of these traits in gastric cancer risk. Multitrait analysis of GWAS…

—

7 A deep adversarial network model for multi-task analysis of single-cell omics data

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用单细胞多组学计算方法论文，聚焦技术挑战，但与 AML 基础研究无直接关联，且提供的文本极度有限，无法评估真实性能。；置信度：低

摘要: Single-cell multi-omics data reveal complex cellular states and deepen our understanding of tissue cell phenotypes and functions. However, data analysis remains challenging due to the discrete nature and high noise level of the data, as well as the lack of modality. Here, we propose scMultiNet, a multi-task deep adversarial neural network that can integrate different tasks to analyze single-cell multi-modal data. In particular, we achieve joint training of multi-modal integration and cross-modal…

—

8 Impact of non-genetic heterogeneity of BRAF-mutant colon cancer organoids on growth kinetics, drug sensitivity and Wnt dynamics

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100

- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇结肠癌领域聚焦非遗传参数的探索性研究，与 AML 基础研究无任何直接或逻辑关联，借鉴价值极低。；置信度：低

摘要: Patient-derived organoid (PDO) models are powerful systems for studying tumor biology and drug response. By retaining genetic, histological, and functional characteristics of the original tumor, including intra- and interpatient heterogeneity, they provide powerful tools to investigate therapy resistance. While genomic profiling is well established in organoid-based screening, non-genetic parameters such as organoid size, seeding density, and morphology remain underexplored, despite their…

—

9 Experimental mechanician for plate lattice metamaterial discovery

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇材料科学领域的工程方法论文，与 AML 或任何血液学/造血生物学基础研究完全无关，学术价值仅限于其自身领域。；置信度：低

摘要: Plate lattice metamaterials offer higher strength-to-weight efficiency than traditional truss lattices. Their optimization, however, is challenging when having more parameters and multiple objectives. Here, we present an experimental mechanician (ExMech) to identify lightweight and high-strength plate lattices. ExMech integrates a robotic platform for standardized data generation and a multi-objective active learning framework for guided exploration of structure-property relationships. It…

—

10 INSIG1/2 succination mediated by the moonlighting function of ADSL promotes lipogenesis and liver tumorigenesis

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦肝癌（HCC）代谢感知与脂质重编程的机制研究，与 AML 基础研究无直接或可迁移的逻辑关联。；置信度：低

摘要: Aerobic glycolysis supports tumor growth, but how tumor cells sense glucose to coordinate biosynthesis remains largely unclear. Here we show that in hepatocellular carcinoma cells, glucose-activated PKCε phosphorylates the purine synthesis enzyme ADSL, triggering its translocation to the endoplasmic reticulum. ADSL then promotes succination

of INSIG1/2, which disrupts the interaction between INSIG proteins and SCAP, leading to the translocation of the SCAP-SREBP complex to the Golgi, the…

—

11 EPInformer: scalable and integrative prediction of gene expression from promoter-enhancer sequences with multimodal epigenomic profiles

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦通用转录调控的计算建模，未涉及任何 AML、造血或白血病相关内容，基于极有限摘要无法评估模型性能、验证或实际生物学意义。；置信度：低

摘要: Transcriptional regulation, critical for cellular differentiation and adaptation to environmental changes, involves coordinated interactions among DNA sequences, regulatory proteins, and chromatin architecture. Despite extensive chromatin profiles and gene expression data from consortia, understanding the dynamics of cis-regulatory elements in gene expression remains challenging. Deep learning is a powerful tool for learning gene expression and epigenomic profiles from DNA sequences, exhibiting…

—

12 Unpacking sources of transmission in HIV prevention trials with deep-sequence pathogen data

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 HIV 流行病学与预防策略的统计建模论文，与 AML（急性髓系白血病）基础研究完全无关。；置信度：低

摘要: To develop effective HIV prevention strategies to guide public health policy the main sources of infection in HIV prevention studies must be identified. Accordingly, we devised a statistical approach that estimates the relative contribution of different sources of infection in community-randomized trials of infectious disease prevention using deep- (or next generation) sequenced pathogen data. We applied this approach to the Botswana Combination Prevention Project (BCPP) and estimated that 90%…

—

13 Joint control of precipitation and CO₂ on global long-term patterns of plant nitrogen availability

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 3 / 30

- **表达与可复用性**: 5 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML、白血病干细胞、造血机制或肿瘤微环境完全无关的植物生态学论文，仅摘要信息无法支持任何 AML 迁移。；置信度：低

摘要: Plant nitrogen availability critically limits plant growth and thus constrains terrestrial carbon sequestration. Growing evidence points to a recent decline in plant nitrogen availability, yet the global consistency of this trend and its underlying drivers remain unclear. Here, we reconstruct the spatiotemporal trajectory of plant nitrogen availability (1980-2020) using four machine-learning algorithms applied to a global database of 37,268 foliar stable nitrogen isotopes measurements and…

—

14 Revealing multiscale competing processes in the solid-state synthesis of single-crystalline layered oxide positive electrodes

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇典型的材料科学领域技术驱动论文，与 AML、白血病干细胞、造血机制或肿瘤微环境毫无关联，仅在电池材料合成领域具有一定价值。；置信度：低

摘要: Solid-state synthesis involves a web of coupled chemical reactions and physical changes that unfold across multiple scales. Efforts to fine-tune its parameters have historically followed heuristic, trial-driven workflows that demand significant time and resources. In this study, we aimed to open this black box by employing multiscale in situ synchrotron imaging and diffraction. Using LiNi(0.5)Mn(0.3)Co(0.2)O(2) battery positive electrode material as a model system and Ba-based sintering aids, we…

—

15 Mining antibody functionality via AI-guided structural landscape profiling

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯计算生物学方法开发论文，聚焦抗体功能预测工具的构建，与 AML 基础研究完全无关。；置信度：低

摘要: Despite substantial progress in single-cell screening techniques, antibody (Ab) repertoires still remain enigmatic. Here we show that Ab sequences can be linked to their

functionality by using big data obtained from high-throughput sequencing. Using the expansive SARS-CoV-2 pandemic data, we develop an AI-based method to reveal the neutralization potential of Ab repertoires. We employ machine learning to process public 3D structural data of Ab-RBD complexes and create a comprehensive tool,…

—

16 Candesartan cilexetil disrupts methicillin-resistant Staphylococcus aureus membrane and potentiates gentamicin and polymyxin B activity

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于极有限摘要信息提出药物膜靶向作用，但完全脱离 AML 领域，与血液学/白血病基础研究无任何逻辑关联。；置信度：低

摘要: Methicillin-resistant Staphylococcus aureus (MRSA) is a leading cause of bacterial infections, but treatment options are limited due to MRSA multi-drug resistance. The anti-hypertensive drug Candesartan cilexetil (CC) exhibits potent anti-MRSA activity. It permeabilizes the membrane of MRSA cells and potentiates the activity of aminoglycoside antibiotics. Here, we used a variety of methods to elucidate the mechanism by which CC disrupts membrane homeostasis. We show that CC binds to bilayer…

—

17 High-Fidelity quantum teleportation mediated by hole transfer in an acceptor-donor-radical molecular triad

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇量子信息科学领域的纯物理化学机制研究，与 AML 或任何血液学/白血病基础研究完全无关。；置信度：低

摘要: Quantum teleportation transfers quantum states between nodes in a quantum network through entanglement distribution and subsequent Bell-state measurements. Here, we report electron spin teleportation through an unexplored hole-transfer mechanism within an ensemble of covalently linked acceptor-donor-stable radical (A-D-R^(·)) molecules. Photoexcitation of A after spin state preparation on R^(·) results in ultrafast hole transfer to produce an entangled pair ^1(A^(·-)-D^(·+)). A subsequent…

—

18 Dopaminergic processes predict temporal distortions in event memory

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30

- **表达与可复用性**: 6 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇认知神经科学领域的初步探索性研究，与 AML 基础研究（包括 LSC、克隆演化、分化阻滞、耐药、微环境、表观遗传等核心问题）无任何逻辑或生物学关联。；置信度：低

摘要: Our memories do not simply keep time - they distort it, stretching and compressing the past to reflect the structure of experience. Here, we combined functional magnetic resonance imaging (fMRI; n = 32) with eye-tracking (n = 28) to test whether activation of the dopaminergic system, known to influence encoding and time perception, expands mnemonic representations of time between contextually distinct events. Participants encoded item sequences while listening to tones that typically repeated…

—

19 Complex marine ecological response during the Eocene-Oligocene revealed by global foraminiferal record

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的古气候-古生态论文，仅基于标题和摘要片段，无法判断其在自身领域的完整证据链。；置信度：低

摘要: The Eocene-Oligocene transition was the crucial turning point when Earth’s climate shifted to its current icehouse state. Understanding how the marine biosphere responded during this transition is not well-constrained, appearing as a simple extinction pulse in low temporal resolution global compendia. Here we design an artificial-intelligence-inspired metaheuristics algorithm to construct a high-resolution global species richness history across the Eocene-Oligocene transition for the rich…

—

20 Real-world effectiveness and safety with nivolumab plus ipilimumab or nivolumab alone in patients with or without melanoma brain metastasis: Results from the German noninterventional NICO study

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇临床肿瘤学真实世界数据报告，与 AML 基础研究（机制、生信、单细胞、造血干/祖细胞等）无任何实质关联。；置信度：低

摘要: The prospective, German NICO study (ClinicalTrials.gov identifier: NCT02990611) evaluated real-world effectiveness and safety with nivolumab plus ipilimumab or nivolumab alone (any-line) in patients with advanced melanoma with/without melanoma brain metastasis (MBM). A total of 755 patients treated with nivolumab plus ipilimumab (n = 486; median follow-up, 46.8 months) or nivolumab alone (n = 269; median follow-up, 38.7 months) were enrolled. Baseline characteristics differed between the…

—

21 Learning data-efficient coarse-grained molecular dynamics from forces and noise

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯计算生物物理/机器学习方法论文，与 AML 或任何血液学、白血病基础研究无任何直接或间接关联，仅讨论通用生物分子模拟的技术瓶颈。；置信度：低

摘要: Molecular dynamics (MD) simulations are essential for elucidating biomolecular function, yet the computational cost of all-atom models often limits their reach. Machine-learned coarse-grained (MLCG) models offer a solution by simplifying the representation while maintaining near-atomistic accuracy. However, the training of MLCG models currently requires vast amounts of force-labeled sample conformations from reference atomistic MD. Here, we overcome this limitation by unifying the training of…

—

22 Integrating integrated circuit wastewater into the metal catalyst supply chain

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇环境工程领域的可持续催化剂制备证明-of-concept 研究，与 AML 基础研究（包括造血干细胞、LSC、表观遗传、代谢重编程等）无任何生物学或医学关联。；置信度：低

摘要: The rapid growth of the integrated circuit (IC) industry has led to the generation of highly complex and challenging-to-treat wastewater streams. This work presents a sustainable paradigm for transforming metal-containing IC wastewater into efficient heterogeneous catalysts. As a proof of concept, Cu-rich IC wastewater is converted into a multifunctional Cu/SiO(2) catalyst via a simple ammonia-evaporation process, achieving up to 99.9% Cu recovery. The catalyst exhibits outstanding activity in…

—

23 The associations between quality of life and a five-day per week peritoneal dialysis prescription: A pilot randomized controlled study

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 19 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇肾脏病临床领域的试点 RCT，与 AML 基础研究（LSC、克隆演化、分化阻滞、耐药、微环境、表观遗传等）无任何主题或机制关联。；置信度：低

摘要: BackgroundIncremental peritoneal dialysis (PD) has been recommended as a patient-focused prescription; however, its safety and potential clinical benefits are based on retrospective studies.MethodsWe conducted a pilot, randomized controlled study comparing 5-day versus 7-day/week PD in incident patients with preserved residual kidney function in three centers in Australia. Modality of PD accorded to patient and clinician preferences for both groups. Participants on 5-day PD were transitioned to…

—

24 tRNA-derived RNA processing in sperm transmits non-genetically inherited phenotypes to offspring in C. elegans

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 18 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦线虫精子 tDRs 介导表观遗传非遗传性传递的机制研究，内容与 AML 基础研究（LSC、表观遗传异常、耐药、微环境等）无直接或可迁移关联。；置信度：低

摘要: The environment encountered by an organism can modulate epigenetic information in gametes to transmit non-genetically inherited phenotypes to offspring. In mouse models, the diet of males regulates specific tRNA-derived RNAs (tDRs) in sperm. After fertilization, tDRs regulate embryonic gene expression and generate metabolic phenotypes in adult offspring through uncharacterized changes during development. Here we demonstrate that in C. elegans, tDRs also accumulate in sperm and similarly transmit…

—

25 Effects of Transradial Catheterisation on Radial Artery Bypass Graft Patency: A Systematic Review and Meta-Analysis

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 18 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的临床心血管外科荟萃分析，仅基于标题及摘要结论片段，无法体现任何血液学、造血干细胞或白血病机制内容。；置信度：低

摘要: CONCLUSIONS: Prior transradial catheterisation is associated with reduced patency of RA bypass grafts. Surgeons should exercise caution when selecting CRA for critical targets, and prospective controlled data are needed to define patient and procedural modifiers of risk.

—

26 Trajectories of quality of life, anxiety and depressive symptomatology, and health-related work productivity after first seizure events

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 16 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一项神经临床领域的心理社会结局纵向描述性研究，与 AML 基础机制（LSC、克隆演化、耐药、微环境、表观遗传等）无任何逻辑关联。；置信度：低

摘要: OBJECTIVE: First seizure events are common and may exert an immediate and profound impact on people’s lives. Less understood are their long-term consequences. This prospective, longitudinal study aimed to measure and compare psychosocial and health-related work productivity trajectories following first seizure events of various etiologies.

—

27 Evaluation of novel assays of non-ceruloplasmin copper to monitor chelation treatment in patients with Wilson disease: Monitoring chelation treatment in Wilson disease

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 16 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文仅提供摘要结论片段，聚焦 WD 铜代谢临床监测，与 AML 基础研究（LSC、克隆演化、代谢重编程等）无任何逻辑关联或可迁移性。；置信度：低

摘要: CONCLUSION: Non-ceruloplasmin by protein speciation and exchangeable copper have potential to guide chelation in WD patients on maintenance therapy. Specific target ranges should be established, and we hypothesize they may include values below normal ranges. Further studies are required to improve our understanding of the responses to dose changes and non-adherence and if standardization of sampling conditions can reduce visit-to-visit variability.

—

28 Cortical representation of multidimensional handwriting movement and implications for neuroprostheses

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 15 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML、白血病、造血干细胞或肿瘤微环境完全无关的神经科学 BCI 研究，基于极有限摘要文本，无法支撑任何 AML 相关结论。；置信度：低

摘要: Handwriting brain-computer interfaces (BCIs) have enabled high performance brain-to-text communication for paralyzed individuals. However, the detailed parameters of handwriting movement and their cortical representations remain incompletely understood. Here, we recorded intracortical neural activity from a paralyzed subject and found distinct neural representations for strokes and pen lifts with respect to two-dimensional (2D) velocity on the writing plane, indicating that 2D kinematics alone…

—

29 Topologically reconstructing Pancharatnam-Berry phase via encircling exceptional point for chiral spin-orbit interaction steering

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 14 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯物理光学领域的拓扑光子学研究，与 AML、白血病干细胞、造血微环境或任何血液学机制完全无关；置信度：低

摘要: Photonic spin-orbit interaction (SOI) is related to Pancharatnam-Berry (PB) phase, which is usually imposed on the cross-polarized components of light incident on a metasurface in broadband manner. However, metasurfaces exhibiting exceptional points (EPs) can block the cross-polarization conversion channel, hinting at the possibility of EPs influencing SOI. Here, we demonstrate that the accumulation of PB phase is equivalent to the encircling of an EP, leading to topologically reconstructed PB…

—

30 Examining the daily life dynamics of anticipation and recovery from work-related stress in frontline medical professionals during pandemic work demands

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 12 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的心理学文章，仅基于标题和摘要结论，无法支撑任何生物医学机制或 AML 借鉴评估。；置信度：低

摘要: CONCLUSIONS: Neuroticism and low self-efficacy may place medical professionals especially at risk in medical crises. Promoting leisure activities to support recovery may replenish resources for handling work-related stress.

—

31 Histone lactylation: A sensor of epigenetic reprogramming mediated by cellular metabolism

- **研究本身价值**: 13 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 12 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文仅提供高度概括的概念性描述，未呈现任何具体实验设计、数据、验证或 AML 相关内容，学术价值与可信度无法实质评估。；置信度：低

摘要: Epigenetic reprogramming plays a critical role in connecting genotype to phenotype, providing the flexibility that allows organisms to adapt to developmental needs, environmental shifts, and disease challenges through the flexible regulation of gene expression. Histone lactylation, a recent modification in the field of epigenetic regulation, directly associates metabolic products-lactate-with chromatin modifications, revealing how cellular metabolism regulates the epigenetic landscape. In this…

—

32 Catalytic enantioselective synthesis of azahelicenes via cascade Pictet-Spengler reaction and dehydrogenative aromatization

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 12 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯有机合成方法学论文，与 AML、白血病或任何生物医学基础研究完全无关。；置信度：低

摘要: The introduction of N-heteroaromatic rings like pyridine and pyrrole into the helicene scaffold can significantly impact the electronic properties, redox potentials and coordination abilities of helicenes, which lead to diverse and intriguing applications for these chiral azahelicenes. However, methods for catalytic enantioselective synthesis of these compounds, especially those with N-heterocyclic moieties, remain largely underdeveloped. Herein, we present an efficient method for catalytic…

—

33 Highly ionic-dispersed oxygen electrode for reversible proton ceramic electrochemical cells

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 11 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇典型的材料科学电化学性能优化论文，与 AML 基础研究（包括 LSC、克隆演化、表观遗传、代谢重编程等）无任何生物学或医学关联。；置信度：低

摘要: The key challenges for commercializing reversible proton ceramic electrochemical cells (R-PCECs) are the insufficient proton conductivity and inferior thermomechanical stability of oxygen electrodes in air with water vapor. We report a multielement micro-doped BaCoO(3-δ)-based perovskite material, in which disorder is induced in the ionic substructure to maximize the oxygen-water reaction activity. Atom probe tomography and density functional theory calculations reveal that reduced proton…

—

34 Solvation sheath reorganization enables fast ion transfer kinetics in lithium-ion battery

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 11 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇材料化学领域的电解液优化机制研究，与血液学、白血病或 AML 基础研究无任何生物学或医学关联。；置信度：低

摘要: The limitations of ion transport kinetics in conventional electrolytes, particularly under extreme operating conditions, arise from suboptimal solvation structures and inefficient charge carrier utilization. Here, we present strategic electrolyte design that reconfigures Li^+ coordination geometry by modulating intermolecular interactions and solvent molecule volume, fundamentally overcoming these transport constraints. By incorporating an optimized moderator with a low dipole moment and small…

—

35 Urinary Albumin-to-Creatinine Associates More Closely than Protein-to-Creatinine Ratio with Kidney Outcomes in Pediatric Chronic Kidney Disease

- **研究本身价值**: 16 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 10 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇儿科 CKD 预后标志物比较的观察性研究，与 AML 基础研究（LSC、克隆演化、耐药、微环境、表观遗传等）无任何逻辑关联。；置信度：低

摘要: CONCLUSION: Both uACR and uPCR were independently associated with CKD progression in this pediatirc cohort. In direct comparison, uACR demonstrated stronger associations with adverse kidney outcomes, suggesting a comparatively greater prognostic signal and supporting current guideline emphasis on albuminuria for risk stratification and intervention outcome monitoring.

—

36 Dynamically assembled photochromic cages operational in water with visible light

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 8 / 100
- **发布日期**: 2026-03-16
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯化学纳米技术领域的初步描述性工作，与血液学、白血病或 AML 基础研究无任何逻辑关联或可迁移性。；置信度：低

摘要: Producing chemical nanostructures that can mimic the efficient adaptability of complicated biological systems to environment changes is among the main goals of nanotechnology. Progress in this area requires understanding of the adaptation mechanisms towards external stimuli at the molecular level. Due to rapid and precise spatiotemporal addressability, light-driven dynamic systems are particularly attractive for such mechanistic studies. Here, we show efficient formation of dynamic covalent…

—

37 Intraindividual development of MR lung perfusion parameters in children after congenital diaphragmatic hernia at 2 and 10 years

- **研究本身价值**: 18 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 8 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；基于极有限的结论片段，文章提示产前重度肺发育不良患儿存在持久且稳定的肺灌注降低，强调早期干预必要性，但全文方法、数据、统计及验证细节完全缺失，无法判断研究质量。；置信度：低

摘要: CONCLUSION: Patients with prenatally more severe lung hypoplasia, particularly those with low o/e FLV, exhibit persistently reduced lung perfusion even in the long term. These ongoing impairments remain stable over time, indicating permanently compromised lung development. Early identification and detailed risk assessment are therefore essential to initiate targeted therapeutic interventions.

—

38 Semi-automated isolation of parenchymal and non-parenchymal liver cells from mice and humans with enhanced stellate cell fraction

- **研究本身价值**: 16 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 8 / 100
- **发布日期**: 2026-03-15
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯方法学/资源型论文，聚焦肝脏细胞分离技术，与 AML（急性髓系白血病）基础研究无任何直接或可迁移关联。；置信度：低

摘要: No abstract

—