追觅 - 2026-03-22

生成时间: 2026-03-22 06:56

分析文章数量: 17

1 Multi-omic identification of key transcriptional regulatory programs during endurance exercise training in rats

- **研究本身价值**: 40 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 43 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇运动生理学领域的描述性多组学资源文章，与 AML 基础研究（LSC、克隆演化、分化阻滞、耐药、微环境等）无任何直接或可迁移关联。；置信度：低

摘要: Transcription factors play a key role in regulating gene expression. We conduct an integrated analysis of chromatin accessibility, DNA methylation, mRNA expression, protein abundance and phosphorylation across eight tissues in fifty rats of equally represented sexes following endurance exercise training to identify coordinated epigenomic and transcriptional changes and determine key transcription factors involved. We uncover tissue-specific endurance exercise training associated changes and…

—

2 Single-cell multiomics uncovers an endothelial mechanosensitive PIEZO1-IL-33 axis driving pulmonary fibrosis

- **研究本身价值**: 34 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 43 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦肺纤维化血管内皮机械生物学的单细胞多组学文章，与 AML 基础研究无直接关联，迁移性极弱。；置信度：低

摘要: Pulmonary fibrosis represents a progressive interstitial lung disease marked by excessive extracellular matrix deposition and architectural distortion. Vascular endothelial cells critically contribute to fibrogenesis through paracrine secretion of pro-fibrotic mediators, yet their mechanobiological regulation remains elusive. Using integrated single-cell multi-omics profiling of human pulmonary fibrosis specimens and experimental fibrosis models induced by bleomycin or silica, we identify…

—

3 Pre-splicing conformation and stepwise circularization of a group I intron in Azoarcus pre-tRNA

- **研究本身价值**: 41 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10

- **最终总分**: 39 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇高质量的 RNA 催化 RNA 结构生物学研究，在其自身领域具有原创性，但与 AML 基础研究完全无关，无法迁移。；置信度：低

摘要: Group I introns are catalytic RNAs capable of self-splicing, yet structural insights into full-length precursor RNAs and post-splicing circularization have been limited. Here, we present cryo-EM structures of the Azoarcus pre-tRNA^(Ile) system across key catalytic stages, including the full-length precursor in Pre-1S conformations, the linear intron, and the circular introns, at 2.8-3.3 Å resolution. These structures reveal a preformed P1 helix, register shifts during splicing, and a…

—

4 DGAT1 mediates sex-specific CD8⁺ T cell antitumour responses

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 33 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文在肿瘤免疫代谢领域有一定新意，但完全不涉及 AML、白血病干细胞、骨髓微环境或任何血液肿瘤模型，与 AML 基础研究无直接或强迁移关联。；置信度：低

摘要: Fatty acid (FA) oxidation plays an important role in T cell responses. However, whether DGAT1-mediated FA esterification to triacylglycerol also regulates T cell function remains unclear. Here we uncover a sexually dimorphic requirement for DGAT1 expression in CD8^(+) tumour-infiltrating lymphocyte function. In female mice, T cell-specific Dgat1 deficiency improves mitochondrial metabolic fitness and expands the pool of progenitor exhausted CD8^(+) T (T(ex)) cells to sustain antitumour…

—

5 Distinct roles of cortical layer 5 subtypes in associative learning

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇神经科学领域探讨皮层神经元亚型功能分化的研究，但与 AML 基础研究（LSC、克隆演化、耐药、微环境等）无任何逻辑关联。；置信度：低

摘要: Adaptive behavior relies on associating sensory cues with rewarding or aversive outcomes. In mammals, the primary sensory cortex processes stimuli and distributes information to cortical and subcortical targets. Layer 5 (L5) contains two major projection neuron classes, intratelencephalic (IT) and extratelencephalic (ET); however, their roles in

associative learning remain unclear. Using transgenic mice, we identified IT and ET neurons in primary somatosensory cortex and tracked their activity…

—

6 Reliving emotional memories: Episodic memory retrieval elicits affective psychophysiological responses

- **研究本身价值**: 35 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇心理学领域针对记忆回忆与情感生理反应的现象级实验研究，与 AML（急性髓系白血病）基础研究在生物学机制、造血系统或肿瘤微环境等方面完全无关。；置信度：低

摘要: Memories of emotional events can guide behavior in the present. One way to fulfill this adaptive function might be through psychophysiological responses that signal desirable and undesirable outcomes. However, it remains unknown whether remembering emotional episodes indeed re-elicits corresponding affective psychophysiological responses. We addressed this question in two experiments (N1 = 48, N2 = 59). Young adults watched positive, negative, and neutral movie clips and recalled these episodes…

—

7 Network model for alignment, stitching and slice-to-volume 3D reconstruction of large-scale spatially resolved slices

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用计算方法学论文，聚焦空间-omics 数据整合的技术挑战，与 AML 基础研究无任何直接关联，仅在通用空间分析流程层面存在极有限的间接可迁移性。；置信度：低

摘要: Advances in spatially resolved technologies enable the characterization of tissues at molecular resolution by preserving spatial information. However, integrating and aligning spatial-omics data across different platforms and modalities remains challenging. Flexible tools for slice alignment, stitching and slice-to-volume 3D reconstruction are still lacking because available spatial-omics datasets are affected by partial overlapping, local non-rigid deformations, and large-scalability. Here we…

—

8 Parvalbumin-positive neurons in the medial septum participate in the formation of hippocampal-dependent spatial memory

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 2 / 30

- **表达与可复用性**: 5 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇神经科学领域的功能机制研究，在其本领域有一定针对性，但与 AML 基础研究（包括 LSC、克隆演化、耐药、微环境、表观遗传等）完全无关，借鉴价值为零。；置信度：低

摘要: GABAergic medial septal neurons play a key role in regulating hippocampus-dependent spatial memory, but the underlying mechanisms are not yet fully understood. In male mice, we establish an object-place recognition impairment model using an acute sleep deprivation protocol. Here we show that parvalbumin-positive neurons in the medial septum regulate object-location memory specificity during encoding by modulating hippocampal place cell population activity. Using in vivo electrophysiology…

—

9 Hijacking ERAD for targeted degradation of transmembrane proteins

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇通用药物发现领域的概念性技术论文，与 AML 基础研究无任何直接或间接关联。；置信度：低

摘要: Targeted protein degradation (TPD) technologies provide huge opportunities for drug discovery, but degrading transmembrane (TM) targets remains challenging. Since TM proteins are canonically folded on the endoplasmic reticulum (ER) membrane, we hypothesized that harnessing ER-associated degradation (ERAD) may enable efficient degradation of TM proteins. Here, we established a TPD technology hijacking ERAD and named it ERAD-engaging chimeras (ERADECs), capable of degrading TM targets with high…

—

10 Ultra-precision deconvolution of spatial transcriptomics decodes immune heterogeneity and fate-defining programs in tissues

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯方法学论文，聚焦空间转录组去卷积技术的精度提升，但全文仅提供标题与摘要，未涉及任何 AML 相关样本、机制或数据。；置信度：低

摘要: Elucidating the spatial organization and functional specialization of immune cells within complex tissues remains challenging. We present UCASpatial, an ultra-precision

spatial transcriptomics deconvolution algorithm utilizing entropy-based weighting to accurately map cell subpopulations. Benchmarking confirms its superiority in identifying low-abundant cell subpopulations and distinguishing transcriptionally heterogeneous cell subpopulations. Applying UCASpatial to human colorectal cancer, we…

—

11 A multi-ancestry genome-wide study of tamoxifen metabolism and breast cancer recurrence

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇聚焦乳腺癌他莫昔芬药代动力学的 GWAS 文章，与 AML 基础研究（LSC、克隆演化、耐药、微环境、表观遗传等）完全无关。；置信度：低

摘要: Tamoxifen’s pharmacokinetics are strongly influenced by the highly polymorphic CYP2D6, while the influence of other genetic variants has been inconclusive. To further delineate this genotypic-phenotypic impact, we conducted a multi-ancestry genome-wide association study in 636 hormone-receptor-positive (HR+) breast cancer (BC) patients treated with 20 mg tamoxifen daily for ⩾8 weeks and validated these genetic determinants in another 869 patients. Association with clinical outcomes was examined…

—

12 Targeting ubiquitin signaling vulnerabilities in KEAP1-inactivated lung cancer

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦非 AML 癌症（肺癌）的 UPS 相关脆弱性筛选研究，提供了一定方法学思路，但对 AML 的直接借鉴价值有限，且本文内容高度不完整。；置信度：低

摘要: Lung cancer cells rely on protein homeostasis regulators, particularly the ubiquitin-proteasome system (UPS), to sustain malignancy. Genetic alterations in UPS components, such as E3 ubiquitin ligases (E3s) and deubiquitinating enzymes (DUBs), are common and create context-dependent therapeutic dependencies. To investigate how these genetic alterations drive tumor formation, we conducted CRISPR screens on metabolically stressed murine lung cancer models and identified specific cancer…

—

13 A single cluster of RNA Polymerase II molecules is stably associated with active genes

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 6 / 30

- **表达与可复用性**: 4 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇针对真核转录调控基础争论的发育生物学观察性研究，与AML基础研究无任何直接或逻辑可迁移关联。；置信度：低

摘要: In eukaryotic nuclei, transcription is associated with the clustering of RNA Polymerase II (RNAPII) molecules. The mechanisms underlying cluster formation, their interactions with genes, and their impact on transcriptional activity remain heavily debated. Here, we take advantage of the naturally occurring increase in transcriptional activity during Zygotic Genome Activation (ZGA) in Drosophila melanogaster embryos to characterize the functional roles of RNAPII clusters in a developmental…

—

14 When bubbles bounce or stick

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇典型的流体力学现象分类研究，与AML或任何血液学/白血病基础研究无逻辑关联。；置信度：低

摘要: Bubble collisions with solid surfaces are critical in both natural and industrial contexts, ranging from gas absorption in bioreactors to marine aerosol production. Despite their ubiquity, the physical criteria underlying bounce dynamics remain unresolved. Here, through experiments and simulations, we map the phase diagram of rising bubbles impacting a wall in Galilei (Ga)-Bond (Bo) space, revealing four dynamic regimes: fully bouncing, underdamped non-bouncing, overdamped non-bouncing, and…

—

15 Optimal DEtection of Atrial fibrillation in Transient Ischaemic Attack: Rationale and design of the ODEA-TIA trial

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯临床试验方案设计论文，与AML基础研究（LSC、克隆演化、表观遗传、代谢重编程等）毫无关联，仅为心血管/神经领域问题。；置信度：低

摘要: The detection of atrial fibrillation (AF) after transient ischaemic attacks (TIA) is crucial for initiating effective secondary prevention using oral anticoagulation. The choice of the best method and the duration of ECG monitoring to detect AF is uncertain. ODEA-TIA

is a multicentre, investigator-initiated, randomised, open-label, blinded-endpoint (PROBE) clinical trial designed to compare three ECG monitoring strategies for detecting AF in patients aged 50 years or older who experienced a TIA…

—

16 The impact of first birth by cesarean delivery on subsequent reproductive outcomes - a population cohort study

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的流行病学关联研究，自身设计基本合理但证据仅停留在相关性层面，对 AML 无任何可迁移价值。；置信度：低

摘要: CONCLUSIONS: First birth by cesarean delivery is associated with an 11% reduced likelihood of a second live birth within the twelve-year study period and a 28% increase in the use of assisted reproductive technologies to achieve a second birth. Factors leading to a cesarean delivery may also be associated with subsequent reproductive outcomes and warrant further study.

—

17 Controlled human influenza infection reveals heterogeneous expulsion of infectious virus into air

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-20
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇流感病毒空气传播方法的描述性研究，与 AML 基础研究（LSC、克隆演化、分化阻滞、耐药、微环境、表观遗传等）无任何直接或可迁移关联。；置信度：低

摘要: Influenza virus is transmitted via respiratory expulsions, but detecting infectious virus in expulsions is challenging. Here, we describe quantification and genotyping of infectious virus in respiratory particles using a modular influenza sampling tunnel (MIST). The particles deposit on cell monolayers, enabling culture, quantification, and sequencing of viruses. Concomitantly, water-sensitive paper and fine particle samplers yield respiratory particle counts over a broad size range. Using the…

—