追觅 - 2026-03-24

生成时间: 2026-03-24 07:05

分析文章数量: 24

1 Lactylation stabilizes PD-L1 to promote tumor immune evasion and cell growth

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 15 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 46 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐（基于当前可用文本）；机制假说具有一定新意且与肿瘤免疫代谢交叉点相关，但当前仅有标题与不完整摘要，证据链无法核验；对 AML 的可迁移性存在但需实证支持。；置信度：低

摘要: Programmed death-ligand 1 (PD-L1) plays a critical role in tumor immune evasion, yet the mechanisms that regulate its expression, specifically the metabolic control of its stability and function, remain elusive. In this study, we demonstrate that lactate, a key metabolite in the tumor microenvironment, upregulates PD-L1 expression via lysine lactylation (Kla) of PD-L1 at residue K280 within its intracellular domain. This modification stabilizes PD-L1 by inhibiting E3 ligase HUWE1 binding,…

—

2 RF-SIRF reveals a replication stress-specific epigenetic code by spatio-temporal mapping of reversed forks

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 16 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 44 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐（基于当前可用文本）；从题目/摘要看方法潜力较大，但当前仅有极不完整摘要，关键证据链（验证、样本、统计、因果机制）均无法核查，现阶段只能把它当作“可能值得关注的技术线索”，不宜采信机制

摘要: DNA replication stress responses are guardians of genomic stability critical during development, hematopoiesis, cancer therapy response, aging and disease suppression. Central to these responses are reversed forks (RF), which are distinct four-way DNA structures formed during DNA replication stalling to protect against toxic DNA lesions. Historically, RF detection relies on specialized electron microscopy, precluding studies within their native cellular context. By harnessing intrinsic…

—

3 Activation of L-histidine biosynthesis as a new antibiotic strategy against Mycobacterium tuberculosis

- **研究本身价值**: 32 / 60

- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 42 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；概念具有一定新颖性，但当前仅凭题目与不完整摘要，关键证据链（体内有效性、安全窗口、机制闭环与广谱性）无法判断；且与 AML 的直接相关性很弱。；置信度：低

摘要: The increasing prevalence of antimicrobial resistance is an important challenge that warrants new approaches to antibiotic development. Currently, all antibiotics inhibit biological processes. To explore whether activation of a biochemical pathway can elicit bactericidal effects we engineered variants of Mycobacterium tuberculosis ATP-phosphoribosyltransferase (ATP-PRT) that are resistant to allosteric inhibition by L-histidine, leading to supraphysiological activation of ATP-PRT and L-histidine…

—

4 Hyperinnervation inhibits organ-level regeneration in mammalian skin

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 42 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐（基于当前可用文本的审稿式评估）；从有限摘要看选题有潜在重要性，但当前文本未提供关键证据链与验证闭环，难以支持强结论，且对 AML 的可迁移性偏间接。；置信度：低

摘要: Some mammalian tissues can replace lost cells within one lineage, but organ-level regeneration-restoring diverse cell types across lineages-remains rare. Here, we show that late embryonic full-thickness skin injuries heal by regenerating epithelial, mesenchymal, neuronal, and vascular tissues with proper connectivity. However, this ability is lost soon after birth, resulting in failure to restore most cell types and hyperinnervation within the wound bed. Single-cell sequencing identified a…

—

5 In vivo metabolic tagging and targeting of circulating red blood cells

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 40 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐（基于当前可用文本的严格审稿结论）；概念上有新意且可能形成通用的体内细胞工程工具，但基于当前仅有的摘要片段，证据链完整性、标记效率/特异性、安全性与可重复性均无法评估，对 AML 的直接借鉴价𝄢

摘要: Engineering red blood cells (RBCs) has been widely explored for drug delivery, imaging, vaccination, and other applications. However, effective strategies to directly

engineer RBCs in vivo are still lacking. Here, we report successful metabolic glycan labeling of RBCs in vivo. We demonstrate that systemically administered azido-sugars can metabolically label circulating RBCs with azido groups, through labeling of both mature RBCs and RBC precursor cells. The surface azido tags on RBCs can…

—

6 The E3-ome gene-centric compendium reveals the human E3 ligase landscape

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 15 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 39 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；作为“目录+注释框架”具有潜在工具价值，但从当前摘要信息看，证据链、验证策略与可复现性细节不足，且对 AML 的直接可迁移结论有限。；置信度：低

摘要: To define and systematically characterize the human E3 ubiquitin ligase (E3) landscape, we generated the E3-ome, a compendium of E3s encoded by the human genome. The E3-ome integrates experimental data, bioinformatics, and published research, revealing 672 high-confidence E3s. We standardized E3 classifications to create a unified framework for annotation and comparative analysis. The E3-ome identified several previously unrecognized domains, motifs, E3 candidates, and relationships, expanding…

—

7 CSDE1-mediated histone lactylation modification of the HOXD9 promoter promotes gastric cancer progression

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 36 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从题目/摘要看概念有一定新意，但当前提供信息极不完整，证据链强度、因果性与可复现性均无法核验；对 AML 的直接借鉴价值有限。；置信度：低

摘要: Recent research has underscored the crucial role of dysregulated RNA-binding proteins (RBP) in the onset and advancement of gastric cancer. Cold shock domain containing E1 (CSDE1), functioning as an RBP, has emerged as an essential contributor to this process, although its specific mechanistic roles in GC are still not fully understood. Immunohistochemistry was performed on human gastric cancer tissue microarrays to assess CSDE1 expression. Extracellular acidification rate, glucose uptake, and…

—

8 Co-expression network multivariate regression

- **研究本身价值**: 18 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 3 / 10

- **最终总分**: 33 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；问题方向重要但当前仅凭截断摘要无法验证方法细节、验证强度与可复现性，尚不足以支持较高可信度与明确可用性结论。；置信度：低

摘要: Accounting for dependence among high-dimensional variables in omics data analysis is critical to obtain accurate and reliable statistical inference. Although latent, omics variables often exhibit structured correlation/co-expression patterns. However, there are few methods explicitly accounting for such structured dependence in the statistical analysis of omics data (e.g. differential expression analysis). To address this methodological gap, we propose a Coexpression network multivariate…

—

9 Exploration of Agonist and Antagonist Binding Sites within the Cytosolic AHR Complex Using Molecular Modeling

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；在结构层面可能有补充价值，但目前提供文本不足且看不到功能/生物学闭环证据，对 AML 的可迁移性主要停留在“靶点提示/药化思路”层面。；置信度：低

摘要: The aryl hydrocarbon receptor (AHR) is a ligand-activated transcription factor involved in metabolism, cell motility, development, and immune responses. Its dysregulation is linked to various diseases, including cancer, in which it can enhance tumor progression and suppress immune responses. High-resolution cryo-electron microscopy (cryo-EM) structures of the human cytosolic AHR complex have recently been solved and have provided insights into its agonist-binding mechanisms. However, our…

—

10 Role of lactylation-induced macrophage failed phenotypic switching in sustaining inflammation of diabetic wounds

- **研究本身价值**: 18 / 60
- **AML 借鉴价值**: 11 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从免疫代谢-表观遗传角度切入的综述选题有启发，但基于当前仅有题目与极简摘要，无法评估其证据链系统性与结论边界；对 AML 的直接借鉴主要停留在“思路层面”。；置信度：低

摘要: The impaired healing of diabetic wounds is closely associated with a persistent inflammatory response, wherein macrophages, as crucial immune effector cells in the local

wound microenvironment, play a vital role in maintaining inflammatory equilibrium. Increasing evidence indicates that lactate, a product of glycolysis, is now recognized as a novel regulator of macrophage function by influencing gene transcription through protein lactylation on histone and non-histone substrates. This review…

—

11 Flat band induced quasi-one-dimensional magnon transport in a two-dimensional spin lattice

- **研究本身价值**: 33 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；对自旋晶格与磁振子输运可能有物理学意义，但就现有摘要信息而言与AML 基础研究几乎无直接可迁移关联，无法支撑对 AML 机制/靶点/微环境等层面的借鉴。；置信度：低

摘要: Flat bands in electronic systems have been widely explored for their ability to enhance electronic correlations and to induce exotic quantum states. However, the impact of flat magnon bands in spintronic devices remains largely unexplored. In this study, we experimentally demonstrate that intrinsic magnonic flat bands in the two-dimensional antiferromagnetic insulator CrOCl enable long-range, quasi-one-dimensional magnon transport within a strongly coupled two-dimensional spin lattice. We…

—

12 H3K18 lactylation promotes POM121 transcription and accelerates gastric cancer progression via the PI3K/AKT pathway

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从题目/结论看是一个“表观修饰-转录-通路”的肿瘤机制叙事，但在缺乏关键实验与数据细节的情况下，证据链强度与可重复性无法评估，且对 AML 仅具概念性启发。；置信度：低

摘要: CONCLUSIONS: H3K18la enhances the malignant behavior of GC cells through activation of the POM121/PI3K/AKT pathway. These findings provide new insights into the role of histone lactylation in GC progression and suggest that targeting the H3K18la-POM121-PI3K/AKT axis may represent a potential therapeutic avenue worthy of further investigation.

—

13 An activated wheat CC_{G10}-NLR immune receptor forms an octameric resistosome

- **研究本身价值**: 31 / 60

- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就原领域（植物 NLR 结构免疫学）可能有价值，但基于当前仅有的题目+不完整摘要，证据链无法评估；且与 AML 基础研究关联度很弱，难以直接借鉴机制结论。；置信度：低

摘要: Nucleotide-binding, leucine-rich repeat (NLR) receptors are widespread intracellular immune sensors across kingdoms. Plant G10-type coiled-coil (CC(G10))-NLRs constitute a distinct phylogenetic clade that remains poorly characterized. Here, we identified a gain-of-function mutant of wheat autoimmunity 3 (WAI3^(GOF)), which encodes a constitutively active CC(G10)-NLR resulting from a residue substitution in the leucine-rich repeat (LRR) domain. Cryo-electron microscopy (cryo-EM) analysis reveals…

—

14 Enhancing cancer classification accuracy with a self-attention network using panel capture sequencing data

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；方法学方向有潜在实用性，但当前仅凭题目与不完整摘要，关键证据链（数据构成、独立验证、混杂控制、与临床基线方法对比）未提供，结论可信度与可迁移性均需大幅降权。；置信度：低

摘要: Cancer classification is pivotal for precision oncology, yet traditional methods struggle with the molecular heterogeneity of tumors. Our study introduces a self-attention based Conv1D machine learning network designed for panel capture sequencing data, which is more commonly used in clinical settings. Combining clinical capture sequencing data and The Cancer Genome Atlas data, we achieved an overall classification accuracy of over 90%, with precision rates reaching 100% for cervical and gastric…

—

15 Deciphering mechanical determinants of morphological evolution

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-21
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从摘要可见是较强的力学生物学/理论建模工作，但当前文本信息极少且与 AML 并不直接相关，对 AML 更多是“方法与概念借鉴”，无法支持机制与证据链的高评分。；置信度：低

摘要: How morphological diversity arises from variations in biomechanical processes remains an open question. Although forces shape tissues, how force-generating systems differ across species to create diverse forms is unclear. Here, we combine comparative morphogenesis and active matter theory across six cnidarian species spanning 500 million years of divergence to identify the mechanical basis of larval shape diversity. We define species-specific configurations of mechanical modules-termed…

—

16 Metabolic pathways and chemotherapy resistance in acute myeloid leukemia (AML): Insights into Enoyl-CoA hydratase domain-containing protein 3 (ECHDC3) as a potential therapeutic target

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；研究问题方向重要，但当前仅有结论性表述、缺乏任何方法与数据支撑，无法验证因果与可转化性，按严格标准不宜采信。；置信度：低

摘要: CONCLUSIONS: ECHDC3 contributes to chemotherapy resistance in AML by regulating mitochondrial function, autophagy, and metabolic pathways. This study highlights ECHDC3 as a potential therapeutic target and offers insights into the development of AML treatments to overcome drug resistance.

—

17 Smallest acyclic tricationic molecule containing a Bis(phosphine)-stabilized low-valent triantimony-based Unit

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就化学结构新颖性而言可能有价值，但与 AML 基础研究几乎无直接关联，且仅凭当前题目+不完整摘要无法评估证据链强度与可复现性。；置信度：低

摘要: Element-element bonded multiply charged cationic species are well known as dimers or small cyclic oligomers in the condensed phase. However, the smallest acyclic version, a trinuclear unit possessing greater than a monocationic charge, has remained elusive. Here we introduce a bis(phosphine) supported low valent triantimony-based tricationic compound as a new entrant in this field. Structural elucidation and electronic understanding reveal a W-shaped tricationic unit comprising of a three-center…

—

18 High-κ KBe₂BO₃F₂ dielectric material with wide bandgap for two-dimensional electronics

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：**不推荐**；从现有摘要看，这是“二维器件介电材料”工作，与 AML 基础研究几乎无直接关联，且当前文本不足以评估其数据与证据链，难以推荐给 AML 研究者作为重点阅读。；置信度：低

摘要: Further transistor scaling requires gate dielectric materials with both high dielectric constant and wide bandgap to improve gate control and suppress leakage current. However, these two properties typically exhibit an inverse relationship, making it challenging to simultaneously optimize both. Here, we report the exfoliation of KBe(2)BO(3)F(2) (KBBF) dielectric nanosheets that overcome this trade-off. The material possesses a wide band gap exceeding 8 eV and a bulk dielectric constant of 63⋯.

—

19 Metal-organic framework-confined Co₃O₄ for humidity-immune ozone decomposition

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：**不推荐**；从现有摘要看属于环境催化材料优化工作，对 AML 基础研究几乎无直接可迁移结论；且目前信息不足以严格评估其数据与机理证据链。；置信度：低

摘要: Ground-level ozone (O(3)) is a major air pollutant, and catalytic decomposition represents a promising strategy for its removal. However, maintaining high catalytic efficiency under humid conditions remains a considerable challenge. In this study, we encapsulate ultrafine metal oxides (UMOs; e.g., Co(3)O(4), NiO) within the nanopores of an Fe(3)O-cluster-based metal-organic framework, PCN-333(Fe), for catalytic O(3) decomposition. The optimized 30% Co(3)O(4)@PCN-333(Fe) catalyst achieves⋯

—

20 Macroscopic particle transport in dissipative long-range bosonic systems

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就其自身领域看可能是严谨的理论工作，但在当前提供的有限摘要信息下证据链无法核查；且与 AML 基础研究几乎无直接可迁移关联。；置信度：低

摘要: Dissipation in quantum many-body systems provides a more general and experimentally realistic perspective on particle transport than closed quantum systems. In this work, we determine the maximal speed of macroscopic particle transport in dissipative bosonic systems featuring both long-range hopping and long-range interactions. By developing a generalized optimal transport theory for open quantum systems, we rigorously establish the relationship between the minimum transport time and the…

—

21 The Inherited Basis of Coronary Artery Disease

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；作为 CAD 遗传学综述可能有科普/领域梳理价值，但就当前提供文本而言缺乏可审查的证据链细节，且对 AML 基础研究的直接借鉴价值很弱。；置信度：低

摘要: Investigations of the genetic basis of coronary artery disease have led to advances in mechanistic insights, therapeutics, prevention, and risk prediction. Indeed, most contemporary medicines for coronary artery disease target pathways that promote atherosclerosis due to underpinning genetic mechanisms. Monogenic causes of coronary artery disease occur in approximately 1 out of 250 people and mostly result in massively elevated lipid levels. At the population level, hundreds of common variants…

—

22 CNN-based detection of pediatric lymphoma on whole body [(18)F]FDG-PET/MRI

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就“影像自动检出”应用问题有一定价值，但当前提供信息仅到训练/测试规模与比较对象层面，证据链与可复现性难评；对 AML 基础研究的直接借鉴价值很弱。；置信度：低

摘要: We assessed the performance of a deep convolutional neural network (CNN) in detecting pediatric lymphoma lesions on [^(18)F]FDG-PET/MRI. We evaluated CNN’s sensitivity, specificity, percentage agreement, and processing time compared to the interpretations of a pediatric radiologist and a second-year radiology resident. In this retrospective study, a CNN was trained on annotated [^(18)F]FDG-PET/MRI scans from 53 pediatric lymphoma patients and tested on 30 additional scans. The CNN and two human…

—

23 The regulation of artificial intelligence in intensive care units: from narrow tools to generalist systems

- **研究本身价值**: 15 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：**不推荐**；框架性讨论对医疗 AI 治理有启发，但缺乏实证数据与可检验命题，对 AML 基础研究的直接借鉴价值很弱。；置信度：低

摘要: Artificial intelligence (AI) is increasingly explored for use in intensive care units. While most approved AI devices use narrow models, research is shifting towards generalist systems based on large language models and agentic AI. In this perspective, we propose a five-paradigm framework that shows how regulatory complexity rises with AI functionality and scale. As current regulatory frameworks are device-centric, new approaches like agentic oversight are needed for orchestrating AI systems.

—

24 Consensus palliative care referral criteria for people with idiopathic pulmonary fibrosis: An international Delphi study

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-22
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；该研究更偏临床流程与共识框架，证据层级主要是专家一致性，难以直接为 AML 基础研究提供机制或可验证的生物学启示。；置信度：低

摘要: CONCLUSION: Consensus referral criteria for specialist palliative care referral in ILD are presented, which emphasise comprehensive needs assessment and open communication between clinicians and patients. Future work is needed to examine the implementation of these criteria in clinical respiratory care.

—