# 追觅 - 2026-03-14

生成时间: 2026-03-14 23:07

分析文章数量: 59

## 1 DNMT3B Controls Enhancer-Linked Chromatin and Cell Cycle Networks in Acute Myeloid Leukemia

- **研究本身价值**: 34 / 60
- **AML 借鉴价值**: 25 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 55 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：一般；文章选题直接切中 AML 表观遗传异常核心，但本次评估仅基于标题+摘要开头（无结果、图表、讨论），证据链完全缺失，难以判断结论可信度。；置信度：低

摘要: Background: DNMT3B is frequently overexpressed in molecular subsets of acute myeloid leukemia (AML) and is associated with poor prognosis. Unlike DNMT3A, DNMT3B is rarely mutated, suggesting dysregulation through epigenetic mechanisms. The regulatory basis and downstream consequences of DNMT3B overexpression in AML remain incompletely defined. Methods: We integrated analyses of BeatAML, TCGA, and BLUEPRINT cohorts with multi-omic profiling (RNA-seq, DNA methylation, ATAC-seq, and proteomics) in…

—

## 2 IDH2 Clonal Hematopoiesis and IKAROS Loss Cooperate in a B-ALL Subtype after Lenalidomide Therapy for Multiple Myeloma

- **研究本身价值**: 34 / 60
- **AML 借鉴价值**: 22 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 55 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：一般；摘要显示 IDH2 突变在 LenB-ALL 中显著富集并形成三大亚群，具有临床相关性，但机制“合作”仅为推断性描述，未见功能证据。；置信度：低

摘要: Lenalidomide, a maintenance treatment in multiple myeloma first-line therapy, increases the risk of secondary malignancies, including B-cell precursor acute lymphoblastic leukemia (B-ALL). We present a comprehensive molecular characterization of 57 patients with lenalidomide-associated B-ALL (LenB-ALL), revealing three mutational subgroups: (1) TP53mt (30%), (2) IDH2mt (p.R140Q) (23%) and (3) other, including NRAS/KRASmt. Remarkably, IDH2 R140Q mutations were highly enriched in LenB-ALL compared…

—

## 3 Defective cerebrovascular development in mice lacking TFPI is restored by activated protein C

- **研究本身价值**: 38 / 60

- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 41 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为凝血平衡在胚胎脑血管发育中的作用提供经典 in vivo 证据，但主题完全属于发育生物学/凝血领域，与 AML 基础研究无直接或可迁移关联。；置信度：低

摘要: Mice lacking Tissue Factor Pathway Inhibitor (Tfpi-/-) succumb to embryonic lethality from excess thrombin production and associated cerebrovascular defects called glomeruloid bodies. A transgene producing hyperactivatable mouse protein C (hMPC) was bred into Tfpi+/- mice to determine if excess activated PC (aPC) would correct the cerebrovascular defects in Tfpi-/- embryos. Tfpi-/-/hMPC+ embryos survived to adulthood. Despite the rescue of embryonic lethality, hMPC reduced glomeruloid body…

—

## 4 Decoding GPX4 regulation in ferroptosis: mechanisms and therapeutic implications

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 38 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦铁死亡关键调控因子的文献综述，非原创机制或生信研究，基于提供的有限摘要无法判断其对 AML 的特异贡献。；置信度：低

摘要: Ferroptosis, a regulated form of cell death, is determined by iron-dependent lipid peroxidation. A selenoenzyme called glutathione peroxidase 4 (GPX4) detoxifies phospholipid hydroperoxides at the heart of this process. As the pivotal gatekeeper of ferroptosis, GPX4 is implicated in a wide range of pathologies, including cancer, neurodegeneration, acute renal failure, and infection. In this review, we discuss how GPX4 transcription and mRNA stability are controlled by transcription factors,…

—

## 5 A Modified-Delphi Consensus on the Management of Patients with FLT3-Mutated AML

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 37 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为临床实践指导性共识，而非 AML 基础机制、生信或功能研究，基于极有限摘要文本，无法评估任何实验证据链。；置信度：低

摘要: Background/Objectives: The emergence of FLT3 inhibitors (FLT3i) has radically transformed the prognostic and therapeutic landscape for FLT3-mutated Acute Myeloid Leukemia, stimulating the need for comprehensive and structured clinical guidance. Methods: We aimed to develop evidence-based recommendations spanning the entire disease continuum of FLT3-mutated AML from leading Italian experts through a modified Delphi consensus process. Results: The panel achieved a high degree of agreement on…

—

## 6 A corticothalamic circuit modulates pain sensitivity and mediates innate fear-induced analgesia in male mice

- **研究本身价值**: 35 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 35 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇典型的神经科学机制研究，与 AML、白血病干细胞、造血微环境或任何血液学核心问题完全无关。；置信度：低

摘要: Fear and pain are two frequently co-occurring states that mammals need to orchestrate to ensure survival. Nevertheless, how the brain dynamically prioritizes between them remains poorly understood. Here, we demonstrate that innate fear suppresses both acute and chronic pain, whereas pain does not reciprocally modulate fear responses in male mice. Using fiber photometry, virus tracing, and electrophysiological approaches, we show that exposure to a fear-inducing odor activates GABAergic neurons…

—

## 7 Lactate Promotes Endothelial-Mesenchymal Transition via Mediating Twist1 Lactylation in Hypoxic Pulmonary Hypertension

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 PH 病理机制的非 AML 研究，基于有限摘要文本，核心结论依赖体外/体内乳酸调控实验，但与 AML 基础研究关联极弱。；置信度：低

摘要: Elevated plasma lactate is a significant risk factor in pulmonary hypertension (PH), and endothelial-mesenchymal transition (EndoMT) is a major contributor to this pathological process, yet its specific role in driving endothelial-mesenchymal transition (EndoMT) remains unclear. Using in vivo and in vitro models, we demonstrate that modulating lactate levels critically influences PH progression. In a hypoxic PH mouse model, inhibition of lactate production ameliorated hemodynamic and vascular…

—

## 8 Butyrate extends health and lifespan in mice with mitochondrial deficiency

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；基于极有限的标题与摘要片段，文章仅呈现了一个线粒体缺陷模型的表型现象及丁酸盐的潜在缓解作用，学术价值与 AML 借鉴价值均无法实质评估。；置信度：低

摘要: Mitochondrial diseases progressively lead to multisystemic failure with treatment options remaining extremely limited. Here, to investigate strategies that alleviate mitochondrial dysfunction, we first generate a ubiquitous and tamoxifen-inducible knockout mouse model of mitochondrial transcription factor A (TFAM), a nuclear-encoded protein involved in mitochondrial DNA (mtDNA) maintenance - Tfam^(fl/fl)Ubc^(Cre-ERT2) (iTfamKO) mice. Systemic TFAM deficiency triggers mitochondrial decline in a…

—

## 9 Saccades orchestrate intraocular glucose dynamics to shape visual responses in birds

- **研究本身价值**: 36 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇视觉神经生理领域的现象-代谢-功能关联研究，但与 AML 基础研究（LSC、代谢重编程、骨髓微环境等）无任何逻辑迁移路径。；置信度：低

摘要: Birds exhibit remarkable vision despite lacking the classical retinal vasculature, raising an enigmatic question about how avascular retinas fulfill energy demands necessary for sight. Avian saccades couple rapid gaze shifts with cyclotorsional oscillations. In pigeons, we show that these saccades orchestrate intraocular metabolic dynamics and visual processing. Using combined measurements of eye movements, intraocular glucose, and neuronal activity in retinorecipient brain regions, we found…

—

## 10 Toward a consensus on the tumor microbiota: Evidence, standards, and interpretation

- **研究本身价值**: 32 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇肿瘤微生物组领域的极化问题共识评论，非原创机制或数据驱动研究，与 AML 基础研究无直接关联。；置信度：低

摘要: Tumors across diverse organs harbor microbial communities that can shape cancer biology and therapeutic responses, yet the field remains polarized by technical and interpretive challenges. In this commentary, we synthesize functional and mechanistic evidence linking intratumoral microbes to cancer hallmarks; critically evaluate current detection approaches; and propose minimal technical and reporting standards to establish microbial presence, viability, and causality. We highlight common…

—

## 11 Targeting Prolyl 3-hydroxylase 1 inhibits pancreatic cancer progression and macrophage immunity

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；该文聚焦 PDAC 中 P3H1 的新潜在功能，但提供的文本仅为标题+摘要片段，无法形成完整证据链。；置信度：低

摘要: Pancreatic ductal adenocarcinoma remains one of the most formidable challenges in oncology, with limited treatment options and a poor prognosis. Understanding the key pathways affecting cancer progression is crucial for the development of therapeutic strategies. Here, we reveal a pivotal role of Prolyl 3-hydroxylase 1 in pancreatic ductal adenocarcinoma using transcriptome sequencing, proteomic analyses and engineered mouse model. Mechanistically, our findings indicate that this effect is, at…

—

## 12 Riboflavin metabolism shapes FSP1-driven ferroptosis resistance

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于 CRISPR 筛选提出核黄素代谢是 FSP1 驱动铁死亡抵抗的新调控因素，但提供的文本仅为标题与摘要片段，未涉及 AML 或造血系统。；置信度：低

摘要: Membrane protection against oxidative insults is achieved by the concerted action of glutathione peroxidase 4 (GPX4) and endogenous lipophilic antioxidants such as ubiquinone and vitamin E. More recently, ferroptosis suppressor protein 1 (FSP1) was identified as a critical ferroptosis inhibitor, acting via the regeneration of membrane-embedded antioxidants. Yet, regulators of FSP1 are largely uncharacterized, and their identification is essential for understanding the mechanisms buffering…

—

## 13 Inflammatory ILC2s migrate to distal tissues during infection using stage-specific S1P receptors

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 32 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇免疫学机制研究，聚焦先天淋巴细胞再循环的分子基础，但与 AML 或造血系统完全无关。；置信度：低

摘要: Tissue-resident lymphocytes can recirculate, but the underlying molecular mechanism is poorly understood. During helminth infection, intestinal group 2 innate lymphoid cells (ILC2s) rapidly proliferate and give rise to inflammatory ILC2s (iILC2s), which migrate from the intestine to distal tissues. Here, we show in mice that the redistribution of iILC2s requires access to lymphatic vessels. Interleukin-25 (IL-25) induces a substantial change in the epigenetic landscape of iILC2s, with…

—

## 14 Non-Fickian diffusion within assemblies of the intrinsically disordered protein β-casein

- **研究本身价值**: 33 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 31 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯生物物理学机制研究，与 AML（急性髓系白血病）或造血系统无任何直接或间接关联。；置信度：低

摘要: The molecular mechanisms governing internal fluctuations in intrinsically disordered protein (IDP) assemblies are crucial to the stability and dynamics of both regulated and aberrant toxic cellular aggregates, but remain poorly understood. By comprehensively combining high-resolution quasi-elastic neutron scattering with all-atom molecular dynamics simulations, we probe the motions of [Formula: see text]-casein, a model IDP, inside its assemblies. We uncover a previously unresolved slow…

—

## 15 Humans with function-disrupting variants in the myostatin gene (MSTN) have increased skeletal muscle mass and strength, and less adiposity

- **研究本身价值**: 33 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；基于提供的有限标题与摘要片段，这是一项大规模人群遗传关联研究，证实 MSTN 失功能变异导致肌肉增加与脂肪减少，但完全未涉及 AML 或任何造血/白血病相关内容。；置信度：低

摘要: Myostatin negatively regulates skeletal muscle size in multiple species, and therefore, myostatin blockade has been therapeutically explored to promote muscle growth in humans, including to counter the muscle loss seen in obese humans using GLP1R agonists. In this study, we present results from a large multi-cohort genetic association analysis, using data from 1.1 million individuals to examine the effects of function-disrupting mutations in the myostatin gene (MSTN) on traits relevant to body…

—

## 16 Mechanical gating of tendon fibrogenic transcription in systemic sclerosis

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；该文在纤维化机械生物学领域提出创新实验模型，但与 AML 基础研究完全无关，无任何可迁移逻辑链条。；置信度：低

摘要: Fibrosis arises from aberrant tissue repair in systemic sclerosis and is characterized by progressive stiffening across multiple tissues, including tendons. How altered tissue mechanics sustain fibrotic remodeling remains poorly understood, in part because of limited experimental models that capture key biophysical features of fibrosis. Here we develop a modular cantilever-based mechano-culture platform that enables tendon-like constructs to be maintained under controlled static tension. We show…

—

## 17 Multi-Omics Characterization of Lactate-Associated Molecular Subtypes in Lung Cancer Suggests a Role for DKK1 in Lactate-Linked Migration, Invasion, and Lactylation Programs

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇针对肺癌代谢异质性的纯生信表征研究，证据链极不完整，仅停留在相关性层面，对 AML 基础研究的直接借鉴价值极低。；置信度：低

摘要: CONCLUSIONS: This study characterizes lactate-associated molecular heterogeneity in lung cancer and provides a lactate-related subtype framework and prognostic risk model for patient stratification. The findings nominate DKK1 as a lactate-responsive candidate linked to migration/invasion phenotypes and lactate/lactylation changes in vitro.

—

## 18 Hepatocyte SLCO4C1 is a cAMP uptake transporter for inhibiting lipogenesis and a therapeutic target for MASLD

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于有限摘要提示 SLCO4C1 可能为 MASLD 提供保护机制，但与 AML 基础研究（LSC、克隆演化、分化阻滞、耐药、骨髓微环境、代谢重编程等）无任何逻辑映射。；置信度：低

摘要: Metabolic dysfunction-associated steatotic liver disease (MASLD), the most prevalent cause of chronic liver disease worldwide, currently lacks precision medicine treatments. SLCO4C1 acts as a transporter for endogenous compounds and xenobiotics. This study aims to investigate whether SLCO4C1 plays a regulatory role in MASLD pathogenesis and to elucidate the underlying mechanisms. Using human, mouse and cellular models, we found that hepatocyte SLCO4C1 is upregulated in MASLD patients, where it…

—

## 19 Microglial cathepsin B is necessary for neuronal efferocytosis in zebrafish and mice during brain development

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 5 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇神经发育生物学机制研究论文，聚焦微胶细胞溶酶体消化机制，与 AML 基础研究无直接或可迁移关联。；置信度：低

摘要: Half of all newborn neurons in the developing brain are removed via efferocytosis - the phagocytic clearance of apoptotic cells. Microglia are brain-resident professional phagocytes that play important roles in neural circuit development including as primary effectors of efferocytosis. While the mechanisms through which microglia recognize potential phagocytic cargo are widely studied, the lysosomal mechanisms that are necessary for efficient digestion are less well defined. Here we show that…

—

## 20 TMEM187 is a novel modulator in the regulation of erythropoiesis

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于有限摘要提出 TMEM187 在正常红系造血中的新作用，但证据链极不完整，且与 AML 核心机制无直接关联。；置信度：低

摘要: Erythropoiesis, the process of red blood cell production, is highly dependent on iron uptake via transferrin and its receptor, transferrin receptor 1 (TfR1), but the mechanisms governing the proper recycling of TfR1 in relation to cellular iron demands remain elusive. Here, we identify human TMEM187, a Golgi transmembrane protein of unknown function, as a novel negative regulator of erythropoiesis. Lack of TMEM187 in a cell model initiates erythropoiesis without the normal induction protocol and…

—

## 21 Focus on Lactate and Lactylation Modification: The Potential Role in Ophthalmic Disease Treatment

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 12 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为新兴 PTM 领域的文献总结性综述，聚焦眼科，未包含任何原创实验数据，与 AML 基础研究关联高度间接且证据缺失。；置信度：低

摘要: Lysine lactylation represents a novel post-translational modification (PTM) involved in cellular functions including glycolysis and macrophage polarisation. It differs in form and mechanism from other PTMs such as acetylation, methylation, phosphorylation, ubiquitination, and SUMOylation. As a recently discovered modification, lactylation has been implicated in the progression of multiple diseases. Recent studies further indicate lactylation’s association with multiple ocular pathologies. This…

—

## 22 Non-invasive imaging of defence responses in plants

- **研究本身价值**: 29 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇植物免疫领域的技术创新论文，聚焦非侵入成像工具开发，与 AML 或任何血液肿瘤/造血基础研究无任何直接或间接关联。；置信度：低

摘要: Jasmonic and salicylic acids are key hormones involved in plant responses to pests and pathogens. Existing fluorescence-based approaches to imaging plant defence hormones are constrained by the need for external illumination and by autofluorescence of plant tissues, while luminescence-based ones require exogenous substrates. Here, we use jasmonate- and salicylate-responsive promoters to engineer autoluminescent plants that report hormone signalling activity with up to a 53-fold contrast. Using…

—

## 23 4-Hydroxyphenanthrene exacerbates obesity by altering gut microbiota and bile acid metabolism

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文聚焦环境污染物与肥胖的微生物-代谢机制，但与 AML 基础研究无任何直接或逻辑可迁移关联，仅为非血液学领域的独立机制探索。；置信度：低

摘要: Elevated exposure to 4-hydroxyphenanthrene (4-OHPhe) is positively associated with obesity risk. However, the specific pathogenic mechanism remains unclear. Here, we use SHapley Additive exPlanations (SHAP) and logistic regression analysis to identify a strong association between 4-OHPhe and obesity. Mechanistically, exposure to 4-OHPhe significantly downregulates the expression of hydroxylamine reductase (hcp) genes in Bacteroides uniformis (B. uniformis), thereby inhibiting the removal of…

—

## 24 Post-infection sequelae of COVID-19 and other infectious diseases-a nationwide Danish study with 40-month follow-up

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一项大规模描述性队列研究，旨在填补 COVID-19 与其他感染长期后遗症对比的空白，但与 AML 基础研究（LSC、克隆演化、耐药、微环境、表观遗传等）无任何逻辑关联。；置信度：低

摘要: Nationwide follow-up studies of long-term post-acute COVID-19 sequelae compared with sequelae of other infections have been lacking. Using nationwide registers, we analyzed all SARS-CoV-2 PCR test results, prescriptions for anti-infective agents, and hospitalizations with COVID-19 or other infections in Denmark from March 2020 to June 2023, including up to 40-month follow-up. We used Cox proportional hazards models with time-varying exposures to estimate the rates of first-time mental disorders…

—

## 25 Pyruvate metabolism enzyme Dlat induces mitochondria protein hyperacetylation to limit fatty acid oxidation in the HFpEF heart

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 25 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文基于有限摘要提出 HFpEF 中线粒体乙酰化新机制，但证据链完全未展示，且与 AML 基础研究无直接关联，仅存在极弱的代谢概念重叠。；置信度：低

摘要: Increased protein acetylation is frequently observed in the failing heart, including in hearts with heart failure with preserved ejection fraction (HFpEF). However, its role in the pathogenesis of HFpEF remains insufficiently investigated. Here, we found that HFpEF hearts displayed significantly protein hyperacetylation, which were predominantly localized to mitochondria and particularly enriched in fatty acid oxidation (FAO) pathway. Notably, Dlat, a pyruvate metabolism enzyme, was identified…

—

## 26 Operando insights on stable Cu<sup>2+</sup> active sites for efficient electrochemical CO<sub>2</sub>-to-C<sub>2</sub>H<sub>4</sub> conversion

- **研究本身价值**: 31 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯材料化学/电催化领域的催化剂优化论文，与 AML、白血病、造血干/祖细胞、肿瘤微环境或任何血液学基础问题无任何逻辑关联。；置信度：低

摘要: Copper catalysts feature unique superiorities for the electrochemical conversion of CO(2) to C(2+) fuels and chemicals. Their surface oxidation states dominantly determine the reaction pathways to various products. However, most Cu-based catalysts inevitably undergo electroreduction from Cu^(2+) to Cu^(1+) or Cu^0 species during the electrochemical CO(2) reduction. Herein, we propose a straightforward strategy to stabilize Cu^(2+) ions by coordinating them with benzobistriazole (H(2)BBTA),…

—

## 27 Acute Toxicities During Proton Therapy with or Without Simultaneous Chemotherapy in Pediatric CNS Tumors: A Retrospective Cohort Study

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇儿科放射肿瘤学临床描述性研究，与 AML 基础研究（包括 LSC、克隆演化、表观遗传、代谢、耐药等核心领域）无任何直接或可迁移关联。；置信度：低

摘要: Background: Proton beam therapy (PBT) is a valuable alternative to photon radiotherapy of CNS tumors in children and adolescents. While most recent studies deal with the outcome or long-term side effects of PBT, the aim of this study was to investigate the feasibility of PBT with a particular focus on the acute toxicity of a simultaneous radiochemotherapy (sPBCT). Patients and methods: We enrolled 199 children [median age 7.4 years (range, 0.9-17.9)], who received altogether 200 courses of…

—

## 28 Machine learning assisted single-molecule sensing towards standard-free quantification of per- and polyfluoroalkyl carboxylic acids

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯环境分析化学/纳米传感技术论文，与 AML、白血病、造血干/祖细胞、肿瘤微环境等任何基础研究领域均无关联。；置信度：低

摘要: Per- and polyfluoroalkyl carboxylic acids (PFCAs) are of global concern for their ubiquitous presence in the environment. However, precise quantification of PFCAs remains challenging due to the shortage of standards. Herein, with the aid of machine learning, a probe-directed nanopore based single-molecule electrochemical sensor is developed towards standard-free digital quantification of PFCAs. To correctly predict the signal without standards, a strict linear relationship (R^2 > 0.9998) is…

—

## 29 Positive Margin Location and Prostate Biopsy Route: A Consecutive Cohort Comparison of Transperineal and Transrectal Techniques

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇针对前列腺癌活检路线临床安全性和手术结局的回顾性比较研究，与 AML 基础研究（包括 LSC、克隆演化、微环境、表观遗传等）完全无关。；置信度：低

摘要: Background: Transperineal (TP) prostate biopsy is increasingly used due to its superior infectious safety and enhanced sampling of anterior and apical prostate regions. Concerns have been raised that TP biopsy may induce greater post-biopsy fibrosis, particularly at the apex, potentially increasing positive surgical margin (PSM) rates at radical prostatectomy (RP). We evaluated whether biopsy technique influences overall or location-specific PSMs at RP. Methods: We performed a retrospective…

—

## 30 Assessment of Mycobacterium tuberculosis dodecin scaffold as a multimerization platform on the immunogenicity of HPV L2 antigens

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 HPV 疫苗开发的蛋白支架评估研究，与 AML（急性髓系白血病）基础研究在主题、机制或实验体系上完全无关。；置信度：低

摘要: Human papillomavirus infection, the primary cause of cervical cancer (the fourth most common cancer in women), is preventable through prophylactic HPV vaccines. As an alternative to current HPV vaccines, all based on a cocktail of L1 virus-like particles (VLPs) from multiple HPV types, the L2 capsid protein offers cross-protection via a conserved epitope residing at the N-terminus between residues 20-38. The Trx-L2((20-38))−8mer (Trx-8mer) is an L2-based antigen composed of a polytope of…

—

## 31 Coenzyme-functionalized photo-redox catalysis for low-energy click labeling

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯化学催化方法开发论文，聚焦生化标记工具的能量优化，与 AML 或造血系统基础研究无任何直接或间接关联。；置信度：低

摘要: Photo-redox catalysis has emerged as a new generation of biochemical protocols in proximity labeling, noninvasive therapy, ultrafast activation. However, the well-developed photocatalysts cannot simultaneously possess low-energy absorption and high excited potentials to initiate photochemical reactions. Herein, we develop new Ru-PCs as bio-photocatalyst exhibiting both green-light absorption and high E(red)*. Ru-PCs cooperatively function with natural coenzyme riboflavin (RF) via an integrated…

—

## 32 Dimer-Projection Contact and the Clock Shift of a Unitary Fermi Gas

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 或任何血液学/白血病基础研究完全无关的物理学论文，仅摘要文本无法支撑任何生物医学迁移。；置信度：低

摘要: Understanding the dynamics of short-range correlations is a central challenge in strongly interacting Fermi gases. In ultracold gases, these correlations are quantified by the contact parameter, yet measurements to date have been limited to equilibrium systems or relatively slow, global dynamics. Here, we introduce a rapid spectroscopic technique based on projection of the interacting state onto an alternate scattering channel with a low-lying dimer state. We demonstrate contact measurements on…

—

## 33 Bioinformatics Profiling of a Paeoniflorin-Associated “Macrophage-Lactylation” Axis in Hepatocellular Carcinoma: An LDHA-Based Prognostic Model and Structural Computational Evidence

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇典型的 HCC 纯生信关联分析论文，试图将中药成分 PF 与 TME 代谢重编程（乳酸化）挂钩构建预后模型，但证据链仅停留在数据库交集与计算层面，无任何实验验证或 AML 相关内容。；置信度：低

摘要: Hepatocellular carcinoma (HCC) progression is shaped by crosstalk between the tumor immune microenvironment (TME) and metabolic reprogramming. This study aims to characterize a macrophage-lactylation molecular axis in HCC and to develop a quantitative prognostic stratification model. Using the TCGA-LIHC cohort, differentially expressed genes were intersected with Paeoniflorin (PF)-related targets, HCC disease targets, and macrophage-/lactylation-related genes to identify candidate genes⋯.

—

## 34 Follow-up care of adults after lung transplantation - an evidence based guideline

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇针对肺移植患者的临床管理指南，与 AML 基础研究、血液学机制或白血病相关领域完全无关。；置信度：低

摘要: The S3 Guideline provides evidence- and consensus-based recommendations for the structured follow-up care of adult patients after lung transplantation. Its goal is to standardize long-term management in order to optimize survival, functional status, and quality of life. The guideline is addressed to professionals in pulmonology, internal medicine, surgery, and general practice, as well as to transplant centers and inpatient institutions in the German-speaking countries of Germany, Austria, and⋯

—

## 35 Electrostatically tunable moiré-mediated Wigner states via interfacial potential engineering in 2D van der Waals heterostructures

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-14

- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇量子物理领域宣称具有前沿创新的物理学研究，与 AML 基础研究（包括 LSC、克隆演化、表观遗传、代谢重编程等任何核心问题）完全无关。；置信度：低

摘要: Electrostatically tunable quantum confinement in nanoscale systems offers unprecedented opportunities for manipulating artificial quantum matter, positioning these platforms at the frontier of quantum science. The strategic integration of distinct confinement mechanisms could revolutionize quantum functionality by enabling novel states with enhanced coherence properties. Here, we demonstrate inherently interfacial potential engineering by combining lateral semiconductor 2D moiré potentials with…

—

## 36 Ultraflexible photoelectrical impedance tomography-based imager for 3-axis robotic tactile sensing

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 19 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇材料与工程技术领域的装置创新论文，与 AML（急性髓系白血病）基础研究、造血机制、LSC、微环境或任何生物学过程完全无关。；置信度：低

摘要: Vision-based robotic triaxial tactile sensing provides superior spatial resolution and rich multimodal data. However, employing rigid CMOS imagers suffers from limitations in mechanical flexibility and large-area scalability. Here we present a large-area ultraflexible photoelectrical impedance tomography (PIT)-based imager that achieves high-fidelity triaxial tactile sensing. The 5-μm-thick PIT imager incorporates a quantum dots/metal-oxide heterojunction layer with 16 peripheral electrodes,…

—

## 37 Spatiotemporal visualization of long-range anisotropic plasmon polaritons in hyperbolic MoOCl<sub>2</sub>

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 19 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML、白血病、造血干细胞或肿瘤微环境完全无关的纯物理学论文，仅基于标题与截断摘要，无法支持任何生物医学结论。；置信度：低

摘要: Manipulating light at the nanoscale with minimal loss remains a central challenge for nanophotonic technologies that can be tackled by using the direction-dependent polariton modes supported by anisotropic materials. Although best known for their highly confined polaritons, hyperbolic materials can also host long-range directional polaritons, whose

direct observation has remained challenging as it requires experimental techniques that combine nanometer and femtosecond spatial and temporal…

—

## 38 Desegregation of neuronal predictive processing

- **研究本身价值**: 22 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 18 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯神经科学领域的理论计算研究，与 AML、造血系统或肿瘤微环境无任何关联，仅基于摘要片段无法判断其实验或机制细节。；置信度：低

摘要: Neural circuits construct internal ‘world-models’ to guide behavior. The predictive processing framework posits that neural activity signaling sensory predictions and concurrently computing prediction-errors is a signature of those internal models. To understand how the brain generates predictions for complex sensorimotor signals, we investigate the emergence of high-dimensional, multi-modal predictive representations in recurrent networks. Contrary to previous proposals of functionally…

—

## 39 Chiral Heptagon-Embedded Double [6]Helicenes via Scholl Reaction

- **研究本身价值**: 26 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 17 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯有机合成化学论文，聚焦新型手性多环芳烃的构建与光物理性质，与 AML、白血病干细胞、造血机制或任何肿瘤微环境/表观遗传研究完全无关。；置信度：低

摘要: A series of heptagon-embedded multiple helicenes was synthesized in which the heptagon subunit was fused to the π-framework through Knoevenagel condensation reactions. By controlling the conditions of the cyclodehydrogenation (Scholl) reactions, different fused chiral and twisted PAHs were accessible. The optical and electronic properties of all products were investigated by UV-vis, fluorescence spectroscopy, and cyclic voltammetry. One member of the series displays an unusual anti-Kasha-like…

—

## 40 A guard cell carbonic anhydrase binds and regulates SLAC1 separate from its catalytic activity

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 16 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A

- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为植物叶片气孔 CO_2 感知的非催化机制研究，与 AML、白血病、造血干/祖细胞、肿瘤微环境或任何血液肿瘤基础研究完全无关。；置信度：低

摘要: Stomata of plant leaves open to enable CO(2) entry for photosynthesis and close when CO(2) in the leaf is elevated. CO(2) is thought to promote stomatal closure in part by activating the SLAC1 anion channel at the guard cell plasma membrane. Carbonic anhydrases (CAs) contribute to this activation, but their contribution as distinct from CO(2)-H(2)CO(3) catalysis remains controversial. Here we show that the β-carbonic anhydrase CA4 binds selectively with the guard-cell anion channel SLAC1 to…

—

## 41 Reassessing the adverse event profiles of levetiracetam and brivaracetam: a systematic review and meta-analysis

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 16 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的临床药物 meta 分析，其科学问题、样本类型、机制探讨均无法迁移至白血病干细胞、表观遗传、代谢或微环境等 AML 核心领域。；置信度：低

摘要: CONCLUSION: LEV and BRV show favorable safety profiles that appear comparable based on meta-analytic findings, despite higher rates of some psychiatric AEs with LEV on a descriptive level. Both remain safe treatment options, though larger head-to-head trials are needed to provide definitive comparative evidence.

—

## 42 Clinical Outcomes of Dual-Beam Particle Therapy in Head and Neck Adenoid Cystic Carcinoma

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 15 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇与 AML 基础研究完全无关的临床放射治疗结局报告，仅描述治疗模式和患者纳入，未提供任何机制、生物学或血液学相关内容。；置信度：低

摘要: Background and Purpose: This study retrospectively evaluates the outcomes of head and neck adenoid cystic carcinomas (ACCs) treated with particle therapy, including carbon ion radiotherapy (CIRT) alone or combined with photon therapy, at a single institution. Methods and Materials: Patients with ACC who underwent CIRT alone or a combination of CIRT and photon therapy at the Marburg Ion Therapy Center between February 2017 and December 2023 were included. Radiation therapy was administered…

—

## 43 Liquid metal dispersed single-atom catalyst with high-temperature stability

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 15 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯材料化学领域的催化剂稳定性研究，与 AML、造血干/祖细胞、白血病机制或肿瘤微环境等生物医学领域完全无关。；置信度：低

摘要: Single-atom catalysts (SACs) enable greener and more economically sustainable chemical production by significantly improving thermocatalysis efficiency and selectivity through maximized atom utilization and highly homogeneous metal coordination environments. Unfortunately, SACs are fundamentally constrained by the stability owing to the severe aggregation of single atoms, especially under the high-temperature thermocatalysis operations, which compromises the overall catalytic performance. Here,…

—

## 44 Passivating pinholes for large-area and high-efficiency silicon solar cells with tunnel oxide passivated contact

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 15 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦光伏材料界面原子结构的非生物学论文，与 AML、白血病、造血干/祖细胞或肿瘤微环境等任何核心问题均无关联。；置信度：低

摘要: The tunnel oxide passivated contact (TOPCon) solar cell is poised to dominate silicon photovoltaics, yet the atomic-scale nature of pinholes-local disruptions in the SiO(x) layer enabling direct conduction-remains unresolved despite its critical importance for device performance. Here, using spherical aberration-corrected transmission electron microscopy, the TOPCon interface is uncovered at the atomic level, revealing two distinct pinhole types: recombinational pinholes with oxygen-depleted…

—

## 45 Data-driven intelligent carbonization unifies diverse biomass into high-performance hard carbon negative electrodes

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 14 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇能源材料科学领域的策略性论文，与 AML、白血病、造血干/祖细胞、肿瘤微环境或任何血液学机制完全无关，无法提供任何可迁移价值。；置信度：低

摘要: Sustainable batteries necessitate high-performance hard carbon negative electrodes derived from abundant biomass. However, realizing their full potential is significantly limited by the inherent diversity of biomass feedstocks, the intricate control over carbonization and resulting microstructures, and the complex interplay between processing, structure, and electrochemical performance. Here, we introduce “intelligent carbonization”, a strategy integrating programmable Joule heating (1000-2000…

—

## 46 Circulating Musclin is associated with skeletal muscle function and subclinical cardiac dysfunction in patients with cancer

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 14 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦癌症患者肌肉-心脏轴生物标志物的观察性研究，与 AML 基础研究（LSC、克隆演化、分化阻滞、耐药、微环境、表观遗传等）无任何直接或可迁移关联。；置信度：低

摘要: CONCLUSION AND IMPLICATIONS: Circulating Musclin may reveal information on skeletal muscle integrity and subclinical cardiac dysfunction, identifying cancer patients at increased functional and prognostic risk. These findings support Musclin as a biologically plausible and clinically relevant biomarker at the interface of muscle-heart signalling.

—

## 47 Mycophenolate Mofetil for Treatment of Ipilimumab-Induced Colitis in Patients with Metastatic Melanoma

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 13 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦黑色素瘤免疫检查点抑制剂副作用临床管理的观察性研究，与 AML 基础机制研究无任何直接或可迁移关联。；置信度：低

摘要: Background/Objectives: Treatment with immune checkpoint inhibitors (ICIs) in patients with advanced melanoma has greatly improved clinical outcomes but can induce immune-related adverse events (irAEs). ICI-induced immune-related (ir) colitis is one of the most common severe irAEs and is primarily managed by treatment with high-dose corticosteroids. Some patients are steroid-refractory and require additional immunosuppressants. Infliximab is commonly used as the additional treatment of choice,…

—

## 48 Residual Absolute Volume of Blastema as a Predictor of Clinical Outcomes in Patients With Wilms Tumor: A Report From the SIOP WT 2001 Study

- **研究本身价值**: 16 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 13 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇聚焦 Wilms 瘤（肾母细胞瘤）临床风险分层的预后研究，与 AML 基础研究（LSC、克隆演化、表观遗传、代谢等）完全无关，借鉴价值极低。；置信度：低

摘要: CONCLUSION: AVB after preoperative chemotherapy is a strong, independent predictor of both EFS and OS in WT across risk groups. An AVB of ⩾20 mL identifies a clinically relevant subset of patients with IRWT who may benefit from intensified therapy with doxorubicin. These findings support the use of AVB as a stratification criterion in future SIOP-RTSG protocols.

—

## 49 Advancing the elimination of cervical cancer across Africa

- **研究本身价值**: 18 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 12 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为宫颈癌公共卫生倡导性讨论，与 AML 基础研究（LSC、克隆演化、表观遗传、代谢、耐药等）无任何直接或间接关联。；置信度：低

摘要: Though cervical cancer is largely preventable, Africa faces a high burden with thousands of deaths. Highlighting the urgent need for action toward the WHO’s 90-70-90 elimination strategy, this study presents the epidemiological burden of HPV and cervical cancer in Africa and advocates for decisive measures to improve screening and implement effective prevention through vaccination. Here, we discuss which is the better option for Africa: a single-dose or a two-dose vaccine regimen. Furthermore,…

—

## 50 Cellulose-based sensors for decentralized monitoring in precision agriculture

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 11 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为农业传感器领域的综述/展望，与 AML 基础研究（包括 LSC、克隆演化、分化阻滞、耐药、表观遗传、代谢或骨髓微环境）无任何逻辑关联或可迁移性。；置信度：低

摘要: Modern agriculture requires rapid, affordable tools to monitor crops and soils directly in the field. Cellulose, the structural polymer of plants, is emerging as a versatile foundation for lightweight, biodegradable sensors that measure nutrients, moisture, stress, and disease without laboratory infrastructure. Spanning simple paper assays to flexible wearables, these platforms enable distributed, real-time insight into plant and soil health. As materials engineering converges with digital…

—

## 51 Return to Sports in Recreational Athletes After Complex Elbow Dislocation

- **研究本身价值**: 19 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 11 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；一篇骨科创伤领域针对罕见损伤后患者中心结局的回顾性研究，与 AML、白血病基础研究、造血机制、LSC 或肿瘤微环境等完全无关；置信度：低

摘要: Background/Objectives: Complex elbow dislocations are rare but severe injuries. Optimal treatment is controversial, and few studies focus on return to sports. The aim of the study is to analyze functional outcomes and return-to-sport and return-to-work metrics in recreational athletes after complex elbow dislocation. Methods: A retrospective single-center study at a level I trauma center was performed from January 2008 to December 2019. Epidemiological data, associated injuries, and treatment of…

—

## 52 Care for uncontrolled asthma in Germany - with focus on Magdeburg and Mannheim regions

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 11 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯临床观察性描述研究，与 AML 基础研究（包括 LSC、克隆演化、表观遗传、代谢重编程、微环境等）无任何关联。；置信度：低

摘要: Description of treatment and prescription patterns among asthma patients in the regions of Magdeburg (MD) and Mannheim (MA) compared nationwide.We analyzed IQVIATM LRx data 1 from the period of July 2022 to June 2023. The prescription database LRx covers about 80% of prescriptions for patients in the statutory health insurance in Germany. A machine learning model was used to assign the indication of asthma to patients with ⩾ 1 redeemed GKV prescription for respiratory diseases.In this analysis,…

—

## 53 Left-sided heart failure determines outcomes in patients with severe tricuspid regurgitation undergoing percutaneous repair

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 10 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯心血管临床预后观察论文，与 AML 基础研究（LSC、克隆演化、耐药、微环境、表观遗传等）完全无关，仅基于标题和结论片段无法支撑任何科学价值评估。；置信度：低

摘要: CONCLUSION: Consideration of left-sided pathologies in patients with significant TR is important as outcomes and predictors for survival differ. Across HF phenotypes, procedural success is associated with survival but the prognostic impact of TR reduction may unfold over time especially in HFpEF.

—

## 54 Targeting MYC in T-Cell lymphoma via epigenetic modulation with a small-molecule inhibitor

- **研究本身价值**: 10 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 8 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文仅提供标题，无摘要、方法、结果、图注或任何正文内容，无法进行实质学术评估，其 AML 借鉴价值极低。；置信度：低

摘要: No abstract

—

## 55 Barrier breakdown: insights into the skin-gut axis in psoriatic arthritis

- **研究本身价值**: 16 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 8 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文为 PsA 领域提供屏障轴的病理生理概述，但作为 AML 审稿专家，本文与白血病干细胞、克隆演化、骨髓微环境或任何 AML 核心机制均无实质关联。；置信度：低

摘要: Psoriatic arthritis (PsA) is a chronic inflammatory disease that primarily affects the skin, joints, and entheses. Growing evidence implicates alterations at the gut and skin barriers in PsA pathogenesis. These sites play a crucial role in tissue homeostasis, and their

disruption can contribute to the translocation of immune cells and bacterial products, resulting in systemic inflammation. Crosstalk between barrier sites and joints can promote or exacerbate disease development, with barriers…

—

## 56 Strong interplay between polar and structural topologies

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 6 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇凝聚态物理领域的初步发现性报道，与 AML（急性髓系白血病）基础研究完全无关，生物学相关性为零。；置信度：低

摘要: Topological structures in condensed matter systems unlock new possibilities for the development of nanoelectronic devices. However, the potential of antiferroelectrics to host topological features remains largely unexplored, constrained by significant energy barriers from antiparallel dipole coupling that suppress polarization rotation and challenges in high-quality film fabrication. Here for the first time, we find that, dislocations, the most common one-dimensional topological structures in…

—

## 57 Templated perpendicular ferroelectricity in textured Aurivillius oxide-based thin films

- **研究本身价值**: 13 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 5 / 10
- **最终总分**: 5 / 100
- **发布日期**: 2026-03-14
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；这是一篇纯材料物理/化学领域的铁电薄膜机制研究，与 AML（急性髓系白血病）或任何血液学/造血系统基础研究无任何逻辑关联。；置信度：低

摘要: The discovery of novel ferroelectric compounds and the modulation of polarization in established ferroelectric materials have persistently represented crucial and highly dynamic areas of research within the field of ferroelectric materials science. Here we demonstrate a structural templating strategy to reconfigure Aurivillius-phase oxides into metastable ferroelectric phases with tailored polarization. By stabilizing a metastable WO(3) phase within a Bi(2)WO(6) framework, we achieve robust…

—

## 58 How the classic computer game Doom became a tool for science

- **研究本身价值**: 1 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 4 / 100
- **发布日期**: 2026-03-14

- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文与 AML（急性髓系白血病）基础研究毫无关联，仅凭标题无法构成任何生物医学研究。；置信度：低

摘要: No abstract

—

## 59 Endomyocardial Biopsy Revisited: Diagnostic Value and Expanding Roles in Cardiac Amyloidosis

- **研究本身价值**: 9 / 60
- **AML 借鉴价值**: 1 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 1 / 100
- **发布日期**: 2026-03-13
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；本文主题与 AML 基础研究（包括 LSC、克隆演化、耐药、微环境、表观遗传等核心问题）完全无关，且仅提供标题、无摘要，研究质量与借鉴价值均无法实质评估。；置信度：低

摘要: No abstract

—