□□ - 2026-03-13

□□□□: 2026-03-13 23:05

□□□□□□: 62

1 VHL synthetic lethality screens uncover CBF-β as a negative regulator of STING

- □□□□□□: 36 / 60
- **AML** □□□□: 15 / 30
- □□□□□□□: 6 / 10
- □□□□: 52 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□ ccRCC □□□□□□□□□□□□□□ CBF-β-STING □□□□□□□□□□□□□□□□□□+□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Clear cell renal cell carcinoma (ccRCC) represents the most common form of kidney cancer and is typified by biallelic inactivation of the von Hippel-Lindau (VHL) tumour suppressor gene. Here, we undertake genome-wide CRISPR/Cas9 screening to reveal synthetic lethal interactors of VHL, and uncover that loss of Core Binding Factor β (CBF-β) causes cell death in VHL-null ccRCC cell lines and impairs tumour establishment and growth in vivo. This synthetic relationship is independent of the elevated…

—

2 Defining the vascular niche of human adipose tissue across metabolic states

- □□□□□□: 38 / 60
- **AML** □□□□: 12 / 30
- □□□□□□□: 6 / 10
- □□□□: 48 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□

—

4 Catalytic Proximal Protein Oligomerization as an Anti-Tumor Strategy Targeting WDR5

- □□□□□□□: 26 / 60
- **AML** □□□□□: 19 / 30
- □□□□□□□□: 5 / 10
- □□□□□: 38 / 100
- □□□□□: 2026-03-13
- **DOI**: N/A
- □□□: □□□□□

□□□□□: □□□□□□□□

modifies histone proteins H2A and H4. However, the molecular details of co-translational NAA40 activity have remained elusive. Here, we show biochemically and by cryo-EM how NAA40 activity is…

—

7 Structure of a pH-sensitive pentameric ligand-gated ion channel from the Sarcoptes scabies mite

- □□□□□□: 33 / 60
- **AML** □□□□: 4 / 30
- □□□□□□□: 6 / 10
- □□□□: 35 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□□□□□□□□□+□□□□□□□□□□□□ AML □□□□□□□□□□□□

□□: Scabies is a skin infestation caused by the mite Sarcoptes scabiei and represents a substantial global health burden exacerbated by emerging resistance to ivermectin. An anionic pentameric ligand-gated ion channel from the mite, SsCl, shows pH-sensitivity and is significantly modulated by ivermectin. Here, we use cryo-EM and electrophysiology to explore the pH-sensing mechanisms of SsCl and the impact of ivermectin on channel activity. Structures of SsCl were resolved in closed (pH 6.5) and…

—

8 Vitamin C inhibits ACSL4 to alleviate ferro-aging in primates

- □□□□□□: 29 / 60
- **AML** □□□□: 10 / 30
- □□□□□□□: 5 / 10
- □□□□: 35 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□“ferro-aging”□□□□□□□□□□□□□□□□□□□

parasitic lifestyle can explain the variability in architectural evolutionary rates, and attempt to answer multiple other evolutionary questions. Double-stranded (genes on both strands)…

—

10 DEUP1 functions as a scaffold for basal foot integrity and planar polarity in multiciliated cells

- □□□□□□: 32 / 60
- **AML** □□□□: 6 / 30
- □□□□□□□: 5 / 10
- □□□□: 34 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□ DEUP1 □□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□

the need for refined interventions. Here, we exploit SnCs' lipid metabolic signature to develop a senotherapeutic strategy. Given the universal lipid accumulation in SnCs and the dual role of lipids as both metabolic liabilities and essential lubricants, we engineer an injectable…

—

13 The COP1-ADA2b module mediates light regulation of DNA double-strand break repair in Arabidopsis

- □□□□□□: 35 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 31 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□/□□□□□□□□□□□□□□□□□

□□: DNA double-strand break (DSB) represents the most severe form of DNA damage, and both defective and hyperactive repair can compromise genome stability. It is known that blue light promotes DSB repair by cryptochromes (CRYs)-enhanced STRUCTURAL MAINTENANCE OF CHROMOSOME 5/6 (SMC5/6) complex recruitment via ALTERATION/DEFICIENCY IN ACTIVATION 2B (ADA2b). Here, we report a negative regulatory module consisting of CONSTITUTIVE PHOTOMORPHOGENIC 1 (COP1), an E3 ubiquitin ligase acting as a central…

—

14 Risk of infection shortly before and after a diagnosis of liver cirrhosis: a nationwide, population-based study in Sweden

- □□□□□□: 35 / 60
- **AML** □□□□: 3 / 30
- □□□□□□□: 5 / 10
- □□□□: 31 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

□□: Copy number variants (CNVs) have been shown to play a significant role in the pathogenesis of various human diseases. Although several tools have been developed for detecting CNVs based on whole-exome sequencing (WES) data, their performance remains suboptimal for small exon-level CNVs (exCNVs). This is primarily due to multiple technical variabilities, including probe capture efficiency, mappability, exon size, batch effects, experimental background noise bias, and control sample selection, all…

—

16 BPX-Net: biomarker-preserved explainable networks for disease diagnosis and prognosis

- □□□□□□: 16 / 60
- **AML** □□□□: 9 / 30
- □□□□□□□: 3 / 10
- □□□□: 28 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

TRAMFIX (TRavelling Across Melbourne for FIXel-based analysis): a multisite dataset of traveling participants (n = 10 healthy adults) scanned across four 3T MRI scanners using a…

—

19 Cortical-limbic circuit dynamics of approach-avoidance conflict in humans

- □□□□□□: 31 / 60
- **AML** □□□□: 2 / 30
- □□□□□□□: 4 / 10
- □□□□: 27 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□ LSC□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Choosing to approach or avoid is common in everyday life and excessive avoidance is a cardinal feature of anxiety disorders. We use intracranial EEG to define a prefrontal-limbic circuit supporting approach and avoidance. Presurgical epilepsy patients (n = 20) performed an approach-avoidance conflict decision-making task inspired by the arcade game Pac-Man, where patients trade off rewards against losses from ghost attack. During approach, theta power increases across a limbic circuit including…

—

20 Pan-tumor activity of olomorasib, a next-generation KRAS G12C inhibitor in KRAS G12C-mutant advanced solid tumors: a first-in-human study

- □□□□□□: 29 / 60
- **AML** □□□□: 5 / 30
- □□□□□□□: 5 / 10
- □□□□: 26 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□ I □□□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□□□□

European coverage contributed by 26 biobanks from 12 countries. This retrospective, multi-center study contains structured and curated clinical data, supporting research on biomarkers for early detection, prognosis, and treatment. A phenotypical/clinical data model has been defined and individual-level…

—

22 Unveiling the hidden regulators: how post-translational modifications influence the progression and treatment of hepatocellular carcinoma

- □□□□□□: 20 / 60
- **AML** □□□□: 11 / 30
- □□□□□□□: 6 / 10
- □□□□: 25 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□ HCC □ PTMs □□□□□□□□ AML □□□□□□□□□□□□□□□□□□

□□: Hepatocellular carcinoma (HCC) is a common and aggressive primary liver cancer. Due to its high incidence and fatality rates, it poses a serious threat to global public health. Protein post-translational modifications (PTMs) are crucial regulatory mechanisms that occur after translation and fine-tune cellular functions. Common PTM types-including phosphorylation, ubiquitination, acetylation, methylation, glycosylation, ubiquitin-like modifications (such as UFMylation and SUMOylation), and…

—

23 Quantitative corrosion framework for anti-corrosive passivation design to extend calendar life in lithium metal batteries

- □□□□□□: 30 / 60
- **AML** □□□□: 2 / 30
- □□□□□□□: 5 / 10
- □□□□: 25 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□ LSC□□□□□□□□□□□□□□□□□□□

- □□□□: 23 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□/□□□□□□□□ RCT □□□□□□□□□ StresSOS □“□□□”□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□

- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□

- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□□□□□□□□□□□□□□□□

- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML[illegible]LSC[illegible]

[illegible]: Large animal models, while valuable, are expensive, time-consuming, and limited to discrete interventional or terminal timepoints, while existing benchtop models do not offer an accurate representation of the esophageal environment. Moreover, current pre-clinical models cannot effectively simulate swallowing dysfunction (dysphagia), restricting progress in understanding motility disorders like achalasia and hindering evidence-based dietary recommendations. In response, we present RoboGullet, a…

—

37 The Development and Acceptability of a Psychology-Based Intervention for Debilitating Symptom Complexes Attributed to Ticks

- [illegible]: 23 / 60
- **AML** [illegible]: 2 / 30
- [illegible]: 4 / 10
- [illegible]: 21 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML[illegible]/[illegible]

[illegible]: CONCLUSIONS: This is a novel and theoretically driven psychotherapeutic intervention for DSCATT, co-produced with patient involvement across four phases. Pilot testing suggested the manualised intervention was feasible and acceptable, supporting future evaluation of feasibility and treatment outcomes with randomised controlled trials.

—

38 Effect of Fresh versus Cryopreserved Grafts on Febrile Neutropenia and Infections in Allogeneic Hematopoietic Cell Transplantation: Factors Determining Mortality

- [illegible]: 17 / 60
- **AML** [illegible]: 12 / 30
- [illegible]: 4 / 10
- [illegible]: 20 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]FN [illegible] HCT-CI [illegible]

[illegible]: CONCLUSIONS: This study demonstrated that, regardless of graft type, prolonged FN duration and a high HCT-CI score are the primary determinants of mortality. Therefore, clinical management of patients should also address these risk factors.

—

39 Continuous assessment of daily-living gait using self-supervised learning of wrist-worn accelerometer data

- [illegible]: 24 / 60
- **AML** [illegible]: 3 / 30

- □□□□□□□: 5 / 10
- □□□□: 20 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML□□□□□□□□□□□□□□□□□□□□□□□□/□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□

□□: Physical activity and mobility are critical for healthy aging and predict diverse health outcomes. While wrist-worn accelerometers are widely used to monitor physical activity, estimating gait metrics from wrist data remains challenging. We extend ElderNet, a self-supervised deep-learning model previously validated for walking-bout detection, to estimate gait metrics from wrist accelerometry. Validation involved 819 older adults (Rush-Memory-and-Aging-Project) and 85 individuals with gait…

—

40 Hospital readmission in children on maintenance dialysis: a multicentre, prospective cohort study

- □□□□□□□: 24 / 60
- **AML □□□□□**: 3 / 30
- □□□□□□□: 5 / 10
- □□□□□: 20 / 100
- □□□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□□

□□□□□: □□□□□□□□□□□□□□□□□□□□

- [redacted]: 5 / 10
- [redacted]: 20 / 100
- [redacted]: 2026-03-12
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]

[redacted]: CONCLUSIONS: These findings show potential evidence of altered WM pathways in children at-risk for mental illness, particularly in emotion regulation related pathways. Additionally, parenting behaviour and stress were associated with WM microstructure in children, underscoring potential influence of early environmental factors on brain development. Longitudinal and replication studies are needed to clarify causal mechanisms and long-term effects.

—

43 Cohort profile: The DIabetes and ST-segment Elevation Myocardial Infarction (DISTEMI) Study

- [redacted]: 20 / 60
- **AML** [redacted]: 5 / 30
- [redacted]: 6 / 10
- [redacted]: 19 / 100
- [redacted]: 2026-03-13
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]

[redacted]: CONCLUSION: The DISTEMI study integrates comprehensive cardiometabolic phenomic with multiomic profiling to identify cardiometabolic STEMI subtypes and predictors of outcomes, and to improve precision risk stratification and targeted prevention.

—

44 Working conditions, job stress and work-related consequences among hospital employees-differences by professional group, working hours and job levels: A cross-sectional study

- [redacted]: 23 / 60
- **AML** [redacted]: 3 / 30
- [redacted]: 5 / 10
- [redacted]: 19 / 100
- [redacted]: 2026-03-12
- **DOI**: N/A
- [redacted]: [redacted]

[redacted]: [redacted] AML [redacted]

[redacted]: CONCLUSIONS: In conclusion, this study provides a comprehensive view of the differing work stressors and consequences faced by hospital staff based on their roles, work hours, and job levels. These insights emphasize the importance of tailoring interventions to target specific groups within hospitals to enhance occupational health and create supportive work environments.

—

45 Chiral orbital lasing in a twisted bilayer metasurface

- [redacted]: 23 / 60
- **AML** [redacted]: 1 / 30
- [redacted]: 6 / 10

- [illegible]: 18 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML[illegible]

[illegible]: Chirality is a fundamental concept in physics, core to many phenomena in nonlinear optics, quantum physics and topological photonics. Photons are intrinsically chiral when carrying spin angular momentum, whereas orbital angular momentum can induce chirality when photons interact with structures with broken mirror symmetry. Here, we observe orbital chiral lasing from a twisted bilayer photonic structure, by leveraging its inherent structural chirality. We design and fabricate a Moiré-type optical…

—

46 Thermal detection of single photons using Dirac fermions

- [illegible]: 25 / 60
- **AML** [illegible]: 1 / 30
- [illegible]: 5 / 10
- [illegible]: 17 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]/[illegible]

[illegible]: Detecting single photons is a crucial process in quantum science, quantum networking, biology, and advanced imaging. To detect the small quantum of energy carried in a photon, conventional mechanisms rely on energy excitation across either a semiconductor bandgap or superconducting gap that hinders their applications to low-energy photons. Here, we detect single near-infrared photons using the thermal properties of Dirac fermions in graphene. By exploiting the extremely low heat capacity of…

—

47 Unintended Consequences of COVID-19 Public Health and Social Measures in Camps and Camp-Like Settings: A Systematic Review and Conceptual Analysis

- [illegible]: 22 / 60
- **AML** [illegible]: 3 / 30
- [illegible]: 5 / 10
- [illegible]: 16 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]/[illegible]LSC[illegible]

[illegible]: CONCLUSION: This study reveals the complex interactions between PHSM and UIC and their broad mostly negative effects on marginalised populations. To reduce UIC in future health emergencies, they must be considered in pandemic planning with all stakeholders. Trust-building should be central in health interventions and PHSM design for more effective and equitable responses.

—

48 Outcomes of SARS-CoV-2 infection post-solid organ transplantation in the cystic fibrosis population

- [illegible]: 19 / 60
- **AML** [illegible]: 4 / 30
- [illegible]: 5 / 10

- □□□□: 16 / 100
- □□□□: 2026-03-12
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□ CF □□□□□□□□□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□

- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□□□□□□□□□□□□ AI □□□□□□□□□□□ AML □□□□□□□□ LSC□□□□□□□□□□□□□□□□□

□□□□: □□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□□ AML □□□□□LSC□□□□□□□□□□□□□□□□□□□

□□: Humic acid (HA), a complex organic compound of natural matter, is harmless on its own but can form carcinogenic disinfection byproducts, such as trihalomethanes and haloacetic acids, during water chlorination and disinfection processes, posing serious risks to aquatic organisms and human health. We present a B-site dual-metal-doping strategy to fabricate ferromagnetic perovskite-type adsorbents for rapid and efficient HA degradation. Perovskite oxides, specifically LaFeO(3), were selected for…

—

58 No such thing as a shark? Genomes shake up ocean predator's family tree

- □□□□□□: 13 / 60
- **AML** □□□□: 2 / 30
- □□□□□□□: 3 / 10
- □□□□: 4 / 100
- □□□□: 2026-03-13
- **DOI**: N/A
- □□: □□□□

□□□□: □□□□□□□

- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible] AML [illegible]

[illegible]: No abstract

—

62 Daily briefing: ‘Virtual cell’ simulates nearly every chemical reaction in the real thing

- [illegible]: 6 / 60
- **AML** [illegible]: 6 / 30
- [illegible]: 4 / 10
- [illegible]: 1 / 100
- [illegible]: 2026-03-13
- **DOI**: N/A
- [illegible]: [illegible]

[illegible]: [illegible]

[illegible]: No abstract

—