追觅 - 2026-03-25

生成时间: 2026-03-25 07:13

分析文章数量: 26

1 An expanded reference catalog of translated open reading frames for biomedical research

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 16 / 30
- **表达与可复用性**: 6 / 10
- **最终总分**: 46 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从资源建设角度具有潜在基础设施价值，但就当前提供的有限摘要信息，证据链、构建标准与验证强度无法评估，且对 AML 的直接可迁移性尚不明确。；置信度：低

摘要: Non-canonical (i.e. unannotated) open reading frames (ncORFs) have until recently been omitted from reference genome annotations, despite evidence of their translation, limiting their incorporation into biomedical research. To address this, in 2022, we initiated the TransCODE consortium and built the first community-driven consensus catalog of human ncORFs, which was openly distributed to the research community via Ensembl-GENCODE. While this catalog represented a starting point for reference…

—

2 Same-Slide Spatial Multi-Omics Integration with IN-DEPTH Reveals Tumor Virus-Linked Spatial Reorganization of the Tumor Microenvironment

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 42 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从题目与摘要片段看方法学目标明确且可能有通用价值，但当前提供信息不足以判断其关键技术指标、基准验证与结论强度，难以支持较高评分。；置信度：低

摘要: Spatial transcriptomics and proteomics have enabled profound insights into tissue organization, yet these technologies remain largely disparate, and emerging same-slide multi-omics approaches are limited in plex, spatial resolution, signal retention, and integrative analytics. We introduce IN-situ DEtailed Phenotyping To High-resolution transcriptomics (IN-DEPTH), a streamlined, resource-efficient, commercially compatible workflow using single-cell spatial proteomics-derived imaging to guide…

—

3 Mitochondrial metabolic imbalance drives diploidization in mouse haploid embryonic stem cells via NADPH overload

- **研究本身价值**: 30 / 60
- **AML 借鉴价值**: 9 / 30

- **表达与可复用性**: 4 / 10
- **最终总分**: 39 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就其原领域（单倍体 ESC 稳定性）可能有意思，但基于当前仅有的截断摘要，证据链无法核验；且与 AML 核心问题距离较远，AML 借鉴价值有限。；置信度：低

摘要: A hallmark of mammals is a diploid genome. Despite constraints from dosage compensation and imprinting, haploid embryonic stem cells can be established. However, rapid diploidization is observed in such cultures from mice, rats, and humans, limiting their use and indicating counterselection of a haploid genome. Here, we use metabolic profiling to discover that diploidization is triggered by an imbalance that arises from a smaller cytoplasmic volume and increased mitochondrial density. Reduced…

—

4 RNA functional modulation by Mitoxantrone via RNA structural ensemble repartitioning

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 13 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 38 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；概念上有潜在新意，但就当前仅有的题目+不完整摘要而言，证据链、适用边界与对 AML 的可迁移性均无法被充分评估，需非常保守对待。；置信度：低

摘要: Targeting RNA with small molecules offers a strategy to modulate gene expression at undruggable targets. Traditional screens favor thermodynamically stable, low-entropy RNA motifs with defined conformations, yet these provide limited energetic leverage for functional modulation. Many RNAs instead sample dynamic structural ensembles that small molecules can repartition. Using group I self-splicing introns as a model, we identified the antineoplastic drug Mitoxantrone as a competitive inhibitor of…

—

5 Asymmetric histone inheritance regulates olfactory stem cell fates during regeneration

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 38 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；问题本身有新意，但在仅有题目与截断摘要的情况下，关键证据链、因果实验与可复现信息均无法核验，对 AML 的可迁移性目前只能停留在“概念启发”层面。；置信度：低

摘要: The olfactory epithelium possesses an adult stem cell population, the horizontal basal cells (HBCs), to permit lifelong tissue regeneration. Here, we show that HBCs exhibit asymmetric inheritance of histone H4 but not H2A-H2B during olfactory epithelium regeneration in mice. Primary HBC cultures further revealed asymmetric histone inheritance for H3 and H3.3. Upon mitotic exit, asymmetric histone inheritance correlates with asynchronous transcription re-initiation and differential enrichment of…

—

6 Enhanced Remission and Survival Outcomes with Decitabine Plus Venetoclax in Additional Sex Comb Like 1 Mutated Acute Myeloid Leukemia

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 14 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 34 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就“特定分子亚型的临床疗效信号”可能有参考意义，但对 AML 基础机制与可迁移生物学结论的支撑非常有限，且当前摘要信息不足以判断关键方法学质量。；置信度：低

摘要: Decitabine (DEC) and Venetoclax (VEN) are approved for elderly adult acute myeloid leukemia (AML) patients with an additional sex comb-like 1 (ASXL1) mutation who cannot tolerate intensive chemotherapy. However, direct comparative data in this population remain limited. A systematic review and meta-analysis were conducted to assess the indirect efficacy of DEC alone and VEN in older AML patients with ASXL1 mutations. A matched cohort was created by comparing outcomes of consecutive adults with…

—

7 Maturase K forms a plastidial splicing complex with a neofunctionalized branching enzyme

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 33 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就原研究领域（植物质体 RNA 剪接）可能有机制意义，但与 AML 基础研究关联极弱，且仅凭当前不完整摘要无法评估关键证据链强度。；置信度：低

摘要: Chloroplast group IIA introns originate from bacterial ribozymes. Their splicing requires the splicing factor Maturase K (MatK). MatK, however, has been difficult to functionally analyze, as it appears essential for plant viability and is encoded in the chloroplast genome. Here we identified a heteromultimeric complex comprising MatK and three other essential, plastid-targeted proteins using co-immunoprecipitation experiments in Arabidopsis and tobacco. Among the MatK interactors is a conserved…

—

8 Prediction of the individual risk for the development of brain metastases in patients with non-oncogene-addicted non-small cell lung cancer: Real-world data from the German prospective CRISP registry (AIO-TRK-0315)

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 30 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；作为 NSCLC 随访管理的风险分层工具有一定应用意义，但当前提供信息不足以核查模型严谨性；且与 AML 基础研究关联极弱，借鉴价值主要停留在“风险建模思路”层面。；置信度：低

摘要: Brain metastases (BM) from non-small cell lung cancer (NSCLC) are a relevant cause of morbidity and mortality. This study aims to identify a population at high risk for BM in order to inform personalized follow-up strategies in the future. We have therefore developed and validated a point-based risk classifier for the risk of BM1 in patients with metastatic non-oncogene-addicted NSCLC without BM at disease onset. A total of 3139 patients were included from the German CRISP lung cancer registry…

—

9 Dual Regulation of Post-Translational Modification of Proteins: Bottlenecks and Breakthroughs in NK Cell Therapy for Glioblastoma

- **研究本身价值**: 21 / 60
- **AML 借鉴价值**: 9 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 29 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就现有题目与摘要片段看，属于领域综述性梳理，对 AML 只能提供“概念层面启发”，缺乏可直接迁移的证据与方法闭环。；置信度：低

摘要: Glioblastoma (GBM) is the most malignant primary central nervous system tumor in adults, with strong invasiveness, high recurrence, and poor prognosis. Natural killer (NK) cells, innate immune cells that eliminate glioma stem cells without MHC matching, show promise for GBM immunotherapy, but their efficacy is limited by GBM’s immunosuppressive tumor microenvironment (TME), especially via protein post-translational modifications (PTMs). This review summarizes seven key PTMs’ (phosphorylation,…

—

10 MADRe: Strain-level Metagenomic Classification Through Assembly-Driven Database Reduction

- **研究本身价值**: 23 / 60
- **AML 借鉴价值**: 10 / 30
- **表达与可复用性**: 3 / 10

- **最终总分**: 29 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从提供的题目与摘要片段看是偏工程与算法整合的宏基因组方法学工作，但关键信据（基准测试、外部验证、误差来源分析等）在当前文本中缺失，且与 AML 基础研究关联度弱。；置信度：低

摘要: Strain-level metagenomic classification is essential for understanding microbial diversity and functional potential, yet remains challenging, particularly when sample composition is unknown and reference databases are large and redundant. Here we present MADRe, a modular and scalable pipeline for long-read strain-level metagenomic classification based on Metagenome Assembly-Driven Database Reduction. Beyond system-level integration, MADRe introduces statistical strategies that leverage…

—

11 Efficient Pt₁Ni single-atom alloy catalyst for hydrogen-free catalytic fractionation of lignocellulose

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 28 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：**不推荐**；就催化/生物质精炼方向看起来是“性能导向的材料催化工作”，但与 AML 基础研究几乎无直接可迁移链条，作为 AML 文献不建议投入精读。；置信度：低

摘要: The production of monophenols from lignocellulose with cellulose intact without exogenous hydrogen via self-hydrogen supplied fractionation is promising, yet requires high metal loadings and yields saturated products. Herein, we report a single-atom alloyed Pt(1)Ni catalyst that achieves 50.9 wt% yield of phenolic monomers from birch sawdust with about 50% selectivity to valuable propenyl side-chained products under mild conditions (140 °C, 1 atm N(2)), while preserving cellulose intact….

—

12 Unveiling the biodiversity of large DNA viruses in intertidal mudflats via metagenomics

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 27 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就原研究领域（环境病毒组/宏基因组）可能是较大规模资源工作，但就“AML 基础研究可迁移价值”目前文本证据极弱，难支撑任何与 AML 直接相关的结论或启发。；置信度：低

摘要: Large DNA viruses (LDVs) are unique members of the Earth’s virosphere, remarkable for their extra-large genome sizes and broad metabolic potential. However, our knowledge of this viral group remains very limited, particularly in complex dynamic habitats. In this study, 237 metagenome-assembled LDV genomes are comprehensively recovered from intertidal mudflats using multiple sampling and sequencing strategies totaling 5.3 TB data. A phylogenetically distinct subgroup within Imitervirales is…

—

13 Intrinsically chiral exciton polaritons in an atomically-thin semiconductor

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就光子学方向可能有一定新意，但基于当前摘要信息证据链无法核查；对 AML 基础研究的直接借鉴价值极低。；置信度：低

摘要: Photonic bound states in the continuum (BICs) have emerged as a versatile tool for enhancing light-matter interactions by strongly confining light fields. Chiral BICs are photonic resonances with a high degree of circular polarisation, which hold great promise for spin-selective applications in quantum optics and nanophotonics. Here, we demonstrate a novel application of a chiral BIC for inducing strong coupling between the circularly polarised photons and spin-polarised (valley) excitons (bound…

—

14 Governing principles of hydration of mixed proton conducting Co-based double perovskites

- **研究本身价值**: 28 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 26 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就摘要可见其更偏材料体系的规律总结，但与 AML 基础研究几乎无直接关联，且仅凭当前截断摘要无法核验其“主导原则”的证据链强度。；置信度：低

摘要: Proton ceramic electrochemical cells PCECs hold promise for efficient, sustainable production and use of hydrogen. The positive electrodes are mixed proton conducting perovskites that facilitate water splitting and oxygen reduction, but the factors that determine the protonation are poorly understood. Here, we establish the governing principles of protonation through a study of hydration of 45 double perovskites with the general formula AIAIICo2O6-δ, having Ba or Ba+Sr on A^(I) and a mix of rare…

—

15 Multi-layered marble for hydrogel encapsulation

- **研究本身价值**: 23 / 60

- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 4 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从现有摘要信息看是界面材料学问题，潜在工程意义存在，但与 AML 基础研究关联极弱，且证据链与数据细节在当前文本中几乎无法评估。；置信度：低

摘要: Hydrophobic skins on fruits play a crucial role in preventing dehydration and shielding against environmental stress. Inspired by this natural structure, hydrophobic encapsulation is widely employed to preserve the functionality and long-term stability of hydrogels. However, instability at the interface between hydrophobic shell and hydrogel remains a critical challenge. Here, we present a hydrogel encapsulation strategy to stabilize the incompatible interface using a concept of liquid marble,…

—

16 From Breakthroughs to Blueprints: Evolving Evidence and Future Directions in Relapsed and Refractory Large B-Cell Lymphoma

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 7 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 24 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就 LBCL 临床证据与策略框架可能有参考价值，但从目前仅有的题目与摘要片段看，缺乏可审查的数据与方法细节，且与 AML 基础研究的可迁移性总体偏弱。；置信度：低

摘要: The therapeutic landscape for relapsed or refractory large B-cell lymphoma (R/R LBCL) has undergone rapid and profound change, driven by cellular therapies, bispecific antibodies, and next-generation antibody-drug conjugates (ADCs). These advances have redefined historical standards while exposing persistent gaps in trial design, biological insight, and therapeutic sequencing. Recent randomized studies show that ADC- and bispecific-anchored regimens can outperform legacy chemotherapy…

—

17 How strong is a mitochondrial targeting signal?

- **研究本身价值**: 18 / 60
- **AML 借鉴价值**: 6 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；作为线粒体蛋白导入领域的概念性/方法学亮点提示有一定价值，但就“研究证据链”与“AML 可迁移性”而言，基于当前文本几乎无法支撑强结论或直接借鉴到 AML 机制研究。；置信度：低

摘要: In this issue, Yan et al. show that mitochondrial targeting signals (presequences) vary widely in import strength. Using the quantitative MitoLuc and PotLuc assays, they dissect multiple parameters of protein import and reveal how presequence features influence mitochondrial targeting efficiency and stress sensitivity.

—

18 Negative CO₂ emissions for long-term mitigation of extremes in land hydrological cycle

- **研究本身价值**: 27 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就其本领域（气候水文）可能是重要问题，但当前提供文本极少、方法与证据链无法核验；且与 AML 基础研究几乎无直接可迁移关联。；置信度：低

摘要: Global warming has profound effects on the terrestrial hydrological cycle, leading to alterations in regional extreme weather patterns. While terrestrial precipitation responses under continued greenhouse gas emissions are well established, the responses of terrestrial precipitation and vegetation feedbacks under climate mitigation scenarios remain uncertain. Here, we investigate terrestrial precipitation changes under idealized negative and zero CO(2) emissions scenarios using the Community…

—

19 AI-based hematological malignancy prediction from peripheral blood smears in a large diagnostic laboratory cohort

- **研究本身价值**: 17 / 60
- **AML 借鉴价值**: 11 / 30
- **表达与可复用性**: 1 / 10
- **最终总分**: 23 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；仅凭标题无法评估其模型性能、证据链与可复现性；对 AML 的可迁移价值“可能存在但未被证明”。；置信度：低

摘要: No abstract

—

20 Medium-scale integrated circuits based on p-type 2D semiconducting MoTe₂

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 3 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就器件/材料领域看可能有一定工程推进意义，但与 AML 基础研究几乎无直接关联，且当前仅凭不完整摘要不足以支撑对证据链与可重复性的高可信评价。；置信度：低

摘要: Two-dimensional (2D) semiconductors hold promise for next-generation electronics, yet the lack of scalable p-type counterparts remains a major bottleneck, with prior studies largely limited to discrete devices or simple circuits. Here we report the realization of medium-scale integrated circuits (MSICs) based on wafer-scale p-type 2D semiconductors, enabled by the controlled synthesis of uniform 4-inch 2H-MoTe(2) films. A precursor-engineering strategy that integrates thickness-tunable Mo…

—

21 Mid-infrared detection through ligand-driven local heating in lanthanide-doped nanoparticles

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 4 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 22 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；从现有摘要片段看更偏工程方法学创新，但当前文本不足以评估证据链强度；且与 AML 基础研究直接关联极弱。；置信度：低

摘要: Mid-infrared photothermal detection, leveraging well-established visible/near-infrared detection techniques, holds significant potential for a wide range of scientific and industrial applications. Conventional mid-infrared photothermal methods show limited response at weak mid-infrared radiation due to insufficient thermal sensitivity. Here, we overcome this challenge by employing ligand-capped lanthanide nanoparticles, where organic molecules with strong and broadband mid-infrared absorption…

—

22 The expanding role of mitochondria-lipid droplet contacts in liver and their disruption by MASLD

- **研究本身价值**: 12 / 60
- **AML 借鉴价值**: 8 / 30
- **表达与可复用性**: 1 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；仅凭标题无法核验证据链与结论边界，现阶段只能将其视为“代谢细胞器接触”方向的背景线索，对 AML 的可迁移价值尚未被文本证明。；置信度：低

摘要: No abstract

—

23 Selective conversion of syngas to C₄₊ long-chain alcohols

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 2 / 10

- **最终总分**: 21 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就现有摘要信息看，这是偏工业催化的研究，证据链与可复现细节未呈现，且与 AML 基础研究几乎无直接可迁移关联。；置信度：低

摘要: The selective conversion of syngas to C(4+) long-chain alcohols holds significant industrial and scientific interest, but challenges in product selectivity and process efficiency remain. Here, we report a precisely catalytic strategy for C(4+) alcohol synthesis with a selectivity of 80% at 17% CO conversion. The reaction channel involves: (i) the development Cs(2)O-Co(2)C-Co catalysts, capable of catalyzing CO hydrogenation to long-chain oxygenates/olefins; and (ii) complete conversion to C(4+)…

—

24 Evidence from Buhais Rockshelter for human settlement in Arabia between 60,000 and 16,000 years ago

- **研究本身价值**: 25 / 60
- **AML 借鉴价值**: 2 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 21 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就“AML 基础研究借鉴”而言几乎不相关；就“研究本身”而言仅凭当前摘要片段无法核验其关键证据链与方法细节，无法支持高可信度评价。；置信度：低

摘要: Several significant milestones in human evolution date to the period between 70,000 and 12,000 years ago, including the replacement of archaic humans, the global dispersal of Homo sapiens and the introduction of Upper Palaeolithic traditions. The Arabian Peninsula provides only sparse records illuminating this period. We introduce here the Buhais Rockshelter archaeological sequence and paleoenvironmental records from the Faya Palaeolandscape in the Emirate of Sharjah (UAE). Buhais Rockshelter…

—

25 Engineered microbial hydrogels with confined architecture and binary microbes for efficient hydrogen production

- **研究本身价值**: 20 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-24
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就摘要有限信息看属于工程优化型研究，但证据链细节无法核查；且与 AML 基础研究直接关联极弱，最多借鉴“限域共培养水凝胶”这一方法学思路。；置信度：低

摘要: Renewable energy sources, particularly hydrogen, offer a promising solution to address global energy crisis and carbon emissions. Microalgae-driven hydrogen production has attracted immense interest in both scientific and industrial fields. However, challenges

such as high oxygen sensitivity, substantial water demand, and low hydrogen production efficiency limit their potential. Here, we develop a core-shell symbiotic hydrogel system for enhanced hydrogen production via leveraging coaxial 3D…

—

26 The effect of Polyethylene Glycol on constipation prophylaxis after benign gynaecological laparoscopy: A randomized controlled trial

- **研究本身价值**: 24 / 60
- **AML 借鉴价值**: 3 / 30
- **表达与可复用性**: 2 / 10
- **最终总分**: 20 / 100
- **发布日期**: 2026-03-23
- **DOI**: N/A
- **链接**: 查看原文

审稿结论: 推荐等级：不推荐；就其自身临床问题而言属于常规 RCT 结论较克制，但与 AML 基础研究几乎无直接可迁移关系。；置信度：低

摘要: CONCLUSION: Short-term use of PEG 3350 following benign gynecological laparoscopy did not significantly reduce overall constipation rates at one week, but was well tolerated and associated with improvements in specific constipation symptoms.

—