

HIGHER DERIVATIVES OF THE GRAM FUNCTION

MARIUS DUNDULIS, RAMŪNAS GARUNKŠTIS, ANTANAS LAURINČIKAS,
AND RAIVYDAS ŠIMĖNAS

ABSTRACT. We find the asymptotics of higher-order derivatives of the Gram function t_u . We also prove the uniform distribution and continuous uniform distribution modulo one of t_u^n for $n = 1, 2, \dots$ and make a conjecture that the Riemann zeta function shifted by powers of the Gram points has discrete universality. We use AI to prove our results in Lean.

1. INTRODUCTION AND MAIN RESULTS

The concept of Gram points was introduced by the Danish mathematician Gram. They are points on the critical line where the Riemann-Siegel theta function $\vartheta(t)$ is a multiple of π . Using a heuristic argument, Gram [3] reasoned that there is precisely one zero of the Riemann zeta function between two neighboring Gram points. Later on, Hutchinson [4] found counterexamples to this conjecture. Nevertheless, the Gram points remain an active area of research.

The Riemann-Siegel theta function $\vartheta(t)$ gives us an increment of the continuous branch of the argument of $\pi^{-s/2}\Gamma(s/2)$ as s goes along the straight line segment of the complex plane joining $s = 1/2$ and $s = 1/2 + it$; note that $\vartheta(t)$ is increasing for $t \geq 7$ (Ivić [5, Section 1.3] and Korolev [10, p. 3]). The function ϑ appears in the symmetric functional equation involving the Riemann zeta function ζ

$$e^{i\vartheta(t)}\zeta(1/2 + it) = e^{-i\vartheta(t)}\zeta(1/2 - it).$$

The function ϑ is related to the function S . The function $S(t)$ is defined as

$$S(t) = \frac{1}{\pi} \arg \zeta(1/2 + it).$$

Here, the argument is taken along the continuous branch of $\arg \zeta(s)$ as we move from $s = 2$ to $s = 2 + it$ and $s = 1/2 + it$. If s is a zero of ζ , then we set

$$S(t) = \lim_{\delta \rightarrow +0} \frac{1}{2} (S(t + \delta) + S(t - \delta)).$$

The Riemann-von Mangoldt formula (see, e.g. Edwards [1, Section 6.6], Ivić [5, formula (1.45)], Karatsuba and Korolev [6]) leads to the following relation between the functions ϑ and S

$$(1) \quad N(t) = \frac{1}{\pi} \vartheta(t) + 1 + S(t).$$

Here $N(t)$ is the number of ζ zeros in the rectangle $0 < \Im s \leq t$, $0 \leq \Re s \leq 1$, with the exception that at points of discontinuity, which coincide with the ordinates of

the zeros γ_n , it is defined as the limit

$$N(t) = \frac{1}{2} \lim_{h \rightarrow 0} (N(t+h) + N(t-h)).$$

The function S is related to the function N by the Riemann-von Mangoldt formula.

$$N(t) = \frac{t}{2\pi} \log \frac{t}{2\pi} - \frac{t}{2\pi} + \frac{7}{8} + S(t) + \frac{1}{\pi} \delta(t)$$

(see Karatsuba and Korolev, [6, Theorem 1]).

From the above equation, the function S can be expressed as

$$(2) \quad S(t) = -\frac{t}{2\pi} \log \frac{t}{2\pi} + \frac{t}{2\pi} - \frac{7}{8} - \frac{1}{\pi} \delta(t) + N(\gamma + 0)$$

with $\gamma < t < \gamma'$, where γ and γ' are neighboring ordinates of the zeros of ζ (see Karatsuba and Korolev [6, Proof of Theorem 2]). The function δ is

$$(3) \quad \delta(t) = \frac{t}{4} \log \left(1 + \frac{1}{4t^2} \right) + \frac{1}{4} \arctan \frac{1}{2t} - \frac{t}{2} \int_0^\infty \frac{\rho(u) du}{(u + 1/4)^2 + (t/2)^2}$$

and $\rho(u) = 1/2 - \{u\}$ (see Karatsuba and Korolev [6, Theorem 1]).

The first derivative of S is

$$(4) \quad S'(t) = -\frac{1}{2\pi} \log \frac{t}{2\pi} + O(t^{-2}), \quad t \rightarrow +\infty$$

except for its points of discontinuity (see Karatsuba and Korolev [6, Theorem 2]).

We give our theorem about the n -th derivative of the S function.

Theorem 1. *The n -th derivative of the S function when $n \geq 2$ is*

$$(5) \quad S^{(n)}(t) = (-1)^{n-1} \frac{(n-2)!}{2\pi} t^{1-n} + O(t^{-n-1}), \quad t \rightarrow +\infty.$$

except for its discontinuity points.

From the above theorem, equation (1), equation (4), and the fact that ϑ is differentiable infinitely many times, we get

Corollary 2. *Suppose $n \geq 2$. Then, the n -th derivative of the Riemann-Siegel theta function is*

$$(6) \quad \vartheta^{(n)}(t) = (-1)^n \frac{(n-2)!}{2} t^{1-n} + O(t^{-n-1}), \quad t \rightarrow +\infty.$$

It is known that for $t \geq 7$, the Riemann-Siegel theta function $\vartheta(t)$ is monotonically increasing. Thus for $u \geq \vartheta(7)/\pi + \pi$, we can implicitly define the Gram function t_u by

$$(7) \quad \vartheta(t_u) = (u-1)\pi,$$

with $t_u \geq 7$. Note that for positive integers k , t_k correspond to Gram points (see Edwards [1, pp. 125–226]).

The asymptotic expression for the Gram function is (see Lavrik [13, Lemma 2])

$$(8) \quad t_u = \frac{2\pi u}{\log u} \left(1 + (1 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty.$$

Korolev [10, Lemma 1.1] gave the asymptotics for the first two derivatives of the Gram function. The first derivative of the Gram function is

$$(9) \quad \frac{dt_u}{du} = \frac{2\pi}{\log u} \left(1 + (1 + o(1)) \frac{\log \log u}{\log u} \right),$$

as $u \rightarrow +\infty$.

We calculate the n -th derivative of the Gram function t_u , which we denote by $t_u^{(n)}$.

Theorem 3. *The n -th derivative of the Gram function is asymptotically*

$$t_u^{(n)} = \frac{(-1)^{n+1} 2\pi(n-2)!}{u^{n-1} \log^2 u} \left(1 + (2 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty,$$

for $n = 2, \dots$

Here, we corrected the constant in [10, Lemma 1.1] for $n = 2$.

The proof of the following theorem of the uniform distribution modulo one of the sequence $\{t_k^n\}_{k=1}^\infty$ is analogous to Pańkowski [15, Proof of Theorem 1].

Theorem 4. *The sequence $\{t_k^n\}_{k=1}^\infty$ is uniformly distributed modulo one for $n = 1, 2, 3, \dots$*

We have the following corollary about the continuous uniform distribution of the n -th degree of the Gram function. Laurinćikas proved this result [12, Lemma 5.7] by a different method. In our case, it follows directly from Theorem 4.

Corollary 5. *The function t_u^n , $n = 1, 2, 3, \dots$, is continuously uniformly distributed modulo one.*

The function t_u can be applied in the universality theory of zeta functions. As an example, we will consider the Riemann zeta function. Recall that the universality of $\zeta(s)$ in the Voronin sense means that a broad class of analytic functions can be approximated by shifts $\zeta(s + iu)$, $u \in \mathbb{R}$. This approximation property of $\zeta(s)$ was discovered by Voronin [18] and has various theoretical and practical applications. For results, see Matsumoto [14].

Now we discuss the discrete universality for the Riemann zeta function using its shifts by the degrees of the Gram points $\zeta(s + it_k^n)$. Korolev and Laurinćikas have extensively studied this problem. In [7], they proved it for the degree 1. They proved the result for short intervals in [8]. The continuous version of joint universality was proved in [9].

By $\#A$ we denote the cardinality of the set A . We assume about the second moment of the Riemann zeta function shifted by the Gram function: For all $\epsilon > 0$ and sufficiently large y and N , we have

$$(10) \quad \frac{1}{N} \sum_{k=2}^N |\zeta(s + it_k^n) - \zeta_{\{p:p \leq y\}}(s + it_k^n)|^2 < \epsilon$$

with $s \in K$ where K is any compact set with connected complement in the right part of the critical strip. In addition, $\zeta_{\{p:p \leq y\}}(s) := \prod_{p \leq y} (1 - 1/p^s)^{-1}$ is a truncated Euler product for the Riemann zeta function, where p is prime. Pańkowski [15, p. 192, Proposition 1] incorrectly thought that this follows from Gallagher's lemma applied to Dirichlet L -functions with shifts $\alpha t^a (\log t)^b$ for $\alpha \neq 0$, $a > 0$, and b real numbers. Sourmelidis [17] partially remedied this error. The proof of universality follows from our assumption (10) and Theorem 4 using the method by Pańkowski [15, pp. 192–194, proof of Theorem 2]. We formulate our statement as follows: *Suppose $K \subset \{s \in \mathbb{C} : 1/2 < \sigma < 1\}$ is a compact set with connected complement, f is a continuous non-vanishing function on K , analytic inside of K . Then, for*

any $\epsilon > 0$,

$$\liminf_{N \rightarrow \infty} \frac{1}{N} \# \left\{ 2 \leq k \leq N : \max_{s \in K} |\zeta(s + it_k^n) - f(s)| < \epsilon \right\} > 0.$$

In the last section of the article, we discuss the proof of our results using Lean. In doing so, we employed the AI models Claude Opus 4.7, Opus 4.8, and Fable 5. Our results can be found at <https://github.com/raivydasiminas/gram-derivatives.git>. Our axiomatization is minimal, consisting of the results in the theory of zeta functions and uniform distribution.

2. PROOFS

Proof of Theorem 1. Our goal is to compute the n -th derivative of $S(t)$. Recall the formula (2). We split the computation into two parts.

First, let us compute

$$\left(-\frac{t}{2\pi} \log \frac{t}{2\pi} + \frac{t}{2\pi} - \frac{7}{8} \right)^{(n)} = \left(-\frac{1}{2\pi t} \right)^{(n-2)} = (-1)^{n-1} \frac{(n-2)!}{2\pi t^{n-1}}.$$

It remains to estimate the error term arising from differentiating the function δ . The terms

$$(11) \quad \frac{t}{4} \log \left(1 + \frac{1}{4t^2} \right) + \frac{1}{4} \arctan \frac{1}{2t}$$

can be expanded into

$$\frac{1}{16t} + \frac{1}{8t} + \sum_{n=3}^{\infty} \frac{a_n}{t^n}$$

with real coefficients a_n . Differentiating the above expression n times term-by-term yields

$$(12) \quad (-1)^n \frac{3n!}{16t^{n+1}} + O(t^{-n-3}), \quad t \rightarrow +\infty.$$

For our purposes, the accuracy $O(t^{-n-1})$ is sufficient.

Continuing our computation of the n -th derivative of $\delta(t)$, let us denote

$$j(t) := \int_0^\infty \frac{\rho(u) du}{(u + 1/4)^2 + (t/2)^2},$$

$$\sigma(u) := \int_0^u \rho(z) dz.$$

We compute the n -th derivative of j . We follow with integration by parts

$$j(t) = \int_0^\infty \frac{d\sigma(u)}{(u + 1/4)^2 + (t/2)^2} = 2 \int_0^\infty \frac{\sigma(u)(u + 1/4) du}{((u + 1/4)^2 + (t/2)^2)^2}.$$

Note that $0 \leq \sigma(u) \leq 1/8$.

We have to differentiate the function $j(t)$ with respect to t n times, which amounts to differentiation under the integral sign. We use the Leibniz rule

$$\frac{d}{dt} \int_a^\infty f(u, t) du = \int_a^\infty \frac{\partial}{\partial t} f(u, t) du.$$

The following conditions must be satisfied: $f(u, t)$ and $\partial f / \partial t$ be continuous for $u \geq a$, and there exists an integrable $g(u)$ such that $|\partial f(u, t) / \partial t| \leq g(u)$ (see Rudin [16, pp. 180–182]). Evidently, they hold in our case.

Applying the Leibniz rule yields

$$\begin{aligned} \frac{d^n}{dt^n} \int_0^\infty \frac{(u+1/4)\sigma(u) du}{((u+1/4)^2 + (t/2)^2)^2} &= \int_0^\infty \frac{\partial^n}{\partial t^n} \frac{(u+1/4)\sigma(u) du}{((u+1/4)^2 + (t/2)^2)^2} \\ &\ll \int_0^\infty \frac{t^n(4u+1)\sigma(u) du}{((4u+1)^2 + 4t^2)^{n+2}}, \quad t \rightarrow +\infty. \end{aligned}$$

Let us split the above integral into two

$$\int_0^t \frac{t^n(4u+1)\sigma(u) du}{((4u+1)^2 + 4t^2)^{n+2}} + \int_t^\infty \frac{t^n(4u+1)\sigma(u) du}{((4u+1)^2 + 4t^2)^{n+2}}.$$

Let us integrate with respect to u using the facts that $u < t$ in the first integral and $u > t$ in the second. Combining this with the bounds $0 \leq \sigma(u) \leq 1/8$, gives the estimate

$$\begin{aligned} j^{(n)}(t) &\ll \int_0^t \frac{t^n(4u+1)\sigma(u) du}{((4u+1)^2 + 4u^2)^{n+2}} + \int_t^\infty \frac{u^n(4u+1)\sigma(u) du}{(4u+1)^{2n+4}} \\ &\ll t^{-n-2}, \quad t \rightarrow +\infty. \end{aligned}$$

We need to find the n -th derivative of the last term of the function $\delta(t)$ (see (3))

$$-\frac{t}{2} \int_0^\infty \frac{\rho(u) du}{(u+1/4)^2 + (t/2)^2} = -\frac{t}{2} j(t).$$

We apply the Leibniz rule to differentiate products. If u and v are n times differentiable functions of x , then (see Gradshteyn and Ryzhik [2, Formula 0.42])

$$\frac{d^n(uv)}{dx^n} = u \frac{d^n v}{dx^n} + \binom{n}{1} \frac{du}{dx} \frac{d^{n-1} v}{dx^{n-1}} + \binom{n}{2} \frac{d^2 u}{dx^2} \frac{d^{n-2} v}{dx^{n-2}} + \dots + \frac{d^n u}{dx^n} v.$$

We get

$$-\frac{1}{2} \left(t j^{(n)}(t) + \binom{n}{1} j^{(n-1)}(t) \right) = O(t^{-n-1}).$$

Collecting this and (12) completes the proof of Theorem 1. $\square$

Proof of Theorem 3. We prove this by induction. Let us make the inductive hypothesis

$$(13) \quad t_u^{(k)} = \frac{(-1)^{k+1} 2\pi(k-2)!}{u^{k-1} \log^2 u} \left(1 + (2 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty.$$

First, we prove that the theorem statement holds for $k = 2$. Differentiating the equation (7) gives

$$\vartheta''(t_u) t_u'^2 + \vartheta'(t_u) t_u'' = 0.$$

We calculate the term $\vartheta''(t_u) t_u'^2$ using Corollary 2, equations (9) and (8)

$$\begin{aligned} \vartheta''(t_u) t_u'^2 &= \left(\frac{1}{2t_u} + O(t_u^{-3}) \right) \left(\frac{2\pi}{\log u} \left(1 + (1 + o(1)) \frac{\log \log u}{\log u} \right) \right)^2 \\ &= \frac{2\pi}{2u \log u} \left(1 + (1 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty. \end{aligned}$$

Differentiating (7) and using the expression (9) yields

$$\frac{1}{\vartheta'(t_u)} = \frac{2}{\log u} \left(1 + (1 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty.$$

Combining the above two results gives

$$t_u'' = -\frac{2\pi}{u \log^2 u} \left(1 + (2 + o(1)) \frac{\log \log u}{\log u} \right), \quad u \rightarrow +\infty.$$

Suppose our hypothesis (13) holds for $2 \leq k \leq n-1$. We will show that it does for $k = n$. By Faà di Bruno's formula (see Gradshteyn and Ryzhik [2, formula (0.430)]), the n -th derivative of a composite function $h(x) = g(f(x))$ is

$$(14) \quad \frac{d^n}{dx^n} g(f(x)) = \sum \frac{n!}{m_1! 1!^{m_1} m_2! 2!^{m_2} \dots m_n! n!^{m_n}} g^{(m_1+m_2+\dots+m_n)}(f(x)) \\ \times \prod_{j=1}^n (f^{(j)}(x))^{m_j},$$

here, the summation is taken over all partitions

$$(15) \quad m_1 + 2m_2 + \dots + nm_n = n.$$

We will need the shorthand notation $\sum m = m_1 + m_2 + \dots + m_n$ for our purposes.

We plug $g(s) = \vartheta(s)$, $f(u) = t_u$ into Faà di Bruno's formula. Then, the relation (7) yields that the $d^n \vartheta(t_u)/du^n$ equals 0 and the formula (14) becomes

$$(16) \quad 0 = \vartheta'(t_u) t_u^{(n)} + \dots + \vartheta^{(n)}(t_u) (t_u')^n.$$

The first member in the above equation corresponds to the case $m_n = 1$ and $m_i = 0$ for $i = 1, \dots, n-1$. The last member - to the case $m_1 = n$ and $m_i = 0$ for $i = 2, \dots, n$.

In the summation of Faà di Bruno's formula, the first term in the above equation (16) is the only one containing $t_u^{(n)}$, and the last one is asymptotically the largest of the remaining ones as $u \rightarrow \infty$. Let us prove this fact. We will show that $\vartheta^{(n)}(t_u) (t_u')^n$ is asymptotically larger than the terms obtained using Faà di Bruno's formula, except possibly the first one. Hence, we can ignore the constants. Any term is uniquely determined by the values $m_1, m_2, \dots, m_n$, which come from the partition (15). For this calculation, we use the asymptotics for the derivatives of $\vartheta(t)$ (Corollary 2), the asymptotics for t_u' and t_u , and our inductive step (13). We skip the term $\vartheta'(t_u) t_u^{(n)}$. In all the remaining terms, we have $m_n = 0$. The arbitrary term in Faà de Bruno's formula, except for the partition $m_n = 1, m_i = 0, i = 1, \dots, n-1$, is

$$(17) \quad \vartheta^{(m_1+m_2+\dots+m_{n-1})}(t_u) \prod_{j=1}^{n-1} (t_u^{(j)})^{m_j} = \Theta \left(\frac{1}{t_u^{\sum m-1}} \cdot \frac{1}{\log^{m_1} u} \prod_{j=2}^{n-1} \left(\frac{1}{u^{j-1} \log^2 u} \right)^{m_j} \right) \\ = \Theta \left(\frac{\log^{\sum m-1} u}{u^{\sum m-1} \log^{m_1} u} \cdot \frac{1}{u^{\sum_{j=2}^{n-1} j m_j - \sum m + m_1} \log^2 \sum m - 2m_1 u} \right) \\ = \Theta \left(\frac{1}{u^{n-1} \log^{\sum m - m_1 + 1} u} \right).$$

We say that a function $f(x)$ is $\Theta(g(x))$ if there exist positive constants c_1, c_2 and a real number x_0 such that $c_1 g(x) \leq f(x) \leq c_2 g(x)$, $x \geq x_0$. In the last step above, we used the equation (15). The last expression shows that among all the terms in Faà de Bruno's formula, except $\vartheta'(t_u) t_u^{(n)}$, the asymptotically largest one

is $\vartheta^{(n)}(t_u)(t'_u)^n$, corresponding to the partition $m_1 = n$ and $m_i = 0$, $i = 2, \dots, n$. This, together with (16), leads to

$$\vartheta'(t_u)t_u^{(n)} = -\vartheta^{(n)}(t_u)(t'_u)^n + o\left(\vartheta^{(n)}(t_u)(t'_u)^n\right) \quad (u \rightarrow \infty).$$

$$\vartheta'(t_u)t_u^{(n)} = -\vartheta^{(n)}(t_u)(t'_u)^n + O\left(\frac{1}{u^{n-1} \log^2 u}\right) \quad (u \rightarrow \infty),$$

It follows

$$t_u^{(n)} = \frac{-\vartheta^{(n)}(t_u)(t'_u)^n}{\vartheta'(t_u)} + O\left(\frac{1}{\vartheta'(t_u)u^{n-1} \log^2 u}\right).$$

Corollary 2 together with the asymptotics for t'_u and t_u leads to

$$t_u^{(n)} = \frac{(-1)^{n+1} 2\pi(n-2)!}{u^{n-1} \log^2 u} \left(1 + (2 + o(1)) \frac{\log \log u}{\log u}\right).$$

This completes the induction and yields the theorem. $\square$

Proof of Theorem 4. We have theorem [11, Chapter 1, Theorem 3.5] stating: if there exists $l \in \mathbb{N}$, the function $f(x)$ is defined for $x \geq 1$, the derivative $f^{(l)}(x)$ tends monotonically to 0 as $x \rightarrow \infty$, and if $x|f^{(l)}(x)| \rightarrow \infty$, $x \rightarrow \infty$, then the sequence $\{f(n)\}_{n \in \mathbb{N}}$ is uniformly distributed modulo one.

Consider the n -th degree of the Gram function t_u^n . By Faà di Bruno's formula, we have

$$(18) \quad \frac{d^n t_u^n}{du^n} = \sum \frac{n!}{m_1! 1!^{m_1} m_2! 2!^{m_2} \dots m_n! n!^{m_n}} \frac{d^{m_1+m_2+\dots+m_n} t_u^n}{dt_u^{m_1+m_2+\dots+m_n}} \prod_{j=1}^n (t_u^{(j)})^{m_j},$$

where we sum over all partitions

$$(19) \quad m_1 + 2m_2 + \dots + nm_n = n.$$

Let us denote $\sum m := m_1 + m_2 + \dots + m_n$. An arbitrary term on the right-hand side of the equation (18) depends on the value of the vector $(m_1, m_2, \dots, m_n)$. Ignoring the constant, recalling the asymptotic of t_u given in the formula (8), and the asymptotic of its derivative in Theorem 3 yields the following asymptotic of an arbitrary term in the formula (18)

$$\begin{aligned} & \frac{u^{n-\sum m}}{\log^{n-\sum m} u} \cdot \frac{1}{\log^{m_1} u} \prod_{j=2}^n \frac{1}{(u^{j-1} \log^2 u)^{m_j}} = \frac{u^{n-\sum m}}{\log^{n-\sum m} u} \cdot \frac{1}{u^{n-\sum m} \log^{-m_1+2\sum m} u} \\ & = \frac{1}{\log^{n-m_1+\sum m} u}, \end{aligned}$$

where we used the formula (19). This value is maximized when we minimize $-m_1 + \sum m$, which happens when $m_1 = n$ and $m_2 = \dots = m_n = 0$.

To prove monotonicity, we calculate the $n+1$ -st derivative of the Gram function. Again, we use Faà di Bruno's formula. Summing over partitions, we have

$$m'_1 + 2m'_2 + \dots + nm'_n + (n+1)m'_{n+1} = n+1,$$

where the integers $m'_1, m'_2, \dots, m'_{n+1}$ appear in our formula now. Let us denote $\sum m' := m'_1 + 2m'_2 + \dots + m'_{n+1}$. It follows

$$\begin{aligned} & \frac{u^{n-\sum m'}}{\log^{n-\sum m'} u} \cdot \frac{1}{\log^{m'_1} u} \prod_{j=2}^{n+1} \frac{1}{(u^{j-1} \log^2 u)^{m'_j}} = \frac{u^{n-\sum m'}}{\log^{n-\sum m'} u} \cdot \frac{1}{u^{n+1-\sum m'} \log^{-m'_1+2\sum m'} u} \\ &= \frac{1}{u \log^{n-m'_1+\sum m'} u} \end{aligned}$$

Again, this value is maximized by setting $m'_1 = n+1, m'_2 = m'_3 = \dots = m'_{n+1} = 0$.

All in all, we have

$$\frac{d^n t_u^n}{du^n} = \Theta\left(\frac{1}{\log^n u}\right)$$

and

$$\frac{d^{n+1} t_u^n}{du^{n+1}} = \Theta\left(\frac{1}{u \log^n u}\right).$$

Based on the continuity of the $n+1$ -st derivative, there exists some value u_0 for which $d^{n+1} t_u^n / du^{n+1}$ does not change its sign provided $u > u_0$. Hence, the n -th derivative of t_u^n will monotonically approach 0 for $u > u_0$. In addition, $u |d^n t_u^n / du^n|$ will monotonically approach $+\infty$. Therefore, by the criterion stated at the beginning of the proof, the sequence $\{t_k^n\}_{k=1}^\infty$ is uniformly distributed modulo one. $\square$

Proof of Corollary 5. By Kuipers and Niederreiter [11, Chapter 1, Theorem 9.6], the function $f(t), t \geq 0$, is continuously uniformly distributed modulo one if 1) the sequence $f(k+t), k = 0, 1, \dots$, is uniformly distributed modulo one for almost all $t \in [0, 1)$ and 2) the sequence $f(kt), k = 1, 2, \dots$, is uniformly distributed modulo one for almost all $t \in [0, 1)$. If we take f to be the n -th degree of the Gram function, both of these facts follow from the fact that there exists real number u_0 such that $d^n t_u^n / du^n$ monotonically approaches zero for $u > u_0$, $u |d^n t_u^n / du^n|$ monotonically approaches $+\infty$ for $u > u_0$, and the criterion for the uniform distribution modulo 1 of sequences, formulated at the beginning of the proof of Theorem 4. $\square$

3. USING AI AND LEAN

Lean has emerged as the definitive programming language for proving mathematical theorems. We used it to formalize our results. To write the code, we employed Claude Opus 4.7, Opus 4.8, and Fable 5 LLMs. The overall amount of code is substantial, more than 10000 lines. It is available at <https://github.com/raivydasiminas/gram-derivatives.git>. The proofs of our theorems and corollaries are located in the folder `GramDerivatives`. There is one file per theorem or corollary. The file `Axioms.md` in the top directory describes the axioms used to prove our theorems. The project builds cleanly, with no "sorry" statements.

Here, for the reader's convenience, we provide the formalizations of our theorems and corollaries in Lean

- Theorem 1

```
theorem theorem1 (n : Nat) (hn : 2 <= n) :
  IsO
    (fun t =>
      iteratedDeriv n S t
      - ((-1 : Real) ^ (n - 1) * (n - 2).factorial
        * (1 / (2 * Real.pi)) * t ^ (1 - (n : Real))))
```



```
(fun t => t ^ (-(n : Real) - 1))
N^\infty_0
```

The predicate `is0(f g)` means that $f = O(g)$. The function `iteratedDeriv n S t` is $S^{(n)}(t)$. The notation `N^\infty_0` essentially means that $f(x) = O(g(x))$, $n \rightarrow \infty$. In Theorem 1, our result does not depend on the analytic nature of the piecewise constant function $N(t)$. The important fact is that $N(t)$ is constant everywhere except at a countable set of jumps. We formalized this in an assumption.

- Corollary 2 is an immediate consequence of Theorem 1. We formalize it as `theorem corollary2 (n : Nat) (hn : 2 <= n) :`

```
Is0
(fun t =>
  iteratedDeriv n theta t
  - ((-1 : Real) ^ n * (n - 2).factorial
    * (1 / 2) * t ^ (1 - (n : Real))))
(fun t => t ^ (-(n : Real) - 1))
N^\infty_0
```

- Theorem 3

```
theorem theorem3 (n : Nat) (hn : 2 <= n) :
```

```
Iso
(fun u : Real =>
  iteratedDeriv n gram u
  - gramLeading n u
  - 2 * gramLeading n u * Real.log (Real.log u) / Real.log u)
(fun u : Real => gramLeading n u * Real.log (Real.log u) / Real.log u)
N^\infty_0
```

For this theorem, we axiomatically define the Gram function as `gram` and its leading term as `gramLeading`.

- Theorem 4

```
theorem theorem4 (n : Nat) (hn : 1 <= n) :
```

```
Gram.UD.IsUDModOne (fun k : Nat => gramPow n (k : Real))
```

Here, we axiomatize the theory of uniform distribution, as it is not yet available in Mathlib.

- Corollary 5

```
lemma corollary5 (n : Nat) (hn : 1 <= n) :
```

```
Gram.UD.IsCUDModOne (fun u : Real => Gram.Theorem4.gramPow n u)
```

Here we axiomatized the theorem about continuous uniform distribution modulo one.

Acknowledgments. This project has received funding from the European Social Fund (project No 09.3.3-LMT-K-712-19-0030) under a grant agreement with the Research Council of Lithuania (LMTLT).

REFERENCES

- [1] H. M. Edwards. *Riemann's zeta function*. Academic Press, New York, 1974. 1, 1
- [2] I. S. Gradshteyn and I. M. Ryzhik. *Table of integrals, series, and products*. Elsevier, 2007. 2, 2
- [3] J.-P. Gram. Sur les zéros de la fonction $\zeta(s)$ de Riemann. *Acta Math.*, (27):289–304, 1903. 1
- [4] J. I. Hutchinson. On the roots of the Riemann zeta-function. *Trans. Amer. Math. Soc.*, 27:49–60, 1925. 1

- [5] A. Ivić. *The theory of Hardy's Z-function*, volume 196 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2013. 1
- [6] A. A. Karatsuba and M. A. Korolev. The argument of the Riemann zeta function. *Russian Math. Surveys*, 60(3):433–488, 2005. 1, 1, 1, 1, 1
- [7] M. Korolev and A. Laurinčikas. A new application of the Gram points. *Aequat. Math.*, 93:859–873, 2019. 1
- [8] M. Korolev and A. Laurinčikas. A new application of the Gram points. II. *Aequat. Math.*, 94:1171–1187, 2020. 1
- [9] M. Korolev and A. Laurinčikas. Joint approximation of analytic functions by shifts of the Riemann zeta-function twisted by the Gram function. *Carpathian J. Math.*, 39(1):175–187, 2023. 1
- [10] M. A. Korolev. Gram's law in the theory of the Riemann zeta-function. Part 1. *Proc. Steklov Inst. Math.*, 292(2):1–146, 2016. 1, 1, 1
- [11] L. Kuipers and H. Niederreiter. *Uniform distribution of sequences*. Courier Corporation, 2012. 2, 2
- [12] A. Laurinčikas. On the extension of the Voronin universality theorem for the Riemann zeta-function. *Quaest. Math.*, 47(4):735–750, 2024. 1
- [13] A. A. Lavrik. Titchmarsh's problem in the discrete theory of the Riemann zeta-function. *Tr. Mat. Inst. Steklova*, 207:197–230, 1994. 1
- [14] K. Matsumoto. A survey on the theory of universality for zeta and L -functions. In *Number theory*, volume 11 of *Ser. Number Theory Appl.*, pages 95–144. World Sci. Publ., Hackensack, NJ, 2015. 1
- [15] Ł. Pańkowski. Joint universality for dependent L -functions. *Ramanujan J.*, 45(1):181–195, 1 2018. 1, 1
- [16] W. Rudin. *Principles of Mathematical Analysis*. McGraw-Hill, New York, 3rd edition, 1976. 2
- [17] A. Sourmelidis. Estimates for weyl sums in the theory of discrete universality. *Ramanujan J.*, 57:1399–1421, 2022. 1
- [18] S. M. Voronin. Theorem on the "universality" of the Riemann zeta-function. *Izv. Akad. Nauk SSSR Ser. Mat.*, 39(3):475–486, 1975. 1

(Marius Dundulis) EUROMONITOR INTERNATIONAL, JOGAILOS G. 4, LT-01116 VILNIUS, LITHUANIA

Email address: marius.dundulis@euromonitor.com

(Ramūnas Garunkštis) INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND INFORMATICS, VILNIUS UNIVERSITY, NAUGARDUKO 24, 03225 VILNIUS, LITHUANIA

Email address: ramunas.garunkstis@mif.vu.lt

URL: <https://mif.vu.lt/~garunkstis/>

(Antanas Laurinčikas) INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND INFORMATICS, VILNIUS UNIVERSITY, NAUGARDUKO 24, 03225 VILNIUS, LITHUANIA

Email address: antanas.laurincikas@mif.vu.lt

(Raivydas Šimėnas) INSTITUTE OF MATHEMATICS, FACULTY OF MATHEMATICS AND INFORMATICS, VILNIUS UNIVERSITY, NAUGARDUKO 24, 03225 VILNIUS, LITHUANIA

Email address: raivydas.simenas@mif.vu.lt

URL: <https://web.vu.lt/mif/r.simenas/>