

Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

# Logistic Regression:

- Classification problem

Eq: Dataset be:

Study hour	Playhour	O/P
------------	----------	-----

2	8	fail
---	---	------


3	7	fail
---	---	------

6	3	pass
---	---	------

Outliers  $\rightarrow$  1                          4                          pass

Here, we cannot perform regression, we need

to perform classification into pass / fail.





Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /


Why we use logistic regression when we can solve classification problem using linear regression?

→ Due to outliers best fit line gets changed and results will be wrong.

⇒ We cannot remove outliers always.

⇒ Threshold can't be changed, once fixed.

→ In logistic, we squash ('cut') the line, we will not change the line.



$$h_{\theta}(x) = \theta_0 + \theta_1 x$$



Sigmoid function ⇒


output will range  
bet<sup>n</sup> 0 and 1.

Sigmoid Function:-

$$\text{Sigmoid function} = \frac{1}{1 + e^{-x}} \rightarrow \text{bet}^n 0 to 1$$

① Create a best fit line

② Squashing → sigmoid function



Mo Tu We Th Fr Sa Su


Memo No. \_\_\_\_\_

Date / /

## Linear Regression Cost function:

$$J(\theta_0, \theta_1) = \frac{1}{m} \sum_{i=1}^m \{h_{\theta}(x^i) - y^i\}^2$$

MSE



## Logistic Regression cost function

Steps:

1) Create a best fit line

2) Apply squashing using sigmoid function.


$$J(\theta_0, \theta_1) = \frac{1}{m} \sum_{i=1}^m \{h_{\theta}(x^i) - y^i\}^2$$

sigmoid function

$$h_{\theta}(x) = \sigma(\theta_0 + \theta_1 x)$$

$$\text{let } z = \theta_0 + \theta_1 x$$

$$\therefore h_{\theta}(x) = \sigma(z)$$



Mo Tu We Th Fr Sa Su


Memo No. \_\_\_\_\_

Date / /

$$h_{\theta}(x) = \frac{1}{1+e^{-\theta^T x}} \Rightarrow h_{\theta}(x) = \frac{1}{1+e^{-(\theta_0 + \theta_1 x)}}$$

$$\therefore h_{\theta}(x) = \frac{1}{1+e^{-(\theta_0 + \theta_1 x)}}$$

But after applying sigmoid function, cost function will become non convex function and have a chance to get local minima.




- change the cost function to solve the convexity problem.

Logloss function:

$$h_{\theta}(x) = \frac{1}{1+e^{-(\theta_0 + \theta_1 x)}}$$

$$\text{Cost Function} = \begin{cases} -\log(h_{\theta}(x)), & \text{if } y=1 \\ -\log(1-h_{\theta}(x)), & \text{if } y=0 \end{cases}$$

Convex function



Mo Tu We Th Fr Sa Su

This will never  
give local minima  
Memo No. \_\_\_\_\_  
Date / /

$$\text{cost}(h_\theta(x^i), y^i) = -y \log(h_\theta(x)) - (1-y) \log(1-h_\theta(x))$$

If  $y=1$ ,

$$\text{cost}(h_\theta(x^i), y^i) = -\log(h_\theta(x))$$

If  $y=0$

$$\text{cost}(h_\theta(x^i), y^i) = -\log(1-h_\theta(x))$$

Minimize cost function  $J(\theta_0, \theta_1)$  by changing

$\theta_0, \theta_1$ .

Convergence algorithm:

Repeat convergence

{

$$\theta_j: \theta_j - \alpha \frac{\partial J(\theta_0, \theta_1)}{\partial \theta_j}$$

}

for  $j=0$  and  $j=1$

By default take threshold = 0.5



Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

Using ROC and ROC, we can define threshold.

## Performance Matrices

⇒ Confusion Matrix

⇒ Accuracy

⇒ Precision

⇒ Recall

⇒ F-Beta Score

### dataset

	$f_1$	$f_2$	$d_P$	$\hat{y}$
	-	-	0	1
	-	-	1	1
	-	-	0	0
	-	-	1	1
	-	-	1	1
	-	-	0	1
	-	-	1	0



Memo No. \_\_\_\_\_

Date / /

## Confusion Matrix

	1	0	$\rightarrow y$
1	TP	FP	
0	FN	TN	
$\uparrow y$			

TP : True positive } correct  
TN : True negative } match

FP : False positive } wrong  
FN : False negative } match


Confusion matrix  
for the previous dataset:

	1	0	$\leftarrow \text{actual}(y)$
1	3	2	
0	1	1	

### Accuracy

Accuracy =  $\frac{\text{TP} + \text{TN}}{\text{TP} + \text{FP} + \text{FN} + \text{TN}}$  → total dataset.  
Correctly predicted

Accuracy for above dataset =  $\frac{3+1}{7} = 4/7 = 57\%$




Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

## Imbalance dataset

1000 datapoint (Binary classification)



dumb model → If our model predict 1 everytime, we still get 90% accuracy.

So, 90% accuracy don't ensure the model goodness. To overcome this problem we use Recall and Precision.


Precision      we have to reduce false positive

		0	1
0	FN	TN	
1	FP	TP	

precision =  $\frac{TP}{TP + FP}$

So, precision deals with correct true prediction among all true prediction.

Hence, precision help us to visualize the reliability of the machine learning model in classifying the model as "Positive."



Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

### Example:

1) Mail  $\rightarrow$  spam or not

TP: Actual  
↓  
spam

Predicted  
↓  
spam  $\Rightarrow$  good

FP: Actual  
↓  
ham

Predicted  
↓  
spam  $\Rightarrow$  bad.

It is true positive rate.

### Recall

Recall =  $\frac{TP}{TP + FN}$



Recall is calculated as the ratio bet'n the number of positive samples correctly classified as positive to the total number of positive samples.

$\uparrow$  the Recall

$\downarrow$   
more positive

sample detected.

We don't care about negative sample are correctly classified or not.



Mo Tu We Th Fr Sa Su

Memo No.

Date

## Difference bet<sup>n</sup> Precision and Recall in ML:

Precision	Recall
→ Helps to measure the ability to classify positive sample in the model. el.	It helps to measure how many positive sample were correctly classified by model

### Using Situation

- a) If there is a requirement of classifying all positive as well as negative sample as positive, whether they are classified correctly or incorrectly then use Precision.
- b) If our goal is detect only all positive sample, then use Recall.



Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

## F-beta Score

$$F\text{-beta score} = \frac{(1+\beta^2) \times \text{Precision} \times \text{Recall}}{\beta^2 \cdot \text{Precision} + \text{Recall}}$$

① If FP and FN are both important:

$$\beta = 1$$

$$F_1 \text{ score} = \frac{2 \times P \times R}{P + R}$$

② If FP is more important than FN:


$$\beta = 0.5$$

$$F_{0.5} = \frac{(1+0.25) \times P \times R}{0.25 \times P + R}$$

③ If FN >> FP, FP is less important than FN.

$$\beta = 2$$

$$F_2 \text{ score} = \frac{(1+4) \times P \times R}{4 \times P + R}$$



Mo Tu We Th Fr Sa Su

Memo No. \_\_\_\_\_

Date / /

## Issue with High Dimensionality:

- When a categorical variable has high number of labels, it leads to quasi complete separation
  - when the dependent variable separates an independent variable or combination of several independent variables to a certain degree.
- It can affect the convergence of the model and can lead to incorrect decision.

Solution :- Collapsing categories based on reduction in chi-square.