

Deaths: Final Data for 2021

by Sherry L. Murphy, B.S., Kenneth D. Kochanek, M.A., Jiaquan Xu, M.D., and Elizabeth Arias, Ph.D.

Abstract

Objectives—This report presents final 2021 data on U.S. deaths, death rates, life expectancy, infant and maternal mortality, and trends by selected characteristics such as age, sex, Hispanic origin and race, state of residence, and cause of death.

Methods—Information reported on death certificates is presented in descriptive tabulations. The original records are filed in state registration offices. Statistical information is compiled in a national database through the Vital Statistics Cooperative Program of the National Center for Health Statistics. Causes of death are processed according to the *International Classification of Diseases, 10th Revision*. Beginning in 2018, all states and the District of Columbia were using the 2003 revised certificate of death for the entire year, which includes the 1997 Office of Management and Budget revised standards for race. Data based on these revised standards are not completely comparable to previous years.

Results—In 2021, a total of 3,464,231 deaths were reported in the United States. The age-adjusted death rate was 879.7 deaths per 100,000 U.S. standard population, an increase of 5.3% from the 2020 rate. Life expectancy at birth was 76.4 years, a decrease of 0.6 year from 2020. Age-specific death rates increased from 2020 to 2021 for every age group. In 2021, 9 of the 10 leading causes of death remained the same as in 2020. Heart disease remained the top leading cause, followed by cancer and COVID-19. The infant mortality rate of 5.44 infant deaths per 1,000 live births in 2021 did not change significantly from the rate in 2020 (5.42).

Conclusions—In 2021, the age-adjusted death rate increased and life expectancy at birth decreased for the total, male, and female populations, primarily due to the influence of deaths from COVID-19.

Keywords: mortality • cause of death • life expectancy • National Vital Statistics System

Highlights

Mortality experience in 2021

- In 2021, a total of 3,464,231 resident deaths were registered in the United States, an increase of 80,502 deaths compared with 2020 (3,383,729). The 1-year increase in the number of deaths was primarily driven by the COVID-19 pandemic.
- The crude death rate was 1,043.8 deaths per 100,000 population. The age-adjusted death rate, which accounts for the aging of the population, was 879.7 deaths per 100,000 U.S. standard population.
- The age-adjusted death rate for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population (1,109.2) was 1.2 times greater than for the White non-Hispanic (subsequently, White) population (893.9).
- The age-adjusted death rate for the Black non-Hispanic (subsequently, Black) population (1,118.0) was 1.3 times greater than for the White population (893.9).
- The age-adjusted death rate for the White population (893.9) was 1.9 times greater than for the Asian non-Hispanic (subsequently, Asian) population (461.7) and 1.2 times greater than for the Hispanic population (724.7).
- Life expectancy at birth was 76.4 years.
- From 2020 to 2021, life expectancy at birth decreased by 0.7 year for males (from 74.2 to 73.5) and by 0.6 year for females (79.9 to 79.3).
- The 15 leading causes of death in 2021 were:
 1. Diseases of heart (heart disease)
 2. Malignant neoplasms (cancer)
 3. COVID-19
 4. Accidents (unintentional injuries)
 5. Cerebrovascular diseases (stroke)
 6. Chronic lower respiratory diseases
 7. Alzheimer disease
 8. Diabetes mellitus (diabetes)
 9. Chronic liver disease and cirrhosis

10. Nephritis, nephrotic syndrome and nephrosis (kidney disease)
 11. Intentional self-harm (suicide)
 12. Essential hypertension and hypertensive renal disease (hypertension)
 13. Influenza and pneumonia
 14. Septicemia
 15. Parkinson disease
- In 2021, the infant mortality rate (IMR) was 5.44 infant deaths per 1,000 live births.
 - The 10 leading causes of infant death were:
 1. Congenital malformations, deformations and chromosomal abnormalities (congenital malformations)
 2. Disorders related to short gestation and low birth weight, not elsewhere classified (low birth weight)
 3. Sudden infant death syndrome (SIDS)
 4. Accidents (unintentional injuries)
 5. Newborn affected by maternal complications of pregnancy (maternal complications)
 6. Newborn affected by complications of placenta, cord and membranes (cord and placental complications)
 7. Bacterial sepsis of newborn
 8. Respiratory distress of newborn
 9. Diseases of the circulatory system
 10. Intrauterine hypoxia and birth asphyxia

Comparison with previous year

- The age-adjusted death rate increased 5.3% from 835.4 per 100,000 standard population in 2020 to 879.7 in 2021.
- Life expectancy for the total population decreased 0.6 year from 77.0 in 2020 to 76.4 in 2021.
- Life expectancy for females (79.3) was 5.8 years higher than for males (73.5), an increase of 0.1 year from 2020.
- The difference in life expectancy between the Black and White populations decreased 0.4 year from 5.9 years in 2020 to 5.5 years in 2021.
- From 2020 to 2021, life expectancy decreased for American Indian and Alaska Native males (1.6 years), American Indian and Alaska Native females (1.5 years), White males (0.8 year), White females (0.6 year), Black females (0.4 year), Asian females (0.3 year), Hispanic females (0.2 year), and Black males (0.2 year).
- The 15 leading causes of death remained the same in 2021 as in 2020, although some causes changed ranks.
- Age-adjusted death rates increased significantly in 2021 from 2020 for 11 of the 15 leading causes of death: heart disease, cancer, COVID-19, unintentional injuries, stroke, diabetes, Chronic liver disease and cirrhosis, kidney disease, suicide, hypertension, and Septicemia. Significant decreases occurred in 2021 from 2020 for 3 of the 15 leading causes of death: Chronic lower respiratory diseases, Alzheimer disease, and Influenza and pneumonia.

- Age-adjusted death rates increased from 2020 to 2021 for drug-induced causes (13.9%) and for alcohol-induced causes (9.9%).
- The decrease in life expectancy at birth for the total population in 2021 was mainly due to increases in mortality from COVID-19, unintentional injuries, Chronic liver disease and cirrhosis, suicide, and homicide.
- Among external causes of injury death, unintentional poisoning has been the leading mechanism of injury mortality since 2011.
- IMR did not change significantly in 2021 from 2020.
- Nine of the 10 leading causes of infant death remained the same in 2021 as in 2020. Intrauterine hypoxia and birth asphyxia replaced Neonatal hemorrhage as the 10th leading cause of infant death.

Introduction

This report presents detailed 2021 data on deaths and death rates according to demographic and medical characteristics. These data provide information on mortality patterns among residents of the United States by such variables as age, sex, Hispanic origin and race, state of residence, and cause of death. Information on these mortality patterns is key to understanding changes in the health and well-being of the U.S. population (1). Companion reports present additional details on leading causes of death and life expectancy in the United States (2,3).

Cause-of-death statistics presented in this report are classified according to the *International Classification of Diseases, 10th Revision* (ICD-10) (4–6). Detail on cause-of-death classification is provided in Technical Notes of this report. The COVID-19 pandemic that began in 2020 continued to have a substantial impact on the mortality profile of the U.S. population for 2021.

Mortality data can be used to monitor and evaluate the health status of the United States in terms of current mortality levels and long-term mortality trends, and to identify segments of the U.S. population at greater risk of death from specific diseases and injuries. Differences in death rates among various demographic subpopulations, including racial and ethnic groups, may reflect subpopulation differences in factors such as socioeconomic status, access to medical care, and the prevalence of specific risk factors in a particular subpopulation.

The 2003 revision of the U.S. Standard Certificate of Death uses the revised 1997 Office of Management and Budget standards for the collection of race and Hispanic ethnicity (7,8). The 1997 standards allow individuals to report more than one race and increase the race categories from four to five by separating the Asian and Pacific Islander groups. Beginning with the 2018 data year, all 50 states and the District of Columbia reported deaths based on the 2003 revision for the entire year, so the revised standards became the official standards for presenting mortality data by race and ethnicity (9). The Hispanic category did not change, remaining consistent with reports before 2018.

The race and ethnicity categories in this report follow the 1997 OMB revised race and Hispanic-origin standards. These categories differ from the bridged-race categories used before

2018 (10). The new categories include Hispanic, single-race American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native), single-race Asian non-Hispanic (subsequently, Asian), single-race Black or African American non-Hispanic (subsequently, Black), single-race Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander), and single-race White non-Hispanic (subsequently, White). Because single-race data are not available for the entire United States before 2018, data by race for 2018 through 2021 are not completely comparable with bridged-race data used in earlier years, so comparisons should be made with this consideration (11).

In addition to the tabulations included in this report, more detailed analysis is possible by using the annual mortality public-use file. The data file may be downloaded from: https://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm (12). Data file documentation is available from: <https://www.cdc.gov/nchs/data/dvs/Multiple-Cause-Record-Layout-2021.pdf>. The public-use file does not include geographic detail, but a file with this information may be available upon request (13). Death data also may be accessed from the Centers for Disease Control and Prevention's (CDC) WONDER, a web-based system that makes the agency's information resources available to public health professionals and the general public (14).

Methods

Data in this report are based on information from all resident death certificates filed in the 50 states and the District of Columbia.

This report provides detailed death data in [Tables 1–25](#). Tables showing data by state also provide information for the Commonwealth of the Northern Mariana Islands (Northern Marianas), Guam, Puerto Rico, and U.S. Virgin Islands.

Mortality data on specific demographic and medical characteristics cover all 50 states and the District of Columbia. Measures of mortality in this report include the number of deaths; crude, age-specific, and age-adjusted death rates; infant, neonatal, postneonatal, and maternal mortality rates; life expectancy; and rate ratios. Changes in death rates in 2021 compared with 2020 and differences in death rates across demographic groups in 2021 were tested for statistical significance. Unless otherwise specified, reported differences are statistically significant. Additional information on these statistical methods, random variation and relative standard error, the computation of derived statistics and rates, population denominators, and the definition of terms are presented in Technical Notes.

In accordance with the revised standards issued by the Office of Management and Budget in 1997, the 2003 revision of the U.S. Standard Certificate of Death provided for the reporting of more than one race (multiple races) and increased the race categories from four to five by separating the Asian and Pacific Islander groups (7,8). Starting in 2018, all 50 states and the District of Columbia reported deaths using the 2003 revision for the entire year.

The race and Hispanic-origin groups in this report follow the 1997 standards and differ from the race categories used in

reports for data years before 2018 (8,10,15). The categories include Hispanic, single-race American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native), single-race Asian non-Hispanic (subsequently, Asian), single-race Black or African American non-Hispanic (subsequently, Black), single-race Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander), and single-race White non-Hispanic (subsequently, White). For brevity, text references to race refer to single race in this report. Because the number of deaths reported with more than one race in 2021 is relatively small (0.6%), these deaths are included in totals but are shown separately in only one report table ([Table 2](#)). Some comparisons between race and ethnicity groups in this report are limited to the following groups based on population size: American Indian and Alaska Native, Asian, Black, Hispanic, and White.

Data presented in this report by the revised Hispanic-origin and race categories for 2018–2021 are not completely comparable with data by bridged race shown in earlier reports, and comparisons should be made with this consideration. The Hispanic-origin category is a separate item on the death certificate and was not affected by the revised standards; as a result, data by Hispanic origin for 2021 and earlier years are comparable.

Death rates by race and ethnicity for the American Indian and Alaska Native, Asian, Hispanic, and Native Hawaiian or Other Pacific Islander populations are affected by inconsistencies in reporting Hispanic origin and race on death certificates as compared with censuses and surveys (16). Death rates for the American Indian and Alaska Native population are underestimated by about 33% due to misclassification (16). Death rates for the Asian and Hispanic populations are underestimated by about 3.0% (16). This should be considered when making rate comparisons across racial and ethnic groups. At this time, information about the prevalence of misclassification for the Native Hawaiian or Other Pacific Islander population is not available.

To maintain consistency with data reported by the jurisdictions and data in the mortality data file, numbers of deaths and death rates in this report are not adjusted for misclassification of race and ethnicity unless otherwise indicated. Specifically, [Tables 4 and 5](#) present life expectancies by Hispanic origin and race that are produced using methods based on death rates adjusted for Hispanic-origin and race misclassification on death certificates. For additional detail, see *Quality of race and Hispanic-origin data* in Technical Notes.

The population data used to calculate death rates for 2021 shown in this report are estimated as of July 1, 2021, based on the Blended Base produced by the U.S. Census Bureau (Technical Notes), and are available from the CDC WONDER website: <https://wonder.cdc.gov/single-race-population.html> (17).

Data presented in this report and other mortality tabulations are available from the National Vital Statistics System website: <https://www.cdc.gov/nchs/nvss/deaths.htm>. The availability of mortality microdata is described in Technical Notes.

Results and Discussion

Deaths and death rates

In 2021, a total of 3,464,231 resident deaths were registered in the United States—80,502 more deaths than in 2020. The crude death rate for 2021 (1,043.8 deaths per 100,000 population) was 1.6% higher than the 2020 rate (1,027.0) (Tables A, 1, 2, 6, 8, and 10). In 2021, 416,893 deaths were identified with COVID-19 as the underlying cause of death (Tables B, 7, and 9). COVID-19 was a contributing cause but was not considered the underlying cause of an additional 45,300 deaths (18).

The age-adjusted death rate in 2021 was 879.7 deaths per 100,000 U.S. standard population—5.3% higher than the rate of 835.4 in 2020 (Tables A and 1). The age-adjusted death rates increased for males (5.0%) and females (5.5%). Age-adjusted death rates should be viewed as relative indexes rather than as actual measures of mortality risk. They are constructs that show what the level of mortality would be if no changes occurred in the age composition of the population from year to year. (For a discussion of age-adjusted death rates, see Technical Notes.) Thus, age-adjusted death rates are better indicators than unadjusted (crude) death rates for examining changes in the risk of death over a period of time when the age distribution of the population is changing. Age-adjusted death rates are also better indicators of relative risk when comparing mortality across geographic areas or between sex or race and ethnicity

subgroups of the population that have different age distributions; see Technical Notes. Since 1980, the age-adjusted death rate decreased significantly every year except for 1983, 1985, 1988, 1993, 1999, 2005, 2010, 2013, 2015, 2017, 2020, and 2021 (Figure 1) (14).

Death rates by Hispanic origin and race

In 2021, age-adjusted death rates by ethnicity and race groups (Table 1) were:

- Hispanic population: 724.7 deaths per 100,000 U.S. standard population
- American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population: 1,109.2
- Asian non-Hispanic (subsequently, Asian) population: 461.7
- Black non-Hispanic (subsequently, Black) population: 1,118.0
- Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population: 924.3
- White non-Hispanic (subsequently, White) population: 893.9

In 2021, the age-adjusted death rate for the American Indian and Alaska Native population was 1.2 times that for the White population. The rate for the Black population was 1.3 times that for the White population. The rate for the White population was 1.9 times the rate for the Asian population and 1.2 times the rate for the Hispanic population (Table B).

Figure 1. Crude and age-adjusted death rates: United States, 1960–2021




Table A. Percent change in death rates and age-adjusted death rates in 2021 from 2020, by age, Hispanic origin and race, and sex: United States

[Based on death rates on an annual basis per 100,000 population and age-adjusted rates per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards; see Technical Notes. Data for some Hispanic-origin or race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes]

Age group (years)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All ages																					
Crude rate	1.6	2.5	0.6	1.1	0.6	1.5	8.2	8.3	8.0	-0.3	-0.8	0.2	-0.9	-1.2	-0.7	15.3	14.0	16.8	2.6	3.9	1.1
Age-adjusted rate	5.3	5.0	5.5	0.2	-2.1	2.2	7.0	6.4	7.4	0.9	-0.4	2.1	-0.1	-1.8	1.3	12.5	10.0	15.2	7.1	7.2	6.9
Younger than 1 year ⁴	6.6	5.2	8.3	8.2	5.2	11.9	3.8	-2.6	12.2	28.7	29.8	27.4	2.7	3.4	1.7	7.0	36.3	-25.2	6.3	5.0	7.9
1–4	10.1	5.9	16.2	10.3	1.0	22.0	-11.0	-2.2	-22.3	11.2	16.0	4.0	17.6	10.0	28.4	1.4	-44.9	129.1	6.3	5.2	7.9
5–14	4.4	1.9	8.0	5.9	6.2	6.6	-6.2	-15.5	4.6	9.0	18.5	-1.3	10.0	6.7	14.8	-27.5	-30.7	-19.0	0.8	-2.6	4.9
15–24	5.6	4.1	10.0	6.1	3.2	16.1	20.5	17.8	26.3	8.8	12.1	2.1	5.4	4.1	10.0	47.8	49.7	43.2	4.3	3.3	7.3
25–34	13.4	13.6	13.5	20.9	22.3	18.4	16.5	20.2	10.6	15.4	16.3	13.8	14.5	12.4	19.9	22.8	24.0	20.6	10.5	10.9	9.9
35–44	16.1	14.7	17.9	19.8	18.3	21.7	20.4	19.5	21.2	7.5	4.5	12.1	14.2	12.8	15.7	45.9	43.4	49.6	15.7	14.2	17.7
45–54	12.1	11.1	12.9	9.6	7.7	12.6	17.6	16.7	18.1	5.0	4.4	5.1	8.7	7.2	10.3	29.1	28.2	29.8	14.5	13.8	14.8
55–64	7.5	5.6	9.7	2.3	-0.6	6.6	13.1	14.0	11.1	-0.2	-2.9	3.1	1.9	-0.7	5.0	11.3	6.6	17.3	10.4	8.8	12.1
65–74	3.8	2.3	5.5	-3.1	-7.3	2.4	2.5	-3.3	9.7	-1.1	-4.2	2.7	-2.6	-5.7	0.7	7.6	6.5	8.9	6.3	5.4	7.0
75–84	2.4	1.6	3.2	-4.8	-7.2	-2.5	-3.1	-3.6	-2.8	0.6	-1.4	2.7	-5.0	-7.8	-2.5	2.5	-7.0	12.8	4.5	4.0	4.8
85 and older	3.5	5.1	2.4	-1.8	-3.8	-0.4	-1.5	-2.6	-0.9	-0.5	-0.3	-0.7	-4.3	-4.0	-4.5	2.6	2.9	2.4	5.3	7.3	4.0

¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes people of Hispanic origin of any race; see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Death rates for younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table B. Number of deaths, percentage of total deaths, death rate, and age-adjusted death rate for 2021, percent change in age-adjusted death rates in 2021 from 2020, and ratio of age-adjusted death rates. by sex and Hispanic origin and race for the 15 leading causes of death for the total population in 2021: United States

[Crude death rates on an annual basis per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards]

Rank ¹	Cause of death (based on ICD-10)	Number	Percent of total deaths, 2021 ²	Crude death rate, 2021	2021	Percent change from 2020 to 2021	Age-adjusted death rate					
							Ratio					
							Non-Hispanic, single race ³					
							Male to female	American Indian and Alaska Native to White	Black to White	Native Hawaiian or Other Pacific Islander to White	White to Asian	White ³ to Hispanic ⁴
...	All causes.	3,464,231	100.0	1,043.8	879.7	5.3	1.4	1.2	1.3	1.0	1.9	1.2
1	Diseases of heart. (I00-I09,I11,I13,I20-I51)	695,547	20.1	209.6	173.8	3.3	1.6	0.9	1.3	1.0	2.1	1.5
2	Malignant neoplasms. (C00-C97)	605,213	17.5	182.4	146.6	1.7	1.3	0.8	1.1	0.9	1.7	1.5
3	COVID-19. (U07.1)	416,893	12.0	125.6	104.1	22.5	1.6	2.0	1.5	2.0	1.5	0.6
4	Accidents (unintentional injuries). (V01-X59,Y85-Y86)	224,935	6.5	67.8	64.7	12.3	2.2	1.8	1.1	0.8	3.7	1.5
5	Cerebrovascular diseases. (I60-I69)	162,890	4.7	49.1	41.1	5.9	1.0	0.9	1.5	1.2	1.2	1.1
6	Chronic lower respiratory diseases. (J40-J47)	142,342	4.1	42.9	34.7	-4.7	1.2	0.8	0.7	0.5	4.4	2.7
7	Alzheimer disease. (G30)	119,399	3.4	36.0	31.0	-4.3	0.7	0.6	0.9	0.6	1.9	1.2
8	Diabetes mellitus. (E10-E14)	103,294	3.0	31.1	25.4	2.4	1.6	2.3	2.1	2.4	1.2	0.8
9	Chronic liver disease and cirrhosis. (K70,K73-K74)	56,585	1.6	17.0	14.5	9.0	1.8	5.1	0.7	0.6	3.6	0.9
10	Nephritis, nephrotic syndrome and nephrosis. (N00-N07,N17-N19,N25-N27)	54,358	1.6	16.4	13.6	7.1	1.4	1.3	2.2	1.7	1.5	1.0
11	Intentional self-harm (suicide). (*U03,X60-X84,Y87.0)	48,183	1.4	14.5	14.1	4.4	4.0	1.6	0.5	0.7	2.6	2.2
12	Essential hypertension and hypertensive renal disease. (I10,I12,I15)	42,816	1.2	12.9	10.7	5.9	1.2	1.1	2.1	1.2	1.0	1.0
13	Influenza and pneumonia. (J09-J18)	41,917	1.2	12.6	10.5	-19.2	1.5	1.4	1.3	1.0	1.3	1.1
14	Septicemia. (A40-A41)	41,281	1.2	12.4	10.2	5.2	1.2	1.2	1.8	1.1	2.2	1.4
15	Parkinson disease. (G20-G21)	38,536	1.1	11.6	9.8	-1.0	2.2	0.5	0.5	0.4	1.8	1.6
...	All other causes. (residual)	670,042	19.3	201.9	...	...	...	...	...	...	...	...

... Category not applicable.

¹Based on number of deaths; see Technical Notes.

²Percentages may not add to 100 due to rounding.

³Includes only one race reported on the death certificate.

⁴Includes people of Hispanic origin of any race; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

For the Native Hawaiian or Other Pacific Islander population, the age-adjusted rate increased 12.5%, from 821.3 in 2020 to 924.3 in 2021. During the same time period, rates increased 7.1% (from 834.7 to 893.9) for the White population and 7.0% (from 1,036.2 to 1,109.2) for the American Indian and Alaska Native population (Tables A and 1). The changes in age-adjusted rates were not statistically significant for the Asian, Black, and Hispanic populations.

From 2020 to 2021, the age-adjusted death rate increased 15.2% for Native Hawaiian or Other Pacific Islander females, 10.0% for Native Hawaiian or Other Pacific Islander males, 7.4% for American Indian and Alaska Native females, 7.2% for White males, 6.9% for White females, 6.4% for American Indian and Alaska Native males, 2.2% for Hispanic females, 2.1% for Asian females, and 1.3% for Black females (Tables A and 1). The rate decreased 2.1% for Hispanic males and 1.8% for Black males. The change in the age-adjusted rate for Asian males was not statistically significant.

Hispanic subgroups—Mortality data for 2021 by specified Hispanic subgroup for the United States are presented in Table 3. Hispanic subgroups shown in the table include Central American,

Cuban, Dominican, Mexican, Puerto Rican, South American, and Other Hispanic populations.

In 2021, age-adjusted rates among the Hispanic subgroups ranged from a low of 400.7 for the South American population to a high of 767.4 for the Mexican population. Differences between subgroups are likely a function of large variation in age-specific death rates for some of the Hispanic subgroups, reflecting their relatively small population sizes. Data aggregated over several years confirm differences among the Hispanic subgroups (16).


Death rates by age and sex

For the total population, age-specific death rates increased significantly from 2020 to 2021 for every age group (Tables 6 and 8; Figure 2).

The age-adjusted death rate for males was 1.4 times the rate for females in 2021 (Table B). The male-to-female death rate ratio was unchanged from the ratio in 2020.

Death rates for males increased significantly for age groups younger than 1, 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. Changes in rates for males ages

Figure 2. Death rate, by age and sex: United States, 1955–2021



1–4 and 5–14 were not statistically significant. Death rates for females increased significantly for every age group.

Race and ethnicity by sex—For the total Hispanic population, age-specific death rates increased from 2020 to 2021 for age groups younger than 1, 15–24, 25–34, 35–44, 45–54, and 55–64, and decreased for age groups 65–74, 75–84, and 85 and older. Rates for Hispanic males increased for age groups 25–34, 35–44, and 45–54, and decreased for age groups 65–74, 75–84, and 85 and older. For Hispanic females, rates increased for age groups younger than 1, 1–4, 15–24, 25–34, 35–44, 45–54, 55–64, and 65–74, and decreased for the age group 75–84.

For the total and male American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) populations, age-specific death rates increased from 2020 to 2021 for age groups 15–24, 25–34, 35–44, 45–54, and 55–64. For American Indian and Alaska Native females, age-specific death rates increased for age groups 15–24, 35–44, 45–54, 55–64, and 65–74.

For the total Asian non-Hispanic (subsequently, Asian) population, age-specific rates increased from 2020 to 2021 for age groups younger than 1, 25–34, 35–44, and 45–54. For Asian males, rates increased for age groups younger than 1, 15–24, and 25–34, and decreased for the age group 65–74. Rates for Asian females increased for age groups younger than 1, 25–34, 35–44, and 75–84.

For the total Black non-Hispanic (subsequently, Black) population, age-specific death rates increased from 2020 to 2021 for age groups 1–4, 5–14, 15–24, 25–34, 35–44, 45–54, and 55–64, and decreased for age groups 65–74, 75–84, and 85 and older. Rates for Black males increased for age groups 15–24, 25–34, 35–44, and 45–54, and decreased for age groups 65–74, 75–84, and 85 and older. For Black females, rates increased for age groups 1–4, 5–14, 15–24, 25–34, 35–44, 45–54, and 55–64, and decreased for age groups 75–84 and 85 and older.

For the total Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population, age-specific death rates increased for age groups 15–24, 25–34, 35–44, 45–54, and 55–64. For Native Hawaiian or Other Pacific Islander males, rates increased for age groups 15–24, 25–34, 35–44, and 45–54. For Native Hawaiian or Other Pacific Islander females, rates increased for age groups 35–44, 45–54, and 55–64.

For the total, male, and female White non-Hispanic (subsequently, White) populations, age-specific death rates increased from 2020 to 2021 for age groups younger than 1, 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older (Tables A and 2).

Other observed changes from 2020 to 2021 in age-specific rates by race and ethnicity and sex were not statistically significant.

Expectation of life at birth and at specified ages

Life expectancy at birth represents the average number of years that a group of infants would live if the group was to experience throughout life the age-specific death rates present in the year of birth.

Life table data shown in this report for 2010–2021 are based on a revised methodology first presented with final data reported for 2008. The life table methodology was revised by changing the smoothing technique used to estimate the life table functions at the oldest ages. This revision improves on the methodologies used previously; see Technical Notes.

The methods used to produce life expectancies by Hispanic origin and race are based on death rates adjusted for Hispanic-origin and race misclassification on death certificates (Technical Notes). As noted in the Methods section, the age-specific and age-adjusted death rates shown in this report (with the exception of Table IV) are not adjusted for misclassification of Hispanic origin and race on death certificates. For further information on the effects of Hispanic-origin and race misclassification on death rates, see Technical Notes.


Life tables were generated for both sexes and by each sex for the following populations:

- Total U.S.
- Hispanic
- American Indian and Alaska Native non-Hispanic
- Asian non-Hispanic
- Black non-Hispanic
- White non-Hispanic

In 2021, life expectancy at birth for the U.S. population was 76.4 years, 0.6 year lower than 2020 (Tables 4 and 5). The single-year decreases in life expectancy observed in 2020 (1.8 years) and 2021 were the largest decreases since 1943. These decreases were mostly due to the effects of the COVID-19 pandemic and increases in unintentional injuries. Before this time, the general trend in U.S. life expectancy had been one of improvement since 1900. In 2021, life expectancy for males (73.5 years) was 0.7 year lower than in 2020. Life expectancy for females (79.3 years) was 0.6 year lower than in 2020. From 1900 through the late 1970s, the gap in life expectancy between the sexes widened (3) from 2.0 to 7.8 years. The gap between sexes has narrowed since its peak in the 1970s (Figure 3 and Table 5). In 2021, the difference in life expectancy between males and females was 5.8 years, an increase of 0.1 year from 2020.

Life expectancy figures for the Hispanic, Black non-Hispanic (subsequently, Black), and White non-Hispanic (subsequently, White) populations have been available starting with data for 2006 (19). Before 2006, life tables were produced for the Black and White populations, regardless of Hispanic origin. Life tables for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) and Asian non-Hispanic (subsequently, Asian) populations were added to the Life Table Program beginning with data for 2019 (20).

Life expectancy for the American Indian and Alaska Native population decreased 1.5 years (from 67.1 years in 2020 to 65.6 in 2021).

Figure 3. Life expectancy at birth, by sex: United States, 1970–2021

Life expectancy decreased by 0.7 year for the White population (from 77.4 years in 2020 to 76.7 in 2021).

Life expectancy decreased by 0.3 year for the Black population (from 71.5 years in 2020 to 71.2 in 2021). The difference in life expectancy between the White and Black populations decreased 0.4 year from 5.9 years in 2020 to 5.5 years in 2021 (Table 5).

Life expectancy for the Hispanic population decreased by 0.1 year (from 77.9 years in 2020 to 77.8 in 2021) (Table 5). The difference in life expectancy between the Hispanic and White populations increased 0.6 year from 0.5 year in 2020 to 1.1 years in 2021 (Table 5).

Life expectancy declined by 0.1 year for the Asian population (from 83.6 years in 2020 to 83.5 in 2021).

Among the 10 major Hispanic-origin–race–sex groups in 2021, Asian females had the highest life expectancy at birth (85.6 years), followed by Asian males (81.2), Hispanic females (81.1), White females (79.5), Black females (75.0), Hispanic males (74.6), White males (74.0), American Indian and Alaska Native females (69.2), Black males (67.6), and American Indian and Alaska Native males (62.2) (Tables 4 and 5).

Life tables shown in this report may be used to compare life expectancies at selected ages from birth to 100 years. For

example, based on mortality experienced in 2021 for the total population, a person age 50 could expect to live an average of 30.1 more years, for a total of 80.1 years. A person age 65 could expect to live an average of 18.4 more years, for a total of 83.4 years, and a person age 85 could expect to live an average of 6.3 more years, for a total of 91.3 years (Table 4). Life expectancy decreased at ages from birth through age 65 and increased at age 80 and older from 2020 to 2021 (Table 4) (3).

Leading causes of death

The 15 leading causes of death in 2021 accounted for 80.7% of all deaths in the United States (Table B). In 2021, the 15 leading causes of death remained the same as in 2020, although some changed ranks. Heart disease, cancer, and COVID-19 remained the top three leading causes in 2021. Of the remaining leading causes in 2020, six causes changed rank. Chronic liver disease and cirrhosis, the 11th leading cause in 2020, became the 9th leading cause in 2021. Suicide, the 12th leading cause in 2020, became the 11th. Hypertension, the 13th leading cause in 2020, became the 12th. Influenza and pneumonia, the 9th leading cause in 2020, became the 13th. Septicemia, the 15th

leading cause in 2020, became the 14th. Parkinson disease, the 14th leading cause in 2020, became the 15th. Causes of death are ranked according to the number of deaths; for ranking procedures, see Technical Notes. By rank, the 15 leading causes of death in 2021 were:


1. Diseases of heart (heart disease)
2. Malignant neoplasms (cancer)
3. COVID-19
4. Accidents (unintentional injuries)
5. Cerebrovascular diseases (stroke)
6. Chronic lower respiratory diseases
7. Alzheimer disease
8. Diabetes mellitus (diabetes)
9. Chronic liver disease and cirrhosis
10. Nephritis, nephrotic syndrome and nephrosis (kidney disease)
11. Intentional self-harm (suicide)
12. Essential hypertension and hypertensive renal disease (hypertension)
13. Influenza and pneumonia

14. Septicemia
15. Parkinson disease

Death rates vary greatly by age. As a result, the shifting age distribution of a population can significantly influence changes in crude death rates over time. Age-adjusted death rates, in contrast, eliminate the influence of such differences in the population age structure. Consequently, while causes of death are ranked according to the number of deaths, age-adjusted death rates are used to depict trends for leading causes of death in this report because they are better than crude rates for showing changes in mortality over time and among causes of death (Figure 4; Tables B and 6).

From 2020 through 2021, age-adjusted death rates increased significantly for 11 of the 15 leading causes of death and decreased for 3 of the 15 leading causes (Tables B and 6). The rate for the top leading cause of death, heart disease, increased 3.3% in 2021 from 2020 (Figure 4; Tables B and 6) (14). The rate for the second leading cause of death, cancer, increased for the first time since 1993, by 1.7% from 2020 to 2021. Deaths from these two diseases combined accounted for 37.5% of deaths in the United States in 2021 (Table B).

Figure 4. Age-adjusted death rate for selected leading causes of death: United States, 1958–2021



COVID-19, the third leading cause of death in 2021, accounted for 12.0% of deaths in the United States. In 2021, the age-adjusted death rate for COVID-19 increased 22.5%, from 85.0 per 100,000 standard population in 2020 to 104.1 in 2021. The rate for males (131.3) was 1.6 times higher than the rate for females (81.7) (Tables 1, 2, and 11). Among race and ethnicity groups, the rate was highest for the Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population (185.4), followed by the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) (184.0), Hispanic (151.8), Black non-Hispanic (subsequently, Black) (136.4), White non-Hispanic (subsequently, White) (93.5), and Asian non-Hispanic (subsequently, Asian) (61.9) populations (Tables 1 and 11).

Other leading causes of death that showed significant increases in 2021 from 2020 were unintentional injuries (12.3%), stroke (5.9%), diabetes (2.4%), Chronic liver disease and cirrhosis (9.0%), kidney disease (7.1%), suicide (4.4%), hypertension (5.9%), and Septicemia (5.2%).

The age-adjusted rate decreased significantly in 2021 from 2020 for Chronic lower respiratory diseases (4.7%), Alzheimer disease (4.3%), and Influenza and pneumonia (19.2%).

The age-adjusted death rate for Parkinson disease in 2021 was not significantly different from the rate in 2020.

The relative risk of death in one population group compared with another can be expressed as a ratio. Ratios based on age-adjusted death rates show that males have higher rates than females for 13 of the 15 leading causes of death (Table B), with rates for males being at least twice as great as those for females for 3 of these leading causes. The largest ratio was for suicide (4.0). Other high ratios were observed for unintentional injuries and Parkinson disease (2.2 each); Chronic liver disease and cirrhosis (1.8); heart disease, COVID-19, and diabetes (1.6 each); Influenza and pneumonia (1.5); kidney disease (1.4); cancer (1.3); and Chronic lower respiratory diseases, hypertension, and Septicemia (1.2 each). Age-adjusted rates were lower for males than for females for one leading cause, Alzheimer disease (0.7).

Age-adjusted death rates for the American Indian and Alaska Native population were higher than for the White population for 9 of the 15 leading causes of death (Tables B and 6). The largest ratio was for Chronic liver disease and cirrhosis (5.1). Other causes for which the ratio was high include diabetes (2.3), COVID-19 (2.0), unintentional injuries (1.8), suicide (1.6), Influenza and pneumonia (1.4), kidney disease (1.3), Septicemia (1.2), and hypertension (1.1). Age-adjusted rates for the American Indian and Alaska Native population were lower than for the White population for 6 of the 15 leading causes of death. The smallest American Indian and Alaska Native-to-White ratio was for Parkinson disease (0.5); that is, the risk of dying from Parkinson disease was two times greater for the White population than for the American Indian and Alaska Native population. Other causes with a low American Indian and Alaska Native-to-White ratio were Alzheimer disease (0.6), Chronic lower respiratory diseases and cancer (0.8 each), and heart disease and stroke (0.9 each).

Age-adjusted death rates for the Black population were higher than for the White population for 10 of the 15 leading causes of death (Tables B and 6). The largest ratio was for kidney disease (2.2). Other causes for which the ratio was high include diabetes and hypertension (2.1 each), Septicemia (1.8), COVID-19 and stroke (1.5 each), heart disease and Influenza and pneumonia (1.3 each), and cancer and unintentional injuries (1.1 each). For five of the leading causes, age-adjusted rates were lower for the Black population than for the White population. The smallest Black-to-White ratios were for suicide and Parkinson disease (0.5 each). Other causes with low Black-to-White ratios were Chronic lower respiratory diseases and Chronic liver disease and cirrhosis (0.7 each) and Alzheimer disease (0.9).

Age-adjusted death rates for the White population were higher than for the Hispanic population for 10 of the 15 leading causes of death (Tables B and 6). The largest ratio was for Chronic lower respiratory diseases (2.7). Other causes with high ratios include suicide (2.2); Parkinson disease (1.6); heart disease, cancer, and unintentional injuries (1.5 each); Septicemia (1.4); Alzheimer disease (1.2); and stroke and Influenza and pneumonia (1.1 each). Age-adjusted rates were lower for the White population than for the Hispanic population for 3 of the 15 leading causes. The smallest White-to-Hispanic ratio was for COVID-19 (0.6), followed by diabetes (0.8) and Chronic liver disease and cirrhosis (0.9).

Age-adjusted death rates for the White population were higher than for the Asian population for 14 of the 15 leading causes of death (Tables B and 6). The highest ratio was for Chronic lower respiratory diseases (4.4), followed by unintentional injuries (3.7), Chronic liver disease and cirrhosis (3.6), suicide (2.6), Septicemia (2.2), heart disease (2.1), Alzheimer disease (1.9), Parkinson disease (1.8), cancer (1.7), COVID-19 and kidney disease (1.5 each), Influenza and pneumonia (1.3), and stroke and diabetes (1.2 each).

Assault (homicide), the 16th leading cause of death in 2021, dropped from among the 15 leading causes of death in 2010. In 2021, the age-adjusted rate for homicide increased 5.1%, from 7.8 in 2020 to 8.2 per 100,000 standard population in 2021 (Table 11). Homicide remains a major issue for some age groups. Homicide was among the 15 leading causes of death in 2021 for age groups 1–4 (3rd), 5–14 (4th), 15–24 (2nd), 25–34 (3rd), 35–44 (7th), 45–54 (10th), and 55–64 (15th) (For leading causes of infant death, see “Infant mortality”) (14).

Although Human immunodeficiency virus (HIV) disease has not been among the 15 leading causes of death since 1997 (21), it is still considered a major public health problem for some age groups. The age-adjusted death rate for HIV disease decreased 7.1% from 1.4 deaths per 100,000 U.S. standard population in 2020 to 1.3 in 2021 (Table 11). Historically, for all ages combined, HIV disease mortality reached its highest level in 1995 after a period of increase from 1987 through 1994. Subsequently, the rate for this disease decreased an average of 33.0% per year from 1995 through 1998, and 6.1% per year from 1999 through 2021 (14). In 2021, HIV disease was among the 15 leading causes of death for age groups 25–34 (11th), 35–44 (13th), and 45–54 (15th) (14). The greatest influence was among the age group 25–34 years, where HIV disease is ranked 9th for males, 8th for Black males, 9th for Black females, and 10th for Hispanic males.

Enterocolitis due to *Clostridium difficile* (*C. difficile*)—A predominantly antibiotic-associated inflammation of the intestines caused by *C. difficile*, a gram-positive, anaerobic, spore-forming bacillus—is often acquired in hospitals or other health care facilities with long-term patients or residents (22,23). The number of deaths from *C. difficile* climbed from 793 deaths in 1999 to a high of 8,085 deaths in 2011 (14). Since 2011, the number of deaths from this cause has trended downward. In 2021, the number of deaths from *C. difficile* was 4,105. The age-adjusted death rate in 2021 was unchanged from 2020, 1.0 deaths per 100,000 standard population. About 84.2% of deaths from *C. difficile* occurred among people age 65 and older (Table 7).

Leading causes of death in 2021 for the total population and for specific subpopulations are detailed further in a companion *National Vital Statistics Report* on leading causes by age, Hispanic origin and race, and sex (2).

Other selected causes

Dementia-related mortality

In 2021, 279,704 people died of dementia-related causes in the United States (Tables 7, 9, and 22). Deaths from dementia-related causes were presented for the first time in this report series in 2018 to provide a more comprehensive estimate of the burden of mortality from Alzheimer disease and other dementias in the United States.

Dementia-related causes include conditions with similar physical signs and symptoms that, collectively, are considered to be a good indicator of dementia mortality (24). Dementia is characterized by memory impairment and cognitive decline (25–27). Causes of death attributable to dementia-related mortality include ICD–10 codes F01, Vascular dementia; F03, Unspecified dementia; G30, Alzheimer disease; and G31, Other degenerative diseases of nervous system, not elsewhere classified. Alzheimer disease, the sixth leading cause of death, is the most common cause of dementia, but other dementias, including Lewy body dementia, frontotemporal degeneration, vascular dementia, and mixed dementias, are often indistinguishable from Alzheimer disease in their symptoms and outcomes and may coexist with Alzheimer disease (24–26).

Certification and coding rule changes can impact data analysis of component causes of dementia. In 2021, Alzheimer disease accounted for 42.7% of all dementia deaths; Unspecified dementia for 32.8%; Other degenerative diseases of nervous system, not elsewhere classified for 17.6%; and Vascular dementia for 6.9%. For detailed information, see CDC WONDER (14). Changes in the percentage of deaths assigned to individual causes comprising dementia may be the result of many factors (27). Combining the types of dementia provides a more comprehensive and stable measure of dementia mortality.

The age-adjusted death rate for dementia-related causes decreased 1.2% in 2021 from 2020 for the total population (from 73.3 to 72.4). The rate for males decreased 1.3% (60.5 to 59.7) and decreased for females 1.1% (81.1 to 80.2) (Tables 11 and 22).

Among race-ethnicity groups—Age-adjusted rates for the total, male, and female Black non-Hispanic (subsequently,

Black) populations decreased 3.9% (74.7 to 71.8), 5.6% (66.4 to 62.7), and 2.9% (78.2 to 75.9), respectively. The rate for Hispanic males decreased 4.1% (46.1 to 44.2), but rates did not change significantly for the total and female Hispanic populations. Changes in age-adjusted death rates for dementia-related causes were not statistically significant for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native), Asian non-Hispanic (subsequently, Asian), Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander), and White non-Hispanic (subsequently, White) populations.

Drug-induced mortality

In 2021, a total of 111,219 people died of drug-induced causes in the United States (Tables 7, 9, and 23). The category of drug-induced causes includes deaths from drug overdose as well as from other medical conditions caused by use of legal or illegal drugs. In 2021, drug-overdose deaths accounted for 95.9% of all drug-induced deaths (Tables 7 and 9). The drug-induced category excludes deaths indirectly related to drug use, as well as newborn deaths due to the mother's drug use. (For a list of all drug-induced causes, including those specifically classified as drug-overdose causes, see Technical Notes.)

The age-adjusted death rate for drug-induced causes increased 13.9% for the total population from 29.5 in 2020 to 33.6 in 2021 (Tables 6, 11, and 23). For males in 2021, the age-adjusted death rate for drug-induced causes was 2.3 times the rate for females. The rate increased by 13.9% for males and by 14.0% for females from 2020 to 2021 (Tables 11 and 23).

Among race-ethnicity groups—Age-adjusted rates increased from 2020 and 2021 by 33.9% for the American Indian and Alaska Native population (36.2% for American Indian and Alaska Native males and 29.1% for American Indian and Alaska Native females), 22.8% for the Black population (21.8% for Black males and 23.9% for Black females), 19.0% for the Hispanic population (17.5% for Hispanic males and 22.8% for Hispanic females), 47.9% for the Native Hawaiian or Other Pacific Islander population (48.0% for Native Hawaiian or Other Pacific Islander males), and 10.7% for the White population (10.6% for White males and 10.7% for White females) (Tables 11 and 23). Changes in age-adjusted rates from 2020 to 2021 were not significant for the Asian population or Native Hawaiian or Other Pacific Islander females.

Alcohol-induced mortality

In 2021, a total of 54,258 people died of alcohol-induced causes in the United States (Tables 7, 9, and 24). This category includes deaths from dependent and nondependent use of alcohol, and deaths from accidental poisoning by alcohol. It excludes unintentional injuries, homicides, and other causes indirectly related to alcohol use, and deaths due to fetal alcohol syndrome. For a list of alcohol-induced causes, see Technical Notes.

The age-adjusted death rate for alcohol-induced causes increased 9.9%, from 13.1 in 2020 to 14.4 in 2021

(Tables 6, 11, and 24). For males in 2021, the age-adjusted death rate for alcohol-induced causes was 2.5 times the rate for females. The rate increased 8.9% for males and 10.7% for females from 2020 to 2021 (Tables 11 and 24).

Among race-ethnicity groups—Age-adjusted rates increased from 2020 to 2021 by 23.9% for the American Indian and Alaska Native population (30.6% for American Indian and Alaska Native males and 13.9% for American Indian and Alaska Native females), 15.5% for the Black population (10.7% for Black males and 21.8% for Black females), 7.9% for the Hispanic population (7.2% for Hispanic males), and 9.1% for the White population (7.9% for White males and 9.2% for White females) (Tables 11 and 24). The age-adjusted rate for alcohol-induced death did not change significantly from 2020 to 2021 for the Asian populations, Hispanic females, and Native Hawaiian or Other Pacific Islander populations.

Firearm-related mortality

In 2021, 48,830 people died from firearm-related injuries in the United States (Tables 7, 9, and 25). The age-adjusted death rate for firearm-related injuries for the total, male, and female populations increased significantly from 2020 to 2021 by 7.4%, 6.3%, and 10.5%, respectively (Tables 6, 11, and 25). For males in 2021, the age-adjusted death rate for firearm-related injuries was 6.0 times the rate for females.

Among race-ethnicity groups—The age-adjusted death rate increased from 2020 to 2021 by 8.8% for the Black population (6.8% for Black males and 23.3% for Black females), 12.7% for the Hispanic population (10.9% for Hispanic males and 21.1% for Hispanic females), and 6.0% for the White population (5.5% for White males and 5.4% for White females) (Tables 11 and 25). The age-adjusted death rates for firearm-related injuries did not change significantly from 2020 to 2021 for the American Indian and Alaska Native, Asian, and Native Hawaiian or Other Pacific Islander populations.

Effect on life expectancy of changes in mortality by age and cause of death

Changes in mortality by age and cause of death can have a major effect on life expectancy. In other words, year-to-year changes in life expectancy may be influenced by changes in age-specific rates for certain causes, particularly for younger age groups. Life expectancy at birth for the total population decreased by 0.6 year (from 77.0 years in 2020 to 76.4 years in 2021) primarily because of increases in mortality from COVID-19, unintentional injuries, Chronic liver disease and cirrhosis, suicide, and homicide (Table 4). The decrease in life expectancy for the total population was slightly offset by decreases in mortality from Influenza and pneumonia, Chronic lower respiratory diseases, Alzheimer disease, and Parkinson disease. Life expectancy at birth for males decreased by 0.7 year (from 74.2 years in 2020 to 73.5 years in 2021) due to increases in mortality from COVID-19, unintentional injuries, Chronic liver disease and cirrhosis, suicide, and homicide. These increases were offset somewhat by decreases in mortality from cancer,

Influenza and pneumonia, Chronic lower respiratory diseases, Alzheimer disease, and heart disease. For the female population, life expectancy decreased by 0.6 year (from 79.9 years in 2020 to 79.3 years in 2021) due to increases in mortality from COVID-19, unintentional injuries, Chronic liver disease and cirrhosis, stroke, and heart disease, which were offset by decreases in mortality from Influenza and pneumonia, Alzheimer disease, Chronic lower respiratory diseases, cancer, and Parkinson disease. (For a discussion of the major causes contributing to the change in life expectancy, see Technical Notes.)

Life expectancy for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population decreased by 1.5 years to 65.6 years. This decrease, the largest among race and ethnicity groups, was primarily due to increases in mortality due to unintentional injuries, COVID-19, Chronic liver disease and cirrhosis, suicide, and heart disease. The decline in life expectancy was offset by decreases in mortality due to Influenza and pneumonia, homicide, congenital malformations, Certain conditions originating in the perinatal period, and Alzheimer disease. Life expectancy for the American Indian and Alaska Native male population decreased 1.6 years to 62.2 years. This decrease, primarily due to increases in mortality from unintentional injuries, Chronic liver disease and cirrhosis, COVID-19, suicide, and heart disease, was offset somewhat by decreases in mortality from homicide, Certain conditions originating in the perinatal period, Influenza and pneumonia, diabetes, and congenital malformations.

Life expectancy for the White non-Hispanic (subsequently, White) population decreased 0.7 year in 2021 to 76.7 years. This decrease was due to increases in mortality from COVID-19, unintentional injuries, heart disease, Chronic liver disease and cirrhosis, and suicide. These increases in mortality were offset to some extent by decreases from Influenza and pneumonia, Chronic lower respiratory diseases, Alzheimer disease, cancer, and Parkinson disease. Life expectancy for the White male population decreased 0.8 year in 2021 to 74.0 years due to increases in mortality from COVID-19, unintentional injuries, Chronic liver disease and cirrhosis, suicide, and diabetes, which was offset somewhat by decreases in mortality from cancer, Chronic lower respiratory diseases, Influenza and pneumonia, Alzheimer disease, and Certain conditions originating in the perinatal period. Life expectancy for White females decreased 0.6 year in 2021 to 79.5 years due to increases in mortality from COVID-19, unintentional injuries, heart disease, stroke, and Chronic liver disease and cirrhosis, which were offset somewhat by decreases in mortality from Influenza and pneumonia, Alzheimer disease, Chronic lower respiratory diseases, Parkinson disease, and homicide.

Life expectancy for the Black non-Hispanic (subsequently, Black) population in 2021 decreased 0.3 year to 71.2 years. This decrease was due to increases in mortality from unintentional injuries, COVID-19, homicide, suicide, and Chronic liver disease and cirrhosis, which were offset by decreases in mortality from heart disease, cancer, Influenza and pneumonia, Chronic lower respiratory diseases, and Certain conditions originating in the perinatal period. Life expectancy for the Black male population in 2020 decreased 0.2 year to 67.6 years. This decrease was due

to increases in mortality from unintentional injuries, homicide, COVID-19, suicide, and Chronic liver disease and cirrhosis, which were offset by decreases in mortality from cancer, heart disease, Influenza and pneumonia, Chronic lower respiratory diseases, and diabetes. Life expectancy for the Black female population decreased 0.4 year in 2021 to 75.0 years due to increases in mortality from COVID-19, unintentional injuries, homicide, Chronic liver disease and cirrhosis, and stroke, which were offset somewhat by decreases from heart disease, cancer, Influenza and pneumonia, Chronic lower respiratory diseases, and Alzheimer disease.

Life expectancy for the Asian non-Hispanic (subsequently, Asian) population decreased 0.1 year to 83.5 years. The decrease, primarily due to increases in mortality due to cancer, unintentional injuries, Certain conditions originating in the perinatal period, stroke, and hypertension, was offset by decreases in mortality due to Influenza and pneumonia, Chronic lower respiratory diseases, diabetes, heart disease, and Alzheimer disease. Life expectancy for the Asian male population increased 0.1 year in 2021 to 81.2 years due to decreases in mortality from heart disease, Influenza and pneumonia, diabetes, Chronic lower respiratory diseases, and Alzheimer disease, which were offset somewhat by increases in mortality from Certain conditions originating in the perinatal period, unintentional injuries, suicide, hypertension, and COVID-19. Life expectancy for the Asian female population decreased by 0.3 year in 2021 to 85.6 years due to increases in mortality from cancer, COVID-19, Certain conditions originating in the perinatal period, stroke, and unintentional injuries, which were offset somewhat by decreases in mortality from Influenza and pneumonia, congenital malformations, Chronic lower respiratory diseases, atherosclerosis, and acute bronchitis.

Life expectancy for the Hispanic population decreased by 0.1 year to 77.8 years. This decrease was due to increases in mortality from unintentional injuries, COVID-19, homicide, Chronic liver disease and cirrhosis, and suicide, which were offset by decreases from heart disease, Influenza and pneumonia, Alzheimer disease, diabetes, and Chronic lower respiratory diseases. Life expectancy for the Hispanic male population in 2021 remained unchanged at 74.6 years due to increases in mortality from unintentional injuries, homicide, Chronic liver disease and cirrhosis, suicide, and Complications of medical and surgical care, which were offset by decreases in mortality from heart disease, Influenza and pneumonia, cancer, diabetes, and Chronic lower respiratory diseases. Life expectancy for the Hispanic female population decreased by 0.2 year in 2021 to 81.1 years due to increases in mortality from COVID-19, unintentional injuries, cancer, stroke, and Pregnancy, childbirth and the puerperium, which were offset somewhat by decreases in mortality from heart disease, Influenza and pneumonia, Alzheimer disease, diabetes, and Chronic lower respiratory diseases.

The difference in life expectancy between the male and female populations increased 0.1 year in 2021 to 5.8 years (Table 4). The widening in the male–female life expectancy gap was due primarily to greater increases in mortality for the male population for unintentional injuries, COVID-19, Alzheimer disease, suicide, and homicide (data not shown).

Injury mortality by mechanism and intent

In 2021, a total of 306,086 deaths were classified as injury-related (Table 12). Injury data are presented using the external cause-of-injury mortality matrix for ICD–10, as jointly conceived by the International Collaborative Effort on Injury Statistics and the Injury Control and Emergency Health Services section of the American Public Health Association (28,29). The ICD codes for injuries have two essential dimensions: the mechanism of the injury and its manner or intent. The mechanism involves the circumstances of the injury (such as fall, motor-vehicle traffic, or poisoning). The manner or intent involves whether the injury was purposefully inflicted (where it can be determined) and, when intentional, whether the injury was self-inflicted (suicide) or inflicted upon another person (assault). In other report tables showing cause of death, the focus is on manner or intent, with subcategories showing selected mechanisms. The matrix has two distinct advantages for the analysis of injury mortality data: It contains a comprehensive list of mechanisms, and data can be displayed by mechanism with subcategories of intent, or vice versa. Four major mechanisms of injury in 2021—poisoning, firearm, fall, and motor-vehicle traffic—accounted for 82.3% of all injury deaths (Table 12). A total of 111,830 deaths occurred as the result of poisonings in 2021, accounting for 36.5% of all injury deaths (Table 12). The age-adjusted death rate for poisoning increased significantly, by 13.4% from 29.8 deaths per 100,000 U.S. standard population in 2020 to 33.8 in 2021. Most poisoning deaths were either unintentional (91.2%) or suicides (5.0%). However, 3.6% of poisoning deaths were of undetermined intent. The age-adjusted death rate for unintentional poisoning increased 15.2%, from 26.9 in 2020 to 31.0 in 2021.

Firearm injuries resulted in 48,830 deaths in 2021 (Table 12), accounting for 16.0% of all injury deaths. The age-adjusted death rate for firearm injuries (all intents) in 2021 increased 7.4% from 2020. The two major component causes of firearm injury deaths in 2021 were suicide (53.9%) and homicide (42.9%). The age-adjusted death rate for firearm suicide increased 7.1%, from 7.0 in 2020 to 7.5 in 2021. The age-adjusted rate for firearm homicide increased 8.1%, from 6.2 in 2020 to 6.7 in 2021. A total of 45,988 people died as the result of falls in 2021, accounting for 15.0% of all injury deaths (Table 12). The age-adjusted death rate for falls in 2021 increased 11.3%, from 10.6 in 2020 to 11.8 in 2021. The overwhelming majority of fall-related deaths (97.2%) were unintentional. Motor vehicle traffic-related injuries in 2021 resulted in 45,404 deaths, accounting for 14.8% of all injury deaths (Table 12). The age-adjusted death rate for these injuries increased 10.8%, from 12.0 in 2020 to 13.3 in 2021.

Marital status

For those age 15 and older, the number of deaths in 2021 among people who were married was 1,279,442; widowed, 1,028,227; divorced, 601,216; and never married, 496,695 (Table 13); see Technical Notes. Those who were widowed had the highest age-adjusted death rate (2,036.2 per 100,000 U.S. standard population), followed by never-married people (1,729.3), divorced people (1,643.7), and married people

(898.6). Never-married people had an age-adjusted death rate 39.4% higher than those ever-married (92.4% higher than currently married people and 5.2% higher than divorced people). Widowed people had a rate more than twice that of married people, 23.9% higher than divorced people, and 17.7% higher than never-married people. Divorced people had a rate 1.8 times the rate of those who were married at the time of death.

For all age groups 15 and older, age-specific death rates for married people were much lower than those for never-married people. For those ages 15–24, divorced people had the highest death rate, whereas for those ages 25–34, 35–44, and 75 and older, widowed people had the highest death rate. Never-married people had the highest death rate among those ages 45–54, 55–64, and 65–74.

For each marital status group in 2021, males had higher age-adjusted death rates than females, ranging from 48.4% greater for those never married to 71.2% higher for those married at the time of death.

Educational attainment

Age-specific and age-adjusted death rates are shown by educational attainment for age groups in the range of 25–64 years (Table 14). In 2021, a total of 409,553 decedents ages 25–64 had received a high school diploma or equivalent, compared with 312,389 who had completed some college or collegiate degree and 159,086 who had achieved less than a high school diploma or equivalent. In 2021, age-adjusted death rates for those with less than a high school diploma or GED (778.2 per 100,000 U.S. standard population) and for those with a high school diploma or GED (817.1) were more than 3 times the rate for those with some college or collegiate degree (253.9).

Injury at work

Deaths, crude death rates, and age-adjusted death rates for injury at work are shown for those age 15 and older (Tables 15 and 16). Age-adjusted death rates for injury at work were computed using age-specific death rates and the 2000 U.S. standard population for those age 15 and older; see “Computing rates” in Technical Notes. Information on deaths attributed to injuries at work is derived from a separate item on the death certificate that asks the medical certifier whether the death resulted from an injury sustained at work. This item is on the death certificate of all states.

State of residence

Mortality patterns varied considerably by state (Tables 17 and 20). The state with the highest age-adjusted death rate in 2021 was West Virginia (1,229.1 deaths per 100,000 U.S. standard population), with a rate 39.7% above the national rate (879.7). The state with the lowest age-adjusted death rate was Hawaii (630.0), with a rate 28.4% below the national rate. The age-adjusted death rate for West Virginia was 95.1% higher than the rate for Hawaii.

Variations in mortality by state were associated with differences in socioeconomic status, racial and ethnic composition, as well as with differences in risk of specific causes of death (30).

Infant mortality

In 2021, a total of 19,920 deaths occurred in children younger than age 1 year (Tables C, D, 19, and 20). This number represents 338 more infant deaths in 2021 than in 2020. The ratio of male-to-female IMRs was 1.2, the same as in 2020. IMR was 5.44 infant deaths per 1,000 live births, the neonatal

Table C. Number of infant, neonatal, and postneonatal deaths and mortality rate, by sex: United States, 2020–2021

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days–11 months) deaths per 1,000 live births in specified group]

Age and sex	2021		2020		Percent change ¹ from 2020 to 2021
	Number	Rate	Number	Rate	
Infant					
Total.....	19,920	5.44	19,582	5.42	0.4
Male.....	10,909	5.82	10,859	5.88	-1.0
Female.....	9,011	5.03	8,723	4.94	1.8
Neonatal					
Total.....	12,768	3.48	12,856	3.56	-2.2
Male.....	6,899	3.68	7,098	3.84	-4.2
Female.....	5,869	3.28	5,758	3.26	0.6
Postneonatal					
Total.....	7,152	1.95	6,726	1.86	4.8
Male.....	4,010	2.14	3,761	2.04	4.9
Female.....	3,142	1.75	2,965	1.68	4.2

¹Based on a comparison of 2021 and 2020 mortality rates.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table D. Number of infant deaths, percentage of total infant deaths, and infant mortality rate for 2021, and percent change in infant mortality rates from 2020 to 2021 for the 10 leading causes of infant death in 2021: United States

[Rates are infant deaths per 100,000 live births]

Rank ¹	Cause of death (based on <i>International Classification of Diseases, 10th Revision</i>)	Number	Percent of total deaths ²	Rate	Percent change ³ from 2020 to 2021
...	All causes.	19,920	100.0	543.6	0.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	3,963	19.9	108.2	-3.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified. (P07)	2,946	14.8	80.4	-7.5
3	Sudden infant death syndrome (R95)	1,459	7.3	39.8	3.6
4	Accidents (unintentional injuries). (V01–X59)	1,306	6.6	35.6	7.9
5	Newborn affected by maternal complications of pregnancy. (P01)	1,113	5.6	30.4	-1.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	672	3.4	18.3	-5.7
7	Bacterial sepsis of newborn (P36)	557	2.8	15.2	1.3
8	Respiratory distress of newborn (P22)	414	2.1	11.3	5.6
9	Diseases of the circulatory system. (I00–I99)	402	2.0	11.0	2.8
10	Intrauterine hypoxia and birth asphyxia (P20–P21)	358	1.8	9.8	15.3
...	All other causes (residual)	6,730	33.8	183.7	...

... Category not applicable.

¹Based on number of deaths; see Technical Notes in this report.²Percentages may not add to 100 due to rounding.³Based on a comparison of the 2021 infant mortality rate with the 2020 infant mortality rate.

NOTE: Due to rounding, percent changes based on rates per 100,000 live births may differ from those computed using rates per 1,000 live births.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

mortality rate (deaths of infants ages 0–27 days per 1,000 live births) was 3.48, and the postneonatal mortality rate (deaths of infants ages 28 days–11 months per 1,000 live births) was 1.95 in 2021 (Figure 5 and Tables C and 18; see Technical Notes for information on alternative data sources).

The postneonatal mortality rate increased 4.8% from 2020 to 2021. Changes in the infant and neonatal mortality rates from 2020 to 2021 were not statistically significant.

The 10 leading causes of infant death in 2021 accounted for 66.2% of all infant deaths in the United States (Table D). By rank, the 10 leading causes were:

1. Congenital malformations, deformations and chromosomal abnormalities (congenital malformations)
2. Disorders related to short gestation and low birth weight, not elsewhere classified (low birth weight)
3. Sudden infant death syndrome (SIDS)
4. Accidents (unintentional injuries)
5. Newborn affected by maternal complications of pregnancy (maternal complications)
6. Newborn affected by complications of placenta, cord and membranes (cord and placental complications)
7. Bacterial sepsis of newborn
8. Respiratory distress of newborn
9. Diseases of the circulatory system
10. Intrauterine hypoxia and birth asphyxia

The rankings of the top nine leading causes of infant death remained unchanged in 2021 from 2020 (31). The 10th leading cause in 2021, Intrauterine hypoxia and birth asphyxia, replaced Neonatal hemorrhage, which dropped from the list. Among the


10 leading causes, IMR decreased 7.5% from 2020 for low birth weight. Changes in rates among the other leading causes of infant death were not statistically significant (Table D).

IMRs by race for non-Hispanic origin that are based on the mortality file may be somewhat understated and are better measured using data from the linked file of live births and infant deaths (32); see Technical Notes. Infant mortality data presented in this report use the general mortality file, not the linked file of live births and infant deaths. IMRs for the Hispanic population are not adjusted for misclassification; see Technical Notes. Because these rates are not adjusted, misclassification of Hispanic origin should be considered when interpreting rate disparities between Hispanic and non-Hispanic populations (16).

In 2021, the IMR for Asian non-Hispanic infants was 2.94 per 1,000 live births, an increase of 25.1% from 2.35 in 2020. In 2021, the IMR was 5.03 for Hispanic infants, 7.46 for American Indian and Alaska Native non-Hispanic infants, 10.97 for Black non-Hispanic infants, 7.24 for Native Hawaiian or Other Pacific Islander non-Hispanic infants, and 4.20 for White non-Hispanic infants (Table 18). The changes in IMRs in 2021 from 2020 for these groups were not statistically significant.

Maternal mortality

In 2021, a total of 1,205 women died of maternal causes in the United States—344 more deaths than in 2020 (Table 21) (31). The overall maternal mortality rate increased 38.2%, from 23.8 deaths per 100,000 live births in 2020 to 32.9 in 2021. COVID-19, listed on the death certificate as a contributing cause of death for 35.6% of maternal deaths in 2021, likely contributed to the rise in maternal mortality in 2021 (14). The maternal mortality rate in 2021 was highest for American Indian and Alaska Native non-Hispanic (subsequently, American

Figure 5. Infant, neonatal, and postneonatal mortality rates: United States, 1960–2021

Indian and Alaska Native) women (118.7), followed by Black non-Hispanic (subsequently, Black) (69.9), Hispanic (28.0), White non-Hispanic (subsequently, White) (26.6), and Asian non-Hispanic (subsequently, Asian) (16.8) women. The maternal mortality rate increased 53.8% for Hispanic women (from 18.2 in 2020 to 28.0 in 2021), 26.4% for Black women (from 55.3 to 69.9), and 39.3% for White women (from 19.1 to 26.6). The change in maternal mortality rates for Asian women was not significant. For American Indian and Alaska Native women, the maternal mortality rate in 2021 is not comparable to the rate in 2020 because the number of deaths in 2020 did not meet National Center for Health Statistics standards of reliability. In 2021, the maternal mortality rate for Native Hawaiian or Other Pacific Islander women did not meet standards of reliability because the number of deaths is too low.

Maternal deaths and death rates shown in this report are based on a revised method for coding maternal deaths that was adopted by the National Center for Health Statistics starting with the 2018 data year, and a new coding change that was adopted in 2021 (33) (see Technical Notes). Maternal deaths include deaths of women, while pregnant or within 42 days of being pregnant,

from any cause related to or aggravated by the pregnancy but exclude deaths from external causes (that is, accidents, homicides, and suicides); for more information, see “Maternal Mortality in the United States: Changes in Coding, Publication, and Data Release, 2018” (33) and Technical Notes.

References

1. Hoyert DL, Singh GK, Rosenberg HM. Sources of data on socio-economic differential mortality in the United States. *J Off Stat* 11(3):233–60. 1995.
2. Curtin SC, Tejada-Vera B, Bastian BA. Deaths: Leading causes for 2021. National Vital Statistics Reports; vol 73 no 4. Hyattsville, MD: National Center for Health Statistics. 2024. DOI: <https://dx.doi.org/10.15620/cdc/147882>.
3. Arias E, Xu JQ, Kochanek KD. United States life tables, 2021. National Vital Statistics Reports; vol 72 no 12. Hyattsville, MD: National Center for Health Statistics. 2023. DOI: <https://dx.doi.org/10.15620/cdc/132418>.

4. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. 2008 ed. 2009.
5. National Vital Statistics System. Volume 1. ICD–10, International statistical classification of diseases and related health problems. Tabular list. (Modified by NCHS for use in the classification and analysis of medical mortality data in the U.S.) NCHS Instruction Manual; part 2e, vol 1. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
6. National Vital Statistics System. Volume 1. ICD–10, International statistical classification of diseases and related health problems. Alphabetical index. (Modified by NCHS for use in the classification and analysis of medical mortality data in the U.S.) NCHS Instruction Manual; part 2e, vol 3. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
7. National Center for Health Statistics. 2003 revision of the U.S. Standard Certificate of Death. 2003. Available from: <https://www.cdc.gov/nchs/data/dvs/DEATH11-03final-acc.pdf>.
8. Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. Fed Regist 62FR58782. 1997. Available from: <https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>.
9. Murphy SL, Xu JQ, Kochanek KD, Arias E, Tejada-Vera B. Deaths: Final data for 2018. National Vital Statistics Reports; vol 69 no 13. Hyattsville, MD: National Center for Health Statistics. 2020. Available from: <https://www.cdc.gov/nchs/data/nvsr/nvsr69/nvsr69-13-508.pdf>.
10. Office of Management and Budget. Race and ethnic standards for federal statistics and administrative reporting. Statistical Policy Directive no 15. 1977. Available from: <https://wonder.cdc.gov/wonder/help/populations/bridged-race/directive15.html>.
11. Heron M. Comparability of race-specific mortality data based on 1977 versus 1997 reporting standards. National Vital Statistics Reports; vol 70 no 3. Hyattsville, MD: National Center for Health Statistics. 2021. DOI: <https://dx.doi.org/10.15620/cdc:103476>.
12. National Center for Health Statistics. Vital statistics online data portal: Mortality multiple cause files. Hyattsville, MD: National Center for Health Statistics. Published annually. Available from: https://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.
13. National Center for Health Statistics. National Vital Statistics System: Data release policy for vital statistics micro-data files. Available from: https://www.cdc.gov/nchs/nvss/dvs_data_release.htm.
14. Centers for Disease Control and Prevention. CDC Wide-ranging online data for epidemiologic research (WONDER). Available from: <https://wonder.cdc.gov/>.
15. Tolson GC, Barnes JM, Gay GA, Kowaleski JL. The 1989 revision of the U.S. standard certificates and reports. National Center for Health Statistics. Vital Health Stat 4(28). 1991. Available from: https://www.cdc.gov/nchs/data/series/sr_04/sr04_028.pdf.
16. Arias E, Heron M, Hakes JK. The validity of race and Hispanic-origin reporting on death certificates in the United States: An update. National Center for Health Statistics. Vital Health Stat 2(172). 2016. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_172.pdf.
17. Centers for Disease Control and Prevention. CDC WONDER. Single-race population estimates, United States, 2020–2021. July 1st resident population by state, age, sex, single-race, and Hispanic origin. Data source: Vintage 2021 estimates released by U.S. Census Bureau on June 30, 2022. Available from: <https://wonder.cdc.gov/single-race-population.html>.
18. National Center for Health Statistics. Multiple cause of death public use file, Control Total Table 1. 2023. Available from: <https://www.cdc.gov/nchs/data/dvs/2021-mortality-control-total-table-1.pdf>.
19. Arias E. United States life tables by Hispanic origin. National Center for Health Statistics. Vital Health Stat 2(152). 2010. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_152.pdf.
20. Arias E, Xu JQ. United States life tables, 2019. National Vital Statistics Reports; vol 70 no 19. Hyattsville, MD: National Center for Health Statistics. 2022. DOI: <https://dx.doi.org/10.15620/cdc:113096>.
21. Hoyert DL, Kochanek KD, Murphy SL. Deaths: Final data for 1997. National Vital Statistics Reports; vol 47 no 19. Hyattsville, MD: National Center for Health Statistics. 1999. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr47/nvs47_19.pdf.
22. Sunenshine RH, McDonald LC. *Clostridium difficile*-associated disease: New challenges from an established pathogen. Cleve Clin J Med 73(2):187–97. 2006.
23. Redelings MD, Sorvillo FJ, Mascola L. Increase in *Clostridium difficile*-related mortality rates, United States, 1999–2004. Emerg Infect Dis 13(9). 2007. Available from: https://wwwnc.cdc.gov/eid/article/13/9/06-1116_article.
24. World Health Organization. Dementia. Available from: <https://www.who.int/news-room/fact-sheets/detail/dementia>.
25. National Institute on Aging. Alzheimer's disease fact sheet. Available from: <https://www.nia.nih.gov/health/alzheimers-and-dementia/alzheimers-disease-fact-sheet>.
26. U.S. Department of Health and Human Services. National plan to address Alzheimer's disease: 2019 update. 2019. Available from: <https://aspe.hhs.gov/system/files/pdf/262601/NatlPlan2019.pdf>.
27. Kramarow EA, Tejada-Vera B. Dementia mortality in the United States, 2000–2017. National Vital Statistics Reports; vol 68 no 2. Hyattsville, MD: National Center for Health Statistics. 2019. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68_02-508.pdf.
28. National Center for Health Statistics. Proceedings of the international collaborative effort on injury statistics; vol 1. 1995.

- Available from: https://stacks.cdc.gov/view/cdc/11212/cdc_11212_DS1.pdf?download-document-submit=Download.
29. Fingerhut LA, Cox CS, Warner M, participants of the International Collaborative Effort on Injury Statistics. International comparative analysis of injury mortality: Findings from the ICE on Injury Statistics. *Advance Data from Vital and Health Statistics*; no 303. Hyattsville, MD: National Center for Health Statistics. 1998. Available from: <https://www.cdc.gov/nchs/data/ad/ad303.pdf>.
 30. Pamuk ER, Makuc DM, Heck KE, Reuben C, Lochner K. Socioeconomic status and health chartbook. Health, United States, 1998. Hyattsville, MD: National Center for Health Statistics. 1998. Available from: <https://www.cdc.gov/nchs/data/hs/hs98.pdf>.
 31. Kochanek KD, Murphy SL, Xu JQ, Arias E. Deaths: Final data for 2020. *National Vital Statistics Reports*; vol 72 no 10. Hyattsville, MD: National Center for Health Statistics. 2023. DOI: <https://dx.doi.org/10.15620/cdc:131355>.
 32. National Center for Health Statistics. User guide to the 2021 period/2020 cohort linked birth/infant death public use file. 2023. Available from: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/DVS/period-cohort-linked/.
 33. Hoyert DL, Miniño AM. Maternal mortality in the United States: Changes in coding, publication, and data release, 2018. *National Vital Statistics Reports*; vol 69 no 2. Hyattsville, MD: National Center for Health Statistics. 2020. Available from: <https://www.cdc.gov/nchs/data/nvsr/nvsr69/nvsr69-02-508.pdf>.
 34. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. 1992.
 35. National Center for Health Statistics. Comparability of cause-of-death between ICD revisions. 2015. Available from: https://www.cdc.gov/nchs/nvss/mortality/comparability_icd.htm.
 36. National Center for Health Statistics. Updated comparability ratios (ICD-10 and ICD-9). 2004. Available from: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/Comparability/icd9_icd10/.
 37. Anderson RN, Miniño AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: Preliminary estimates. *National Vital Statistics Reports*; vol 49 no 2. Hyattsville, MD: National Center for Health Statistics. 2001. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_02.pdf.
 38. Faust MM, Dolman AB. Comparability of mortality statistics for the sixth and seventh revisions: United States, 1958. *Vital Statistics—Special Reports* 51(4). National Center for Health Statistics. 1965. Available from: https://www.cdc.gov/nchs/data/spec_rpt51_04.pdf.
 39. Klebba AJ, Dolman AB. Comparability of mortality statistics for the seventh and eighth revisions of the International Classification of Diseases, United States. *National Center for Health Statistics. Vital Health Stat* 2(66). 1975. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_066.pdf.
 40. Klebba AJ, Scott JH. Estimates of selected comparability ratios based on dual coding of 1976 death certificates by the eighth and ninth revisions of the International Classification of Diseases. *Monthly Vital Statistics Report*; vol 28 no 11. Hyattsville, MD: National Center for Health Statistics. 1980. Available from: https://www.cdc.gov/nchs/data/mvss/supp/mv28_11s.pdf.
 41. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2a: Instructions for classifying the underlying cause of death. Published annually. Available from: <https://www.cdc.gov/nchs/nvss/manuals/2a-2021.htm>.
 42. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2b: Instructions for classifying the multiple causes of death. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
 43. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2c: ICD-10 ACME decision tables for classifying underlying causes of death. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
 44. National Center for Health Statistics. Instruction manuals. Historic manuals—Prior to ICD-10. Part 2g: Data entry instructions for the mortality medical indexing, classification, and references system (MICAR), 1996–1997. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
 45. National Center for Health Statistics. Instruction manuals. Historic manuals—ICD-10. Part 2s: SuperMICAR data entry instructions. Available from: https://www.cdc.gov/nchs/data/dvs/2011_2s.pdf.
 46. Chamblee RF, Evans MC. TRANSAX: The NCHS system for producing multiple cause-of-death statistics, 1968–78. *National Center for Health Statistics. Vital Health Stat* 1(20). 1986. Available from: https://www.cdc.gov/nchs/data/series/sr_01/sr01_020acc.pdf.
 47. Israel RA, Rosenberg HM, Curtin LR. Analytical potential for multiple cause-of-death data. *Am J Epidemiol* 124(2):161–79. 1986. Available from: <https://aje.oxfordjournals.org/content/124/2/161.full.pdf>.
 48. National Center for Health Statistics. Instruction manual, part 9: ICD-10 cause-of-death lists for tabulating mortality statistics (updated October 2020 to include WHO updates to ICD-10 for data year 2020). 2020. Available from: <https://www.cdc.gov/nchs/data/dvs/Part9InstructionManual2020-508.pdf>.
 49. Hoyert DL, Arias E, Smith BL, Murphy SL, Kochanek KD. Deaths: Final data for 1999. *National Vital Statistics Reports*; vol 49 no 8. Hyattsville, MD: National Center for Health Statistics. 2001. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_08.pdf.
 50. National Center for Health Statistics. Excess deaths associated with COVID-19. 2023. Available from: https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm.
 51. Ahmad FB, Cisewski JA, Anderson RN. Provisional mortality data—United States, 2021. *MMWR Morb Mortal Wkly Rep* 71:597–600. 2021. DOI: <https://dx.doi.org/10.15585/mmwr.mm7117e1>.

52. National Center for Health Statistics, National Vital Statistics System. Computer edits for mortality data, including separate section for fetal deaths effective 2020. NCHS Instruction Manual; part 11. Hyattsville, MD. 2020. Available from: https://www.cdc.gov/nchs/data/dvs/manuals/2020_PT11_B.pdf.
53. National Center for Health Statistics. Instruction manual, part 20: ICD-10 cause-of-death querying, 2013. 2013. Available from: https://www.cdc.gov/nchs/data/dvs/Instruction_Manual_revise20_2013.pdf.
54. Arias E, Eschbach K, Schauman WS, Backlund EL, Sorlie PD. The Hispanic mortality advantage and ethnic misclassification on US death certificates. *Am J Public Health* 100 Suppl 1:S171–7. 2010. Available from: <https://ajph.aphapublications.org/doi/pdf/10.2105/AJPH.2008.135863>.
55. Rosenberg HM, Maurer JD, Sorlie PD, Johnson NJ, MacDorman MF, Hoyert DL, et al. Quality of death rates by race and Hispanic origin: A summary of current research, 1999. National Center for Health Statistics. *Vital Health Stat* 2(128). 1999. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_128.pdf.
56. Sorlie PD, Rogot E, Johnson NJ. Validity of demographic characteristics on the death certificate. *Epidemiology* 3(2):181–4. 1992.
57. Arias E, Xu JQ, Curtin S, Bastian B, Tejada-Vera B. Mortality profile of the non-Hispanic American Indian or Alaska Native population, 2019. National Vital Statistics Reports; vol 70 no 12. Hyattsville, MD: National Center for Health Statistics. 2021. DOI: <https://dx.doi.org/10.15620/cdc:110370>.
58. Mulry M. Summary of accuracy and coverage evaluation for Census 2000. Research Report Series Statistics #2006–3. Washington, DC: U.S. Census Bureau. 2006. Available from: <https://www.census.gov/library/working-papers/2006/adrm/rrs2006-03.html>.
59. Poe GS, Powell-Griner E, McLaughlin JK, Placek PJ, Thompson GB, Robinson K. Comparability of the death certificate and the 1986 National Mortality Followback Survey. National Center for Health Statistics. *Vital Health Stat* 2(118). 1993. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_118.pdf.
60. U.S. Census Bureau. DSSD 2010 census coverage measurement memorandum series 2010–G–01. 2012. Available from: <https://www2.census.gov/programs-surveys/decennial/2010/technical-documentation/methodology/g-series/g01.pdf>.
61. Osterman MJK, Hamilton BE, Martin JA, Driscoll AK, Valenzuela CP. Births: Final data for 2021. National Vital Statistics Reports; vol 72 no 1. Hyattsville, MD: National Center for Health Statistics. 2023. DOI: <https://dx.doi.org/10.15620/cdc:122047>.
62. Sirken MG. Comparison of two methods of constructing abridged life tables by reference to a “standard” table. National Center for Health Statistics. *Vital Health Stat* 2(4). 1966. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_004.pdf.
63. Anderson RN. Method for constructing complete annual U.S. life tables. National Center for Health Statistics. *Vital Health Stat* 2(129). 1999. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_129.pdf.
64. National Center for Health Statistics. U.S. decennial life tables for 1989–91. Methodology of the National and State Life Tables; vol 1 no 2. 1998. Available from: https://www.cdc.gov/nchs/data/lifetables/life89_1_2.pdf.
65. Wei R, Curtin LR, Arias E, Anderson RN. U.S. decennial life tables for 1999–2001: Methodology of the United States life tables. National Vital Statistics Reports; vol 57 no 4. Hyattsville, MD: National Center for Health Statistics. 2008. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_04.pdf.
66. Miniño AM, Murphy SL, Xu JQ, Kochanek KD. Deaths: Final data for 2008. National Vital Statistics Reports; vol 59 no 10. Hyattsville, MD: National Center for Health Statistics. 2011. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_10.pdf.
67. Arias E. United States life tables, 2008. National Vital Statistics Reports; vol 61 no 3. Hyattsville, MD: National Center for Health Statistics. 2012. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_03.pdf.
68. Kochanek KD, Maurer JD, Rosenberg HM. Causes of death contributing to changes in life expectancy: United States, 1984–89. National Center for Health Statistics. *Vital Health Stat* 20(23). 1994. Available from: https://www.cdc.gov/nchs/data/series/sr_20/sr20_023.pdf.
69. Arriaga EE. Changing trends in mortality decline during the last decades. In: Ruzicka L, Wunsch G, Kane P, editors. *Differential mortality: Methodological issues and biosocial factors*. Oxford, England: Clarendon Press. 1989.
70. Arriaga EE. Measuring and explaining the change in life expectancies. *Demography* 21(1):83–96. 1984.
71. Miniño AM, Anderson RN, Fingerhut LA, Boudreault MA, Warner M. Deaths: Injuries, 2002. National Vital Statistics Reports; vol 54 no 10. Hyattsville, MD: National Center for Health Statistics. 2006. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_10.pdf.
72. Sorlie PD, Johnson NJ. Validity of education information on the death certificate. *Epidemiology* 7(4):437–9. 1996.
73. U.S. Census Bureau, American Community Survey, 2021 1-year ACS Public Use Microdata Sample. Available from https://www2.census.gov/programs-surveys/acs/tech_docs/pums/2021ACS_PUMS_User_Guide.pdf.
74. U.S. Census Bureau. International data base. Population by single years of age and sex, 2021. 2022. Available from: https://www.census.gov/data-tools/demo/idb/#/dashboard?COUNTRY_=&COUNTRY_YEAR=2021&COUNTRY_YR_ANIM=2024.
75. Anderson RN, Rosenberg HM. Age standardization of death rates: Implementation of the Year 2000 standard. National Vital Statistics Reports; vol 47 no 3. Hyattsville, MD: National Center for Health Statistics. 1998. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_03.pdf.
76. Brillinger DR. The natural variability of vital rates and associated statistics. *Biometrics* 42(4):693–734. 1986.

77. Fay MP, Feuer EJ. Confidence intervals for directly standardized rates: A method based on the gamma distribution. *Stat Med* 16(7):791–801. 1997.
78. National Center for Health Statistics. Vital statistics of the United States: Mortality, 1999. Technical appendix. 2004. Available from: <https://www.cdc.gov/nchs/data/statab/techap99.pdf>.
79. Schenker N, Gentleman JF. On judging the significance of differences by examining the overlap between confidence intervals. *Am Stat* 55(3):182–6. 2001. Available from: https://www.jstor.org/stable/2685796?seq=1#page_scan_tab_contents.
80. Arnold SF. Mathematical statistics. Englewood Cliffs, NJ: Prentice Hall. 1990.

List of Detailed Tables

1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2021	22	13. Number of deaths, death rate, and age-adjusted death rate for age 15 and older, by marital status and sex: United States, 2021	86
2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2021	24	14. Number of deaths, death rate, and age-adjusted death rate for ages 25–64, by educational attainment and sex: United States, 2021	87
3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2021	28	15. Number of deaths, death rate, and age-adjusted death rate for injury at work for age 15 and older, by Hispanic origin and race and sex: United States, 2021	88
4. Life expectancy at selected ages, by Hispanic origin and race and sex: United States, 2021	30	16. Number of deaths, death rate, and age-adjusted death rate for injury at work, by Hispanic origin and race and sex: United States, 1997–2021	89
5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2021	31	17. Number of deaths, death rate, and age-adjusted death rate for major causes of death: United States, each state, Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, and Northern Marianas, 2021	91
6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021	33	18. Infant, neonatal, and postneonatal mortality rates, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000–2021	98
7. Number of deaths from 113 selected causes, Enterocolitis due to <i>Clostridium difficile</i> , COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021	43	19. Number of infant deaths and infant mortality rate for 130 selected causes, by Hispanic origin and race: United States, 2021	100
8. Death rate for 113 selected causes, Enterocolitis due to <i>Clostridium difficile</i> , COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021	49	20. Number of infant deaths and infant mortality rate, by Hispanic origin and race for the United States, each state, Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, and Northern Marianas, and by sex for the United States, 2021	107
9. Number of deaths from 113 selected causes, Enterocolitis due to <i>Clostridium difficile</i> , COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021	54	21. Number of maternal deaths and maternal mortality rate for selected causes, by Hispanic origin and race: United States, 2021	109
10. Death rate for 113 selected causes, Enterocolitis due to <i>Clostridium difficile</i> , COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021	66	22. Number of deaths, death rate, and age-adjusted death rate for dementia-related causes, by Hispanic origin and race and sex: United States, 1999–2021	111
11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to <i>Clostridium difficile</i> , COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021	75	23. Number of deaths, death rate, and age-adjusted death rate for drug-induced causes, by Hispanic origin and race and sex: United States, 1999–2021	113
12. Number of deaths, death rate, and age-adjusted death rate for injury deaths, by mechanism and intent of death: United States, 2021	84	24. Number of deaths, death rate, and age-adjusted death rate for alcohol-induced causes, by Hispanic origin and race and sex: United States, 1999–2021	115
		25. Number of deaths, death rate, and age-adjusted death rate for injury by firearms, by Hispanic origin and race and sex: United States, 1999–2021	117

Table 1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2021

[Excludes deaths of nonresidents of the United States. Data for specified race or Hispanic-origin groups other than non-Hispanic White and non-Hispanic Black should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys, although misclassification is very minor for the Hispanic and non-Hispanic Asian or Pacific Islander populations; see Technical Notes in this report]

Hispanic origin and race and year	Number			Crude death rate ¹			Age-adjusted death rate ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All origins and races³									
2021.....	3,464,231	1,838,108	1,626,123	1,043.8	1,118.2	970.8	879.7	1,048.0	733.3
2020.....	3,383,729	1,769,884	1,613,845	1,027.0	1,090.8	965.1	835.4	998.3	695.1
2019.....	2,854,838	1,473,823	1,381,015	869.7	911.7	829.0	715.2	846.7	602.7
2018.....	2,839,205	1,458,469	1,380,736	867.8	905.2	831.6	723.6	855.5	611.3
2017.....	2,813,503	1,439,111	1,374,392	863.8	897.2	831.4	731.9	864.5	619.7
2016.....	2,744,248	1,400,232	1,344,016	849.3	880.2	819.3	728.8	861.0	617.5
2015.....	2,712,630	1,373,404	1,339,226	844.0	868.0	820.7	733.1	863.2	624.2
2014.....	2,626,418	1,328,241	1,298,177	823.7	846.4	801.7	724.6	855.1	616.7
2013.....	2,596,993	1,306,034	1,290,959	821.5	839.1	804.4	731.9	863.6	623.5
2012.....	2,543,279	1,273,722	1,269,557	810.2	824.5	796.4	732.8	865.1	624.7
2011.....	2,515,458	1,254,978	1,260,480	807.3	818.7	796.3	741.3	875.3	632.4
2010.....	2,468,435	1,232,432	1,236,003	799.5	812.0	787.4	747.0	887.1	634.9
2000.....	2,403,351	1,177,578	1,225,773	854.0	853.0	855.0	869.0	1,053.8	731.4
1990.....	2,148,463	1,113,417	1,035,046	863.8	918.4	812.0	938.7	1,202.8	750.9
1980.....	1,989,841	1,075,078	914,763	878.3	976.9	785.3	1,039.1	1,348.1	817.9
1970.....	1,921,031	1,078,478	842,553	945.3	1,090.3	807.8	1,222.6	1,542.1	971.4
1960.....	1,711,982	975,648	736,334	954.7	1,104.5	809.2	1,339.2	1,609.0	1,105.3
1950.....	1,452,454	827,749	624,705	963.8	1,106.1	823.5	1,446.0	1,674.2	1,236.0
1940.....	1,417,269	791,003	626,266	1,076.4	1,197.4	954.6	1,785.0	1,976.0	1,599.4
Hispanic⁴									
2021.....	315,664	181,195	134,469	503.9	571.3	434.7	724.7	884.9	582.7
2020.....	305,708	175,585	130,123	498.6	567.8	428.2	723.6	903.8	570.1
2019.....	212,397	117,683	94,714	350.7	384.9	315.7	523.8	633.2	430.7
2018.....	204,719	113,045	91,674	341.9	373.9	309.3	524.1	633.1	431.7
2017.....	197,249	108,579	88,670	334.6	364.6	304.0	524.7	631.8	434.2
2016.....	188,254	103,532	84,722	327.6	356.8	297.7	525.8	631.8	436.4
2015.....	179,457	98,170	81,287	317.1	343.2	290.4	525.3	628.9	438.3
2014.....	169,387	92,474	76,913	305.8	330.1	281.0	523.3	626.8	437.5
2013.....	163,241	88,880	74,361	301.9	323.7	279.4	535.4	639.8	448.6
2012.....	156,419	85,238	71,181	295.0	316.5	272.7	539.1	643.9	452.5
2011.....	149,635	81,887	67,748	287.5	309.7	264.6	540.7	647.3	452.8
2010.....	144,490	79,622	64,868	286.2	310.8	260.9	558.6	677.7	463.4
2000.....	107,254	60,172	47,082	303.8	331.3	274.6	665.7	818.1	546.0
Non-Hispanic, single race⁵									
American Indian and Alaska Native:									
2021.....	26,972	14,724	12,248	1,100.0	1,216.9	986.2	1,109.2	1,282.7	946.6
2020.....	24,725	13,431	11,294	1,016.5	1,123.4	913.2	1,036.2	1,205.9	881.5
2019.....	18,057	9,732	8,325	741.6	812.1	673.3	782.5	901.9	673.3
2018.....	17,790	9,678	8,112	735.9	813.5	660.8	790.8	918.7	673.1
Asian:									
2021.....	92,432	48,386	44,046	469.5	511.7	430.5	461.7	554.9	386.3
2020.....	91,175	47,699	43,476	470.8	515.8	429.6	457.7	557.4	378.5
2019.....	70,532	35,914	34,618	373.1	398.7	349.8	372.8	442.4	317.2
2018.....	68,768	35,089	33,679	367.2	393.4	343.3	381.2	454.1	324.1
Black:									
2021.....	449,764	238,599	211,165	1,074.5	1,185.8	971.4	1,118.0	1,374.0	917.2
2020.....	449,213	237,703	211,510	1,084.3	1,200.0	978.4	1,119.0	1,399.0	905.2
2019.....	346,677	181,363	165,314	842.5	921.8	769.9	884.0	1,092.8	724.9
2018.....	341,408	177,958	163,450	834.7	909.8	765.9	892.6	1,102.8	733.7
Native Hawaiian or Other Pacific Islander:									
2021.....	5,223	2,906	2,317	834.0	917.5	748.6	924.3	1,042.8	806.0
2020.....	4,439	2,489	1,950	723.5	804.6	641.1	821.3	947.9	699.8
2019.....	3,491	1,938	1,553	585.8	646.5	524.4	679.0	769.0	589.5
2018.....	3,277	1,786	1,491	558.9	605.4	511.8	675.7	758.1	597.3

See footnotes at end of table.

Table 1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2021—Con.

[Excludes deaths of nonresidents of the United States. Data for specified race or Hispanic-origin groups other than non-Hispanic White and non-Hispanic Black should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys, although misclassification is very minor for the Hispanic and non-Hispanic Asian or Pacific Islander populations; see Technical Notes in this report]

Hispanic origin and race and year	Number			Crude death rate ¹			Age-adjusted death rate ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Non-Hispanic, single race ⁵ —Con.									
White:									
2021	2,548,809	1,337,385	1,211,424	1,294.9	1,368.8	1,222.1	893.9	1,055.6	751.4
2020	2,484,072	1,278,612	1,205,460	1,262.4	1,317.7	1,208.6	834.7	985.0	703.1
2019	2,183,844	1,115,767	1,068,077	1,106.8	1,146.6	1,068.1	739.9	868.8	627.4
2018	2,182,552	1,108,848	1,073,704	1,104.8	1,138.2	1,072.3	748.7	878.0	636.5

¹Rates are on an annual basis per 100,000 population in specified group; see Technical Notes.

²Age-adjusted rates are per 100,000 U.S. standard population. For method of computation, see Technical Notes.

³Includes origins and races not shown separately; see Technical Notes.

⁴Includes people of Hispanic origin of any race. The Hispanic-origin category is consistent with 1997 Office of Management and Budget (OMB) standards; see Technical Notes.

⁵Only one race was reported on the death certificate. Hispanic-origin and race categories are consistent with 1997 OMB standards; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2021

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	Age group (years)												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
		Number												
Total	3,464,231	19,920	3,816	5,975	38,307	82,274	124,939	216,037	478,171	724,266	829,653	940,780	93	...
Male	1,838,108	10,909	2,105	3,494	27,911	57,910	81,589	135,774	292,961	418,941	432,389	374,055	70	...
Female	1,626,123	9,011	1,711	2,481	10,396	24,364	43,350	80,263	185,210	305,325	397,264	566,725	23	...
Single race ³	3,444,013	18,975	3,651	5,766	37,283	80,827	123,373	214,033	475,074	720,570	826,379	937,989	93	...
Male	1,826,819	10,409	2,008	3,383	27,215	56,905	80,587	134,528	291,125	416,861	430,763	372,965	70	...
Female	1,617,194	8,566	1,643	2,383	10,068	23,922	42,786	79,505	183,949	303,709	395,616	565,024	23	...
Two or more races ⁴	20,218	945	165	209	1,024	1,447	1,566	2,004	3,097	3,696	3,274	2,791	–	...
Male	11,289	500	97	111	696	1,005	1,002	1,246	1,836	2,080	1,626	1,090	–	...
Female	8,929	445	68	98	328	442	564	758	1,261	1,616	1,648	1,701	–	...
Hispanic, total ⁵	315,664	4,453	804	1,347	8,421	14,924	21,252	32,603	51,588	59,784	60,039	60,445	4	...
Male	181,195	2,421	416	750	6,273	11,248	15,205	22,205	33,385	34,921	30,574	23,793	4	...
Female	134,469	2,032	388	597	2,148	3,676	6,047	10,398	18,203	24,863	29,465	36,652	–	...
Hispanic, single race ³	312,795	4,270	776	1,315	8,238	14,677	21,029	32,311	51,189	59,277	59,620	60,089	4	...
Male	179,564	2,329	402	734	6,153	11,063	15,056	22,020	33,149	34,637	30,361	23,656	4	...
Female	133,231	1,941	374	581	2,085	3,614	5,973	10,291	18,040	24,640	29,259	36,433	–	...
American Indian and Alaska Native	1,878	41	9	13	89	156	181	206	345	380	236	222	–	...
Male	1,090	26	5	4	62	106	122	128	195	221	137	84	–	...
Female	788	15	4	9	27	50	59	78	150	159	99	138	–	...
Asian	961	35	5	12	38	49	69	108	118	169	198	160	–	...
Male	515	20	3	6	30	34	47	68	73	89	96	49	–	...
Female	446	15	2	6	8	15	22	40	45	80	102	111	–	...
Black	5,232	257	55	64	297	426	423	514	737	843	817	799	–	...
Male	2,923	137	33	41	217	299	275	322	451	476	410	262	–	...
Female	2,309	120	22	23	80	127	148	192	286	367	407	537	–	...
Native Hawaiian or Other Pacific Islander	372	10	1	5	22	36	51	52	59	62	41	33	–	...
Male	241	7	–	3	18	31	34	40	45	28	20	15	–	...
Female	131	3	1	2	4	5	17	12	14	34	21	18	–	...
White	304,352	3,927	706	1,221	7,792	14,010	20,305	31,431	49,930	57,823	58,328	58,875	4	...
Male	174,795	2,139	361	680	5,826	10,593	14,578	21,462	32,385	33,823	29,698	23,246	4	...
Female	129,557	1,788	345	541	1,966	3,417	5,727	9,969	17,545	24,000	28,630	35,629	–	...
Hispanic, two or more races ⁴	2,869	183	28	32	183	247	223	292	399	507	419	356	–	...
Male	1,631	92	14	16	120	185	149	185	236	284	213	137	–	...
Female	1,238	91	14	16	63	62	74	107	163	223	206	219	–	...

See footnotes at end of table.

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	Age group (years)												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
		Number—Con.												
Non-Hispanic, single race ³	3,123,200	14,495	2,867	4,446	28,983	66,002	102,035	181,074	422,162	659,006	765,229	876,854	47	...
Male	1,642,000	7,973	1,603	2,645	21,019	45,740	65,300	112,056	256,678	380,628	399,474	348,850	34	...
Female	1,481,200	6,522	1,264	1,801	7,964	20,262	36,735	69,018	165,484	278,378	365,755	528,004	13	...
American Indian and Alaska Native	26,972	195	55	83	700	1,876	2,582	3,373	5,344	5,506	4,401	2,856	1	...
Male	14,724	105	34	41	465	1,204	1,574	2,030	3,140	2,886	2,165	1,079	1	...
Female	12,248	90	21	42	235	672	1,008	1,343	2,204	2,620	2,236	1,777	—	...
Asian	92,432	628	120	192	902	1,672	2,783	5,480	10,364	17,702	22,964	29,624	1	...
Male	48,386	352	65	111	622	1,158	1,821	3,453	6,477	10,155	12,175	11,996	1	...
Female	44,046	276	55	81	280	514	962	2,027	3,887	7,547	10,789	17,628	—	...
Black	449,764	5,682	1,064	1,514	10,211	19,442	25,938	41,975	85,176	105,401	83,750	69,602	9	...
Male	238,599	3,153	571	923	7,757	13,588	16,320	25,204	49,838	58,065	39,561	23,613	6	...
Female	211,165	2,529	493	591	2,454	5,854	9,618	16,771	35,338	47,336	44,189	45,989	3	...
Native Hawaiian or Other Pacific Islander	5,223	69	15	15	123	265	475	762	1,039	1,156	816	488	—	...
Male	2,906	46	6	10	91	189	293	460	583	640	389	199	—	...
Female	2,317	23	9	5	32	76	182	302	456	516	427	289	—	...
White	2,548,809	7,921	1,613	2,642	17,047	42,747	70,257	129,484	320,239	529,241	653,298	774,284	36	...
Male	1,337,385	4,317	927	1,560	12,084	29,601	45,292	80,909	196,640	308,882	345,184	311,963	26	...
Female	1,211,424	3,604	686	1,082	4,963	13,146	24,965	48,575	123,599	220,359	308,114	462,321	10	...
Non-Hispanic, two or more races ⁴	17,316	760	136	176	841	1,195	1,341	1,707	2,693	3,184	2,849	2,434	—	...
Male	9,636	407	82	94	576	815	852	1,058	1,599	1,793	1,408	952	—	...
Female	7,680	353	54	82	265	380	489	649	1,094	1,391	1,441	1,482	—	...
Not stated or not classifiable origin ⁶	8,051	212	9	6	62	153	311	653	1,728	2,292	1,536	1,047	42	...
Male	5,277	108	4	5	43	107	232	455	1,299	1,599	933	460	32	...
Female	2,774	104	5	1	19	46	79	198	429	693	603	587	10	...

See footnotes at end of table.

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	Age group (years)												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
								Rate ⁷						
Total ⁸	1,043.8	558.8	25.0	14.3	88.9	180.8	287.9	531.0	1,117.1	2,151.3	5,119.4	15,743.3	...	879.7
Male	1,118.2	598.9	27.0	16.4	126.9	251.2	373.3	668.4	1,397.5	2,640.0	6,031.8	17,190.5	...	1,048.0
Female	970.8	517.0	22.9	12.2	49.3	108.6	201.2	393.9	848.0	1,715.6	4,395.7	14,914.5	...	733.3
Single race ³	1,069.2	570.5	25.6	14.7	90.5	183.3	290.8	535.2	1,124.7	2,163.3	5,147.3	15,834.6	...	885.7
Male	1,145.2	612.3	27.5	16.8	129.4	254.5	376.8	673.3	1,406.6	2,654.7	6,065.4	17,296.1	...	1,055.2
Female	994.7	526.8	23.5	12.4	50.0	110.1	203.4	397.3	853.8	1,725.1	4,419.0	14,998.1	...	738.0
Two or more races ⁵	206.5	396.2	16.8	8.5	53.6	102.7	160.1	287.7	550.2	1,032.6	2,162.8	5,357.7	...	390.1
Male	232.2	411.1	19.3	8.9	71.9	144.0	213.6	375.1	688.5	1,251.5	2,444.9	5,564.6	...	461.5
Female	181.2	380.8	14.1	8.2	34.8	62.2	110.9	208.0	425.7	842.9	1,941.7	5,233.0	...	327.6
Hispanic, total ⁵	503.9	472.4	20.3	12.5	81.7	155.7	236.3	433.1	932.7	1,894.0	4,328.7	11,953.4	...	724.7
Male	571.3	502.9	20.6	13.7	119.3	227.1	324.2	579.3	1,213.4	2,374.1	5,208.9	12,927.3	...	884.9
Female	434.7	440.6	20.0	11.3	42.6	79.3	140.4	281.5	654.9	1,475.0	3,683.0	11,396.2	...	582.7
Hispanic, single race ³	515.8	477.5	20.6	12.8	83.0	158.0	240.1	438.5	944.0	1,912.5	4,368.0	12,048.3	...	732.1
Male	584.6	509.9	21.0	14.0	121.4	230.3	329.3	586.3	1,227.9	2,396.6	5,254.2	13,025.3	...	893.9
Female	445.1	443.7	20.3	11.6	42.9	80.5	142.6	284.8	662.6	1,489.5	3,717.4	11,488.8	...	588.4
American Indian and Alaska Native	98.0	97.1	*	*	29.6	54.8	66.2	93.3	218.0	475.5	823.6	2,534.0	...	163.7
Male	109.5	121.5	*	*	40.4	70.6	82.0	108.3	236.6	556.4	1,055.6	2,548.5	...	189.7
Female	85.6	*	*	*	18.4	37.2	47.4	76.0	197.8	395.6	631.6	2,525.2	...	138.4
Asian	144.6	224.3	*	*	36.3	49.7	76.7	151.9	236.5	621.0	1,827.4	4,604.3	...	264.6
Male	154.6	250.6	*	*	55.8	69.0	102.9	193.2	302.1	720.6	2,140.5	4,032.9	...	297.6
Female	134.6	*	*	*	*	*	49.7	111.5	174.9	538.2	1,606.3	4,911.5	...	232.9
Black	163.4	364.6	20.1	10.0	57.1	86.7	95.9	157.7	309.2	638.3	1,487.4	4,406.1	...	268.3
Male	184.8	381.3	23.8	12.6	81.6	120.2	128.5	210.1	402.7	797.6	1,805.1	4,168.7	...	318.7
Female	142.5	347.4	16.4	7.3	31.5	52.4	65.2	111.2	226.3	506.9	1,263.3	4,532.0	...	224.0
Native Hawaiian or Other Pacific Islander	157.1	*	*	*	61.5	97.8	140.0	203.5	348.4	764.5	1,336.4	3,552.2	...	272.3
Male	197.0	*	*	*	*	158.1	168.2	302.2	531.0	731.1	1,534.9	*	...	338.1
Female	114.5	*	*	*	*	*	*	*	*	794.4	1,189.8	*	...	209.3
White	557.1	516.4	21.8	13.4	86.9	167.2	256.4	467.3	1,006.9	2,027.3	4,602.0	12,595.4	...	773.1
Male	631.4	550.3	21.9	14.6	127.3	244.3	351.8	624.6	1,309.7	2,543.7	5,536.9	13,636.6	...	945.1
Female	480.8	480.9	21.7	12.2	44.8	84.5	151.7	303.0	705.7	1,576.3	3,916.1	11,997.7	...	620.1
Hispanic, two or more races ⁴	143.5	378.0	14.2	6.5	48.3	84.1	94.9	184.3	367.5	889.4	1,898.9	5,131.9	...	320.9
Male	162.8	372.5	*	*	62.4	124.5	127.1	238.6	458.1	1,107.7	2,335.8	5,621.7	...	390.2
Female	124.1	383.8	*	*	33.8	42.8	62.8	132.3	285.6	710.9	1,591.2	4,866.7	...	264.3

See footnotes at end of table.

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	Age group (years)												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
							Rate ⁷ —Con.							
Non-Hispanic, single race ³	1,194.5	596.1	27.3	15.3	92.7	189.7	303.1	555.1	1,146.6	2,181.5	5,209.3	16,163.4	...	903.0
Male	1,274.8	641.4	29.8	17.8	131.7	260.6	388.3	690.7	1,426.2	2,669.6	6,123.0	17,665.6	...	1,072.2
Female	1,116.6	548.7	24.6	12.7	52.1	117.5	218.0	420.9	879.3	1,745.2	4,479.2	15,303.6	...	756.3
American Indian and Alaska Native	1,100.0	722.6	47.5	24.2	193.9	498.0	819.4	1,191.9	1,771.6	2,613.2	4,938.0	10,070.5	...	1,109.2
Male	1,216.9	761.1	57.7	23.4	254.0	630.5	1,002.0	1,456.6	2,182.4	2,963.9	5,450.8	9,996.3	...	1,282.7
Female	986.2	682.4	37.0	24.9	132.1	361.9	637.8	935.0	1,397.0	2,311.9	4,525.7	10,116.1	...	946.6
Asian	469.5	328.1	13.9	8.5	38.4	53.1	87.0	198.5	468.8	1,082.0	3,014.3	9,945.8	...	461.7
Male	511.7	357.1	14.5	9.6	52.7	74.8	119.7	265.5	632.5	1,400.2	3,636.7	10,593.4	...	554.9
Female	430.5	297.4	13.1	7.4	24.0	32.1	57.3	138.9	327.6	828.6	2,526.3	9,548.6	...	386.3
Black	1,074.5	1,129.2	50.0	26.4	171.7	296.8	465.3	827.8	1,681.3	3,100.0	6,003.1	14,435.6	...	1,118.0
Male	1,185.8	1,231.7	53.0	31.7	257.9	416.3	608.6	1,050.9	2,118.4	3,935.6	7,251.9	15,908.6	...	1,374.0
Female	971.4	1,023.0	47.0	20.9	83.5	178.1	332.5	627.6	1,302.4	2,459.4	5,201.3	13,780.5	...	917.2
Native Hawaiian or Other Pacific Islander	834.0	846.7	*	*	143.1	262.2	488.9	1,012.4	1,546.9	2,665.6	4,469.3	8,062.1	...	924.3
Male	917.5	1,096.8	*	*	206.2	364.3	587.2	1,204.5	1,753.5	3,076.6	4,565.7	8,112.5	...	1,042.8
Female	748.6	581.5	*	*	76.6	154.5	385.1	814.5	1,344.3	2,286.6	4,384.9	8,027.8	...	806.0
White	1,294.9	465.4	21.9	12.8	75.7	173.7	287.0	529.9	1,097.8	2,123.9	5,257.7	16,793.9	...	893.9
Male	1,368.8	495.9	24.5	14.7	104.7	236.9	365.1	655.3	1,361.5	2,587.2	6,168.8	18,352.7	...	1,055.6
Female	1,222.1	433.4	19.1	10.8	45.3	108.5	206.8	401.9	839.1	1,697.7	4,511.3	15,883.6	...	751.4
Non-Hispanic, two or more races ⁴	222.3	399.8	17.3	9.0	54.9	107.1	180.5	317.1	592.8	1,058.1	2,203.2	5,390.2	...	406.0
Male	249.6	419.9	20.4	9.4	74.3	148.3	242.1	415.5	743.3	1,275.6	2,453.5	5,550.7	...	479.3
Female	195.4	378.9	14.0	8.6	35.1	67.1	125.1	228.8	457.4	867.4	2,003.4	5,291.9	...	341.5

... Category not applicable.

— Quantity zero.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

¹Death rates for ages younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

²For method of computation, see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Two or more races were reported on the death certificate; see Technical Notes.

⁵Includes people of Hispanic origin of any race; see Technical Notes.

⁶Includes origin not stated or not classifiable; see Technical Notes.

⁷Data for age not stated included in all ages category but not distributed among age groups.

⁸Includes deaths with origin not stated and origin not classifiable; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2021

[Rates are per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Populations used for computing death rates for Hispanic total are postcensal estimates based on the 2010 census estimated as of July 1, 2021; populations used for computing death rates for Mexican, Puerto Rican, Cuban, Central American, South American, Dominican, and Other and unknown Hispanic are estimates based on the 2021 1-year American Community Survey adjusted to control totals. The control totals are 2010-based postcensal estimates for the United States for July 1, 2021; see Technical Notes. Hispanic origin is reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and sex	Age group (years)												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
	Number													
Hispanic	315,664	4,453	804	1,347	8,421	14,924	21,252	32,603	51,588	59,784	60,039	60,445	4	...
Male	181,195	2,421	416	750	6,273	11,248	15,205	22,205	33,385	34,921	30,574	23,793	4	...
Female	134,469	2,032	388	597	2,148	3,676	6,047	10,398	18,203	24,863	29,465	36,652	–	...
Central American	21,205	446	94	139	854	1,396	1,997	2,561	3,612	3,842	3,244	3,020	–	...
Male	12,339	255	51	74	705	1,128	1,517	1,862	2,318	2,096	1,442	891	–	...
Female	8,866	191	43	65	149	268	480	699	1,294	1,746	1,802	2,129	–	...
Cuban	21,675	99	18	25	152	310	405	933	2,330	3,191	5,587	8,625	–	...
Male	11,255	59	8	14	119	231	293	653	1,603	1,977	2,929	3,369	–	...
Female	10,420	40	10	11	33	79	112	280	727	1,214	2,658	5,256	–	...
Dominican	7,598	88	21	36	169	366	431	590	1,048	1,494	1,642	1,713	–	...
Male	4,003	48	11	18	133	268	302	367	596	851	823	586	–	...
Female	3,595	40	10	18	36	98	129	223	452	643	819	1,127	–	...
Mexican	174,828	2,566	426	785	5,094	8,690	12,799	20,272	30,378	34,076	31,238	28,503	1	...
Male	103,658	1,393	219	456	3,745	6,608	9,212	13,910	19,856	20,221	16,197	11,840	1	...
Female	71,170	1,173	207	329	1,349	2,082	3,587	6,362	10,522	13,855	15,041	16,663	–	...
Puerto Rican	35,727	403	75	120	643	1,448	2,195	3,303	5,719	6,903	7,789	7,128	1	...
Male	19,598	220	38	66	445	1,059	1,538	2,214	3,626	3,893	3,848	2,650	1	...
Female	16,129	183	37	54	198	389	657	1,089	2,093	3,010	3,941	4,478	–	...
South American	14,523	119	21	46	219	413	572	918	1,917	2,812	3,628	3,858	–	...
Male	7,590	68	11	20	163	297	397	606	1,199	1,568	1,767	1,494	–	...
Female	6,933	51	10	26	56	116	175	312	718	1,244	1,861	2,364	–	...
Other and unknown Hispanic	40,108	732	149	196	1,290	2,301	2,853	4,026	6,584	7,466	6,911	7,598	2	...
Male	22,752	378	78	102	963	1,657	1,946	2,593	4,187	4,315	3,568	2,963	2	...
Female	17,356	354	71	94	327	644	907	1,433	2,397	3,151	3,343	4,635	–	...

See footnotes at end of table.

Table 3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2021—Con.

[Rates are per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Populations used for computing death rates for Hispanic total are postcensal estimates based on the 2010 census estimated as of July 1, 2021; populations used for computing death rates for Mexican, Puerto Rican, Cuban, Central American, South American, Dominican, and Other and unknown Hispanic are estimates based on the 2021 1-year American Community Survey adjusted to control totals. The control totals are 2010-based postcensal estimates for the United States for July 1, 2021; see Technical Notes. Hispanic origin is reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and sex	All ages	Age group (years)											Age not stated	Age-adjusted rate ²
		Younger than 1 year ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
		Rate ³												
Hispanic	503.9	472.4	20.3	12.5	81.7	155.7	236.3	433.1	932.7	1,894.0	4,328.7	11,953.4	...	724.7
Male	571.3	502.9	20.6	13.7	119.3	227.1	324.2	579.3	1,213.4	2,374.1	5,208.9	12,927.3	...	884.9
Female	434.7	440.6	20.0	11.3	42.6	79.3	140.4	281.5	654.9	1,475.0	3,683.0	11,396.2	...	582.7
Central American	336.2	402.4	21.2	13.1	83.2	133.4	192.2	347.2	734.4	1,580.7	4,047.7	11,588.6	...	646.0
Male	382.0	425.2	22.0	13.8	133.2	207.6	267.4	482.4	974.8	2,047.4	5,127.1	13,500.0	...	823.8
Female	288.1	375.5	20.2	12.4	30.0	53.3	101.7	198.8	509.4	1,241.1	3,464.1	10,940.4	...	514.2
Cuban	903.1	400.4	*	9.7	57.0	90.6	129.5	269.0	704.7	1,549.8	4,196.5	12,811.9	...	636.0
Male	925.1	492.5	*	*	86.8	128.5	180.2	354.9	921.8	2,015.9	5,164.4	13,956.1	...	777.3
Female	880.4	313.8	*	*	25.5	48.7	74.6	171.9	463.8	1,125.8	3,478.1	12,172.3	...	511.6
Dominican	317.4	226.4	12.8	9.4	45.4	92.7	128.8	212.7	468.1	1,091.1	3,051.5	11,543.1	...	502.4
Male	355.0	256.5	*	*	68.9	140.6	194.6	305.1	637.8	1,522.0	4,259.8	12,070.0	...	641.4
Female	284.0	198.4	*	*	20.1	48.0	71.9	141.9	346.5	793.7	2,374.6	11,286.9	...	409.1
Mexican	469.5	467.6	17.9	11.5	77.4	152.4	243.0	463.0	1,018.0	2,056.3	4,531.2	12,709.0	...	767.4
Male	545.6	500.6	18.0	13.1	111.1	223.7	335.5	617.8	1,304.0	2,556.2	5,464.7	13,596.7	...	931.9
Female	390.3	433.6	17.7	9.9	41.9	75.8	142.3	299.1	720.0	1,599.6	3,827.1	12,145.5	...	616.4
Puerto Rican	616.2	503.3	21.2	12.5	73.6	166.2	273.8	483.9	996.5	1,865.3	4,636.4	11,714.1	...	752.1
Male	678.3	553.3	21.0	13.2	99.6	237.8	382.7	659.2	1,311.6	2,275.7	5,611.8	12,538.4	...	918.7
Female	554.5	454.0	21.5	11.7	46.4	91.4	164.4	314.0	703.6	1,512.6	3,963.8	11,275.3	...	610.9
South American	334.0	256.7	9.7	8.1	38.2	65.0	79.7	144.9	384.7	1,007.3	2,791.1	7,971.1	...	400.7
Male	360.9	264.8	*	7.0	57.0	95.7	111.2	200.9	513.1	1,309.2	3,296.0	8,593.6	...	483.6
Female	308.8	246.6	*	9.1	19.5	35.7	48.5	94.1	271.4	780.4	2,436.7	7,622.1	...	336.1
Other and unknown Hispanic	991.3	1,099.8	56.8	28.1	204.2	416.1	537.7	906.8	1,635.8	2,703.2	5,249.7	15,334.0	...	1,111.3
Male	1,102.6	1,126.3	56.6	28.3	290.8	564.4	665.0	1,148.8	2,200.7	3,428.4	6,496.1	17,491.1	...	1,383.6
Female	875.4	1,072.9	57.0	27.9	108.8	248.2	381.3	656.5	1,129.4	2,096.1	4,357.4	14,213.4	...	874.1

... Category not applicable.

— Quantity zero.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

¹Death rates for ages younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

²For method of computation, see Technical Notes.

³Figures for age not stated are included in all ages category but not distributed among age groups.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 4. Life expectancy at selected ages, by Hispanic origin and race and sex: United States, 2021

[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards]

Exact age	Non-Hispanic, single race																	
	Total ¹			Hispanic ^{2,3}			American Indian or Alaska Native ^{2,4}			Asian ^{2,4}			Black ^{2,4}			White ^{2,4}		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
0.....	76.4	73.5	79.3	77.8	74.6	81.1	65.6	62.2	69.2	83.5	81.2	85.6	71.2	67.6	75.0	76.7	74.0	79.5
1.....	75.8	73.0	78.7	77.2	74.0	80.5	65.1	61.7	68.8	82.8	80.5	84.9	71.0	67.4	74.7	76.0	73.4	78.8
5.....	71.9	69.1	74.8	73.3	70.1	76.5	61.2	57.8	64.9	78.8	76.5	80.9	67.1	63.5	70.8	72.1	69.4	74.8
10.....	66.9	64.1	69.8	68.3	65.1	71.5	56.3	52.9	59.9	73.9	71.6	75.9	62.2	58.6	65.9	67.1	64.5	69.9
15.....	62.0	59.2	64.9	63.4	60.2	66.6	51.3	48.0	55.0	68.9	66.6	71.0	57.3	53.7	61.0	62.1	59.5	64.9
20.....	57.1	54.4	60.0	58.5	55.4	61.7	46.7	43.4	50.3	64.0	61.7	66.0	52.7	49.2	56.1	57.3	54.7	60.0
25.....	52.5	49.8	55.2	53.8	50.8	56.9	42.4	39.2	45.8	59.2	56.9	61.1	48.2	44.9	51.4	52.6	50.1	55.2
30.....	47.8	45.4	50.4	49.2	46.3	52.1	38.3	35.3	41.5	54.3	52.1	56.2	43.8	40.7	46.8	47.9	45.6	50.4
35.....	43.3	41.0	45.7	44.6	41.9	47.3	34.5	31.7	37.5	49.5	47.3	51.3	39.5	36.6	42.3	43.4	41.2	45.7
40.....	38.8	36.6	41.1	40.1	37.5	42.6	31.0	28.4	33.7	44.7	42.6	46.4	35.3	32.5	37.8	38.9	36.8	41.1
45.....	34.4	32.3	36.5	35.6	33.2	37.9	27.5	25.1	30.0	39.9	37.9	41.6	31.1	28.6	33.5	34.5	32.5	36.5
50.....	30.1	28.2	32.1	31.2	28.9	33.3	24.5	22.3	26.7	35.2	33.3	36.8	27.1	24.8	29.3	30.2	28.3	32.1
55.....	26.0	24.2	27.8	27.0	24.9	28.9	21.5	19.6	23.4	30.7	28.8	32.1	23.3	21.1	25.3	26.0	24.3	27.8
60.....	22.1	20.5	23.7	23.0	21.1	24.7	18.8	17.2	20.3	26.2	24.6	27.5	19.8	17.8	21.6	22.1	20.5	23.7
65.....	18.4	17.0	19.7	19.3	17.6	20.6	16.3	15.0	17.3	21.9	20.5	23.0	16.7	14.9	18.1	18.4	17.0	19.7
70.....	14.9	13.7	16.0	15.8	14.4	16.8	13.6	12.6	14.4	17.9	16.7	18.7	13.7	12.3	14.9	14.9	13.7	15.9
75.....	11.6	10.6	12.5	12.4	11.3	13.2	11.2	10.4	11.7	14.0	13.0	14.6	11.0	9.8	11.8	11.5	10.6	12.4
80.....	8.7	7.9	9.4	9.4	8.5	9.9	9.1	8.5	9.4	10.5	9.7	10.9	8.5	7.6	9.1	8.7	7.8	9.3
85.....	6.3	5.6	6.7	6.9	6.1	7.1	7.2	6.8	7.3	7.5	6.9	7.7	6.4	5.6	6.7	6.2	5.6	6.6
90.....	4.4	3.9	4.6	4.8	4.3	4.9	5.6	5.4	5.6	5.0	4.7	5.1	4.7	4.2	4.9	4.3	3.8	4.5
95.....	3.0	2.7	3.2	3.4	3.0	3.3	4.5	4.3	4.3	3.4	3.1	3.3	3.4	3.1	3.5	3.0	2.6	3.1
100.....	2.2	1.9	2.2	2.4	2.2	2.3	3.6	3.5	3.4	2.3	2.2	2.2	2.6	2.4	2.6	2.1	1.9	2.2

¹Includes origins and races not shown separately; see Technical Notes in this report.

²Based on death rates adjusted for Hispanic-origin and race misclassification on death certificates; see Technical Notes.

³Includes people of Hispanic origin of any race; see Technical Notes.

⁴Only one race was reported on the death certificate; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2021

Hispanic origin and race and year	Both sexes	Male	Female
All origins and races ¹			
2021 ²	76.4	73.5	79.3
2020 ²	77.0	74.2	79.9
2019 ^{2,3}	78.8	76.3	81.4
2018 ²	78.7	76.2	81.2
2017 ²	78.6	76.1	81.1
2016 ²	78.7	76.2	81.1
2015 ²	78.7	76.3	81.1
2014 ²	78.9	76.5	81.3
2013 ²	78.8	76.4	81.2
2012 ²	78.8	76.4	81.2
2011 ²	78.7	76.3	81.1
2010 ²	78.7	76.2	81.0
2009 ²	78.5	76.0	80.9
2008 ²	78.2	75.6	80.6
2007.....	78.1	75.5	80.6
2006.....	77.8	75.2	80.3
2005.....	77.6	75.0	80.1
2004.....	77.6	75.0	80.1
2003.....	77.2	74.5	79.7
2002.....	77.0	74.4	79.6
2001.....	77.0	74.3	79.5
2000.....	76.8	74.1	79.3
1990.....	75.4	71.8	78.8
1980.....	73.7	70.0	77.4
1970.....	70.8	67.1	74.7
1960.....	69.7	66.6	73.1
1950.....	68.2	65.6	71.1
1940.....	62.9	60.8	65.2
Hispanic ^{4–6}			
2021 ²	77.8	74.6	81.1
2020 ²	77.9	74.6	81.3
2019 ^{2,3}	81.9	79.1	84.4
2018 ²	81.8	79.1	84.3
2017 ²	81.8	79.1	84.3
2016 ²	81.8	79.1	84.3
2015 ^{2,7}	81.9	79.3	84.3
2014 ²	82.1	79.4	84.5
2013 ²	81.9	79.2	84.2
2012 ²	81.9	79.3	84.3
2011 ²	81.8	79.2	84.2
2010 ²	81.7	78.8	84.3
2009 ²	81.1	78.4	83.5
2008 ²	80.8	78.0	83.3
2007.....	80.7	77.8	83.2
2006.....	80.3	77.5	82.9

See footnotes at end of table.

Table 5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2021—Con.

Hispanic origin and race and year	Both sexes	Male	Female
Non-Hispanic, single race ^{6,8}			
American Indian and Alaska Native:			
2021 ²	65.6	62.2	69.2
2020 ²	67.1	63.8	70.7
2019 ^{2,3}	71.8	68.6	75.0
2018 ²	---	---	---
Asian:			
2021 ²	83.5	81.2	85.6
2020 ²	83.6	81.1	85.9
2019 ^{2,3}	85.6	83.5	87.4
2018 ²	---	---	---
Black:			
2021 ²	71.2	67.6	75.0
2020 ²	71.5	67.8	75.4
2019 ^{2,3}	74.8	71.3	78.1
2018 ²	74.7	71.3	78.0
White:			
2021 ²	76.7	74.0	79.5
2020 ²	77.4	74.8	80.1
2019 ^{2,3}	78.8	76.3	81.3
2018 ²	78.6	76.2	81.1

--- Data not available.

¹Includes origins and races not shown separately; see Technical Notes in this report.

²Life expectancies for 2008–2021 were calculated using newly revised life table methodology described in the Technical Notes.

³Life expectancies were updated using final data; therefore, data may differ from preliminary data previously published; see Technical Notes.

⁴Includes people of Hispanic origin of any race. The Hispanic-origin category is consistent with 1997 Office of Management and Budget (OMB) standards; see Technical Notes.

⁵Data by Hispanic origin available beginning with data year 2006.

⁶Based on death rates adjusted for Hispanic-origin and race misclassification on death certificates; see Technical Notes.

⁷Life expectancies were revised using updated Medicare data; as a result, data may differ from those previously published; see Technical Notes.

⁸Only one race was reported on the death certificate. Hispanic-origin and race categories are consistent with 1997 OMB standards; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
All causes													
2021.....	1,043.8	558.8	25.0	14.3	88.9	180.8	287.9	531.0	1,117.1	2,151.3	5,119.4	15,743.3	879.7
2020.....	1,027.0	524.3	22.7	13.7	84.2	159.5	248.0	473.5	1,038.9	2,072.3	4,997.0	15,210.9	835.4
2019.....	869.7	553.0	23.3	13.4	69.7	128.8	199.2	392.4	883.3	1,764.6	4,308.3	13,228.6	715.2
2018.....	867.8	557.8	24.0	13.3	70.2	128.8	194.7	395.9	886.7	1,783.3	4,386.1	13,450.7	723.6
2017.....	863.8	567.0	24.3	13.6	74.0	132.8	195.2	401.5	885.8	1,790.9	4,472.6	13,573.6	731.9
2016.....	849.3	583.4	25.3	13.4	74.9	129.0	192.2	405.5	883.8	1,788.6	4,474.8	13,392.1	728.8
2015.....	844.0	589.6	24.9	13.2	69.5	116.7	180.1	404.0	875.3	1,796.8	4,579.2	13,673.9	733.1
2014.....	823.7	588.0	24.0	12.7	65.5	108.4	175.2	404.8	870.3	1,786.3	4,564.2	13,407.9	724.6
2013.....	821.5	594.7	25.5	13.0	64.8	106.1	172.0	406.1	860.0	1,802.1	4,648.1	13,660.4	731.9
2012.....	810.2	599.3	26.3	12.6	66.4	105.4	170.7	405.4	854.2	1,802.5	4,674.5	13,678.6	732.8
2011.....	807.3	600.1	26.3	13.2	67.7	104.7	172.0	409.8	849.4	1,846.2	4,753.0	13,779.3	741.3
2010.....	799.5	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3	747.0
2009.....	794.5	659.7	27.4	13.8	69.8	104.4	180.0	418.1	856.7	1,888.7	4,820.2	13,660.1	749.6
2008.....	812.9	678.9	29.3	13.9	74.2	105.1	181.0	419.6	867.1	1,958.4	4,998.1	14,332.4	774.9
2007.....	804.6	702.5	29.4	15.2	78.8	107.2	186.0	420.3	866.7	1,976.0	4,987.1	14,160.9	775.3
2006.....	813.1	705.8	29.1	15.2	81.4	109.0	192.0	427.5	881.3	2,031.4	5,096.1	14,426.7	791.8
2005.....	828.4	710.2	29.9	16.3	80.7	106.8	194.9	431.9	898.5	2,109.7	5,251.8	14,982.4	815.0
2004.....	818.8	695.9	30.3	16.7	79.7	104.1	194.9	426.8	903.2	2,141.0	5,267.4	14,777.6	813.7
2003.....	843.9	704.9	31.8	16.9	81.1	105.2	202.6	433.1	937.3	2,235.0	5,451.3	15,401.4	843.5
2002.....	849.5	709.5	31.4	17.4	80.9	105.1	204.2	431.0	948.7	2,300.3	5,543.8	15,589.5	855.9
2001.....	848.0	687.0	33.4	17.2	80.2	105.6	203.5	426.7	972.5	2,344.2	5,573.7	15,432.6	858.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4	869.0
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6	875.6
Diseases of heart (I00-I09,I11,I13,I20-I51)													
2021.....	209.6	7.4	0.8	0.5	2.2	9.1	29.4	84.9	208.7	416.6	1,044.2	4,078.7	173.8
2020.....	211.5	6.5	0.7	0.4	2.0	8.6	28.9	84.6	208.8	414.2	1,017.2	3,822.1	168.2
2019.....	200.8	7.1	0.8	0.4	2.0	7.6	25.2	76.2	190.4	388.8	991.2	3,798.3	161.5
2018.....	200.3	7.5	0.7	0.4	2.1	7.8	25.5	77.4	191.7	392.4	1,008.3	3,844.8	163.6
2017.....	198.8	7.7	0.8	0.4	2.1	8.1	25.4	77.1	190.7	392.9	1,028.4	3,882.9	165.0
2016.....	196.6	7.4	0.7	0.5	2.2	7.7	25.9	79.5	189.6	392.5	1,037.1	3,873.4	165.5
2015.....	197.2	7.3	0.9	0.5	2.3	8.0	25.6	79.3	188.1	389.5	1,071.6	3,986.5	168.5
2014.....	192.7	8.0	0.9	0.5	2.2	7.7	25.6	80.1	185.8	385.2	1,070.2	3,920.9	167.0
2013.....	193.3	7.8	1.1	0.4	2.1	7.6	25.6	80.3	184.6	390.3	1,095.1	4,013.9	169.8
2012.....	191.0	8.5	1.0	0.4	2.2	7.6	25.9	79.7	184.6	388.3	1,103.7	4,046.1	170.5
2011.....	191.5	7.7	1.0	0.5	2.3	7.9	26.2	80.7	183.2	399.0	1,134.7	4,111.6	173.7
2010.....	193.6	8.3	1.0	0.5	2.4	7.8	25.8	81.6	186.6	409.2	1,172.0	4,285.2	179.1
2009.....	195.4	9.6	0.9	0.5	2.4	7.8	26.7	82.3	190.0	422.8	1,210.8	4,316.9	182.8
2008.....	202.8	9.6	1.2	0.6	2.5	8.1	26.9	85.2	195.3	441.4	1,271.7	4,598.4	192.1
2007.....	204.5	10.2	1.1	0.6	2.5	8.1	27.7	85.2	197.8	454.8	1,308.6	4,668.1	196.1
2006.....	211.7	8.6	1.0	0.6	2.5	8.4	28.5	88.0	205.1	483.0	1,378.0	4,877.6	205.5
2005.....	220.7	8.9	0.9	0.6	2.6	8.3	29.2	89.7	212.8	512.3	1,458.5	5,188.3	216.8
2004.....	222.8	10.5	1.2	0.6	2.5	8.1	29.5	90.2	217.1	535.7	1,504.1	5,233.8	221.6
2003.....	236.1	11.0	1.2	0.6	2.7	8.3	30.8	92.4	232.3	579.8	1,607.7	5,570.7	236.3
2002.....	242.3	12.7	1.1	0.6	2.5	8.0	30.7	93.9	240.5	612.0	1,673.2	5,726.3	244.6
2001.....	245.7	11.9	1.5	0.7	2.5	8.0	29.6	92.4	248.9	632.6	1,723.0	5,784.1	249.5
2000.....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1	257.6
1999.....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0	266.5

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)										Age- adjusted rate ³	
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84		85 and older
Malignant neoplasms (C00–C97)													
2021.....	182.4	1.5	1.8	1.9	3.1	7.9	25.8	82.5	252.4	532.1	1,017.2	1,712.9	146.6
2020.....	182.8	1.4	2.0	1.9	3.1	7.8	25.5	85.7	260.0	539.1	990.0	1,538.1	144.1
2019.....	182.7	1.5	1.8	1.9	3.3	7.8	25.7	87.1	263.3	543.3	1,005.9	1,571.0	146.2
2018.....	183.2	1.3	2.0	2.1	3.2	8.1	25.8	89.6	269.6	554.4	1,031.5	1,577.7	149.1
2017.....	183.9	1.4	2.0	2.1	3.2	8.0	26.7	92.7	273.4	567.5	1,060.2	1,600.3	152.5
2016.....	185.1	1.7	2.4	2.1	3.3	8.5	26.9	96.5	280.6	578.3	1,081.7	1,620.3	155.8
2015.....	185.4	1.3	2.2	2.1	3.4	8.4	26.9	99.7	284.1	594.3	1,100.8	1,628.6	158.5
2014.....	185.6	1.3	2.0	2.1	3.6	8.3	27.8	103.2	287.6	603.1	1,125.9	1,632.9	161.2
2013.....	185.0	1.6	2.1	2.2	3.4	8.6	28.1	105.5	288.2	616.9	1,139.4	1,635.4	163.2
2012.....	185.6	1.6	2.4	2.2	3.6	8.7	28.0	108.5	293.2	632.2	1,161.7	1,658.9	166.5
2011.....	185.1	1.8	2.2	2.1	3.7	8.4	28.8	109.3	295.8	647.6	1,179.1	1,676.2	169.0
2010.....	186.2	1.6	2.1	2.2	3.7	8.8	28.8	111.6	300.1	666.1	1,202.2	1,729.5	172.8
2009.....	185.0	1.8	2.2	2.2	3.8	9.0	30.2	112.8	301.7	668.2	1,213.0	1,699.3	173.5
2008.....	186.0	1.7	2.4	2.2	3.8	8.8	30.1	113.4	304.7	688.4	1,230.9	1,724.6	176.4
2007.....	186.9	1.7	2.3	2.4	3.8	8.7	31.0	114.2	311.4	702.9	1,250.1	1,739.4	179.3
2006.....	187.6	1.9	2.4	2.2	3.8	9.3	32.2	116.3	317.7	716.3	1,259.2	1,748.3	181.8
2005.....	189.3	1.9	2.4	2.5	4.0	9.2	33.5	118.6	323.9	733.2	1,272.8	1,778.2	185.1
2004.....	189.2	1.8	2.5	2.5	4.1	9.3	33.6	119.0	330.8	746.8	1,278.6	1,767.4	186.8
2003.....	192.0	1.9	2.5	2.6	4.0	9.5	35.1	122.1	341.6	763.5	1,299.7	1,792.3	190.9
2002.....	193.7	1.9	2.6	2.6	4.2	9.8	36.0	124.1	349.7	787.2	1,308.8	1,812.4	194.3
2001.....	194.3	1.6	2.7	2.4	4.2	10.1	36.8	125.8	359.4	799.7	1,313.7	1,802.9	196.5
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4	199.6
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8	200.8
COVID-19 (U07.1) ⁴													
2021.....	125.6	2.6	0.4	0.3	3.3	13.5	36.9	90.6	172.2	305.1	609.7	1,354.4	104.1
2020.....	106.5	0.9	*	0.1	1.2	4.9	14.4	42.0	99.3	234.3	589.8	1,645.0	85.0
Accidents (unintentional injuries) (V01–X59,Y85–Y86)													
2021.....	67.8	36.6	8.5	4.2	36.7	75.7	84.0	77.2	78.2	64.7	126.8	446.0	64.7
2020.....	61.0	32.0	7.4	3.8	35.5	68.0	73.7	68.9	68.2	57.2	113.3	383.6	57.6
2019.....	52.7	33.5	7.3	3.6	27.5	53.4	57.8	57.1	58.6	54.5	115.5	377.4	49.3
2018.....	51.1	30.4	7.7	3.5	28.0	53.9	54.9	55.4	56.0	52.3	111.3	368.6	48.0
2017.....	52.2	33.4	7.9	3.8	31.1	56.6	55.8	57.7	55.7	50.7	113.3	374.9	49.4
2016.....	49.9	30.7	7.9	4.0	31.9	53.7	51.8	54.6	52.7	49.1	110.7	365.7	47.4
2015.....	45.6	32.5	7.8	3.7	28.5	44.8	43.9	49.8	47.7	47.0	111.5	364.5	43.2
2014.....	42.6	29.4	7.6	3.6	26.8	39.8	39.6	47.4	44.9	45.1	108.7	349.1	40.5
2013.....	41.3	29.3	8.3	3.7	26.4	37.8	38.0	46.5	43.4	43.5	107.4	340.0	39.4
2012.....	40.7	29.6	8.4	3.8	27.1	37.5	37.1	46.1	41.0	44.0	107.8	336.9	39.1
2011.....	40.6	29.1	8.5	4.0	28.2	37.1	37.5	46.4	39.8	44.5	107.0	333.8	39.1
2010.....	39.1	28.1	8.6	4.0	28.3	35.5	36.0	43.7	38.4	43.3	106.1	328.4	38.0
2009.....	38.5	29.5	9.0	4.1	28.6	34.5	36.4	44.5	36.5	42.1	103.5	310.9	37.5
2008.....	40.1	31.8	9.1	4.6	32.5	36.3	38.1	45.8	37.4	43.9	105.7	318.3	39.2
2007.....	41.1	31.0	9.9	5.4	36.8	37.7	39.6	46.2	36.8	44.4	105.0	313.6	40.4
2006.....	40.8	28.4	10.1	5.6	37.9	38.0	40.5	45.5	35.8	43.8	104.7	299.2	40.2
2005.....	39.9	27.0	10.5	5.9	37.1	35.7	38.9	43.2	35.4	45.7	106.0	303.5	39.5
2004.....	38.3	26.2	10.4	6.5	36.8	33.2	37.6	40.7	32.9	43.5	103.6	295.8	38.1
2003.....	37.7	23.8	11.0	6.4	36.9	32.0	38.0	38.8	32.7	43.7	101.6	294.3	37.6
2002.....	37.1	23.9	10.6	6.6	37.7	31.9	37.4	36.7	31.3	44.0	101.1	289.6	37.1
2001.....	35.6	24.3	11.2	6.9	35.8	30.0	35.4	33.9	30.5	42.6	100.7	282.2	35.7
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5	34.9
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4	35.3

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Cerebrovascular diseases (I60–I69)													
2021.....	49.1	2.7	0.4	0.2	0.4	1.4	5.0	14.1	34.2	84.1	274.8	1,111.1	41.1
2020.....	48.6	2.8	0.4	0.2	0.4	1.3	4.8	14.1	33.4	81.0	262.9	1,017.9	38.8
2019.....	45.7	2.7	0.3	0.2	0.4	1.3	4.2	12.6	30.5	76.4	254.2	977.3	37.0
2018.....	45.2	2.5	0.3	0.2	0.3	1.2	4.1	12.3	30.3	76.8	256.0	984.3	37.1
2017.....	44.9	2.5	0.4	0.2	0.4	1.3	4.4	12.3	30.3	76.4	263.1	993.5	37.6
2016.....	44.0	3.1	0.3	0.2	0.3	1.3	4.6	12.5	29.7	76.0	265.5	972.9	37.3
2015.....	43.7	2.2	0.3	0.2	0.4	1.3	4.4	12.3	29.6	75.5	273.0	975.8	37.6
2014.....	41.7	2.4	0.2	0.2	0.4	1.3	4.3	12.3	29.3	74.5	265.7	929.7	36.5
2013.....	40.8	2.7	0.2	0.2	0.3	1.2	4.2	12.4	28.9	74.2	268.9	906.0	36.2
2012.....	40.9	2.6	0.3	0.2	0.4	1.3	4.3	12.8	28.7	75.7	272.2	931.2	36.9
2011.....	41.4	3.4	0.3	0.2	0.4	1.3	4.2	12.8	29.4	78.2	285.4	943.7	37.9
2010.....	41.9	3.3	0.3	0.2	0.4	1.3	4.6	13.1	29.3	81.7	288.3	993.8	39.1
2009.....	42.0	3.7	0.3	0.2	0.4	1.3	4.6	13.7	29.7	82.8	294.9	992.2	39.6
2008.....	44.1	3.4	0.4	0.2	0.4	1.3	4.8	13.7	30.6	87.3	313.3	1,071.0	42.1
2007.....	45.1	3.2	0.3	0.2	0.5	1.3	5.0	14.5	31.7	91.4	320.8	1,110.7	43.5
2006.....	46.0	3.5	0.3	0.2	0.5	1.3	5.1	14.6	32.9	94.9	333.9	1,131.7	44.8
2005.....	48.6	3.1	0.4	0.2	0.5	1.4	5.2	15.0	32.7	99.8	358.4	1,239.7	48.0
2004.....	51.3	3.2	0.3	0.2	0.5	1.4	5.4	14.8	34.0	106.6	385.6	1,331.9	51.2
2003.....	54.4	2.5	0.3	0.2	0.5	1.5	5.6	15.0	35.5	111.9	409.8	1,446.0	54.6
2002.....	56.6	3.0	0.3	0.2	0.4	1.4	5.4	15.1	37.1	119.6	430.0	1,520.1	57.2
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.0	38.3	122.9	443.3	1,532.0	58.4
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2	60.9
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8	61.6
Chronic lower respiratory diseases (J40–J47)													
2021.....	42.9	*	0.2	0.2	0.4	0.8	1.7	7.8	41.2	112.7	285.9	600.6	34.7
2020.....	46.3	*	*	0.4	0.5	0.9	1.9	8.8	44.4	118.5	300.1	612.6	36.4
2019.....	47.8	*	0.2	0.4	0.4	0.8	1.6	8.8	44.2	124.6	318.0	654.3	38.2
2018.....	48.7	*	0.3	0.3	0.4	0.7	1.6	9.1	44.5	128.6	335.7	682.4	39.7
2017.....	49.2	*	0.2	0.3	0.4	0.7	1.7	9.4	44.4	133.8	347.6	700.6	40.9
2016.....	47.8	0.7	0.3	0.3	0.5	0.8	1.7	10.1	43.0	134.1	347.2	676.9	40.6
2015.....	48.2	0.7	0.3	0.4	0.5	0.7	1.7	10.1	42.7	136.6	357.9	705.1	41.6
2014.....	46.1	*	0.3	0.3	0.4	0.8	1.9	10.1	41.2	134.9	349.0	670.5	40.5
2013.....	47.2	0.6	0.4	0.4	0.4	0.7	1.9	10.6	40.5	141.2	367.0	699.3	42.1
2012.....	45.7	0.5	0.3	0.3	0.3	0.7	1.8	10.2	39.4	140.0	364.0	687.8	41.5
2011.....	45.9	0.8	0.3	0.3	0.4	0.6	1.8	10.4	39.5	144.3	374.9	697.9	42.5
2010.....	44.7	0.9	0.3	0.3	0.3	0.7	1.7	9.9	39.0	146.3	369.9	690.7	42.2
2009.....	44.8	0.7	0.4	0.3	0.4	0.7	1.8	10.4	40.0	147.5	376.4	684.9	42.7
2008.....	46.4	0.8	0.3	0.3	0.4	0.6	1.9	9.9	41.1	155.9	395.4	722.7	44.7
2007.....	42.5	1.0	0.4	0.3	0.3	0.7	1.9	9.5	38.6	145.5	367.1	652.0	41.4
2006.....	41.8	0.7	0.3	0.3	0.4	0.6	1.9	9.1	38.8	147.0	362.0	641.3	41.0
2005.....	44.3	0.8	0.4	0.3	0.3	0.7	2.0	9.4	41.6	158.4	385.0	691.9	43.9
2004.....	41.7	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.1	152.1	366.2	643.2	41.6
2003.....	43.6	0.8	0.4	0.3	0.5	0.7	2.2	8.7	43.1	161.7	382.2	670.2	43.7
2002.....	43.4	1.0	0.4	0.3	0.5	0.8	2.3	8.7	42.2	162.0	385.8	670.3	43.9
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.4	44.5	167.3	379.3	658.3	43.9
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6	44.2
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0	45.4

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Alzheimer disease (G30)													
2021.....	36.0	*	*	*	*	*	*	0.3	3.2	26.4	214.3	1,243.6	31.0
2020.....	40.7	*	*	*	*	*	*	0.2	3.3	28.6	229.3	1,287.3	32.4
2019.....	37.0	*	*	*	*	*	*	0.3	3.0	24.9	210.2	1,191.3	29.8
2018.....	37.3	*	*	*	*	*	*	0.3	2.9	24.7	213.9	1,225.3	30.5
2017.....	37.3	*	*	*	*	*	*	0.2	2.8	24.5	219.7	1,244.7	31.0
2016.....	35.9	*	*	*	*	*	*	0.2	2.7	23.6	214.1	1,216.9	30.3
2015.....	34.4	*	*	*	*	*	*	0.2	2.4	22.4	211.9	1,174.2	29.4
2014.....	29.3	*	*	*	*	*	*	0.2	2.1	19.6	185.6	1,006.8	25.4
2013.....	26.8	*	*	*	*	*	*	0.2	2.2	18.1	171.6	929.5	23.5
2012.....	26.6	*	*	*	*	*	*	0.2	2.2	17.9	175.4	936.1	23.8
2011.....	27.3	*	*	*	*	*	*	0.2	2.2	19.2	183.9	967.1	24.7
2010.....	27.0	*	*	*	*	*	*	0.3	2.1	19.8	184.5	987.1	25.1
2009.....	25.8	*	*	*	*	*	*	0.2	2.0	19.4	179.1	945.3	24.2
2008.....	27.1	*	*	*	*	*	*	0.2	2.2	21.1	192.5	1,002.2	25.8
2007.....	24.8	*	*	*	*	*	*	0.2	2.2	20.2	175.8	928.7	23.8
2006.....	24.3	*	*	*	*	*	*	0.2	2.1	19.9	175.0	923.4	23.7
2005.....	24.2	*	*	*	*	*	*	0.2	2.1	20.2	177.0	935.5	24.0
2004.....	22.5	*	*	*	*	*	*	0.2	1.8	19.5	168.5	875.3	22.6
2003.....	21.9	*	*	*	*	*	*	0.2	2.0	20.7	164.1	846.8	22.1
2002.....	20.5	*	*	*	*	*	*	0.1	1.9	19.6	157.7	790.9	20.8
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.6	147.2	725.4	19.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7	18.1
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3	16.5
Diabetes mellitus (E10–E14)													
2021.....	31.1	*	*	0.1	0.8	2.8	6.8	18.7	43.5	83.0	160.4	309.9	25.4
2020.....	31.0	*	*	0.1	0.7	2.5	6.9	18.7	42.5	83.6	154.7	293.4	24.8
2019.....	26.7	*	*	0.1	0.6	1.9	5.3	15.5	36.5	73.3	138.9	259.5	21.6
2018.....	26.0	*	*	0.1	0.6	1.8	5.5	15.4	35.3	72.1	137.5	260.4	21.4
2017.....	25.7	*	*	0.1	0.6	1.8	5.2	15.1	35.5	71.9	140.8	262.4	21.5
2016.....	24.8	*	*	0.1	0.5	1.8	5.1	14.6	34.4	69.9	137.9	263.6	21.0
2015.....	24.7	*	*	0.1	0.4	1.8	4.9	14.4	34.7	70.6	143.0	267.0	21.3
2014.....	24.0	*	*	0.1	0.4	1.6	4.9	13.9	33.3	69.0	141.8	268.6	20.9
2013.....	23.9	*	*	0.1	0.4	1.6	4.8	13.5	33.2	68.5	145.7	279.5	21.2
2012.....	23.6	*	*	0.1	0.4	1.5	4.6	13.0	32.5	69.7	145.8	285.7	21.2
2011.....	23.7	*	*	0.1	0.4	1.6	4.5	13.4	33.3	72.0	148.8	289.5	21.6
2010.....	22.4	*	*	0.1	0.4	1.5	4.4	12.5	32.0	67.6	144.1	285.5	20.8
2009.....	22.4	*	*	0.1	0.4	1.5	4.5	12.8	32.1	69.6	145.8	282.6	21.0
2008.....	23.2	*	*	0.1	0.5	1.4	4.4	12.6	33.3	74.7	153.2	298.9	22.0
2007.....	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.1	76.7	161.9	302.2	22.8
2006.....	24.3	*	*	0.1	0.4	1.7	4.8	13.1	35.8	80.6	166.2	310.4	23.6
2005.....	25.4	*	*	0.1	0.5	1.6	4.7	13.4	36.9	85.7	177.0	338.8	24.9
2004.....	25.0	*	*	0.1	0.4	1.5	4.6	13.4	36.8	86.2	176.6	328.2	24.7
2003.....	25.6	*	*	0.1	0.4	1.7	4.6	13.9	38.3	90.0	180.7	335.1	25.5
2002.....	25.5	*	*	0.1	0.4	1.6	4.8	13.7	37.5	90.9	182.4	337.0	25.6
2001.....	25.0	*	*	0.1	0.4	1.5	4.3	13.6	38.1	91.0	181.1	328.6	25.4
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7	25.0
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2	25.0

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Chronic liver disease and cirrhosis (K70,K73–K74)													
2021.....	17.0	*	*	*	0.1	4.0	13.4	25.8	41.3	38.4	36.4	31.0	14.5
2020.....	15.7	*	*	*	0.1	3.5	11.7	23.5	38.1	36.5	34.5	27.0	13.3
2019.....	13.5	*	*	*	0.1	2.4	8.2	19.8	33.9	33.1	32.2	26.5	11.3
2018.....	13.1	*	*	*	0.1	2.2	7.5	19.6	33.0	32.5	32.5	25.5	11.1
2017.....	12.8	*	*	*	0.1	2.0	7.3	19.6	32.7	31.7	31.3	26.8	10.9
2016.....	12.5	*	*	*	0.1	2.1	7.0	19.5	32.4	30.7	31.9	24.5	10.7
2015.....	12.5	*	*	*	0.1	1.9	7.0	20.5	32.5	30.5	31.9	25.1	10.8
2014.....	12.0	*	*	*	0.1	1.7	6.4	19.9	31.9	29.6	30.4	23.4	10.4
2013.....	11.5	*	*	*	0.1	1.6	6.2	20.1	30.4	28.1	29.9	23.0	10.2
2012.....	11.1	*	*	*	0.1	1.4	6.1	20.1	29.1	27.6	29.3	21.4	9.9
2011.....	10.8	*	*	*	0.1	1.2	6.0	19.8	28.2	26.3	29.3	22.1	9.7
2010.....	10.3	*	*	*	0.1	1.2	5.9	19.2	26.8	26.3	27.7	21.8	9.4
2009.....	10.0	*	*	*	0.1	1.1	6.0	18.7	25.9	25.4	27.2	21.1	9.1
2008.....	9.9	*	*	*	0.1	1.1	6.1	18.5	25.0	26.3	28.0	21.9	9.2
2007.....	9.7	*	*	*	0.1	1.0	6.0	18.7	24.2	26.2	28.2	21.7	9.1
2006.....	9.2	*	*	*	0.1	0.8	5.9	17.8	22.6	25.6	28.9	21.1	8.8
2005.....	9.3	*	*	*	0.1	0.8	6.2	17.7	23.3	26.8	28.9	21.3	8.9
2004.....	9.2	*	*	*	*	0.8	6.4	18.0	22.4	27.4	28.7	21.1	9.0
2003.....	9.5	*	*	*	*	0.9	6.8	18.3	22.9	29.2	29.9	21.2	9.3
2002.....	9.5	*	*	*	0.1	1.0	7.1	18.0	22.8	29.3	31.3	22.5	9.4
2001.....	9.5	*	*	*	0.1	1.0	7.4	18.4	22.9	29.8	30.2	22.7	9.5
2000.....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1	9.5
1999.....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2	9.6
Nephritis, nephrotic syndrome and nephrosis (N00–N07,N17–N19,N25–N27)													
2021.....	16.4	1.3	*	*	0.1	0.8	2.2	6.2	14.9	36.4	96.1	271.1	13.6
2020.....	15.9	1.2	*	*	0.2	0.7	2.0	5.8	14.7	35.2	89.3	248.1	12.7
2019.....	15.7	1.6	*	*	0.2	0.7	1.8	5.6	13.8	34.7	92.2	250.9	12.7
2018.....	15.7	2.0	*	*	0.1	0.6	1.8	5.4	13.6	35.6	94.3	257.9	12.9
2017.....	15.5	2.0	*	*	0.1	0.6	1.7	5.2	13.5	34.7	95.8	267.1	13.0
2016.....	15.5	1.6	*	*	0.1	0.6	1.8	5.0	13.6	34.6	98.1	270.1	13.1
2015.....	15.5	2.1	*	*	0.1	0.6	1.7	4.9	13.3	35.1	99.7	281.8	13.4
2014.....	15.1	2.3	*	*	0.2	0.5	1.7	4.7	12.6	34.3	98.6	282.4	13.2
2013.....	14.9	2.2	*	*	0.1	0.6	1.5	4.6	12.6	33.8	99.0	285.4	13.2
2012.....	14.5	2.1	*	*	0.2	0.5	1.6	4.7	12.3	33.3	99.9	280.0	13.1
2011.....	14.6	1.9	*	*	0.2	0.5	1.6	4.4	12.5	34.2	101.4	292.1	13.4
2010.....	16.3	2.7	*	0.1	0.2	0.6	1.8	4.9	13.9	39.3	115.7	333.8	15.3
2009.....	16.0	2.8	*	*	0.2	0.7	2.0	5.2	13.5	38.7	115.1	321.4	15.1
2008.....	15.9	3.5	*	*	0.2	0.6	1.8	5.0	14.1	39.9	113.3	325.6	15.1
2007.....	15.4	3.5	0.1	0.1	0.2	0.7	1.8	5.1	13.4	39.4	112.4	317.9	14.9
2006.....	15.2	4.0	*	*	0.2	0.7	1.8	5.2	13.7	38.8	111.0	316.2	14.8
2005.....	14.9	4.0	*	0.1	0.2	0.7	1.7	4.8	13.5	38.8	110.2	313.1	14.7
2004.....	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.5	38.1	108.2	306.4	14.5
2003.....	14.6	4.6	*	0.1	0.2	0.7	1.8	4.9	13.6	39.7	109.3	309.3	14.7
2002.....	14.2	4.4	*	0.1	0.2	0.7	1.7	4.7	12.9	39.0	108.9	303.4	14.4
2001.....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.1	40.0	104.0	293.8	14.1
2000.....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8	13.5
1999.....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9	13.0

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)													
2021.....	14.5	...	...	1.5	15.2	19.5	18.1	18.2	17.0	15.3	19.6	22.4	14.1
2020.....	14.0	...	...	1.5	14.2	18.4	17.4	18.0	16.9	14.5	18.4	20.9	13.5
2019.....	14.5	...	...	1.3	13.9	17.5	18.1	19.6	19.4	15.5	18.6	20.1	13.9
2018.....	14.8	...	...	1.5	14.5	17.6	18.2	20.0	20.2	16.3	18.7	19.1	14.2
2017.....	14.5	...	...	1.3	14.5	17.5	17.9	20.2	19.0	15.6	18.0	20.1	14.0
2016.....	13.9	...	...	1.1	13.2	16.5	17.4	19.7	18.7	15.4	18.2	19.0	13.5
2015.....	13.7	...	...	1.0	12.5	15.7	17.1	20.3	18.9	15.2	17.9	19.4	13.3
2014.....	13.4	...	...	1.0	11.6	15.1	16.6	20.2	18.8	15.6	17.5	19.3	13.0
2013.....	13.0	...	...	1.0	11.1	14.8	16.2	19.7	18.1	15.0	17.1	18.6	12.6
2012.....	12.9	...	...	0.8	11.1	14.7	16.7	20.0	18.0	14.0	16.8	17.8	12.6
2011.....	12.7	...	...	0.7	11.0	14.6	16.2	19.8	17.1	14.1	16.5	16.9	12.3
2010.....	12.4	...	...	0.7	10.5	14.0	16.0	19.6	17.5	13.7	15.7	17.6	12.1
2009.....	12.0	...	...	0.6	10.0	13.1	16.1	19.2	16.4	13.7	15.8	16.4	11.8
2008.....	11.8	...	...	0.5	9.9	13.2	15.9	18.6	16.0	13.6	16.1	16.4	11.6
2007.....	11.5	...	...	0.5	9.6	13.3	15.7	17.7	15.3	12.4	16.2	17.0	11.3
2006.....	11.2	...	...	0.5	9.8	12.7	15.2	17.2	14.4	12.4	15.8	17.3	11.0
2005.....	11.0	...	...	0.7	9.9	12.7	15.1	16.5	13.7	12.4	16.8	18.3	10.9
2004.....	11.1	...	...	0.7	10.3	12.9	15.2	16.6	13.7	12.2	16.3	17.6	11.0
2003.....	10.9	...	...	0.6	9.6	12.9	15.0	15.9	13.7	12.6	16.4	17.9	10.8
2002.....	11.0	...	...	0.6	9.8	12.8	15.3	15.8	13.5	13.4	17.7	18.9	10.9
2001 ⁵	10.7	...	...	0.7	9.9	12.8	14.7	15.1	13.2	13.2	17.4	17.8	10.7
2000.....	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6	10.4
1999.....	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3	10.5
Essential hypertension and hypertensive renal disease (I10,I12,I15)													
2021.....	12.9	*	*	*	0.1	0.4	1.5	4.8	12.1	24.9	63.2	270.8	10.7
2020.....	12.7	*	*	*	0.1	0.3	1.4	4.5	11.8	23.9	60.3	250.3	10.1
2019.....	11.1	*	*	*	*	0.3	1.2	3.6	9.8	20.9	54.3	227.1	8.9
2018.....	11.0	*	*	*	*	0.3	1.2	3.7	9.7	20.2	54.4	229.4	8.9
2017.....	10.8	*	*	*	*	0.3	1.3	3.9	9.5	19.7	55.8	231.4	9.0
2016.....	10.3	*	*	*	*	0.3	1.1	3.8	9.0	19.0	53.7	222.3	8.6
2015.....	10.0	*	*	*	*	0.3	1.2	3.4	8.8	18.1	54.1	222.7	8.5
2014.....	9.5	*	*	*	0.0	0.2	1.1	3.3	8.4	16.9	51.3	217.0	8.2
2013.....	9.7	*	*	*	0.1	0.3	1.0	3.5	8.0	17.3	53.7	231.6	8.5
2012.....	9.3	*	*	*	*	0.2	0.8	3.0	7.8	16.1	51.7	230.7	8.2
2011.....	8.9	*	*	*	*	0.2	1.0	3.1	7.0	16.6	51.4	222.7	8.1
2010.....	8.6	*	*	*	0.0	0.3	1.0	3.1	7.3	16.7	51.8	212.0	8.0
2009.....	8.4	*	*	*	0.1	0.3	1.0	3.1	7.1	16.3	51.0	208.0	7.8
2008.....	8.5	*	*	*	0.1	0.3	1.0	3.0	7.2	16.5	51.9	215.3	8.0
2007.....	8.0	*	*	*	0.1	0.2	0.9	2.8	6.4	15.9	49.2	209.1	7.6
2006.....	8.0	*	*	*	0.0	0.3	0.9	3.0	6.8	16.5	50.8	206.1	7.7
2005.....	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.5	55.5	228.0	8.3
2004.....	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	16.9	52.5	212.2	7.9
2003.....	7.6	*	*	*	0.1	0.2	0.8	2.5	6.3	16.8	51.6	199.4	7.6
2002.....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	15.9	48.1	189.6	7.1
2001.....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.4	47.6	175.6	6.9
2000.....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9	6.5
1999.....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1	6.2

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Influenza and pneumonia (J09–J18)													
2021.....	12.6	3.5	0.3	0.1	0.3	0.9	1.9	4.4	11.3	25.3	69.9	231.8	10.5
2020.....	16.3	3.3	0.5	0.3	0.4	1.3	2.7	6.2	14.8	32.1	84.1	273.6	13.0
2019.....	15.2	4.1	0.8	0.3	0.4	1.0	2.3	5.1	12.5	27.3	77.2	294.7	12.3
2018.....	18.1	4.6	0.8	0.3	0.5	1.0	2.3	5.6	13.9	31.7	94.2	377.6	14.9
2017.....	17.1	4.0	0.7	0.3	0.4	0.9	1.9	4.8	12.0	29.6	93.8	375.3	14.3
2016.....	15.9	4.2	0.6	0.2	0.4	1.0	2.2	5.0	12.1	28.5	88.5	340.3	13.5
2015.....	17.8	4.4	0.6	0.2	0.4	0.9	1.7	4.7	11.3	29.5	101.6	421.4	15.2
2014.....	17.3	4.7	0.7	0.2	0.5	1.3	2.8	6.3	13.4	29.8	96.4	385.9	15.1
2013.....	18.0	4.5	0.6	0.3	0.4	1.0	2.2	5.1	12.2	29.5	103.7	441.0	15.9
2012.....	16.1	4.0	0.6	0.2	0.3	0.8	1.7	4.1	10.2	26.1	98.2	408.4	14.4
2011.....	17.3	5.2	0.7	0.3	0.5	1.2	2.1	5.0	11.0	28.9	104.0	439.2	15.7
2010.....	16.2	4.9	0.6	0.2	0.4	0.9	1.9	4.3	9.9	27.9	102.4	426.2	15.1
2009.....	17.5	6.3	0.9	0.6	1.0	2.0	3.2	6.5	11.7	29.5	107.0	433.8	16.5
2008.....	18.5	5.5	0.9	0.2	0.5	0.9	2.1	5.1	10.9	30.5	118.6	512.3	17.6
2007.....	17.5	5.4	0.7	0.3	0.4	0.8	1.8	4.3	9.5	28.2	113.5	506.7	16.8
2006.....	18.9	6.5	0.8	0.2	0.4	0.9	1.9	4.6	9.9	31.6	127.3	547.0	18.4
2005.....	21.3	6.6	0.7	0.3	0.4	0.9	2.1	5.1	11.2	35.1	142.0	644.9	21.0
2004.....	20.4	6.8	0.8	0.2	0.4	0.8	2.0	4.6	10.8	34.2	139.1	622.8	20.4
2003.....	22.5	8.1	1.0	0.4	0.5	1.0	2.2	5.2	11.2	36.9	150.8	703.0	22.6
2002.....	22.8	6.7	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.2	156.6	732.4	23.2
2001.....	21.8	7.5	0.7	0.2	0.5	0.9	2.2	4.6	10.8	36.2	148.3	700.1	22.2
2000.....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1	23.7
1999.....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8	23.5
Septicemia (A40–A41)													
2021.....	12.4	3.2	0.2	0.1	0.2	0.9	2.6	6.0	15.1	31.1	67.4	153.3	10.2
2020.....	12.2	3.1	0.3	0.1	0.2	0.8	2.3	6.2	14.7	29.7	63.5	143.1	9.7
2019.....	11.7	3.5	0.3	0.2	0.3	0.8	1.9	5.3	13.4	28.0	64.5	150.2	9.5
2018.....	12.4	3.9	0.3	0.1	0.2	0.9	2.0	5.7	14.1	30.0	69.4	167.4	10.2
2017.....	12.6	3.7	0.3	0.1	0.3	0.9	2.1	5.8	13.9	30.9	72.4	173.3	10.6
2016.....	12.6	4.8	0.4	0.2	0.3	0.9	2.2	5.8	14.3	30.6	73.9	174.2	10.7
2015.....	12.7	4.5	0.3	0.2	0.3	0.9	2.0	5.9	14.1	31.4	75.3	185.8	11.0
2014.....	12.2	4.0	0.3	0.2	0.3	0.8	2.1	5.8	14.2	31.1	73.1	176.9	10.7
2013.....	12.1	3.9	0.3	0.1	0.3	0.8	2.0	5.6	13.6	30.5	76.4	179.6	10.7
2012.....	11.4	4.5	0.4	0.1	0.3	0.8	1.9	5.3	12.9	29.2	73.9	173.4	10.3
2011.....	11.5	4.5	0.4	0.2	0.3	0.8	2.0	5.5	13.0	29.5	74.4	179.7	10.5
2010.....	11.3	5.5	0.4	0.2	0.3	0.8	1.9	5.2	12.6	30.1	76.0	179.0	10.6
2009.....	11.6	5.5	0.4	0.2	0.3	0.9	2.2	5.4	13.1	31.4	79.2	182.4	11.0
2008.....	11.8	7.0	0.6	0.2	0.3	0.9	2.1	5.7	13.3	31.4	82.0	189.8	11.3
2007.....	11.6	6.8	0.5	0.2	0.4	0.7	2.1	5.5	12.8	32.2	79.5	190.8	11.2
2006.....	11.5	6.7	0.6	0.2	0.3	0.7	2.0	5.2	12.6	31.6	82.1	193.0	11.2
2005.....	11.6	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.8	32.2	81.3	203.4	11.4
2004.....	11.4	6.8	0.5	0.2	0.3	0.8	1.9	5.4	12.8	32.1	81.5	199.6	11.3
2003.....	11.7	7.0	0.5	0.2	0.4	0.8	2.1	5.3	13.0	32.3	84.8	213.7	11.8
2002.....	11.8	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.5	86.3	213.4	11.9
2001.....	11.3	7.8	0.7	0.2	0.3	0.7	1.8	5.0	12.4	32.6	82.2	210.3	11.5
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7	11.3
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7	11.3

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Parkinson disease (G20–G21)													
2021.....	11.6	*	*	*	*	*	*	0.2	2.0	18.6	102.5	246.0	9.8
2020.....	12.2	*	*	*	*	*	*	0.2	2.1	19.0	105.3	237.4	9.9
2019.....	10.8	*	*	*	*	*	*	0.2	1.9	16.8	93.3	215.9	8.8
2018.....	10.3	*	*	*	*	*	*	0.2	1.8	16.2	91.0	214.6	8.7
2017.....	9.8	*	*	*	*	*	*	0.1	1.7	15.7	89.6	206.3	8.4
2016.....	9.2	*	*	*	*	*	*	0.2	1.6	14.3	85.5	198.6	8.0
2015.....	8.7	*	*	*	*	*	*	0.2	1.5	13.8	82.4	190.6	7.7
2014.....	8.2	*	*	*	*	*	*	0.2	1.4	13.0	79.2	182.0	7.4
2013.....	8.0	*	*	*	*	*	*	0.2	1.5	12.7	78.5	178.2	7.3
2012.....	7.6	*	*	*	*	*	*	0.1	1.4	12.3	76.2	172.3	7.0
2011.....	7.4	*	*	*	*	*	*	0.1	1.3	12.8	76.0	168.1	7.0
2010.....	7.1	*	*	*	*	*	*	0.2	1.3	11.8	74.8	165.9	6.8
2009.....	6.7	*	*	*	*	*	*	0.2	1.3	11.2	70.8	157.0	6.5
2008.....	6.7	*	*	*	*	*	*	0.2	1.2	12.3	71.2	157.4	6.6
2007.....	6.7	*	*	*	*	*	*	0.1	1.2	11.7	71.5	157.0	6.5
2006.....	6.6	*	*	*	*	*	*	0.2	1.2	12.0	69.5	157.6	6.5
2005.....	6.6	*	*	*	*	*	*	0.2	1.4	12.8	71.1	156.0	6.6
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	11.9	67.4	145.1	6.2
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.6	67.6	145.8	6.3
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.1	63.8	142.2	6.0
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.5	137.0	5.9
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9	5.7
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4	5.4
Dementia-related causes ⁶													
2021.....	84.3	*	*	0.1	*	0.0	0.2	0.9	8.6	65.0	477.6	2,949.3	72.4
2020.....	92.3	*	*	*	*	0.0	0.1	0.8	8.7	68.1	495.1	2,946.9	73.3
2019.....	82.8	*	*	0.1	*	0.1	0.1	0.9	8.2	59.3	448.6	2,688.9	66.6
2018.....	81.6	*	*	*	*	0.1	0.1	0.8	7.8	57.9	447.0	2,700.3	66.6
2017.....	80.4	*	*	0.0	*	*	0.1	0.7	7.5	56.9	450.9	2,707.3	66.7
2016.....	77.2	*	*	0.1	*	0.1	0.1	0.8	7.3	54.7	441.6	2,626.4	64.9
2015.....	76.5	*	*	0.1	*	*	0.1	0.8	6.8	53.0	447.2	2,637.4	65.2
2014.....	75.2	*	0.1	0.1	*	*	0.1	0.8	6.9	52.7	450.6	2,611.3	64.9
2013.....	74.1	*	*	*	*	*	0.1	0.9	7.4	52.2	449.7	2,601.8	64.8
2012.....	71.2	*	0.2	*	*	*	0.1	0.9	6.8	50.3	445.2	2,532.7	63.3
2011.....	68.3	*	0.1	0.1	*	*	0.1	0.8	6.5	50.4	436.6	2,458.6	61.8
2010.....	63.6	*	*	*	*	*	0.1	0.9	6.3	48.3	412.1	2,352.4	58.8
2009.....	57.9	*	0.1	0.1	*	0.1	0.2	0.8	5.7	45.2	383.3	2,151.3	54.2
2008.....	58.9	*	0.2	0.1	*	*	0.1	0.9	6.0	46.4	396.7	2,213.2	55.9
2007.....	51.8	*	0.2	0.1	*	*	0.1	0.8	5.5	42.3	350.0	1,976.0	49.8
2006.....	50.8	*	0.1	0.1	*	*	0.1	0.8	5.5	42.4	346.2	1,967.0	49.5
2005.....	43.8	*	0.2	*	*	*	0.1	0.6	4.5	36.3	302.1	1,735.4	43.4
2004.....	39.0	*	0.2	0.1	*	*	0.1	0.6	3.9	33.1	275.4	1,556.9	39.1
2003.....	38.1	*	0.2	0.1	*	*	0.1	0.6	4.0	34.1	269.5	1,523.1	38.4
2002.....	35.5	*	0.2	0.1	*	*	0.1	0.4	3.6	31.9	254.9	1,428.9	36.1
2001.....	32.5	*	0.2	0.0	*	0.1	0.1	0.5	3.8	30.5	234.6	1,299.3	33.1
2000.....	29.7	*	0.2	0.1	0.1	0.1	0.1	0.5	3.6	29.4	218.5	1,180.9	30.5
1999.....	26.7	*	0.2	0.1	*	*	0.1	0.4	3.4	27.0	197.9	1,062.5	27.5

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes]

Cause of death (based on ICD–10) and year	Age group (years)												Age- adjusted rate ³
	All ages ¹	Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Drug-induced causes ⁶													
2021.....	33.5	2.1	0.8	0.3	17.5	54.2	64.0	56.2	48.0	18.5	6.0	5.3	33.6
2020.....	29.2	1.7	0.4	0.3	17.0	48.8	55.8	49.1	39.9	14.7	4.9	5.1	29.5
2019.....	22.7	1.3	0.3	0.2	11.5	36.9	42.3	38.9	32.7	12.6	5.2	4.6	22.8
2018.....	21.7	0.8	0.2	0.1	11.0	36.8	40.0	37.3	30.4	11.4	4.8	5.1	21.8
2017.....	22.7	0.9	0.2	0.2	13.0	39.8	40.6	39.8	30.0	10.5	4.5	5.3	22.8
2016.....	20.8	0.9	0.3	0.1	12.8	35.9	36.6	36.5	27.7	9.2	4.1	5.3	20.8
2015.....	17.2	0.7	0.4	0.1	10.0	28.0	29.6	31.9	23.3	8.1	4.4	5.6	17.2
2014.....	15.6	0.6	0.3	0.1	8.9	24.0	26.2	29.8	21.7	7.6	4.4	5.0	15.5
2013.....	14.7	0.8	0.3	0.1	8.6	21.7	24.1	29.0	20.6	7.1	4.4	5.3	14.6
2012.....	14.0	0.8	0.2	0.1	8.3	20.9	23.1	28.3	17.9	6.5	4.0	5.1	13.8
2011.....	14.0	0.6	0.2	0.1	8.9	20.9	23.4	28.2	17.1	6.0	4.0	4.9	13.9
2010.....	13.1	0.6	0.3	0.2	8.4	19.2	21.7	26.5	16.2	5.2	4.0	5.5	12.9
2009.....	12.8	0.8	0.2	0.1	8.0	17.8	21.5	26.9	14.9	5.4	4.5	5.1	12.6
2008.....	12.7	0.5	0.3	0.1	8.3	17.4	22.2	26.8	14.0	5.2	4.0	5.0	12.6
2007.....	12.7	0.8	0.3	0.2	8.5	17.5	22.6	26.8	13.4	4.6	3.9	5.2	12.6
2006.....	12.9	1.1	0.2	0.1	8.5	17.2	23.5	26.7	12.1	5.2	6.0	8.8	12.8
2005.....	11.3	0.9	0.2	0.1	7.3	14.6	21.5	23.6	10.6	4.7	5.4	8.3	11.3
2004.....	10.5	0.7	0.2	0.2	6.9	12.9	21.1	21.7	9.0	4.2	4.8	6.7	10.5
2003.....	9.9	0.6	0.2	0.1	6.3	12.3	20.7	20.0	8.0	4.1	4.2	6.3	9.9
2002.....	9.1	0.7	0.2	0.1	5.4	11.3	19.8	18.0	6.8	3.6	3.8	6.0	9.1
2001.....	7.6	0.5	0.2	0.1	4.5	9.5	17.0	14.7	5.4	3.0	3.5	5.2	7.6
2000.....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7	7.0
1999.....	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8	6.8
Alcohol-induced causes ⁶													
2021.....	16.3	*	*	*	0.5	6.7	18.0	30.3	42.0	29.6	15.4	7.8	14.4
2020.....	14.9	*	*	*	0.4	6.0	16.0	27.5	39.3	27.1	14.4	6.5	13.1
2019.....	11.9	*	*	*	0.4	4.2	11.1	22.0	32.5	22.5	12.6	6.4	10.4
2018.....	11.4	*	*	*	0.3	3.7	10.0	21.6	31.5	22.2	12.4	6.1	9.9
2017.....	11.0	*	*	*	0.3	3.4	9.4	21.8	30.2	20.9	11.7	6.4	9.6
2016.....	10.8	*	*	*	0.4	3.6	9.2	21.4	29.7	20.3	11.8	6.3	9.5
2015.....	10.3	*	*	*	0.4	3.2	8.7	21.6	28.2	19.1	11.2	5.8	9.1
2014.....	9.6	*	*	*	0.3	2.8	8.0	20.4	26.8	17.6	10.5	5.6	8.5
2013.....	9.2	*	*	*	0.3	2.5	7.7	20.1	25.3	16.6	10.3	4.9	8.2
2012.....	8.8	*	*	*	0.4	2.4	7.4	20.0	24.1	15.8	10.3	5.0	8.0
2011.....	8.6	*	*	*	0.4	2.1	7.6	19.8	22.7	15.2	9.6	5.1	7.7
2010.....	8.3	*	*	*	0.3	2.2	7.5	19.1	21.9	15.8	9.6	5.3	7.6
2009.....	8.0	*	*	*	0.4	1.8	7.6	18.7	20.8	15.1	9.2	4.8	7.4
2008.....	8.0	*	*	*	0.4	2.0	7.6	18.6	20.7	15.3	9.4	5.2	7.4
2007.....	7.7	*	*	*	0.4	1.9	7.3	18.2	19.9	15.2	9.6	5.0	7.2
2006.....	7.4	*	*	*	0.3	1.6	7.5	17.5	19.2	14.5	9.7	5.3	7.0
2005.....	7.3	*	*	*	0.4	1.4	7.5	17.6	19.4	14.9	9.2	5.0	7.0
2004.....	7.2	*	*	*	0.3	1.6	7.7	17.3	18.6	15.5	9.2	4.6	7.0
2003.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.0	9.2	4.3	7.0
2002.....	7.0	*	*	*	0.3	1.5	8.1	16.9	18.3	15.4	9.3	4.6	6.9
2001.....	7.1	*	*	*	0.3	1.6	8.3	17.1	18.3	15.5	9.6	5.1	7.0
2000.....	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4	7.0
1999.....	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5	7.1

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2021, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group (years)											Age- adjusted rate ³
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Injury by firearms ⁶													
2021.....	14.7	*	0.9	1.6	23.5	24.8	18.1	14.5	12.1	11.7	16.2	18.3	14.6
2020.....	13.7	*	0.8	1.5	22.2	22.7	16.7	13.3	11.8	10.9	15.3	16.2	13.6
2019.....	12.1	*	0.5	1.0	17.4	18.1	14.6	12.7	12.3	11.1	15.0	16.1	11.9
2018.....	12.1	*	0.6	1.1	17.2	17.7	14.6	12.8	12.7	12.0	15.4	14.7	11.9
2017.....	12.2	*	0.5	1.1	17.7	18.5	14.4	13.1	12.3	11.4	14.8	15.6	12.0
2016.....	12.0	*	0.6	0.9	17.2	18.2	14.5	12.8	11.9	11.4	14.7	14.3	11.8
2015.....	11.3	*	0.5	0.9	15.7	16.8	13.1	12.4	11.7	11.3	14.5	14.5	11.1
2014.....	10.5	*	0.4	0.9	14.0	14.7	12.1	12.2	11.4	11.5	13.9	15.0	10.3
2013.....	10.6	*	0.4	0.8	14.1	15.3	12.3	12.3	11.5	11.3	14.1	13.9	10.4
2012.....	10.7	*	0.4	0.8	14.7	15.3	12.4	12.4	11.6	10.8	14.1	13.6	10.5
2011.....	10.4	*	0.5	0.8	14.4	15.0	11.7	12.2	11.0	10.9	13.7	13.1	10.2
2010.....	10.3	*	0.4	0.7	14.2	15.0	11.7	12.0	11.1	10.7	12.7	13.2	10.1
2009.....	10.2	*	0.4	0.7	14.4	14.5	11.9	11.8	10.8	10.9	13.3	12.5	10.1
2008.....	10.4	*	0.5	0.7	15.4	15.4	11.8	11.5	10.8	10.7	13.2	12.5	10.3
2007.....	10.4	*	0.4	0.8	16.0	15.9	12.0	11.1	10.1	9.8	13.1	12.7	10.3
2006.....	10.4	*	0.4	0.9	16.7	15.7	11.6	11.2	9.7	9.9	12.9	12.5	10.3
2005.....	10.4	*	0.4	0.8	16.1	16.1	11.7	11.2	9.7	10.2	13.6	13.0	10.3
2004.....	10.1	*	0.3	0.7	15.6	15.3	11.4	11.0	9.8	10.1	13.3	12.7	10.0
2003.....	10.4	*	0.3	0.8	16.5	15.8	11.6	11.1	10.0	10.3	13.4	13.2	10.3
2002.....	10.5	*	0.4	0.8	16.6	15.6	12.2	10.8	10.2	10.8	14.4	13.2	10.5
2001.....	10.4	*	0.5	0.8	16.6	15.5	11.7	10.5	10.1	10.9	14.3	13.1	10.3
2000.....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2	10.2
1999.....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5	10.3

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Figures for age not stated included in all ages category but not distributed among age groups.

²Death rates for younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

³For method of computation, see Technical Notes.

⁴COVID-19 became an official cause of death in 2020; rates for years before 2020 are not applicable.

⁵Figures include September 11, 2001, related deaths for which death certificates were filed as of October 24, 2002; see Technical Notes from "Deaths: Final Data for 2001. *National Vital Statistics Reports*, vol 53, no 3."

⁶For the list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021

[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
All causes	3,464,231	19,920	3,816	5,975	38,307	82,274	124,939	216,037	478,171	724,266	829,653	940,780	93
Salmonella infections (A01-A02)	72	9	3	2	—	—	2	4	9	22	12	9	—
Shigellosis and amebiasis (A03,A06)	7	—	—	1	—	—	—	—	3	1	2	—	—
Certain other intestinal infections. (A04,A07-A09)	7,036	160	9	13	15	33	112	261	789	1,585	2,070	1,989	—
Tuberculosis. (A16-A19)	602	2	—	2	3	15	38	52	114	165	127	84	—
Respiratory tuberculosis. (A16)	435	—	—	1	3	12	25	31	80	120	88	75	—
Other tuberculosis (A17-A19)	167	2	—	1	—	3	13	21	34	45	39	9	—
Whooping cough (A37)	4	1	—	1	—	—	—	1	—	—	1	—	—
Scarlet fever and erysipelas (A38,A46)	—	—	—	—	—	—	—	—	—	—	—	—	—
Meningococcal infection (A39)	21	2	1	—	1	2	5	2	3	2	1	2	—
Septicemia (A40-A41)	41,281	113	34	45	107	426	1,108	2,426	6,477	10,466	10,919	9,160	—
Syphilis (A50-A53)	43	5	—	—	—	—	2	3	8	14	8	3	—
Acute poliomyelitis. (A80)	—	—	—	—	—	—	—	—	—	—	—	—	—
Arthropod-borne viral encephalitis. (A83-A84,A85.2)	6	—	—	—	—	—	1	—	2	1	2	—	—
Measles (B05)	—	—	—	—	—	—	—	—	—	—	—	—	—
Viral hepatitis. (B15-B19)	3,589	2	—	—	4	33	175	453	1,332	1,138	346	106	—
Human immunodeficiency virus (HIV) disease (B20-B24)	4,977	1	—	—	33	510	738	1,058	1,549	831	217	40	—
Malaria. (B50-B54)	6	—	—	—	—	—	—	2	1	1	—	2	—
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44, A48-A49,A54-A79,A81-A82, A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99,U07.1)	427,811	242	168	231	1,478	6,311	16,411	37,642	75,512	105,487	101,412	82,913	4
Malignant neoplasms. (C00-C97)	605,213	52	282	796	1,323	3,615	11,194	33,567	108,023	179,145	164,850	102,359	7
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	11,394	—	—	2	10	52	216	886	2,675	3,587	2,515	1,450	1
Malignant neoplasm of esophagus. (C15)	15,743	—	1	—	3	37	235	999	3,665	5,369	3,761	1,673	—
Malignant neoplasm of stomach. (C16)	10,894	—	1	1	17	131	403	951	2,053	3,012	2,680	1,644	1
Malignant neoplasms of colon, rectum and anus (C18-C21)	54,121	—	—	4	42	356	1,672	4,846	10,478	14,008	12,631	10,082	2
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	28,719	4	15	17	26	108	346	1,401	6,488	10,556	6,743	3,014	1
Malignant neoplasm of pancreas (C25)	47,906	—	—	4	5	59	495	2,358	8,907	15,643	13,543	6,892	—
Malignant neoplasm of larynx (C32)	3,887	—	—	1	—	4	22	245	1,004	1,340	858	413	—
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	134,592	—	—	1	17	119	774	4,859	25,908	45,284	40,331	17,297	2
Malignant melanoma of skin. (C43)	8,224	—	—	1	19	81	287	576	1,444	2,208	2,075	1,533	—
Malignant neoplasm of breast (C50)	42,810	—	—	—	7	371	1,873	4,169	8,474	10,892	9,419	7,605	—
Malignant neoplasm of cervix uteri. (C53)	4,366	—	—	—	—	234	649	852	1,004	853	523	251	—
Malignant neoplasms of corpus uteri and uterus, part unspecified. (C54-C55)	12,208	—	—	—	4	50	232	741	2,480	4,313	2,948	1,440	—
Malignant neoplasm of ovary (C56)	13,430	—	—	2	28	95	286	1,050	2,754	4,099	3,444	1,672	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Malignant neoplasm of prostate (C61)	32,563	—	—	—	1	3	18	335	2,783	8,224	11,011	10,188	—
Malignant neoplasms of kidney and renal pelvis (C64-C65)	14,163	1	14	24	17	61	218	832	2,625	4,171	3,791	2,409	—
Malignant neoplasm of bladder (C67)	16,840	—	—	2	1	8	73	345	1,723	3,854	5,429	5,405	—
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	17,857	12	71	308	242	419	834	1,735	4,003	5,286	3,558	1,389	—
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	57,375	14	77	209	346	569	942	2,160	6,867	14,978	18,576	12,637	—
Hodgkin disease (C81)	1,003	—	—	1	17	46	49	60	156	259	271	144	—
Non-Hodgkin lymphoma (C82-C85)	20,020	—	7	25	53	154	307	766	2,440	5,088	6,622	4,558	—
Leukemia (C91-C95)	23,271	14	70	183	274	363	503	875	2,522	5,996	7,263	5,208	—
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	12,933	—	—	—	1	5	82	446	1,739	3,606	4,368	2,686	—
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	148	—	—	—	1	1	1	13	10	29	52	41	—
All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)	78,121	21	103	220	538	858	1,619	4,227	12,688	21,468	21,014	15,365	—
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16,237	26	37	54	83	126	240	547	1,440	3,439	5,079	5,166	—
Anemias (D50-D64)	5,987	12	10	19	61	148	239	320	583	1,086	1,426	2,083	—
Diabetes mellitus (E10-E14)	103,294	2	2	48	345	1,285	2,961	7,597	18,603	27,945	25,989	18,517	—
Nutritional deficiencies (E40-E64)	17,505	12	5	2	25	53	98	279	890	2,198	4,308	9,635	—
Malnutrition (E40-E46)	17,106	11	4	1	24	51	91	267	821	2,112	4,238	9,486	—
Other nutritional deficiencies (E50-E64)	399	1	1	1	1	2	7	12	69	86	70	149	—
Meningitis (G00,G03)	534	55	13	9	7	25	40	56	97	86	106	40	—
Parkinson disease (G20-G21)	38,536	—	—	—	2	4	13	99	850	6,252	16,615	14,701	—
Alzheimer disease (G30)	119,399	—	—	—	—	2	13	108	1,353	8,886	34,724	74,312	1
Major cardiovascular diseases (I00-I78)	925,923	379	174	299	1,202	5,147	16,178	43,496	112,152	182,530	230,520	333,820	26
Diseases of heart (I00-I09,I11,I13,I20-I51)	695,547	265	116	198	944	4,155	12,754	34,535	89,342	140,250	169,231	243,733	24
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	3,907	2	1	2	12	84	124	172	411	755	1,003	1,341	—
Hypertensive heart disease (I11)	67,774	—	—	2	66	569	2,160	5,049	10,443	12,760	12,985	23,736	4
Hypertensive heart and renal disease (I13)	13,918	—	—	—	5	51	166	396	989	2,087	3,505	6,717	2
Ischemic heart diseases (I20-I25)	375,476	15	6	13	115	1,057	5,335	18,382	53,321	83,934	94,650	118,636	12
Acute myocardial infarction (I21-I22)	109,097	7	3	3	44	437	2,025	6,729	18,297	27,039	27,224	27,288	1
Other acute ischemic heart diseases (I24)	4,748	2	—	2	3	19	88	312	826	1,136	1,171	1,188	1
Other forms of chronic ischemic heart disease (I20,I25)	261,631	6	3	8	68	601	3,222	11,341	34,198	55,759	66,255	90,160	10
Atherosclerotic cardiovascular disease, so described (I25.0)	78,175	—	—	1	23	321	1,822	6,084	16,797	20,676	15,986	16,456	9

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.

[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
All other forms of chronic ischemic heart disease. (I20,I25.1-I25.9)	183,456	6	3	7	45	280	1,400	5,257	17,401	35,083	50,269	73,704	1
Other heart diseases. (I26-I51)	234,472	248	109	181	746	2,394	4,969	10,536	24,178	40,714	57,088	93,303	6
Acute and subacute endocarditis. (I33)	1,767	—	—	1	15	144	191	214	329	381	327	165	—
Diseases of pericardium and acute myocarditis. (I30-I31,I40)	1,237	12	5	14	23	38	66	125	187	250	281	236	—
Heart failure. (I50)	85,037	16	9	9	36	220	702	2,032	5,767	12,444	21,777	42,024	1
All other forms of heart disease. (I26-I28, I34-I38,I42-I49,I51)	146,431	220	95	157	672	1,992	4,010	8,165	17,895	27,639	34,703	50,878	5
Essential hypertension and hypertensive renal disease. (I10,I12,I15)	42,816	1	—	1	24	171	668	1,965	5,187	8,369	10,244	16,185	1
Cerebrovascular diseases. (I60-I69)	162,890	97	55	88	190	624	2,189	5,755	14,634	28,327	44,533	66,397	1
Atherosclerosis. (I70)	4,214	3	—	—	—	4	16	89	335	693	1,082	1,992	—
Other diseases of circulatory system. (I71-I78)	20,456	13	3	12	44	193	551	1,152	2,654	4,891	5,430	5,513	—
Aortic aneurysm and dissection. (I71)	10,037	—	1	10	25	121	395	713	1,382	2,422	2,680	2,288	—
Other diseases of arteries, arterioles and capillaries. (I72-I78)	10,419	13	2	2	19	72	156	439	1,272	2,469	2,750	3,225	—
Other disorders of circulatory system. (I80-I99)	5,655	23	6	3	61	223	455	779	1,076	1,140	943	946	—
Influenza and pneumonia. (J09-J18)	41,917	125	47	49	128	400	842	1,783	4,845	8,518	11,327	13,852	1
Influenza. (J09-J11)	608	5	1	4	8	11	17	32	94	149	157	130	—
Pneumonia. (J12-J18)	41,309	120	46	45	120	389	825	1,751	4,751	8,369	11,170	13,722	1
Other acute lower respiratory infections. (J20-J22,U04)	189	27	15	3	1	3	10	7	15	35	28	45	—
Acute bronchitis and bronchiolitis. (J20-J21)	128	27	15	2	—	3	5	4	10	22	13	27	—
Other and unspecified acute lower respiratory infections. (J22,U04)	61	—	—	1	1	—	5	3	5	13	15	18	—
Chronic lower respiratory diseases. (J40-J47)	142,342	13	31	99	158	360	733	3,174	17,620	37,932	46,328	35,892	2
Bronchitis, chronic and unspecified. (J40-J42)	335	6	6	1	3	3	8	18	36	75	69	110	—
Emphysema. (J43)	7,336	—	1	—	1	7	25	170	999	2,133	2,364	1,636	—
Asthma. (J45-J46)	3,517	3	23	92	140	293	374	453	626	579	481	453	—
Other chronic lower respiratory diseases. (J44,J47)	131,154	4	1	6	14	57	326	2,533	15,959	35,145	43,414	33,693	2
Pneumoconioses and chemical effects. (J60-J66, J68,U07.0)	564	—	—	1	4	2	7	8	37	123	193	189	—
Pneumonitis due to solids and liquids. (J69)	19,969	5	6	8	52	133	296	597	1,971	3,758	5,480	7,663	—
Other diseases of respiratory system. (J00-J06, J30-J39,J67,J70-J98)	45,888	173	91	72	129	284	635	1,679	5,089	10,773	14,452	12,511	—
Peptic ulcer. (K25-K28)	4,008	2	2	1	6	52	119	280	668	928	1,002	948	—
Diseases of appendix. (K35-K38)	449	3	6	10	10	12	18	29	69	96	102	94	—
Hernia. (K40-K46)	2,488	15	3	4	2	7	50	92	279	478	655	902	1
Chronic liver disease and cirrhosis. (K70,K73-K74)	56,585	2	2	2	59	1,833	5,833	10,501	17,664	12,934	5,900	1,853	2
Alcoholic liver disease. (K70)	33,098	—	—	—	46	1,536	4,600	7,636	11,348	6,140	1,541	250	1

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Other chronic liver disease and cirrhosis (K73-K74)	23,487	2	2	2	13	297	1,233	2,865	6,316	6,794	4,359	1,603	1
Cholelithiasis and other disorders of gallbladder (K80-K82)	4,490	—	—	—	4	21	58	131	370	958	1,300	1,648	—
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54,358	48	5	7	54	369	972	2,516	6,374	12,245	15,569	16,199	—
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	781	7	—	1	1	10	8	34	72	155	235	258	—
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	290	—	1	1	2	14	13	31	43	71	76	38	—
Renal failure (N17-N19)	53,261	41	4	5	51	342	951	2,448	6,256	12,013	15,252	15,898	—
Other disorders of kidney (N25,N27)	26	—	—	—	—	3	—	3	3	6	6	5	—
Infections of kidney (N10-N12,N13.6,N15.1)	1,248	4	4	—	7	12	57	81	208	291	322	262	—
Hyperplasia of prostate (N40)	715	—	—	—	—	—	—	4	35	96	218	362	—
Inflammatory diseases of female pelvic organs (N70-N76)	214	—	—	—	3	9	23	10	36	51	45	37	—
Pregnancy, childbirth and the puerperium (O00-O99)	1,687	...	...	4	214	797	654	17	1	—	—	—	—
Pregnancy with abortive outcome (O00-O07)	48	...	...	1	11	25	10	1	—	—	—	—	—
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	1,639	...	...	3	203	772	644	16	1	—	—	—	—
Certain conditions originating in the perinatal period (P00-P96)	9,579	9,434	68	35	19	7	3	6	4	1	1	—	1
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9,664	3,963	412	350	419	459	513	765	1,249	734	450	349	1
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	34,503	2,714	273	140	603	1,374	1,851	2,165	3,816	4,813	5,878	10,860	16
All other diseases (residual)	397,499	611	375	725	2,028	5,707	11,321	20,164	42,223	67,115	95,068	152,157	5
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224,935	1,306	1,299	1,742	15,792	34,452	36,444	31,407	33,471	21,797	20,557	26,649	19
Transport accidents (V01-V99,Y85)	49,845	108	389	1,020	7,654	9,235	7,607	6,649	7,352	5,289	3,124	1,414	4
Motor vehicle accidents (V02-V04,V09.0, V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6, V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1, V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)	46,980	107	373	979	7,425	8,897	7,143	6,228	6,722	4,783	2,943	1,376	4
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9, V10-V11,V15-V18,V19.3,V19.8-V19.9, V80.0-V80.2,V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	1,179	1	14	17	98	166	201	168	216	190	91	17	—
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,686	—	2	24	131	172	263	253	414	316	90	21	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.

[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Nontransport accidents (W00-X59,Y86)	175,090	1,198	910	722	8,138	25,217	28,837	24,758	26,119	16,508	17,433	25,235	15
Falls (W00-W19)	44,686	8	21	35	150	404	714	1,333	3,278	6,409	12,136	20,197	1
Accidental discharge of firearms. (W32-W34)	549	3	51	53	125	81	76	46	44	46	20	4	—
Accidental drowning and submersion. (W65-W74)	4,337	38	475	238	423	561	599	460	587	490	321	145	—
Accidental exposure to smoke, fire and flames. (X00-X09)	3,389	15	100	150	105	188	227	328	662	793	559	261	1
Accidental poisoning and exposure to noxious substances (X40-X49)	102,001	27	34	80	6,929	23,167	26,070	21,011	18,596	5,251	615	212	9
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)	20,128	1,107	229	166	406	816	1,151	1,580	2,952	3,519	3,782	4,416	4
Intentional self-harm (suicide). (*U03,X60-X84,Y87.0)	48,183	...	...	607	6,528	8,862	7,862	7,401	7,267	5,144	3,170	1,338	4
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	26,328	...	...	236	3,590	4,221	3,644	3,717	3,984	3,430	2,470	1,035	1
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	21,855	...	...	371	2,938	4,641	4,218	3,684	3,283	1,714	700	303	3
Assault (homicide). (*U01-U02,X85-Y09,Y87.1)	26,031	267	309	486	6,635	7,571	4,863	2,768	1,828	882	316	105	1
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	20,958	15	72	359	6,207	6,679	3,931	2,024	1,096	413	115	47	—
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-U01.3,*U01.5-U01.9,*U02,X85-X92,X96-Y09,Y87.1)	5,073	252	237	127	428	892	932	744	732	469	201	58	1
Legal intervention (Y35,Y89.0)	669	—	—	2	101	245	170	85	46	11	6	3	—
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	6,259	98	107	80	528	1,205	1,338	1,139	1,041	444	182	95	2
Discharge of firearms, undetermined intent (Y22-Y24)	458	1	11	22	140	98	64	42	31	24	19	6	—
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	5,801	97	96	58	388	1,107	1,274	1,097	1,010	420	163	89	2
Operations of war and their sequelae. (Y36,Y89.1)	9	—	—	—	—	1	—	2	—	4	1	1	—
Complications of medical and surgical care (Y40-Y84,Y88)	6,051	12	17	23	73	139	244	474	1,079	1,685	1,426	879	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.

[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group (years)											Age not stated
		Younger than 1 year	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ¹	4,105	1	—	1	—	11	53	145	438	960	1,317	1,179	—
COVID-19. (U07.1) ¹	416,893	91	54	142	1,401	6,133	16,006	36,881	73,725	102,716	98,807	80,934	3
Dementia-related causes ¹	279,704	4	14	22	16	20	69	351	3,668	21,885	77,408	176,245	2
Drug-induced deaths ¹	111,219	76	126	108	7,527	24,664	27,771	22,864	20,532	6,243	980	318	10
Drug overdose deaths ¹	106,699	75	126	108	7,426	24,070	26,908	21,875	19,399	5,646	800	256	10
Alcohol-induced deaths ¹	54,258	—	—	2	214	3,027	7,797	12,313	17,960	9,978	2,500	464	3
Firearm-related injuries ¹	48,830	19	134	672	10,132	11,278	7,862	5,898	5,189	3,923	2,627	1,095	1

— Quantity zero.
 ... Category not applicable.
¹Included in selected categories above. For the list of ICD-10 codes included, see Technical Notes.
 SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group (years)										
		Younger than 1 year ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
All causes.	1,043.8	558.8	25.0	14.3	88.9	180.8	287.9	531.0	1,117.1	2,151.3	5,119.4	15,743.3
Salmonella infections. (A01-A02)	0.0	*	*	*	*	*	*	*	*	0.1	*	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections. (A04,A07-A09)	2.1	4.5	*	*	*	0.1	0.3	0.6	1.8	4.7	12.8	33.3
Tuberculosis. (A16-A19)	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	0.8	1.4
Respiratory tuberculosis. (A16)	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.5	1.3
Other tuberculosis (A17-A19)	0.1	*	*	*	*	*	*	0.1	0.1	0.1	0.2	*
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40-A41)	12.4	3.2	0.2	0.1	0.2	0.9	2.6	6.0	15.1	31.1	67.4	153.3
Syphilis (A50-A53)	0.0	*	*	*	*	*	*	*	*	*	*	*
Acute poliomyelitis. (A80)	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis. (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis. (B15-B19)	1.1	*	*	*	*	0.1	0.4	1.1	3.1	3.4	2.1	1.8
Human immunodeficiency virus (HIV) disease (B20-B24)	1.5	*	*	*	0.1	1.1	1.7	2.6	3.6	2.5	1.3	0.7
Malaria. (B50-B54)	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05, A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99,U07.1)	128.9	6.8	1.1	0.6	3.4	13.9	37.8	92.5	176.4	313.3	625.8	1,387.5
Malignant neoplasms. (C00-C97)	182.4	1.5	1.8	1.9	3.1	7.9	25.8	82.5	252.4	532.1	1,017.2	1,712.9
Malignant neoplasms of lip, oral cavity and pharynx . . . (C00-C14)	3.4	*	*	*	*	0.1	0.5	2.2	6.2	10.7	15.5	24.3
Malignant neoplasm of esophagus. (C15)	4.7	*	*	*	*	0.1	0.5	2.5	8.6	15.9	23.2	28.0
Malignant neoplasm of stomach. (C16)	3.3	*	*	*	*	0.3	0.9	2.3	4.8	8.9	16.5	27.5
Malignant neoplasms of colon, rectum and anus (C18-C21)	16.3	*	*	*	0.1	0.8	3.9	11.9	24.5	41.6	77.9	168.7
Malignant neoplasms of liver and intrahepatic bile ducts . . . (C22)	8.7	*	*	*	0.1	0.2	0.8	3.4	15.2	31.4	41.6	50.4
Malignant neoplasm of pancreas (C25)	14.4	*	*	*	*	0.1	1.1	5.8	20.8	46.5	83.6	115.3
Malignant neoplasm of larynx (C32)	1.2	*	*	*	*	*	0.1	0.6	2.3	4.0	5.3	6.9
Malignant neoplasms of trachea, bronchus and lung . . . (C33-C34)	40.6	*	*	*	*	0.3	1.8	11.9	60.5	134.5	248.9	289.5
Malignant melanoma of skin. (C43)	2.5	*	*	*	*	0.2	0.7	1.4	3.4	6.6	12.8	25.7
Malignant neoplasm of breast (C50)	12.9	*	*	*	*	0.8	4.3	10.2	19.8	32.4	58.1	127.3
Malignant neoplasm of cervix uteri. (C53)	1.3	*	*	*	*	0.5	1.5	2.1	2.3	2.5	3.2	4.2
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	3.7	*	*	*	*	0.1	0.5	1.8	5.8	12.8	18.2	24.1
Malignant neoplasm of ovary (C56)	4.0	*	*	*	0.1	0.2	0.7	2.6	6.4	12.2	21.3	28.0
Malignant neoplasm of prostate (C61)	9.8	*	*	*	*	*	*	0.8	6.5	24.4	67.9	170.5
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.3	*	*	0.1	*	0.1	0.5	2.0	6.1	12.4	23.4	40.3
Malignant neoplasm of bladder (C67)	5.1	*	*	*	*	*	0.2	0.8	4.0	11.4	33.5	90.4
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	5.4	*	0.5	0.7	0.6	0.9	1.9	4.3	9.4	15.7	22.0	23.2

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group (years)										
		Younger than 1 year ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	17.3	*	0.5	0.5	0.8	1.3	2.2	5.3	16.0	44.5	114.6	211.5
Hodgkin disease (C81)	0.3	*	*	*	*	0.1	0.1	0.1	0.4	0.8	1.7	2.4
Non-Hodgkin lymphoma (C82-C85)	6.0	*	*	0.1	0.1	0.3	0.7	1.9	5.7	15.1	40.9	76.3
Leukemia (C91-C95)	7.0	*	0.5	0.4	0.6	0.8	1.2	2.2	5.9	17.8	44.8	87.2
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.9	*	*	*	*	*	0.2	1.1	4.1	10.7	27.0	44.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	*	*	*	*	*	*	*	*	0.1	0.3	0.7
All other and unspecified malignant neoplasms (C17, C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60, C62-C63,C66,C68-C69,C73-C80,C97)	23.5	0.6	0.7	0.5	1.2	1.9	3.7	10.4	29.6	63.8	129.7	257.1
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.9	0.7	0.2	0.1	0.2	0.3	0.6	1.3	3.4	10.2	31.3	86.4
Anemias (D50-D64)	1.8	*	*	*	0.1	0.3	0.6	0.8	1.4	3.2	8.8	34.9
Diabetes mellitus (E10-E14)	31.1	*	*	0.1	0.8	2.8	6.8	18.7	43.5	83.0	160.4	309.9
Nutritional deficiencies (E40-E64)	5.3	*	*	*	0.1	0.1	0.2	0.7	2.1	6.5	26.6	161.2
Malnutrition (E40-E46)	5.2	*	*	*	0.1	0.1	0.2	0.7	1.9	6.3	26.2	158.7
Other nutritional deficiencies (E50-E64)	0.1	*	*	*	*	*	*	*	0.2	0.3	0.4	2.5
Meningitis (G00,G03)	0.2	1.5	*	*	*	0.1	0.1	0.1	0.2	0.3	0.7	0.7
Parkinson disease (G20-G21)	11.6	*	*	*	*	*	*	0.2	2.0	18.6	102.5	246.0
Alzheimer disease (G30)	36.0	*	*	*	*	*	*	0.3	3.2	26.4	214.3	1,243.6
Major cardiovascular diseases (I00-I78)	279.0	10.6	1.1	0.7	2.8	11.3	37.3	106.9	262.0	542.2	1,422.4	5,586.2
Diseases of heart (I00-I09,I11,I13,I20-I51)	209.6	7.4	0.8	0.5	2.2	9.1	29.4	84.9	208.7	416.6	1,044.2	4,078.7
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.2	*	*	*	*	0.2	0.3	0.4	1.0	2.2	6.2	22.4
Hypertensive heart disease (I11)	20.4	*	*	*	0.2	1.3	5.0	12.4	24.4	37.9	80.1	397.2
Hypertensive heart and renal disease (I13)	4.2	*	*	*	*	0.1	0.4	1.0	2.3	6.2	21.6	112.4
Ischemic heart diseases (I20-I25)	113.1	*	*	*	0.3	2.3	12.3	45.2	124.6	249.3	584.0	1,985.3
Acute myocardial infarction (I21-I22)	32.9	*	*	*	0.1	1.0	4.7	16.5	42.7	80.3	168.0	456.6
Other acute ischemic heart diseases (I24)	1.4	*	*	*	*	*	0.2	0.8	1.9	3.4	7.2	19.9
Other forms of chronic ischemic heart disease (I20,I25)	78.8	*	*	*	0.2	1.3	7.4	27.9	79.9	165.6	408.8	1,508.8
Atherosclerotic cardiovascular disease, so described (I25.0)	23.6	*	*	*	0.1	0.7	4.2	15.0	39.2	61.4	98.6	275.4
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	55.3	*	*	*	0.1	0.6	3.2	12.9	40.7	104.2	310.2	1,233.4
Other heart diseases (I26-I51)	70.6	7.0	0.7	0.4	1.7	5.3	11.4	25.9	56.5	120.9	352.3	1,561.4
Acute and subacute endocarditis (I33)	0.5	*	*	*	*	0.3	0.4	0.5	0.8	1.1	2.0	2.8
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.4	*	*	*	0.1	0.1	0.2	0.3	0.4	0.7	1.7	3.9
Heart failure (I50)	25.6	*	*	*	0.1	0.5	1.6	5.0	13.5	37.0	134.4	703.2

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group (years)										
		Younger than 1 year ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)	44.1	6.2	0.6	0.4	1.6	4.4	9.2	20.1	41.8	82.1	214.1	851.4
Essential hypertension and hypertensive renal disease (I10,I12,I15)	12.9	*	*	*	0.1	0.4	1.5	4.8	12.1	24.9	63.2	270.8
Cerebrovascular diseases (I60-I69)	49.1	2.7	0.4	0.2	0.4	1.4	5.0	14.1	34.2	84.1	274.8	1,111.1
Atherosclerosis (I70)	1.3	*	*	*	*	*	*	0.2	0.8	2.1	6.7	33.3
Other diseases of circulatory system (I71-I78)	6.2	*	*	*	0.1	0.4	1.3	2.8	6.2	14.5	33.5	92.3
Aortic aneurysm and dissection (I71)	3.0	*	*	*	0.1	0.3	0.9	1.8	3.2	7.2	16.5	38.3
Other diseases of arteries, arterioles and capillaries (I72-I78)	3.1	*	*	*	*	0.2	0.4	1.1	3.0	7.3	17.0	54.0
Other disorders of circulatory system (I80-I99)	1.7	0.6	*	*	0.1	0.5	1.0	1.9	2.5	3.4	5.8	15.8
Influenza and pneumonia (J09-J18)	12.6	3.5	0.3	0.1	0.3	0.9	1.9	4.4	11.3	25.3	69.9	231.8
Influenza (J09-J11)	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.2
Pneumonia (J12-J18)	12.4	3.4	0.3	0.1	0.3	0.9	1.9	4.3	11.1	24.9	68.9	229.6
Other acute lower respiratory infections (J20-J22,U04)	0.1	0.8	*	*	*	*	*	*	*	0.1	0.2	0.8
Acute bronchitis and bronchiolitis (J20-J21)	0.0	0.8	*	*	*	*	*	*	*	0.1	*	0.5
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	*	*	*	*	*	*	*	*	*	*	*
Chronic lower respiratory diseases (J40-J47)	42.9	*	0.2	0.2	0.4	0.8	1.7	7.8	41.2	112.7	285.9	600.6
Bronchitis, chronic and unspecified (J40-J42)	0.1	*	*	*	*	*	*	*	0.1	0.2	0.4	1.8
Emphysema (J43)	2.2	*	*	*	*	*	0.1	0.4	2.3	6.3	14.6	27.4
Asthma (J45-J46)	1.1	*	0.2	0.2	0.3	0.6	0.9	1.1	1.5	1.7	3.0	7.6
Other chronic lower respiratory diseases (J44,J47)	39.5	*	*	*	*	0.1	0.8	6.2	37.3	104.4	267.9	563.8
Pneumoconioses and chemical effects (J60-J66,J68,U07.0)	0.2	*	*	*	*	*	*	*	0.1	0.4	1.2	3.2
Pneumonitis due to solids and liquids (J69)	6.0	*	*	*	0.1	0.3	0.7	1.5	4.6	11.2	33.8	128.2
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	13.8	4.9	0.6	0.2	0.3	0.6	1.5	4.1	11.9	32.0	89.2	209.4
Peptic ulcer (K25-K28)	1.2	*	*	*	*	0.1	0.3	0.7	1.6	2.8	6.2	15.9
Diseases of appendix (K35-K38)	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.6	1.6
Hernia (K40-K46)	0.7	*	*	*	*	*	0.1	0.2	0.7	1.4	4.0	15.1
Chronic liver disease and cirrhosis (K70,K73-K74)	17.0	*	*	*	0.1	4.0	13.4	25.8	41.3	38.4	36.4	31.0
Alcoholic liver disease (K70)	10.0	*	*	*	0.1	3.4	10.6	18.8	26.5	18.2	9.5	4.2
Other chronic liver disease and cirrhosis (K73-K74)	7.1	*	*	*	*	0.7	2.8	7.0	14.8	20.2	26.9	26.8
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.4	*	*	*	*	0.0	0.1	0.3	0.9	2.8	8.0	27.6
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16.4	1.3	*	*	0.1	0.8	2.2	6.2	14.9	36.4	96.1	271.1
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	4.3
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	0.1	*	*	*	*	*	*	0.1	0.1	0.2	0.5	0.6
Renal failure (N17-N19)	16.0	1.2	*	*	0.1	0.8	2.2	6.0	14.6	35.7	94.1	266.0
Other disorders of kidney (N25,N27)	0.0	*	*	*	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group (years)										
		Younger than 1 year ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Infections of kidney (N10-N12,N13.6,N15.1)	0.4	*	*	*	*	*	0.1	0.2	0.5	0.9	2.0	4.4
Hyperplasia of prostate (N40)	0.2	*	*	*	*	*	*	*	0.1	0.3	1.3	6.1
Inflammatory diseases of female pelvic organs (N70-N76)	0.1	*	*	*	*	*	0.1	*	0.1	0.2	0.3	0.6
Pregnancy, childbirth and the puerperium (O00-O99)	0.5	...	...	*	0.5	1.8	1.5	*	*	*	*	*
Pregnancy with abortive outcome (O00-O07)	0.0	...	...	*	*	0.1	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.5	...	...	*	0.5	1.7	1.5	*	*	*	*	*
Certain conditions originating in the perinatal period (P00-P96)	2.9	264.7	0.4	0.1	*	*	*	*	*	*	*	*
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2.9	111.2	2.7	0.8	1.0	1.0	1.2	1.9	2.9	2.2	2.8	5.8
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	10.4	76.1	1.8	0.3	1.4	3.0	4.3	5.3	8.9	14.3	36.3	181.7
All other diseases (residual)	119.8	17.1	2.5	1.7	4.7	12.5	26.1	49.6	98.6	199.4	586.6	2,546.2
Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67.8	36.6	8.5	4.2	36.7	75.7	84.0	77.2	78.2	64.7	126.8	446.0
Transport accidents (V01-V99,Y85)	15.0	3.0	2.5	2.4	17.8	20.3	17.5	16.3	17.2	15.7	19.3	23.7
Motor vehicle accidents (V02-V04,V09.0, V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6, V20-V79, V80.3-V80.5,V81.0-V81.1,V82.0-V82.1, V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)	14.2	3.0	2.4	2.3	17.2	19.6	16.5	15.3	15.7	14.2	18.2	23.0
Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2, V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9, V88.9, V89.1,V89.3,V89.9)	0.4	*	*	*	0.2	0.4	0.5	0.4	0.5	0.6	0.6	*
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.5	*	*	0.1	0.3	0.4	0.6	0.6	1.0	0.9	0.6	0.4
Nontransport accidents (W00-X59,Y86)	52.8	33.6	6.0	1.7	18.9	55.4	66.4	60.8	61.0	49.0	107.6	422.3
Falls (W00-W19)	13.5	*	0.1	0.1	0.3	0.9	1.6	3.3	7.7	19.0	74.9	338.0
Accidental discharge of firearms (W32-W34)	0.2	*	0.3	0.1	0.3	0.2	0.2	0.1	0.1	0.1	0.1	*
Accidental drowning and submersion (W65-W74)	1.3	1.1	3.1	0.6	1.0	1.2	1.4	1.1	1.4	1.5	2.0	2.4
Accidental exposure to smoke, fire and flames (X00-X09)	1.0	*	0.7	0.4	0.2	0.4	0.5	0.8	1.5	2.4	3.4	4.4
Accidental poisoning and exposure to noxious substances (X40-X49)	30.7	0.8	0.2	0.2	16.1	50.9	60.1	51.6	43.4	15.6	3.8	3.5
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64, W75-W99,X10-X39,X50-X59,Y86)	6.1	31.1	1.5	0.4	0.9	1.8	2.7	3.9	6.9	10.5	23.3	73.9
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14.5	...	...	1.5	15.2	19.5	18.1	18.2	17.0	15.3	19.6	22.4
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	7.9	...	...	0.6	8.3	9.3	8.4	9.1	9.3	10.2	15.2	17.3

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group (years)										
		Younger than 1 year ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71,X75–X84,Y87.0)	6.6	...	...	0.9	6.8	10.2	9.7	9.1	7.7	5.1	4.3	5.1
Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	7.8	7.5	2.0	1.2	15.4	16.6	11.2	6.8	4.3	2.6	1.9	1.8
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	6.3	*	0.5	0.9	14.4	14.7	9.1	5.0	2.6	1.2	0.7	0.8
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3,*U01.5–*U01.9,*U02,X85–X92,X96–Y09,Y87.1) ³	1.5	7.1	1.6	0.3	1.0	2.0	2.1	1.8	1.7	1.4	1.2	1.0
Legal intervention (Y35,Y89.0)	0.2	*	*	*	0.2	0.5	0.4	0.2	0.1	*	*	*
Events of undetermined intent ¹⁹ (Y10–Y34,Y87.2,Y89.9)	1.9	2.7	0.7	0.2	1.2	2.6	3.1	2.8	2.4	1.3	1.1	1.6
Discharge of firearms, undetermined intent (Y22–Y24)	0.1	*	*	0.1	0.3	0.2	0.1	0.1	0.1	0.1	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21,Y25–Y34,Y87.2,Y89.9)	1.7	2.7	0.6	0.1	0.9	2.4	2.9	2.7	2.4	1.2	1.0	1.5
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40–Y84,Y88)	1.8	*	*	0.1	0.2	0.3	0.6	1.2	2.5	5.0	8.8	14.7
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ³	1.2	*	*	*	*	*	0.1	0.4	1.0	2.9	8.1	19.7
COVID-19 (U07.1) ³	125.6	2.6	0.4	0.3	3.3	13.5	36.9	90.6	172.2	305.1	609.7	1,354.4
Dementia-related causes ³	84.3	*	*	0.1	*	0.0	0.2	0.9	8.6	65.0	477.6	2,949.3
Drug-induced deaths ³	33.5	2.1	0.8	0.3	17.5	54.2	64.0	56.2	48.0	18.5	6.0	5.3
Drug overdose deaths ³	32.1	2.1	0.8	0.3	17.2	52.9	62.0	53.8	45.3	16.8	4.9	4.3
Alcohol-induced deaths ³	16.3	*	*	*	0.5	6.7	18.0	30.3	42.0	29.6	15.4	7.8
Firearm-related injuries ³	14.7	*	0.9	1.6	23.5	24.8	18.1	14.5	12.1	11.7	16.2	18.3

0.0 Quantity more than zero but less than 0.05.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Data for age not stated included in "all ages" but not distributed among age groups.

²Death rates for younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

³Included in selected categories above. For the list of ICD-10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	3,464,231	1,838,108	1,626,123	315,664	181,195	134,469	26,972	14,724	12,248	92,432	48,386	44,046	449,764	238,599	211,165	5,223	2,906	2,317	2,548,809	1,337,385	1,211,424
Salmonella infections (A01-A02)	72	42	30	12	5	7	1	—	1	4	3	1	12	8	4	1	1	—	41	25	16
Shigellosis and amebiasis (A03,A06)	7	1	6	1	—	1	—	—	—	—	—	—	—	—	—	—	—	—	6	1	5
Certain other intestinal infections (A04, A07-A09)	7,036	2,925	4,111	596	282	314	46	23	23	151	69	82	774	353	421	10	4	6	5,408	2,172	3,236
Tuberculosis (A16-A19)	602	392	210	108	71	37	7	4	3	138	94	44	93	62	31	3	—	3	248	158	90
Respiratory tuberculosis (A16)	435	293	142	77	53	24	6	3	3	121	81	40	70	49	21	3	—	3	157	107	50
Other tuberculosis (A17-A19)	167	99	68	31	18	13	1	1	—	17	13	4	23	13	10	—	—	—	91	51	40
Whooping cough (A37)	4	3	1	1	1	—	—	—	—	—	—	—	1	1	—	—	—	—	2	1	1
Scarlet fever and erysipelas (A38,A46)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Meningococcal infection (A39)	21	10	11	4	3	1	1	1	—	1	1	—	5	1	4	—	—	—	9	3	6
Septicemia (A40-A41)	41,281	20,529	20,752	3,163	1,638	1,525	292	153	139	885	453	432	7,159	3,404	3,755	61	32	29	29,455	14,704	14,751
Syphilis (A50-A53)	43	31	12	10	8	2	1	—	1	1	1	—	21	14	7	—	—	—	10	8	2
Acute poliomyelitis (A80)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Arthropod-borne viral encephalitis (A83-A84, A85.2)	6	4	2	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6	4	2
Measles (B05)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Viral hepatitis (B15-B19)	3,589	2,317	1,272	529	370	159	60	35	25	200	126	74	512	314	198	22	15	7	2,217	1,426	791
Human immunodeficiency virus (HIV) disease (B20-B24)	4,977	3,800	1,177	764	641	123	38	26	12	71	60	11	2,438	1,674	764	7	7	—	1,574	1,328	246
Malaria (B50-B54)	6	3	3	1	—	1	—	—	—	—	—	—	2	1	1	—	—	—	3	2	1

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other and unspecified infectious and parasitic diseases and their sequelae. (A00, A05,A20–A36,A42–A44, A48–A49,A54–A79, A81–A82,A85.0–A85.1, A85.8,A86–B04, B06–B09,B25–B49, B55–B99,U07.1)	427,811	242,562	185,249	69,681	42,651	27,030	4,756	2,500	2,256	13,080	7,786	5,294	57,524	28,412	29,112	1,113	609	504	279,037	159,045	119,992
Malignant neoplasms (C00–C97)	605,213	318,670	286,543	45,544	23,330	22,214	3,162	1,599	1,563	19,271	9,528	9,743	70,036	35,235	34,801	840	375	465	462,601	246,599	216,002
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	11,394	8,141	3,253	680	495	185	45	26	19	449	296	153	1,095	804	291	25	19	6	9,013	6,443	2,570
Malignant neoplasm of esophagus (C15)	15,743	12,558	3,185	779	634	145	89	78	11	339	253	86	1,167	805	362	11	7	4	13,278	10,715	2,563
Malignant neoplasm of stomach (C16)	10,894	6,592	4,302	1,964	1,092	872	96	57	39	880	476	404	1,860	1,110	750	30	20	10	5,991	3,792	2,199
Malignant neoplasms of colon, rectum and anus (C18–C21)	54,121	28,911	25,210	4,810	2,695	2,115	324	186	138	1,976	1,010	966	7,135	3,860	3,275	83	45	38	39,425	20,893	18,532
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	28,719	18,828	9,891	3,907	2,500	1,407	263	162	101	1,765	1,118	647	3,505	2,355	1,150	57	39	18	19,016	12,511	6,505
Malignant neoplasm of pancreas. (C25)	47,906	24,912	22,994	3,745	1,844	1,901	213	104	109	1,690	818	872	5,857	2,824	3,033	68	27	41	36,095	19,170	16,925
Malignant neoplasm of larynx (C32)	3,887	3,126	761	235	210	25	17	11	6	59	53	6	612	486	126	2	2	–	2,937	2,343	594
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	134,592	71,601	62,991	5,781	3,208	2,573	668	330	338	3,779	2,058	1,721	14,248	8,003	6,245	133	63	70	109,126	57,487	51,639
Malignant melanoma of skin. (C43)	8,224	5,363	2,861	295	179	116	18	13	5	42	21	21	131	62	69	2	1	1	7,703	5,063	2,640
Malignant neoplasm of breast (C50)	42,810	500	42,310	3,405	26	3,379	201	1	200	1,368	12	1,356	6,326	84	6,242	93	–	93	31,134	374	30,760
Malignant neoplasm of cervix uteri (C53)	4,366	...	4,366	611	...	611	53	...	53	202	...	202	762	...	762	19	...	19	2,680	...	2,680

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	12,208	...	12,208	1,115	...	1,115	66	...	66	412	...	412	2,509	...	2,509	61	...	61	7,974	...	7,974
Malignant neoplasm of ovary (C56)	13,430	...	13,430	1,217	...	1,217	82	...	82	518	...	518	1,343	...	1,343	17	...	17	10,180	...	10,180
Malignant neoplasm of prostate (C61)	32,563	32,563	...	2,284	2,284	...	136	136	...	670	670	...	5,292	5,292	...	43	43	...	23,941	23,941	...
Malignant neoplasms of kidney and renal pelvis (C64–C65)	14,163	9,370	4,793	1,435	917	518	124	84	40	299	189	110	1,291	809	482	9	4	5	10,912	7,297	3,615
Malignant neoplasm of bladder (C67)	16,840	12,196	4,644	844	570	274	50	35	15	300	201	99	1,270	789	481	9	7	2	14,297	10,547	3,750
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	17,857	10,098	7,759	1,551	908	643	66	40	26	477	258	219	1,142	597	545	15	10	5	14,487	8,217	6,270
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	57,375	32,988	24,387	4,721	2,586	2,135	232	125	107	1,680	917	763	5,989	3,159	2,830	63	36	27	44,368	25,981	18,387
Hodgkin disease (C81)	1,003	596	407	130	78	52	6	4	2	21	8	13	125	81	44	1	1	–	713	418	295
Non-Hodgkin lymphoma (C82–C85)	20,020	11,519	8,501	1,739	931	808	80	45	35	731	412	319	1,476	829	647	19	9	10	15,850	9,220	6,630
Leukemia (C91–C95)	23,271	13,540	9,731	1,832	1,012	820	94	46	48	639	342	297	1,964	1,029	935	30	18	12	18,584	11,019	7,565
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	12,933	7,244	5,689	1,011	558	453	50	30	20	286	154	132	2,412	1,213	1,199	12	8	4	9,100	5,250	3,850
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	148	89	59	9	7	2	2	–	2	3	1	2	12	7	5	1	–	1	121	74	47

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																					
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	
All other and unspecified malignant neoplasms (C17, C23–C24,C26–C31, C37–C41,C44–C49, C51–C52,C57–C60, C62–C63,C66, C68–C69, C73–C80,C97)	78,121	40,923	37,198	6,165	3,182	2,983	419	211	208	2,366	1,178	1,188	8,502	4,196	4,306	100	52	48	60,044	31,825	28,219	
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	16,237	8,720	7,517	1,039	529	510	57	37	20	500	262	238	1,520	713	807	18	11	7	13,021	7,122	5,899	
Anemias (D50–D64)	5,987	2,720	3,267	414	194	220	34	17	17	164	84	80	1,240	547	693	6	2	4	4,099	1,859	2,240	
Diabetes mellitus (E10–E14)	103,294	58,628	44,666	12,489	7,029	5,460	1,269	689	580	3,639	1,963	1,676	18,968	9,843	9,125	314	175	139	65,789	38,428	27,361	
Nutritional deficiencies (E40–E64)	17,505	6,605	10,900	1,047	423	624	95	42	53	374	148	226	1,872	747	1,125	10	2	8	14,019	5,206	8,813	
Malnutrition (E40–E46)	17,106	6,428	10,678	1,025	412	613	90	40	50	371	148	223	1,822	726	1,096	10	2	8	13,704	5,065	8,639	
Other nutritional deficiencies (E50–E64)	399	177	222	22	11	11	5	2	3	3	–	3	50	21	29	–	–	–	315	141	174	
Meningitis (G00,G03)	534	314	220	83	45	38	11	7	4	21	11	10	94	51	43	2	1	1	314	193	121	
Parkinson disease. (G20–G21)	38,536	23,567	14,969	2,369	1,365	1,004	123	62	61	1,169	683	486	1,893	1,095	798	21	11	10	32,818	20,268	12,550	
Alzheimer disease. (G30)	119,399	36,975	82,424	9,142	2,775	6,367	370	101	269	3,203	968	2,235	9,638	2,791	6,847	85	29	56	96,522	30,170	66,352	
Major cardiovascular diseases (I00–I78)	925,923	488,914	437,009	66,425	36,411	30,014	4,928	2,757	2,171	25,918	13,422	12,496	124,348	65,454	58,894	1,344	788	556	696,709	366,401	330,308	
Diseases of heart (I00–I09,I11, I13,I20–I51)	695,547	384,886	310,661	47,345	27,211	20,134	3,760	2,234	1,526	17,009	9,383	7,626	90,507	49,490	41,017	1,002	617	385	531,162	292,973	238,189	
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	3,907	1,382	2,525	266	95	171	25	10	15	117	37	80	333	114	219	16	5	11	3,119	1,110	2,009	
Hypertensive heart disease (I11)	67,774	34,520	33,254	4,726	2,622	2,104	509	306	203	1,499	739	760	12,800	7,073	5,727	98	64	34	47,536	23,338	24,198	

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Hypertensive heart and renal disease.(I13)	13,918	6,473	7,445	949	441	508	119	61	58	310	130	180	2,268	1,117	1,151	17	13	4	10,166	4,658	5,508
Ischemic heart diseases(I20–I25)	375,476	226,452	149,024	28,094	17,095	10,999	2,012	1,250	762	10,547	6,305	4,242	44,468	25,543	18,925	543	349	194	287,088	174,148	112,940
Acute myocardial infarction(I21–I22)	109,097	65,673	43,424	8,388	5,051	3,337	601	373	228	3,238	1,942	1,296	12,933	7,295	5,638	147	82	65	83,165	50,529	32,636
Other acute ischemic heart diseases(I24)	4,748	2,704	2,044	269	162	107	26	10	16	112	64	48	738	420	318	5	3	2	3,571	2,029	1,542
Other forms of chronic ischemic heart disease(I20,I25)	261,631	158,075	103,556	19,437	11,882	7,555	1,385	867	518	7,197	4,299	2,898	30,797	17,828	12,969	391	264	127	200,352	121,590	78,762
Atherosclerotic cardiovascular disease, so described(I25.0)	78,175	49,832	28,343	6,670	4,549	2,121	509	327	182	2,051	1,302	749	12,322	7,736	4,586	123	86	37	55,640	35,236	20,404
All other forms of chronic ischemic heart disease. . . . (I20,I25.1–I25.9)	183,456	108,243	75,213	12,767	7,333	5,434	876	540	336	5,146	2,997	2,149	18,475	10,092	8,383	268	178	90	144,712	86,354	58,358
Other heart diseases(I26–I51)	234,472	116,059	118,413	13,310	6,958	6,352	1,095	607	488	4,536	2,172	2,364	30,638	15,643	14,995	328	186	142	183,253	89,719	93,534
Acute and subacute endocarditis(I33)	1,767	1,072	695	166	106	60	27	16	11	33	21	12	243	147	96	2	1	1	1,280	770	510
Diseases of pericardium and acute myocarditis(I30–I31,I40)	1,237	639	598	109	55	54	7	1	6	26	12	14	168	95	73	4	3	1	911	464	447
Heart failure.(I50)	85,037	40,344	44,693	4,647	2,249	2,398	363	211	152	1,603	734	869	10,110	4,902	5,208	82	47	35	67,866	31,993	35,873
All other forms of heart disease(I26–I28, I34–I38,I42–I49,I51)	146,431	74,004	72,427	8,388	4,548	3,840	698	379	319	2,874	1,405	1,469	20,117	10,499	9,618	240	135	105	113,196	56,492	56,704

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Essential hypertension and hypertensive renal disease (I10,I12,I15)	42,816	20,086	22,730	3,718	1,846	1,872	256	129	127	1,828	858	970	7,868	3,875	3,993	65	29	36	28,798	13,214	15,584
Cerebrovascular diseases (I60–I69)	162,890	70,852	92,038	13,776	6,433	7,343	799	351	448	6,428	2,848	3,580	22,837	10,428	12,409	247	124	123	117,809	50,219	67,590
Atherosclerosis. (I70)	4,214	2,008	2,206	247	119	128	17	5	12	136	64	72	448	213	235	3	1	2	3,320	1,580	1,740
Other diseases of circulatory system (I71–I78)	20,456	11,082	9,374	1,339	802	537	96	38	58	517	269	248	2,688	1,448	1,240	27	17	10	15,620	8,415	7,205
Aortic aneurysm and dissection (I71)	10,037	6,039	3,998	580	412	168	27	10	17	319	193	126	1,150	700	450	18	14	4	7,844	4,649	3,195
Other diseases of arteries, arterioles and capillaries . . . (I72–I78)	10,419	5,043	5,376	759	390	369	69	28	41	198	76	122	1,538	748	790	9	3	6	7,776	3,766	4,010
Other disorders of circulatory system (I80–I99)	5,655	2,935	2,720	471	269	202	39	20	19	86	46	40	1,160	590	570	11	2	9	3,848	1,982	1,866
Influenza and pneumonia. (J09–J18)	41,917	22,373	19,544	3,664	2,022	1,642	333	189	144	1,586	880	706	5,310	2,786	2,524	57	39	18	30,673	16,286	14,387
Influenza. (J09–J11)	608	294	314	61	24	37	10	5	5	16	10	6	73	38	35	1	–	1	445	216	229
Pneumonia. (J12–J18)	41,309	22,079	19,230	3,603	1,998	1,605	323	184	139	1,570	870	700	5,237	2,748	2,489	56	39	17	30,228	16,070	14,158
Other acute lower respiratory infections (J20–J22,U04)	189	88	101	26	14	12	2	2	–	4	4	–	30	11	19	–	–	–	123	56	67
Acute bronchitis and bronchiolitis (J20–J21)	128	56	72	19	9	10	2	2	–	3	3	–	23	7	16	–	–	–	77	34	43
Other and unspecified acute lower respiratory infections (J22,U04)	61	32	29	7	5	2	–	–	–	1	1	–	7	4	3	–	–	–	46	22	24
Chronic lower respiratory diseases. (J40–J47)	142,342	67,528	74,814	5,479	2,749	2,730	825	372	453	1,754	1,046	708	11,218	5,586	5,632	90	51	39	122,054	57,231	64,823
Bronchitis, chronic and unspecified (J40–J42)	335	141	194	26	11	15	4	1	3	7	2	5	51	29	22	1	–	1	244	97	147
Emphysema (J43)	7,336	3,945	3,391	226	146	80	50	23	27	87	70	17	504	288	216	3	2	1	6,405	3,376	3,029
Asthma (J45–J46)	3,517	1,430	2,087	366	156	210	19	5	14	133	68	65	1,020	469	551	6	3	3	1,929	711	1,218
Other chronic lower respiratory diseases (J44,J47)	131,154	62,012	69,142	4,861	2,436	2,425	752	343	409	1,527	906	621	9,643	4,800	4,843	80	46	34	113,476	53,047	60,429

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Pneumoconioses and chemical effects(J60–J66,J68,U07.0)	564	527	37	27	23	4	3	3	–	2	1	1	26	24	2	–	–	–	505	475	30
Pneumonitis due to solids and liquids.....(J69)	19,969	11,663	8,306	1,276	726	550	105	63	42	612	356	256	2,399	1,298	1,101	16	7	9	15,447	9,135	6,312
Other diseases of respiratory system(J00–J06, J30–J39,J67,J70–J98)	45,888	24,205	21,683	3,750	2,008	1,742	332	173	159	1,161	625	536	4,775	2,257	2,518	35	16	19	35,586	18,976	16,610
Peptic ulcer(K25–K28)	4,008	2,092	1,916	350	205	145	36	19	17	149	78	71	421	233	188	8	7	1	3,005	1,529	1,476
Diseases of appendix(K35–K38)	449	265	184	53	33	20	1	–	1	15	6	9	48	34	14	1	–	1	327	189	138
Hernia(K40–K46)	2,488	1,132	1,356	197	101	96	16	6	10	37	22	15	208	110	98	2	1	1	2,010	878	1,132
Chronic liver disease and cirrhosis.....(K70, K73–K74)	56,585	35,707	20,878	8,864	6,187	2,677	1,899	1,049	850	881	552	329	4,524	2,707	1,817	51	39	12	39,824	24,824	15,000
Alcoholic liver disease.....(K70)	33,098	22,598	10,500	5,276	4,206	1,070	1,503	855	648	417	325	92	2,624	1,644	980	26	21	5	22,873	15,295	7,578
Other chronic liver disease and cirrhosis(K73–K74)	23,487	13,109	10,378	3,588	1,981	1,607	396	194	202	464	227	237	1,900	1,063	837	25	18	7	16,951	9,529	7,422
Cholelithiasis and other disorders of gallbladder(K80–K82)	4,490	2,362	2,128	452	221	231	51	23	28	158	86	72	430	211	219	7	5	2	3,368	1,806	1,562
Nephritis, nephrotic syndrome and nephrosis(N00–N07, N17–N19,N25–N27)	54,358	28,589	25,769	5,021	2,712	2,309	395	185	210	1,639	856	783	10,434	5,189	5,245	117	57	60	36,444	19,406	17,038
Acute and rapidly progressive nephritic and nephrotic syndrome(N00–N01,N04)	781	380	401	62	35	27	6	3	3	16	7	9	118	51	67	1	1	–	576	282	294
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified(N02–N03, N05–N07,N26)	290	155	135	30	14	16	4	1	3	18	6	12	37	18	19	–	–	–	200	115	85

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Renal failure (N17–N19)	53,261	28,039	25,222	4,926	2,662	2,264	385	181	204	1,605	843	762	10,273	5,117	5,156	116	56	60	35,651	18,998	16,653
Other disorders of kidney (N25,N27)	26	15	11	3	1	2	–	–	–	–	–	–	6	3	3	–	–	–	17	11	6
Infections of kidney (N10–N12, N13.6,N15.1)	1,248	414	834	127	48	79	15	5	10	34	6	28	114	42	72	4	2	2	937	305	632
Hyperplasia of prostate (N40)	715	715	...	61	61	...	–	–	...	34	34	...	78	78	...	1	1	...	533	533	...
Inflammatory diseases of female pelvic organs (N70–N76)	214	...	214	17	...	17	3	...	3	5	...	5	43	...	43	–	...	–	144	...	144
Pregnancy, childbirth and the puerperium (O00–O99)	1,687	...	1,687	352	...	352	42	...	42	52	...	52	519	...	519	6	...	6	687	...	687
Pregnancy with abortive outcome (O00–O07)	48	...	48	11	...	11	1	...	1	2	...	2	18	...	18	–	...	–	15	...	15
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	1,639	...	1,639	341	...	341	41	...	41	50	...	50	501	...	501	6	...	6	672	...	672
Certain conditions originating in the perinatal period (P00–P96)	9,579	5,276	4,303	2,237	1,239	998	57	25	32	390	222	168	2,807	1,559	1,248	27	17	10	3,557	1,941	1,616
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	9,664	5,014	4,650	1,678	871	807	83	43	40	235	127	108	1,424	707	717	17	11	6	5,986	3,126	2,860
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	34,503	17,237	17,266	2,916	1,796	1,120	306	179	127	687	341	346	5,407	2,926	2,481	52	25	27	24,678	11,683	12,995
All other diseases (residual)	397,499	175,191	222,308	27,500	13,171	14,329	3,020	1,530	1,490	8,432	3,674	4,758	45,773	20,443	25,330	397	203	194	309,852	134,877	174,975
Accidents (unintentional injuries) (V01–X59, Y85–Y86)	224,935	149,602	75,333	27,126	20,459	6,667	2,965	1,907	1,058	3,768	2,452	1,316	33,830	24,160	9,670	322	244	78	153,897	98,274	55,623
Transport accidents (V01–V99,Y85)	49,845	36,082	13,763	8,488	6,372	2,116	891	580	311	992	660	332	8,939	6,646	2,293	104	81	23	29,806	21,320	8,486

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motor vehicle accidents (V02–V04,V09.0, V09.2,V12–V14, V19.0–V19.2, V19.4–V19.6, V20–V79, V80.3–V80.5, V81.0–V81.1, V82.0–V82.1, V83–V86, V87.0–V87.8, V88.0–V88.8, V89.0,V89.2)	46,980	33,757	13,223	8,143	6,095	2,048	833	530	303	932	614	318	8,583	6,335	2,248	93	72	21	27,805	19,714	8,091
Other land transport accidents. (V01, V05–V06,V09.1, V09.3–V09.9, V10–V11,V15–V18, V19.3,V19.8–V19.9, V80.0–V80.2, V80.6–V80.9, V81.2–V81.9, V82.2–V82.9, V87.9, V88.9, V89.1,V89.3,V89.9)	1,179	956	223	193	158	35	31	27	4	29	21	8	179	159	20	6	6	–	727	572	155
Water, air and space, and other and unspecified transport accidents and their sequelae (V90–V99,Y85)	1,686	1,369	317	152	119	33	27	23	4	31	25	6	177	152	25	5	3	2	1,274	1,034	240
Nontransport accidents (W00–X59,Y86)	175,090	113,520	61,570	18,638	14,087	4,551	2,074	1,327	747	2,776	1,792	984	24,891	17,514	7,377	218	163	55	124,091	76,954	47,137
Falls (W00–W19)	44,686	22,907	21,779	2,768	1,714	1,054	246	135	111	1,206	670	536	2,160	1,258	902	41	28	13	38,059	18,979	19,080
Accidental discharge of firearms (W32–W34)	549	482	67	65	58	7	7	6	1	7	4	3	150	133	17	1	1	–	310	275	35
Accidental drowning and submersion (W65–W74)	4,337	3,236	1,101	623	490	133	55	45	10	217	153	64	736	582	154	17	13	4	2,610	1,894	716

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental exposure to smoke, fire and flames (X00–X09)	3,389	2,101	1,288	268	172	96	47	25	22	56	28	28	701	428	273	4	3	1	2,273	1,418	855
Accidental poisoning and exposure to noxious substances (X40–X49)	102,001	72,515	29,486	13,071	10,390	2,681	1,463	935	528	862	689	173	18,368	13,396	4,972	126	99	27	66,277	45,689	20,588
Other and unspecified nontransport accidents and their sequelae (W20–W31, W35–W64,W75–W99, X10–X39, X50–X59,Y86)	20,128	12,279	7,849	1,843	1,263	580	256	181	75	428	248	180	2,776	1,717	1,059	29	19	10	14,562	8,699	5,863
Intentional self-harm (suicide). (*U03, X60–X84,Y87.0)	48,183	38,358	9,825	4,907	3,967	940	692	525	167	1,379	962	417	3,692	2,972	720	82	65	17	36,681	29,292	7,389
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	26,328	22,936	3,392	2,037	1,795	242	241	209	32	352	310	42	2,165	1,885	280	27	25	2	21,197	18,448	2,749
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71, X75–X84,Y87.0)	21,855	15,422	6,433	2,870	2,172	698	451	316	135	1,027	652	375	1,527	1,087	440	55	40	15	15,484	10,844	4,640
Assault (homicide) (*U01–*U02, X85–Y09,Y87.1)	26,031	21,084	4,947	4,453	3,694	759	307	241	66	313	205	108	14,313	12,277	2,036	46	40	6	6,215	4,340	1,875
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	20,958	17,596	3,362	3,455	2,992	463	185	147	38	204	146	58	12,721	11,110	1,611	37	34	3	4,064	2,945	1,119

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3, *U01.5–*U01.9, *U02.X85–X92, X96–Y09,Y87.1)	5,073	3,488	1,585	998	702	296	122	94	28	109	59	50	1,592	1,167	425	9	6	3	2,151	1,395	756
Legal intervention (Y35,Y89.0)	669	630	39	151	143	8	13	13	–	9	9	–	163	159	4	1	1	–	317	290	27
Events of undetermined intent (Y10–Y34, Y87.2,Y89.9)	6,259	4,133	2,126	529	394	135	119	67	52	74	50	24	1,502	1,062	440	7	5	2	3,926	2,486	1,440
Discharge of firearms, undetermined intent (Y22–Y24)	458	339	119	55	47	8	20	15	5	6	2	4	134	102	32	2	2	–	231	164	67
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21, Y25–Y34,Y87.2,Y89.9)	5,801	3,794	2,007	474	347	127	99	52	47	68	48	20	1,368	960	408	5	3	2	3,695	2,322	1,373
Operations of war and their sequelae (Y36,Y89.1)	9	9	–	–	–	–	–	–	–	–	–	–	3	3	–	–	–	–	6	6	–
Complications of medical and surgical care (Y40–Y84,Y88)	6,051	3,247	2,804	558	311	247	62	32	30	146	84	62	963	455	508	9	6	3	4,271	2,336	1,935

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	4,105	1,808	2,297	331	162	169	30	15	15	81	40	41	410	189	221	5	2	3	3,226	1,393	1,833
COVID-19..... (U07.1) ⁴	416,893	236,610	180,283	68,656	42,070	26,586	4,624	2,432	2,192	12,757	7,595	5,162	55,994	27,613	28,381	1,087	595	492	271,253	154,806	116,447
Dementia-related causes ⁴	279,704	92,303	187,401	17,876	5,815	12,061	874	264	610	6,793	2,189	4,604	23,685	7,506	16,179	170	61	109	229,199	76,082	153,117
Drug-induced deaths ⁴	111,219	77,309	33,910	13,436	10,513	2,923	1,448	876	572	1,017	757	260	19,944	14,457	5,487	137	106	31	73,225	49,176	24,049
Drug overdose deaths ⁴	106,699	74,301	32,398	12,986	10,156	2,830	1,358	824	534	971	719	252	19,212	13,929	5,283	127	97	30	70,121	47,217	22,904
Alcohol-induced deaths ⁴	54,258	38,700	15,558	7,533	6,148	1,385	2,221	1,377	844	638	504	134	5,023	3,477	1,546	40	33	7	38,117	26,671	11,446
Firearm-related injuries ⁴	48,830	41,866	6,964	5,741	5,014	727	466	390	76	576	469	107	15,290	13,349	1,941	68	63	5	26,054	22,068	3,986

— Quantity zero.

... Category not applicable.

¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes people of Hispanic origin of any race; see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Included in selected categories above. For the list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes.	1,043.8	1,118.2	970.8	503.9	571.3	434.7	1,100.0	1,216.9	986.2	469.5	511.7	430.5	1,074.5	1,185.8	971.4	834.0	917.5	748.6	1,294.9	1,368.8	1,222.1
Salmonella infections (A01–A02)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	*
Shigellosis and amebiasis. (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07–A09)	2.1	1.8	2.5	1.0	0.9	1.0	1.9	1.9	1.9	0.8	0.7	0.8	1.8	1.8	1.9	*	*	*	2.7	2.2	3.3
Tuberculosis. (A16–A19)	0.2	0.2	0.1	0.2	0.2	0.1	*	*	*	0.7	1.0	0.4	0.2	0.3	0.1	*	*	*	0.1	0.2	0.1
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.1	0.2	0.1	*	*	*	0.6	0.9	0.4	0.2	0.2	0.1	*	*	*	0.1	0.1	0.1
Other tuberculosis (A17–A19)	0.1	0.1	0.0	0.0	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	0.0	0.1	0.0
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas. (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40–A41)	12.4	12.5	12.4	5.0	5.2	4.9	11.9	12.6	11.2	4.5	4.8	4.2	17.1	16.9	17.3	9.7	10.1	9.4	15.0	15.0	14.9
Syphilis (A50–A53)	0.0	0.0	*	*	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	*	*	*
Acute poliomyelitis. (A80)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83–A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis. (B15–B19)	1.1	1.4	0.8	0.8	1.2	0.5	2.4	2.9	2.0	1.0	1.3	0.7	1.2	1.6	0.9	3.5	*	*	1.1	1.5	0.8
Human immunodeficiency virus (HIV) disease. (B20–B24)	1.5	2.3	0.7	1.2	2.0	0.4	1.5	2.1	*	0.4	0.6	*	5.8	8.3	3.5	*	*	*	0.8	1.4	0.2
Malaria. (B50–B54)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05,A20–A36, A42–A44,A48–A49,A54–A79, A81–A82,A85.0–A85.1,A85.8, A86–B04,B06–B09,B25–B49, B55–B99,U07.1)	128.9	147.6	110.6	111.2	134.5	87.4	194.0	206.6	181.7	66.4	82.3	51.7	137.4	141.2	133.9	177.7	192.3	162.8	141.8	162.8	121.0
Malignant neoplasms (C00–C97)	182.4	193.9	171.1	72.7	73.6	71.8	129.0	132.2	125.9	97.9	100.8	95.2	167.3	175.1	160.1	134.1	118.4	150.2	235.0	252.4	217.9
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	3.4	5.0	1.9	1.1	1.6	0.6	1.8	2.1	*	2.3	3.1	1.5	2.6	4.0	1.3	4.0	*	*	4.6	6.6	2.6
Malignant neoplasm of esophagus (C15)	4.7	7.6	1.9	1.2	2.0	0.5	3.6	6.4	*	1.7	2.7	0.8	2.8	4.0	1.7	*	*	*	6.7	11.0	2.6

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasm of stomach (C16)	3.3	4.0	2.6	3.1	3.4	2.8	3.9	4.7	3.1	4.5	5.0	3.9	4.4	5.5	3.5	4.8	6.3	*	3.0	3.9	2.2
Malignant neoplasms of colon, rectum and anus. (C18–C21)	16.3	17.6	15.0	7.7	8.5	6.8	13.2	15.4	11.1	10.0	10.7	9.4	17.0	19.2	15.1	13.3	14.2	12.3	20.0	21.4	18.7
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	8.7	11.5	5.9	6.2	7.9	4.5	10.7	13.4	8.1	9.0	11.8	6.3	8.4	11.7	5.3	9.1	12.3	*	9.7	12.8	6.6
Malignant neoplasm of pancreas. (C25)	14.4	15.2	13.7	6.0	5.8	6.1	8.7	8.6	8.8	8.6	8.7	8.5	14.0	14.0	14.0	10.9	8.5	13.2	18.3	19.6	17.1
Malignant neoplasm of larynx (C32)	1.2	1.9	0.5	0.4	0.7	0.1	*	*	*	0.3	0.6	*	1.5	2.4	0.6	*	*	*	1.5	2.4	0.6
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	40.6	43.6	37.6	9.2	10.1	8.3	27.2	27.3	27.2	19.2	21.8	16.8	34.0	39.8	28.7	21.2	19.9	22.6	55.4	58.8	52.1
Malignant melanoma of skin. (C43)	2.5	3.3	1.7	0.5	0.6	0.4	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*	*	3.9	5.2	2.7
Malignant neoplasm of breast (C50)	12.9	0.3	25.3	5.4	0.1	10.9	8.2	*	16.1	6.9	*	13.3	15.1	0.4	28.7	14.9	*	30.0	15.8	0.4	31.0
Malignant neoplasm of cervix uteri (C53)	1.3	...	2.6	1.0	...	2.0	2.2	...	4.3	1.0	...	2.0	1.8	...	3.5	*	...	*	1.4	...	2.7
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	3.7	...	7.3	1.8	...	3.6	2.7	...	5.3	2.1	...	4.0	6.0	...	11.5	9.7	...	19.7	4.1	...	8.0
Malignant neoplasm of ovary (C56)	4.0	...	8.0	1.9	...	3.9	3.3	...	6.6	2.6	...	5.1	3.2	...	6.2	*	...	*	5.2	...	10.3
Malignant neoplasm of prostate (C61)	9.8	19.8	...	3.6	7.2	...	5.5	11.2	...	3.4	7.1	...	12.6	26.3	...	6.9	13.6	...	12.2	24.5	...
Malignant neoplasms of kidney and renal pelvis (C64–C65)	4.3	5.7	2.9	2.3	2.9	1.7	5.1	6.9	3.2	1.5	2.0	1.1	3.1	4.0	2.2	*	*	*	5.5	7.5	3.6
Malignant neoplasm of bladder (C67)	5.1	7.4	2.8	1.3	1.8	0.9	2.0	2.9	*	1.5	2.1	1.0	3.0	3.9	2.2	*	*	*	7.3	10.8	3.8
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	5.4	6.1	4.6	2.5	2.9	2.1	2.7	3.3	2.1	2.4	2.7	2.1	2.7	3.0	2.5	*	*	*	7.4	8.4	6.3
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	17.3	20.1	14.6	7.5	8.2	6.9	9.5	10.3	8.6	8.5	9.7	7.5	14.3	15.7	13.0	10.1	11.4	8.7	22.5	26.6	18.5
Hodgkin disease (C81)	0.3	0.4	0.2	0.2	0.2	0.2	*	*	*	0.1	*	*	0.3	0.4	0.2	*	*	*	0.4	0.4	0.3

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Non-Hodgkin lymphoma (C82–C85)	6.0	7.0	5.1	2.8	2.9	2.6	3.3	3.7	2.8	3.7	4.4	3.1	3.5	4.1	3.0	*	*	*	8.1	9.4	6.7
Leukemia (C91–C95)	7.0	8.2	5.8	2.9	3.2	2.7	3.8	3.8	3.9	3.2	3.6	2.9	4.7	5.1	4.3	4.8	*	*	9.4	11.3	7.6
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.9	4.4	3.4	1.6	1.8	1.5	2.0	2.5	1.6	1.5	1.6	1.3	5.8	6.0	5.5	*	*	*	4.6	5.4	3.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.1	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.0
All other and unspecified malignant neoplasms (C17, C23–C24,C26–C31,C37–C41, C44–C49,C51–C52,C57–C60, C62–C63,C66,C68–C69, C73–C80,C97)	23.5	24.9	22.2	9.8	10.0	9.6	17.1	17.4	16.7	12.0	12.5	11.6	20.3	20.9	19.8	16.0	16.4	15.5	30.5	32.6	28.5
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	4.9	5.3	4.5	1.7	1.7	1.6	2.3	3.1	1.6	2.5	2.8	2.3	3.6	3.5	3.7	*	*	*	6.6	7.3	6.0
Anemias (D50–D64)	1.8	1.7	2.0	0.7	0.6	0.7	1.4	*	*	0.8	0.9	0.8	3.0	2.7	3.2	*	*	*	2.1	1.9	2.3
Diabetes mellitus (E10–E14)	31.1	35.7	26.7	19.9	22.2	17.7	51.8	56.9	46.7	18.5	20.8	16.4	45.3	48.9	42.0	50.1	55.3	44.9	33.4	39.3	27.6
Nutritional deficiencies (E40–E64)	5.3	4.0	6.5	1.7	1.3	2.0	3.9	3.5	4.3	1.9	1.6	2.2	4.5	3.7	5.2	*	*	*	7.1	5.3	8.9
Malnutrition (E40–E46)	5.2	3.9	6.4	1.6	1.3	2.0	3.7	3.3	4.0	1.9	1.6	2.2	4.4	3.6	5.0	*	*	*	7.0	5.2	8.7
Other nutritional deficiencies (E50–E64)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	0.1	0.1	*	*	*	0.2	0.1	0.2
Meningitis (G00,G03)	0.2	0.2	0.1	0.1	0.1	0.1	*	*	*	0.1	*	*	0.2	0.3	0.2	*	*	*	0.2	0.2	0.1
Parkinson disease (G20–G21)	11.6	14.3	8.9	3.8	4.3	3.2	5.0	5.1	4.9	5.9	7.2	4.8	4.5	5.4	3.7	3.4	*	*	16.7	20.7	12.7
Alzheimer disease (G30)	36.0	22.5	49.2	14.6	8.7	20.6	15.1	8.3	21.7	16.3	10.2	21.8	23.0	13.9	31.5	13.6	9.2	18.1	49.0	30.9	66.9
Major cardiovascular diseases (I00–I78)	279.0	297.4	260.9	106.0	114.8	97.0	201.0	227.9	174.8	131.7	142.0	122.1	297.1	325.3	270.9	214.6	248.8	179.6	354.0	375.0	333.2
Diseases of heart (I00–I09, I11,I13,I20–I51)	209.6	234.1	185.5	75.6	85.8	65.1	153.3	184.6	122.9	86.4	99.2	74.5	216.2	246.0	188.7	160.0	194.8	124.4	269.9	299.9	240.3
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	1.2	0.8	1.5	0.4	0.3	0.6	1.0	*	*	0.6	0.4	0.8	0.8	0.6	1.0	*	*	*	1.6	1.1	2.0
Hypertensive heart disease (I11)	20.4	21.0	19.9	7.5	8.3	6.8	20.8	25.3	16.3	7.6	7.8	7.4	30.6	35.2	26.3	15.6	20.2	11.0	24.2	23.9	24.4
Hypertensive heart and renal disease (I13)	4.2	3.9	4.4	1.5	1.4	1.6	4.9	5.0	4.7	1.6	1.4	1.8	5.4	5.6	5.3	*	*	*	5.2	4.8	5.6

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Ischemic heart diseases (I20–I25)	113.1	137.8	89.0	44.8	53.9	35.6	82.1	103.3	61.4	53.6	66.7	41.5	106.2	126.9	87.1	86.7	110.2	62.7	145.9	178.2	113.9
Acute myocardial infarction (I21–I22)	32.9	40.0	25.9	13.4	15.9	10.8	24.5	30.8	18.4	16.4	20.5	12.7	30.9	36.3	25.9	23.5	25.9	21.0	42.3	51.7	32.9
Other acute ischemic heart diseases (I24)	1.4	1.6	1.2	0.4	0.5	0.3	1.1	*	*	0.6	0.7	0.5	1.8	2.1	1.5	*	*	*	1.8	2.1	1.6
Other forms of chronic ischemic heart disease (I20,I25)	78.8	96.2	61.8	31.0	37.5	24.4	56.5	71.7	41.7	36.6	45.5	28.3	73.6	88.6	59.7	62.4	83.4	41.0	101.8	124.4	79.5
Atherosclerotic cardiovascular disease, so described (I25.0)	23.6	30.3	16.9	10.6	14.3	6.9	20.8	27.0	14.7	10.4	13.8	7.3	29.4	38.4	21.1	19.6	27.2	12.0	28.3	36.1	20.6
All other forms of chronic ischemic heart disease (I20,I25.1–I25.9)	55.3	65.8	44.9	20.4	23.1	17.6	35.7	44.6	27.1	26.1	31.7	21.0	44.1	50.2	38.6	42.8	56.2	29.1	73.5	88.4	58.9
Other heart diseases (I26–I51)	70.6	70.6	70.7	21.2	21.9	20.5	44.7	50.2	39.3	23.0	23.0	23.1	73.2	77.7	69.0	52.4	58.7	45.9	93.1	91.8	94.4
Acute and subacute endocarditis (I33)	0.5	0.7	0.4	0.3	0.3	0.2	1.1	*	*	0.2	0.2	*	0.6	0.7	0.4	*	*	*	0.7	0.8	0.5
Diseases of pericardium and acute myocarditis (I30–I31,I40)	0.4	0.4	0.4	0.2	0.2	0.2	*	*	*	0.1	*	*	0.4	0.5	0.3	*	*	*	0.5	0.5	0.5
Heart failure (I50)	25.6	24.5	26.7	7.4	7.1	7.8	14.8	17.4	12.2	8.1	7.8	8.5	24.2	24.4	24.0	13.1	14.8	11.3	34.5	32.7	36.2
All other forms of heart disease (I26–I28, I34–I38,I42–I49,I51)	44.1	45.0	43.2	13.4	14.3	12.4	28.5	31.3	25.7	14.6	14.9	14.4	48.1	52.2	44.2	38.3	42.6	33.9	57.5	57.8	57.2
Essential hypertension and hypertensive renal disease (I10,I12,I15)	12.9	12.2	13.6	5.9	5.8	6.1	10.4	10.7	10.2	9.3	9.1	9.5	18.8	19.3	18.4	10.4	9.2	11.6	14.6	13.5	15.7
Cerebrovascular diseases (I60–I69)	49.1	43.1	54.9	22.0	20.3	23.7	32.6	29.0	36.1	32.7	30.1	35.0	54.6	51.8	57.1	39.4	39.2	39.7	59.9	51.4	68.2
Atherosclerosis (I70)	1.3	1.2	1.3	0.4	0.4	0.4	*	*	*	0.7	0.7	0.7	1.1	1.1	1.1	*	*	*	1.7	1.6	1.8
Other diseases of circulatory system (I71–I78)	6.2	6.7	5.6	2.1	2.5	1.7	3.9	3.1	4.7	2.6	2.8	2.4	6.4	7.2	5.7	4.3	*	*	7.9	8.6	7.3
Aortic aneurysm and dissection (I71)	3.0	3.7	2.4	0.9	1.3	0.5	1.1	*	*	1.6	2.0	1.2	2.7	3.5	2.1	*	*	*	4.0	4.8	3.2
Other diseases of arteries, arterioles and capillaries (I72–I78)	3.1	3.1	3.2	1.2	1.2	1.2	2.8	2.3	3.3	1.0	0.8	1.2	3.7	3.7	3.6	*	*	*	4.0	3.9	4.0

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other disorders of circulatory system (I80–I99)	1.7	1.8	1.6	0.8	0.8	0.7	1.6	1.7	*	0.4	0.5	0.4	2.8	2.9	2.6	*	*	*	2.0	2.0	1.9
Influenza and pneumonia (J09–J11)	12.6	13.6	11.7	5.8	6.4	5.3	13.6	15.6	11.6	8.1	9.3	6.9	12.7	13.8	11.6	9.1	12.3	*	15.6	16.7	14.5
Influenza (J09–J11)	0.2	0.2	0.2	0.1	0.1	0.1	*	*	*	*	*	*	0.2	0.2	0.2	*	*	*	0.2	0.2	0.2
Pneumonia (J12–J18)	12.4	13.4	11.5	5.8	6.3	5.2	13.2	15.2	11.2	8.0	9.2	6.8	12.5	13.7	11.5	8.9	12.3	*	15.4	16.4	14.3
Other acute lower respiratory infections (J20–J22,U04)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	0.1	0.1	0.1
Acute bronchitis and bronchiolitis (J20–J21)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	0.0	0.0	0.0
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	0.0
Chronic lower respiratory diseases (J40–J47)	42.9	41.1	44.7	8.7	8.7	8.8	33.6	30.7	36.5	8.9	11.1	6.9	26.8	27.8	25.9	14.4	16.1	12.6	62.0	58.6	65.4
Bronchitis, chronic and unspecified (J40–J42)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	0.1	0.1	*	*	*	0.1	0.1	0.1
Emphysema (J43)	2.2	2.4	2.0	0.4	0.5	0.3	2.0	1.9	2.2	0.4	0.7	*	1.2	1.4	1.0	*	*	*	3.3	3.5	3.1
Asthma (J45–J46)	1.1	0.9	1.2	0.6	0.5	0.7	*	*	*	0.7	0.7	0.6	2.4	2.3	2.5	*	*	*	1.0	0.7	1.2
Other chronic lower respiratory diseases (J44,J47)	39.5	37.7	41.3	7.8	7.7	7.8	30.7	28.3	32.9	7.8	9.6	6.1	23.0	23.9	22.3	12.8	14.5	11.0	57.7	54.3	61.0
Pneumoconioses and chemical effects (J60–J66,J68,U07.0)	0.2	0.3	0.0	0.0	0.1	*	*	*	*	*	*	*	0.1	0.1	*	*	*	*	0.3	0.5	0.0
Pneumonitis due to solids and liquids (J69)	6.0	7.1	5.0	2.0	2.3	1.8	4.3	5.2	3.4	3.1	3.8	2.5	5.7	6.5	5.1	*	*	*	7.8	9.3	6.4
Other diseases of respiratory system (J00–J06, J30–J39,J67,J70–J98)	13.8	14.7	12.9	6.0	6.3	5.6	13.5	14.3	12.8	5.9	6.6	5.2	11.4	11.2	11.6	5.6	*	*	18.1	19.4	16.8
Peptic ulcer (K25–K28)	1.2	1.3	1.1	0.6	0.6	0.5	1.5	*	*	0.8	0.8	0.7	1.0	1.2	0.9	*	*	*	1.5	1.6	1.5
Diseases of appendix (K35–K38)	0.1	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*	*	0.1	0.2	*	*	*	*	0.2	0.2	0.1
Hernia (K40–K46)	0.7	0.7	0.8	0.3	0.3	0.3	*	*	*	0.2	0.2	*	0.5	0.5	0.5	*	*	*	1.0	0.9	1.1
Chronic liver disease and cirrhosis (K70,K73–K74)	17.0	21.7	12.5	14.1	19.5	8.7	77.4	86.7	68.4	4.5	5.8	3.2	10.8	13.5	8.4	8.1	12.3	*	20.2	25.4	15.1
Alcoholic liver disease (K70)	10.0	13.7	6.3	8.4	13.3	3.5	61.3	70.7	52.2	2.1	3.4	0.9	6.3	8.2	4.5	4.2	6.6	*	11.6	15.7	7.6
Other chronic liver disease and cirrhosis (K73–K74)	7.1	8.0	6.2	5.7	6.2	5.2	16.2	16.0	16.3	2.4	2.4	2.3	4.5	5.3	3.9	4.0	*	*	8.6	9.8	7.5
Cholelithiasis and other disorders of gallbladder (K80–K82)	1.4	1.4	1.3	0.7	0.7	0.7	2.1	1.9	2.3	0.8	0.9	0.7	1.0	1.0	1.0	*	*	*	1.7	1.8	1.6

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Nephritis, nephrotic syndrome and nephrosis (N00–N07, N17–N19, N25–N27)	16.4	17.4	15.4	8.0	8.6	7.5	16.1	15.3	16.9	8.3	9.1	7.7	24.9	25.8	24.1	18.7	18.0	19.4	18.5	19.9	17.2
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01, N04)	0.2	0.2	0.2	0.1	0.1	0.1	*	*	*	*	*	*	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02–N03, N05–N07, N26)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	0.1	0.1	0.1
Renal failure (N17–N19)	16.0	17.1	15.1	7.9	8.4	7.3	15.7	15.0	16.4	8.2	8.9	7.4	24.5	25.4	23.7	18.5	17.7	19.4	18.1	19.4	16.8
Other disorders of kidney (N25, N27)	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10–N12, N13.6, N15.1)	0.4	0.3	0.5	0.2	0.2	0.3	*	*	*	0.2	*	0.3	0.3	0.2	0.3	*	*	*	0.5	0.3	0.6
Hyperplasia of prostate (N40)	0.2	0.4	...	0.1	0.2	...	*	*	...	0.2	0.4	...	0.2	0.4	...	*	*	...	0.3	0.5	...
Inflammatory diseases of female pelvic organs (N70–N76)	0.1	...	0.1	*	...	*	*	...	*	*	...	*	0.1	...	0.2	*	...	*	0.1	...	0.1
Pregnancy, childbirth and the puerperium (O00–O99)	0.5	...	1.0	0.6	...	1.1	1.7	...	3.4	0.3	...	0.5	1.2	...	2.4	*	...	*	0.3	...	0.7
Pregnancy with abortive outcome (O00–O07)	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	0.5	...	1.0	0.5	...	1.1	1.7	...	3.3	0.3	...	0.5	1.2	...	2.3	*	...	*	0.3	...	0.7
Certain conditions originating in the perinatal period (P00–P96)	2.9	3.2	2.6	3.6	3.9	3.2	2.3	2.1	2.6	2.0	2.3	1.6	6.7	7.7	5.7	4.3	*	*	1.8	2.0	1.6
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	2.9	3.1	2.8	2.7	2.7	2.6	3.4	3.6	3.2	1.2	1.3	1.1	3.4	3.5	3.3	*	*	*	3.0	3.2	2.9
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	10.4	10.5	10.3	4.7	5.7	3.6	12.5	14.8	10.2	3.5	3.6	3.4	12.9	14.5	11.4	8.3	7.9	8.7	12.5	12.0	13.1
All other diseases (residual)	119.8	106.6	132.7	43.9	41.5	46.3	123.2	126.4	120.0	42.8	38.9	46.5	109.4	101.6	116.5	63.4	64.1	62.7	157.4	138.0	176.5
Accidents (unintentional injuries) (V01–X59, Y85–Y86)	67.8	91.0	45.0	43.3	64.5	21.6	120.9	157.6	85.2	19.1	25.9	12.9	80.8	120.1	44.5	51.4	77.0	25.2	78.2	100.6	56.1
Transport accidents (V01–V99, Y85)	15.0	21.9	8.2	13.5	20.1	6.8	36.3	47.9	25.0	5.0	7.0	3.2	21.4	33.0	10.5	16.6	25.6	7.4	15.1	21.8	8.6

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motor vehicle accidents (V02-V04,V09.0,V09.2, V12-V14,V19.0-V19.2, V19.4-V19.6,V20-V79, V80.3-V80.5,V81.0-V81.1, V82.0-V82.1,V83-V86, V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)	14.2	20.5	7.9	13.0	19.2	6.6	34.0	43.8	24.4	4.7	6.5	3.1	20.5	31.5	10.3	14.9	22.7	6.8	14.1	20.2	8.2
Other land transport accidents. . . . (V01,V05-V06, V09.1,V09.3-V09.9,V10-V11, V15-V18,V19.3,V19.8-V19.9, V80.0-V80.2,V80.6-V80.9, V81.2-V81.9,V82.2-V82.9, V87.9,V88.9,V89.1, V89.3,V89.9)	0.4	0.6	0.1	0.3	0.5	0.1	1.3	2.2	*	0.1	0.2	*	0.4	0.8	0.1	*	*	*	0.4	0.6	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.5	0.8	0.2	0.2	0.4	0.1	1.1	1.9	*	0.2	0.3	*	0.4	0.8	0.1	*	*	*	0.6	1.1	0.2
Nontransport accidents (W00-X59,Y86)	52.8	69.1	36.8	29.8	44.4	14.7	84.6	109.7	60.1	14.1	19.0	9.6	59.5	87.0	33.9	34.8	51.5	17.8	63.0	78.8	47.6
Falls (W00-W19)	13.5	13.9	13.0	4.4	5.4	3.4	10.0	11.2	8.9	6.1	7.1	5.2	5.2	6.3	4.1	6.5	8.8	*	19.3	19.4	19.2
Accidental discharge of firearms (W32-W34)	0.2	0.3	0.0	0.1	0.2	*	*	*	*	*	*	*	0.4	0.7	*	*	*	*	0.2	0.3	0.0
Accidental drowning and submersion. . . . (W65-W74)	1.3	2.0	0.7	1.0	1.5	0.4	2.2	3.7	*	1.1	1.6	0.6	1.8	2.9	0.7	*	*	*	1.3	1.9	0.7
Accidental exposure to smoke, fire and flames. . . . (X00-X09)	1.0	1.3	0.8	0.4	0.5	0.3	1.9	2.1	1.8	0.3	0.3	0.3	1.7	2.1	1.3	*	*	*	1.2	1.5	0.9
Accidental poisoning and exposure to noxious substances (X40-X49)	30.7	44.1	17.6	20.9	32.8	8.7	59.7	77.3	42.5	4.4	7.3	1.7	43.9	66.6	22.9	20.1	31.3	8.7	33.7	46.8	20.8
Other and unspecified nontransport accidents and their sequelae (W20-W31, W35-W64,W75-W99, X10-X39,X50-X59,Y86)	6.1	7.5	4.7	2.9	4.0	1.9	10.4	15.0	6.0	2.2	2.6	1.8	6.6	8.5	4.9	4.6	*	*	7.4	8.9	5.9

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	14.5	23.3	5.9	7.8	12.5	3.0	28.2	43.4	13.4	7.0	10.2	4.1	8.8	14.8	3.3	13.1	20.5	*	18.6	30.0	7.5
Intentional self-harm (suicide) by discharge of firearms(X72–X74)	7.9	14.0	2.0	3.3	5.7	0.8	9.8	17.3	2.6	1.8	3.3	0.4	5.2	9.4	1.3	4.3	7.9	*	10.8	18.9	2.8
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71, X75–X84,Y87.0)	6.6	9.4	3.8	4.6	6.8	2.3	18.4	26.1	10.9	5.2	6.9	3.7	3.6	5.4	2.0	8.8	12.6	*	7.9	11.1	4.7
Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	7.8	12.8	3.0	7.1	11.6	2.5	12.5	19.9	5.3	1.6	2.2	1.1	34.2	61.0	9.4	7.3	12.6	*	3.2	4.4	1.9
Assault (homicide) by discharge of firearms (*U01.4, X93–X95)	6.3	10.7	2.0	5.5	9.4	1.5	7.5	12.1	3.1	1.0	1.5	0.6	30.4	55.2	7.4	5.9	10.7	*	2.1	3.0	1.1
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3, *U01.5–*U01.9,*U02,X85–X92, X96–Y09,Y87.1)	1.5	2.1	0.9	1.6	2.2	1.0	5.0	7.8	2.3	0.6	0.6	0.5	3.8	5.8	2.0	*	*	*	1.1	1.4	0.8
Legal intervention (Y35,Y89.0)	0.2	0.4	0.0	0.2	0.5	*	*	*	*	*	*	*	0.4	0.8	*	*	*	*	0.2	0.3	0.0
Events of undetermined intent (Y10–Y34,Y87.2,Y89.9)	1.9	2.5	1.3	0.8	1.2	0.4	4.9	5.5	4.2	0.4	0.5	0.2	3.6	5.3	2.0	*	*	*	2.0	2.5	1.5
Discharge of firearms, undetermined intent (Y22–Y24)	0.1	0.2	0.1	0.1	0.1	*	0.8	*	*	*	*	*	0.3	0.5	0.1	*	*	*	0.1	0.2	0.1
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21, Y25–Y34,Y87.2,Y89.9)	1.7	2.3	1.2	0.8	1.1	0.4	4.0	4.3	3.8	0.3	0.5	0.2	3.3	4.8	1.9	*	*	*	1.9	2.4	1.4
Operations of war and their sequelae (Y36,Y89.1)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40–Y84,Y88)	1.8	2.0	1.7	0.9	1.0	0.8	2.5	2.6	2.4	0.7	0.9	0.6	2.3	2.3	2.3	*	*	*	2.2	2.4	2.0

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes			Both sexes			Both sexes			Both sexes			Both sexes			Both sexes			Both sexes		
	Male	Female		Male	Female		Male	Female		Male	Female		Male	Female		Male	Female		Male	Female	
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	1.2	1.1	1.4	0.5	0.5	0.5	1.2	*	*	0.4	0.4	0.4	1.0	0.9	1.0	*	*	*	1.6	1.4	1.8
COVID-19. (U07.1) ⁴	125.6	143.9	107.6	109.6	132.6	86.0	188.6	201.0	176.5	64.8	80.3	50.5	133.8	137.2	130.6	173.6	187.9	159.0	137.8	158.4	117.5
Dementia-related causes ⁴	84.3	56.2	111.9	28.5	18.3	39.0	35.6	21.8	49.1	34.5	23.2	45.0	56.6	37.3	74.4	27.1	19.3	35.2	116.4	77.9	154.5
Drug-induced deaths ⁴	33.5	47.0	20.2	21.4	33.1	9.5	59.1	72.4	46.1	5.2	8.0	2.5	47.6	71.8	25.2	21.9	33.5	10.0	37.2	50.3	24.3
Drug overdose deaths ⁴	32.1	45.2	19.3	20.7	32.0	9.1	55.4	68.1	43.0	4.9	7.6	2.5	45.9	69.2	24.3	20.3	30.6	9.7	35.6	48.3	23.1
Alcohol-induced deaths ⁴	16.3	23.5	9.3	12.0	19.4	4.5	90.6	113.8	68.0	3.2	5.3	1.3	12.0	17.3	7.1	6.4	10.4	*	19.4	27.3	11.5
Firearm-related injuries ⁴	14.7	25.5	4.2	9.2	15.8	2.4	19.0	32.2	6.1	2.9	5.0	1.0	36.5	66.3	8.9	10.9	19.9	*	13.2	22.6	4.0

0.0 Quantity more than zero but less than 0.05.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes people of Hispanic origin of any race; see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Included in selected categories above. For the list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2021

[Age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes.	879.7	1,048.0	733.3	724.7	884.9	582.7	1,109.2	1,282.7	946.6	461.7	554.9	386.3	1,118.0	1,374.0	917.2	924.3	1,042.8	806.0	893.9	1,055.6	751.4
Salmonella infections (A01–A02)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	*
Shigellosis and amebiasis. (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections. (A04,A07–A09)	1.8	1.7	1.8	1.4	1.5	1.4	1.9	2.2	1.7	0.7	0.8	0.7	2.0	2.1	1.8	*	*	*	1.8	1.7	2.0
Tuberculosis. (A16–A19)	0.1	0.2	0.1	0.2	0.3	0.2	*	*	*	0.7	1.1	0.4	0.2	0.3	0.1	*	*	*	0.1	0.1	0.0
Respiratory tuberculosis. (A16)	0.1	0.2	0.1	0.2	0.2	0.1	*	*	*	0.6	0.9	0.3	0.2	0.3	0.1	*	*	*	0.0	0.1	0.0
Other tuberculosis. (A17–A19)	0.0	0.0	0.0	0.1	*	*	*	*	*	*	*	*	0.0	*	*	*	*	*	0.0	0.0	0.0
Whooping cough. (A37)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas. (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection. (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Septicemia. (A40–A41)	10.2	11.4	9.3	7.4	8.5	6.6	11.8	13.2	10.6	4.5	5.3	3.8	17.6	20.2	16.0	10.5	11.6	9.6	10.0	11.2	9.2
Syphilis. (A50–A53)	0.0	0.0	*	*	*	*	*	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
Acute poliomyelitis. (A80)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis. (A83–A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles. (B05)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis. (B15–B19)	0.8	1.1	0.6	1.0	1.5	0.6	2.2	2.7	1.7	0.9	1.3	0.6	1.1	1.5	0.8	3.4	*	*	0.8	1.0	0.6
Human immunodeficiency virus (HIV) disease. (B20–B24)	1.3	2.1	0.6	1.4	2.3	0.5	1.5	2.1	*	0.3	0.6	*	5.5	8.0	3.3	*	*	*	0.6	1.1	0.2
Malaria. (B50–B54)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05,A20–A36, A42–A44,A48–A49,A54–A79, A81–A82,A85.0–A85.1,A85.8, A86–B04,B06–B09,B25–B49, B55–B99,U07.1)	106.9	134.6	84.0	154.0	202.0	113.2	189.4	213.5	167.8	63.5	86.2	45.8	140.1	163.8	123.6	189.9	210.9	168.8	96.2	121.5	75.2
Malignant neoplasms (C00–C97)	146.6	172.0	127.7	105.1	122.1	93.3	124.9	138.5	114.3	92.9	105.4	83.7	167.4	203.9	144.5	144.6	132.1	156.9	153.7	180.1	133.6
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	2.7	4.2	1.4	1.5	2.4	0.8	1.7	1.9	*	2.1	3.1	1.3	2.5	4.3	1.2	4.3	*	*	3.0	4.5	1.6
Malignant neoplasm of esophagus. (C15)	3.7	6.4	1.4	1.7	3.1	0.6	3.3	6.3	*	1.6	2.6	0.7	2.6	4.3	1.5	*	*	*	4.3	7.6	1.5

See footnotes at end of table.