

Deaths: Final Data for 2022

by Jiaquan Xu, M.D., Sherry L. Murphy, B.S., Kenneth D. Kochanek, M.A., and Elizabeth Arias, Ph.D.

Abstract

Objectives—This report presents final 2022 data on U.S. deaths, death rates, life expectancy, infant and maternal mortality, and trends by selected characteristics such as age, sex, Hispanic origin and race, state of residence, and cause of death.

Methods—Information reported on death certificates is presented in descriptive tabulations. The original records are filed in state registration offices. Statistical information is compiled in a national database through the Vital Statistics Cooperative Program of the National Center for Health Statistics. Causes of death are processed according to the *International Classification of Diseases, 10th Revision*. Beginning in 2018, all states and the District of Columbia were using the 2003 revised certificate of death for the entire year, which includes the 1997 Office of Management and Budget revised standards for race. Data based on these revised standards are not completely comparable to previous years.

Results—In 2022, a total of 3,279,857 deaths were reported in the United States. The age-adjusted death rate was 798.8 deaths per 100,000 U.S. standard population, a decrease of 9.2% from the 2021 rate. Life expectancy at birth was 77.5 years, an increase of 1.1 years from 2021. Age-specific death rates decreased from 2021 to 2022 for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older and increased for age groups 1–4 and 5–14. In 2022, the 10 leading causes of death remained the same as in 2021, although four causes changed rank. Heart disease remained the top leading cause, followed by cancer. The infant mortality rate, 5.60 deaths per 1,000 live births in 2022, increased 2.9% from the rate in 2021 (5.44).

Conclusions—In 2022, the age-adjusted death rate decreased and life expectancy at birth increased for the total, male, and female populations, primarily due to the decrease in deaths from COVID-19.

Keywords: mortality • cause of death • life expectancy • National Vital Statistics System

Highlights

Mortality experience in 2022

- In 2022, a total of 3,279,857 resident deaths were registered in the United States, a decrease of 184,374 deaths compared with 2021 (3,464,231). The 1-year decrease in the number of deaths was primarily driven by the decrease in deaths from COVID-19.
- The crude death rate was 984.1 deaths per 100,000 population. The age-adjusted death rate, which accounts for the aging of the population, was 798.8 deaths per 100,000 U.S. standard population.
- The age-adjusted death rate for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population (947.9) was 1.2 times greater than for the White non-Hispanic (subsequently, White) population (822.2).
- The age-adjusted death rate for the Black non-Hispanic (subsequently, Black) population (1,002.8) was 1.2 times greater than for the White population (822.2).
- The age-adjusted death rate for the White population (822.2) was 2.0 times greater than for the Asian non-Hispanic (subsequently, Asian) population (417.5) and 1.3 times greater than for the Hispanic population (614.7).
- The ratio of the age-adjusted rate for the White population (822.2) to that of the Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population (782.0) was 1.1.
- Life expectancy at birth was 77.5 years for the total population, 74.8 for males, and 80.2 for females.
- The 15 leading causes of death in 2022 were:
 1. Diseases of heart (heart disease)
 2. Malignant neoplasms (cancer)
 3. Accidents (unintentional injuries)
 4. COVID-19
 5. Cerebrovascular diseases (stroke)
 6. Chronic lower respiratory diseases

7. Alzheimer disease
 8. Diabetes mellitus (diabetes)
 9. Nephritis, nephrotic syndrome and nephrosis (kidney disease)
 10. Chronic liver disease and cirrhosis
 11. Intentional self-harm (suicide)
 12. Influenza and pneumonia
 13. Essential hypertension and hypertensive renal disease (hypertension)
 14. Septicemia
 15. Parkinson disease
- In 2022, the infant mortality rate (IMR) was 5.60 infant deaths per 1,000 live births.
 - The 10 leading causes of infant death were:
 1. Congenital malformations, deformations and chromosomal abnormalities (congenital malformations)
 2. Disorders related to short gestation and low birth weight, not elsewhere classified (low birth weight)
 3. Sudden infant death syndrome (SIDS)
 4. Accidents (unintentional injuries)
 5. Newborn affected by maternal complications of pregnancy (maternal complications)
 6. Newborn affected by complications of placenta, cord and membranes (cord and placental complications)
 7. Bacterial sepsis of newborn
 8. Respiratory distress of newborn
 9. Intrauterine hypoxia and birth asphyxia
 10. Diseases of the circulatory system

Comparison with previous year

- The age-adjusted death rate decreased 9.2% from 879.7 per 100,000 standard population in 2021 to 798.8 in 2022.
- Life expectancy for the total population increased 1.1 years from 76.4 in 2021 to 77.5 in 2022.
- From 2021 to 2022, life expectancy at birth increased by 1.3 years for males (from 73.5 to 74.8) and by 0.9 year for females (79.3 to 80.2).
- Life expectancy for females (80.2) was 5.4 years higher than for males (74.8), a decrease of 0.4 year from 2021.
- The difference in life expectancy between the Black and White populations decreased 0.8 year from 5.5 years in 2021 to 4.7 years in 2022.
- From 2021 to 2022, life expectancy increased for Hispanic males (2.4 years), American Indian and Alaska Native males (2.3 years), American Indian and Alaska Native females (2.1 years), Hispanic females (1.7 years), Black males and females (1.5 years each), Asian males and White males (1.1 years each), Asian females (0.7 year), and White females (0.6 year).
- The 15 leading causes of death remained the same in 2022 as in 2021, although some causes changed ranks.
- Age-adjusted death rates decreased significantly in 2022 from 2021 for 11 of the 15 leading causes of death: heart

disease, cancer, unintentional injuries, COVID-19, stroke, Chronic lower respiratory diseases, Alzheimer disease, diabetes, Chronic liver disease and cirrhosis, hypertension, and Parkinson disease. Significant increases occurred in 2022 from 2021 for 2 of the 15 leading causes of death: kidney disease and Influenza and pneumonia.

- The age-adjusted death rate decreased in 2022 from 2021 for alcohol-induced causes (6.3%).
- The age-adjusted rate for drug-induced causes did not change significantly in 2022 from 2021.
- The increase in life expectancy at birth for the total population in 2022 was mainly due to decreases in mortality from COVID-19, heart disease, cancer, unintentional injuries, and homicide.
- Among external causes of injury death, unintentional poisoning has been the leading mechanism of injury mortality since 2011.
- IMR increased 2.9% from 5.44 infant deaths per 1,000 live births in 2021 to 5.60 in 2022.
- The 10 leading causes of infant death remained the same in 2022 as in 2021, although two causes, Intrauterine hypoxia and birth asphyxia and Diseases of the circulatory system, changed ranks.

Introduction

This report presents detailed 2022 data on deaths and death rates according to demographic and medical characteristics. These data provide information on mortality patterns among residents of the United States by such variables as age, sex, Hispanic origin and race, state of residence, and cause of death. Information on these mortality patterns is key to understanding changes in the health and well-being of the U.S. population (1). Companion reports present additional details on leading causes of death and life expectancy in the United States (2,3).

Cause-of-death statistics presented in this report are classified according to the *International Classification of Diseases, 10th Revision* (ICD-10) (4–6). Detail on cause-of-death classification is provided in Technical Notes of this report. The COVID-19 pandemic that began in 2020 continued to have a substantial impact on the mortality profile of the U.S. population for 2022.

Mortality data can be used to monitor and evaluate the health status of the United States in terms of current mortality levels and long-term mortality trends, and to identify segments of the U.S. population at greater risk of death from specific diseases and injuries. Differences in death rates among various demographic subpopulations, including racial and ethnic groups, may reflect subpopulation differences in factors such as socioeconomic status, access to medical care, and the prevalence of specific risk factors in a particular subpopulation.

The 2003 revision of the U.S. Standard Certificate of Death uses the revised 1997 Office of Management and Budget (OMB) standards for the collection of race and Hispanic ethnicity (7,8). The 1997 standards allow individuals to report more than one race and increase the race categories from four to five by separating the Asian and Pacific Islander groups. Beginning with the 2018

data year, all 50 states and the District of Columbia reported deaths based on the 2003 revision for the entire year, so the revised standards became the official standards for presenting mortality data by race and ethnicity (9). The Hispanic category did not change, remaining consistent with reports before 2018.

The race and ethnicity categories in this report follow the 1997 OMB revised race and Hispanic-origin standards. These categories differ from the bridged-race categories used before 2018 (10). The new categories include Hispanic, single-race American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native), single-race Asian non-Hispanic (subsequently, Asian), single-race Black or African American non-Hispanic (subsequently, Black), single-race Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander), and single-race White non-Hispanic (subsequently, White). Because single-race data are not available for the entire United States before 2018, data by race for 2018 through 2022 are not completely comparable with bridged-race data used in earlier years, so comparisons should be made with this consideration (11).

In addition to the tabulations included in this report, more detailed analysis is possible by using the annual mortality public-use file. The data file may be downloaded from: https://www.cdc.gov/nchs/data_access/Vitalstatsonline.htm (12). Data file documentation is available from: <https://www.cdc.gov/nchs/data/dvs/2022-Mortality-Public-Use-File-Documentation.pdf>. The public-use file does not include geographic detail, but a file with this information may be available upon request (13). Death data also may be accessed from the Centers for Disease Control and Prevention's (CDC) WONDER, a web-based system that makes the agency's information resources available to public health professionals and the general public (14).

Methods

Data in this report are based on information from all resident death certificates filed in the 50 states and the District of Columbia.

This report provides detailed death data in [Tables 1–25](#). Tables showing data by state also provide information for the Commonwealth of the Northern Mariana Islands (Northern Marianas), Guam, Puerto Rico, and U.S. Virgin Islands.

Mortality data on specific demographic and medical characteristics cover all 50 states and the District of Columbia. Measures of mortality in this report include the number of deaths; crude, age-specific, and age-adjusted death rates; infant, neonatal, postneonatal, and maternal mortality rates; life expectancy; and rate ratios. Changes in death rates in 2022 compared with 2021 and differences in death rates across demographic groups in 2022 were tested for statistical significance. Unless otherwise specified, reported differences are statistically significant. Additional information on these statistical methods, random variation and relative standard error, the computation of derived statistics and rates, population denominators, and the definition of terms are presented in Technical Notes.

In accordance with the revised standards issued by OMB in 1997, the 2003 revision of the U.S. Standard Certificate of Death

provided for the reporting of more than one race (multiple races) and increased the race categories from four to five by separating the Asian and Pacific Islander groups (7,8). Starting in 2018, all 50 states and the District of Columbia reported deaths using the 2003 revision for the entire year.

The race and Hispanic-origin groups in this report follow the 1997 standards and differ from the race categories used in reports for data years before 2018 (8,10,15). The categories include Hispanic, single-race American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native), single-race Asian non-Hispanic (subsequently, Asian), single-race Black or African American non-Hispanic (subsequently, Black), single-race Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander), and single-race White non-Hispanic (subsequently, White). For brevity, text references to race refer to single race in this report. Because the number of deaths reported with more than one race in 2022 is relatively small (0.6%), these deaths are included in totals but are shown separately in only one report table ([Table 2](#)). Some comparisons between race and ethnicity groups in this report are limited to the following groups based on population size: American Indian and Alaska Native, Asian, Black, Hispanic, and White.

Data presented in this report by the revised Hispanic-origin and race categories for 2018–2022 are not completely comparable with data by bridged-race shown in earlier reports, and comparisons should be made with this consideration. The Hispanic-origin category is a separate item on the death certificate and was not affected by the revised standards; as a result, data by Hispanic origin for 2022 and earlier years are comparable.

Death rates by race and ethnicity for the American Indian and Alaska Native, Asian, Hispanic, and Native Hawaiian or Other Pacific Islander populations are affected by inconsistencies in reporting Hispanic origin and race on death certificates as compared with censuses and surveys (16). Death rates for the American Indian and Alaska Native population are underestimated by about 33% due to misclassification (16). Death rates for the Asian and Hispanic populations are underestimated by about 3.0% (16). This should be considered when making rate comparisons across racial and ethnic groups. At this time, information about the prevalence of misclassification for the Native Hawaiian or Other Pacific Islander population is not available.

To maintain consistency with data reported by the jurisdictions and data in the mortality data file, numbers of deaths and death rates in this report are not adjusted for misclassification of race and ethnicity unless otherwise indicated. Specifically, [Tables 4 and 5](#) present life expectancies by Hispanic origin and race that are produced using methods based on death rates adjusted for Hispanic origin and race misclassification on death certificates. For additional detail, see *Quality of race and Hispanic-origin data* in Technical Notes.

The population data used to calculate death rates for 2022 shown in this report are estimated as of July 1, 2022, based on the Blended Base produced by the U.S. Census Bureau (Technical Notes), and are available from the CDC WONDER website: <https://wonder.cdc.gov/single-race-population.html> (17).

Data presented in this report and other mortality tabulations are available from the National Vital Statistics System website: <https://www.cdc.gov/nchs/deaths.htm>. The availability of mortality microdata is described in Technical Notes.

Results and Discussion

Deaths and death rates

In 2022, a total of 3,279,857 resident deaths were registered in the United States—184,374 fewer deaths than in 2021. The crude death rate for 2022 (984.1 deaths per 100,000 population) was 5.7% lower than the 2021 rate (1,043.8) (Tables A, 1, 2, 6, 8, and 10).

The age-adjusted death rate in 2022 was 798.8 deaths per 100,000 U.S. standard population—9.2% lower than the rate of 879.7 in 2021 (Tables A and 1). The age-adjusted death rates decreased for males (8.9%) and females (9.2%). Age-adjusted death rates should be viewed as relative indexes rather than as actual measures of mortality risk. They are constructs that show what the level of mortality would be if no changes occurred in the age composition of the population from year to year. (For a discussion of age-adjusted death rates, see Technical Notes.) Thus, age-adjusted death rates are better indicators than unadjusted (crude) death rates for examining changes in the risk of death over a period of time when the age distribution of the population is changing. Age-adjusted death rates are also better indicators of relative risk when comparing mortality across geographic areas or between sex or race and ethnicity subgroups of the population that have different age distributions; see Technical Notes. Since 1980, the age-adjusted death rate decreased significantly every year except for 1983, 1985, 1988, 1993, 1999, 2005, 2010, 2013, 2015, 2017, 2020, and 2021 (Figure 1) (14).

Death rates by Hispanic origin and race

In 2022, age-adjusted death rates by ethnicity and race groups (Table 1) were:

- Hispanic population: 614.7 deaths per 100,000 U.S. standard population
- American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population: 947.9
- Asian non-Hispanic (subsequently, Asian) population: 417.5
- Black non-Hispanic (subsequently, Black) population: 1,002.8
- Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population: 782.0
- White non-Hispanic (subsequently, White) population: 822.2

In 2022, the age-adjusted death rate for the American Indian and Alaska Native population was 1.2 times that for the White population. The rate for the Black population was 1.2 times that

for the White population. The rate for the White population was 2.0 times the rate for the Asian population, 1.1 times the rate for Native Hawaiian or Other Pacific Islander population, and 1.3 times the rate for the Hispanic population (Table B).

From 2021 to 2022, the age-adjusted rate decreased 15.4% (from 924.3 to 782.0) for the Native Hawaiian or Other Pacific Islander population, 15.2% (from 724.7 to 614.7) for the Hispanic population, 14.5% (from 1,109.2 to 947.9) for the American Indian and Alaska Native population, 10.3% (from 1,118.0 to 1,002.8) for the Black population, 9.6% (461.7 to 417.5) for the Asian population, and 8.0% (893.9 to 822.2) for the White population (Tables A and 1).

From 2021 to 2022, the age-adjusted death rate decreased 17.6% for Native Hawaiian or Other Pacific Islander males, 15.5% for American Indian and Alaska Native males, 15.4% for Hispanic males, 14.5% for Hispanic females, 13.8% for American Indian and Alaska Native females, 12.5% for Native Hawaiian or Other Pacific Islander females, 11.8% for Black females, 9.6% for Asian males, 9.2% for Asian females, 8.5% for Black males, 7.9% for White males, and 7.8% for White females (Tables A and 1).

Hispanic subgroups—Mortality data for 2022 by specified Hispanic subgroup for the United States are presented in Table 3. Hispanic subgroups shown in the table include Central American, Cuban, Dominican, Mexican, Puerto Rican, South American, and Other Hispanic populations.

In 2022, age-adjusted rates among the Hispanic subgroups ranged from a low of 344.9 for the South American population to a high of 688.1 for the Puerto Rican population. Differences between subgroups are likely a function of large variation in age-specific death rates for some of the Hispanic subgroups, reflecting their relatively small population sizes. Data aggregated over several years confirm differences among the Hispanic subgroups (16).

Death rates by age and sex

For the total population, age-specific death rates increased significantly from 2021 to 2022 for age groups 1–4 and 5–14 and decreased for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older (Tables 6 and 8; Figure 2). The rate for younger than 1 year did not change significantly.

The age-adjusted death rate for males was 1.4 times the rate for females in 2022 (Table B). The male-to-female death rate ratio was unchanged from the ratio in 2021.

Death rates for males increased significantly for age group 1–4 and decreased for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. Changes in rates for males younger than 1 and ages 5–14 were not statistically significant. Death rates for females increased significantly for age groups 1–4 and 5–14 and decreased for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. The rate for females younger than 1 did not change significantly.

Race and ethnicity by sex—For the total Hispanic population, Hispanic males, and Hispanic females, age-specific death rates decreased from 2021 to 2022 for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older.

For the total, male, and female American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska

Table A. Percent change in death rates and age-adjusted death rates in 2022 from 2021, by age, Hispanic origin and race, and sex: United States

[Based on death rates on an annual basis per 100,000 population and age-adjusted rates per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards; see Technical Notes. Data for some Hispanic-origin or race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes]

Age group	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All ages																					
Crude rate	-5.7	-7.0	-4.3	-14.1	-15.8	-11.7	-11.3	-12.7	-9.7	-5.9	-7.7	-3.9	-8.9	-8.5	-9.3	-13.4	-16.6	-9.4	-3.7	-5.0	-2.1
Age-adjusted rate	-9.2	-8.9	-9.2	-15.2	-15.4	-14.5	-14.5	-15.5	-13.8	-9.6	-9.6	-9.2	-10.3	-8.5	-11.8	-15.4	-17.6	-12.5	-8.0	-7.9	-7.8
Younger than 1 ⁴	-0.1	0.8	-1.4	1.1	2.2	-0.3	15.3	22.6	6.9	-6.7	-5.3	-8.6	-5.5	-5.0	-6.2	30.0	6.4	76.9	1.6	2.8	0.1
1–4	12.0	14.8	8.3	9.4	13.6	5.0	26.3	2.1	65.1	1.4	10.3	-6.9	5.6	11.7	-1.7	*	*	*	16.0	17.6	14.1
5–14	7.0	4.3	9.0	4.8	6.6	2.7	39.3	56.4	22.9	-1.2	-3.1	0.0	0.8	-3.5	8.1	*	*	*	9.4	6.8	13.0
15–24	-10.6	-11.0	-10.1	-8.9	-8.4	-11.3	-5.7	-4.6	-7.9	-13.0	-13.3	-13.3	-10.4	-9.5	-13.2	-27.9	-24.6	-36.9	-11.2	-12.8	-7.7
25–34	-9.6	-9.9	-9.8	-8.7	-7.7	-12.5	-6.3	-7.8	-3.9	-7.3	-7.6	-7.2	-10.7	-10.2	-12.6	-21.1	-24.7	-12.2	-9.7	-10.8	-8.1
35–44	-11.3	-11.0	-12.2	-16.9	-16.4	-18.8	-10.3	-7.2	-15.6	-9.7	-12.4	-5.1	-11.3	-10.5	-13.2	-28.4	-27.0	-30.9	-9.5	-9.3	-10.3
45–54	-14.6	-15.3	-13.8	-24.3	-25.4	-22.4	-15.8	-17.0	-14.4	-12.5	-14.1	-10.1	-15.2	-14.4	-16.6	-27.8	-32.8	-20.2	-11.7	-12.5	-10.6
55–64	-11.2	-12.0	-10.1	-24.7	-25.3	-23.7	-17.2	-20.2	-12.9	-14.9	-16.0	-13.5	-12.7	-11.9	-13.8	-15.3	-15.0	-15.8	-8.2	-9.4	-6.6
65–74	-8.0	-8.3	-7.5	-19.6	-20.0	-19.1	-15.9	-15.1	-16.7	-11.0	-10.8	-11.4	-10.0	-8.5	-11.6	-12.9	-18.9	-5.4	-6.1	-6.9	-5.1
75–84	-8.0	-8.4	-7.6	-14.9	-16.1	-13.6	-16.3	-19.9	-12.8	-9.3	-10.5	-7.7	-9.6	-7.3	-11.4	-5.7	-5.7	-5.8	-7.1	-7.7	-6.5
85 and older	-8.6	-6.1	-9.9	-7.1	-4.2	-8.5	-19.2	-21.1	-18.0	-7.4	-4.8	-9.0	-7.8	-3.3	-9.7	-16.9	-14.5	-18.5	-8.7	-6.3	-10.0

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.


¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes persons of Hispanic origin of any race; see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Death rates for "Younger than 1" (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Figure 1. Crude and age-adjusted death rates: United States, 1960–2022

Native) populations, age-specific death rates decreased from 2021 to 2022 for age groups 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. For the total and male American Indian and Alaska Native populations, the rates increased for age group 5–14.

For the total Asian non-Hispanic (subsequently, Asian) population, age-specific rates decreased from 2021 to 2022 for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. For Asian males, rates decreased for age groups 15–24, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. For Asian females, rates decreased for age groups 45–54, 55–64, 65–74, 75–84, and 85 and older.

For the total, male, and female Black non-Hispanic (subsequently, Black) populations, age-specific death rates decreased from 2021 to 2022 for age groups younger than 1, 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older.

For the total Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) population, age-specific death rates decreased for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, and 85 and older. For Native Hawaiian or Other Pacific Islander males, rates

decreased for age groups 25–34, 35–44, 45–54, 55–65, and 65–74. For Native Hawaiian or Other Pacific Islander females, rates increased for younger than 1 year and decreased for age groups 35–44, 45–54, 55–64, and 85 and older.

For the total, male, and female White non-Hispanic (subsequently, White) populations, age-specific death rates decreased from 2021 to 2022 for age groups 15–24, 25–34, 35–44, 45–54, 55–64, 65–74, 75–84, and 85 and older. For the total and female White populations, the rates increased for age groups 1–4 and 5–14 and for White males, the rate increased for age group 1–4 ([Tables A and 2](#)).

Other observed changes from 2021 to 2022 in age-specific rates by race and ethnicity and sex were not statistically significant.

Expectation of life at birth and at specified ages

Life expectancy at birth represents the average number of years that a group of infants would live if the group was to experience throughout life the age-specific death rates present in the year of birth.

Table B. Number of deaths, percentage of total deaths, death rate, and age-adjusted death rate for 2022, percent change in age-adjusted death rates in 2022 from 2021, and ratio of age-adjusted death rates, by sex and by Hispanic origin and race for the 15 leading causes of death for the total population in 2022: United States

[Crude death rates on an annual basis per 100,000 population; age-adjusted rates per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards]

Rank ¹	Cause of death (based on ICD-10)	Number	Percent of total deaths, 2022 ²	Crude death rate, 2022	2022	Percent change from 2021 to 2022	Male to female	Age-adjusted death rate				
								Ratio				
								Non-Hispanic, single race ³				
								American Indian and Alaska Native to White	Black to White	White to Asian	White to Native Hawaiian or Other Pacific Islander	White, non-Hispanic ³ to Hispanic ⁴
...	All causes.	3,279,857	100.0	984.1	798.8	-9.2	1.4	1.2	1.2	2.0	1.1	1.3
1	Diseases of heart. (I00-I09,I11,I13,I20-I51)	702,880	21.4	210.9	167.2	-3.8	1.6	0.9	1.3	2.1	1.0	1.5
2	Malignant neoplasms. (C00-C97)	608,371	18.5	182.5	142.3	-2.9	1.3	0.8	1.1	1.7	1.1	1.4
3	Accidents (unintentional injuries). (V01-X59,Y85-Y86)	227,039	6.9	68.1	64.0	-1.1	2.3	1.9	1.2	3.6	1.4	1.4
4	COVID-19. (U07.1)	186,552	5.7	56.0	44.5	-57.3	1.6	1.5	1.2	1.7	0.9	0.9
5	Cerebrovascular diseases. (I60-I69)	165,393	5.0	49.6	39.5	-3.9	1.1	0.8	1.5	1.3	0.7	1.1
6	Chronic lower respiratory diseases. (J40-J47)	147,382	4.5	44.2	34.3	-1.2	1.1	0.7	0.7	4.7	2.0	2.8
7	Alzheimer disease. (G30)	120,122	3.7	36.0	28.9	-6.8	0.7	0.5	0.9	1.9	1.5	1.1
8	Diabetes mellitus. (E10-E14)	101,209	3.1	30.4	24.1	-5.1	1.6	2.2	2.0	1.2	0.4	0.8
9	Nephritis, nephrotic syndrome and nephrosis. (N00-N07,N17-N19,N25-N27)	57,937	1.8	17.4	13.8	1.5	1.4	1.4	2.2	1.3	0.6	1.0
10	Chronic liver disease and cirrhosis. (K70,K73-K74)	54,803	1.7	16.4	13.8	-4.8	1.8	4.4	0.6	3.6	2.0	0.9
11	Intentional self-harm (suicide). (*U03,X60-X84,Y87.0)	49,476	1.5	14.8	14.2	0.7	3.9	1.5	0.5	2.6	1.2	2.2
12	Influenza and pneumonia. (J09-J18)	47,052	1.4	14.1	11.3	7.6	1.4	1.4	1.1	1.4	1.1	1.2
13	Essential hypertension and hypertensive renal disease. (I10,I12,I15)	43,293	1.3	13.0	10.3	-3.7	1.2	1.0	2.0	1.1	0.8	1.0
14	Septicemia. (A40-A41)	42,261	1.3	12.7	10.1	-1.0	1.3	1.3	1.7	2.2	0.9	1.3
15	Parkinson disease. (G20-G21)	39,915	1.2	12.0	9.5	-3.1	2.3	0.4	0.5	1.8	*	1.6
...	All other causes. (residual)	686,172	20.9	205.9	...	...	...	...	...	...	...	...

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.


¹Based on number of deaths; see Technical Notes.

²Percentages may not add to 100 due to rounding.

³Includes only one race reported on the death certificate; see Technical Notes.

⁴Includes people of Hispanic origin of any race; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Figure 2. Death rate, by age and sex: United States, 1955–2022

Life table data shown in this report for 2010–2022 are based on a revised methodology first presented with final data reported for 2008. The life table methodology was revised by changing the smoothing technique used to estimate the life table functions at the oldest ages. This revision improves on the methodologies used previously; see Technical Notes.

The methods used to produce life expectancies by Hispanic origin and race are based on death rates adjusted for Hispanic-origin and race misclassification on death certificates (Technical Notes). As noted in the Methods section, the age-specific and age-adjusted death rates shown in this report (with the exception of Table IV) are not adjusted for misclassification of Hispanic origin and race on death certificates. For further information on the effects of Hispanic origin and race misclassification on death rates, see Technical Notes.


Life tables were generated for both sexes and by each sex for the following populations:

- Total United States
- Hispanic
- American Indian and Alaska Native non-Hispanic
- Asian non-Hispanic

- Black non-Hispanic
- White non-Hispanic

In 2022, life expectancy for the total population (77.5 years) was 1.1 years higher than in 2021. This increase followed the two largest single-year decreases in life expectancy (1.8 years in 2020 and 0.6 year in 2021) since 1943 (Tables 4 and 5). The increase in life expectancy was mostly due to the decrease in deaths from COVID-19. Before the COVID-19 pandemic, the general trend in U.S. life expectancy had been one of improvement since 1900. In 2022, life expectancy for males (74.8 years) was 1.3 years higher than in 2021. Life expectancy for females (80.2 years) was 0.9 year higher than in 2021. From 1900 through the late 1970s, the gap in life expectancy between the sexes widened from 2.0 to 7.8 years (3). The gap between sexes has narrowed since its peak in the 1970s (Figure 3 and Table 5). In 2022, the difference in life expectancy between males and females was 5.4 years, a decrease of 0.4 year from 2021.

Life expectancy figures for the Hispanic, Black non-Hispanic (subsequently, Black), and White non-Hispanic (subsequently, White) populations have been available starting with data for 2006. Before 2006, life tables were produced for the Black

Figure 3. Life expectancy at birth, by sex: United States, 1970–2022

and White populations, regardless of Hispanic origin (18). Life tables for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) and Asian non-Hispanic (subsequently, Asian) populations were added to the Life Table Program beginning with data for 2019 (19).

Life expectancy for the American Indian and Alaska Native population increased 2.2 years (from 65.6 years in 2021 to 67.8 in 2022).

Life expectancy for the Hispanic population increased by 2.2 years (from 77.8 years in 2021 to 80.0 in 2022) (Table 5). The difference in life expectancy between the Hispanic and White populations increased 1.4 years from 1.1 years in 2021 to 2.5 years in 2022.

Life expectancy increased by 1.6 years for the Black population (from 71.2 years in 2021 to 72.8 in 2022). The difference in life expectancy between the White and Black populations decreased 0.8 year from 5.5 years in 2021 to 4.7 years in 2022 (Table 5).

Life expectancy increased by 0.9 year for the Asian population (from 83.5 years in 2021 to 84.4 in 2022) (Table 5).

Life expectancy increased by 0.8 year for the White population (from 76.7 years in 2021 to 77.5 in 2022) (Table 5).

Among the 10 major race-ethnicity-sex groups in 2022, Asian females had the highest life expectancy at birth (86.3 years), followed by Hispanic females (82.8), Asian males (82.3), White females (80.1), Hispanic males (77.0), Black females (76.5), White males (75.1), American Indian and Alaska Native females (71.3), Black males (69.1), and American Indian and Alaska Native males (64.5) (Tables 4 and 5).

Life tables shown in this report may be used to compare life expectancies at selected ages from birth to 100 years. For example, based on mortality experienced in 2022 for the total population, a person age 50 could expect to live an average of 30.9 more years, for a total of 80.9 years. A person age 65 could expect to live an average of 18.9 more years, for a total of 83.9 years, and a person age 85 could expect to live an average of 6.4 more years, for a total of 91.4 years (Table 4). Life expectancy increased at ages from birth through age 85 and was either unchanged or decreased for age 90 and older from 2021 to 2022 (3).

Leading causes of death


The 15 leading causes of death in 2022 accounted for 79.1% of all deaths in the United States (Tables B and 6). In 2022, the 15 leading causes of death remained the same as in 2021, although some changed ranks. Heart disease and cancer remained the top two leading causes in 2022. Of the remaining leading causes in 2022, six causes changed rank. Unintentional injuries, the 4th leading cause in 2021, became the 3rd leading cause in 2022. COVID-19, the 3rd leading cause in 2021, became the 4th. Kidney disease, the 10th leading cause in 2021, became the 9th. Chronic liver disease and cirrhosis, the 9th leading cause in 2021, became the 10th. Influenza and pneumonia, the 13th leading cause in 2021, became the 12th. Hypertension, the 12th leading cause in 2021, became the 13th. Causes of death are ranked according to the number of deaths; for ranking procedures, see Technical Notes. By rank, the 15 leading causes of death in 2022 were:

1. Diseases of heart (heart disease)
2. Malignant neoplasms (cancer)
3. Accidents (unintentional injuries)

4. COVID-19
5. Cerebrovascular diseases (stroke)
6. Chronic lower respiratory diseases
7. Alzheimer disease
8. Diabetes mellitus (diabetes)
9. Nephritis, nephrotic syndrome and nephrosis (kidney disease)
10. Chronic liver disease and cirrhosis
11. Intentional self-harm (suicide)
12. Influenza and pneumonia
13. Essential hypertension and hypertensive renal disease (hypertension)
14. Septicemia
15. Parkinson disease

Death rates vary greatly by age. As a result, the shifting age distribution of a population can significantly influence changes in crude death rates over time. Age-adjusted death rates, in contrast, eliminate the influence of such differences in the population age structure. Consequently, while causes of death are ranked according to the number of deaths, age-adjusted death rates are used to depict trends for leading causes of death in this report

Figure 4. Age-adjusted death rate for selected leading causes of death: United States, 1958–2022



because they are better than crude rates for showing changes in mortality over time and among causes of death (Figure 4; Tables B and 6).

From 2021 through 2022, age-adjusted death rates decreased significantly for 11 of the 15 leading causes of death and increased for 2 of the 15 leading causes (Tables B and 6). The rate for the top leading cause of death, heart disease, decreased 3.8% in 2022 from 2021 (Figure 4; Tables B and 6) (14). The rate for the second leading cause of death, cancer, decreased 2.9% from 2021 to 2022. Deaths from these two diseases combined accounted for 40.0% of deaths in the United States in 2022.

Unintentional injuries, the third leading cause of death in 2022, accounted for 6.9% of deaths in the United States. The age-adjusted death rate for unintentional injuries decreased 1.1% from 2021 (64.7) to 2022 (64.0).

COVID-19, the fourth leading cause of death in 2022, accounted for 5.7% of deaths in the United States. The number of deaths from COVID-19 decreased 55.3% from 416,893 in 2021 to 186,552 in 2022.

In 2022, the age-adjusted death rate for COVID-19 decreased 57.3%, from 104.1 per 100,000 standard population in 2021 to 44.5 in 2022. The rate for males (56.7) was 1.6 times higher than the rate for females (35.4) (Tables B and 11). Among race and ethnicity groups, the rate was highest for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population (65.9), followed by Black non-Hispanic (subsequently, Black) (53.9), Native Hawaiian or Other Pacific Islander non-Hispanic (subsequently, Native Hawaiian or Other Pacific Islander) (48.9), Hispanic (47.2), White non-Hispanic (subsequently, White) (43.9), and Asian non-Hispanic (subsequently, Asian) (25.3) populations (Tables 1 and 11).

Other leading causes of death that showed significant decreases in 2022 from 2021 were stroke (3.9%), Chronic lower respiratory diseases (1.2%), Alzheimer disease (6.8%), diabetes (5.1%), Chronic liver disease and cirrhosis (4.8%), hypertension (3.7%), and Parkinson disease (3.1%).

The age-adjusted rate increased significantly in 2022 from 2021 for kidney disease (1.5%) and Influenza and pneumonia (7.6%).

The age-adjusted death rates for suicide and Septicemia in 2022 were not significantly different from the rates in 2021.

The relative risk of death in one population group compared with another can be expressed as a ratio. Ratios based on age-adjusted death rates show that males have higher rates than females for 14 of the 15 leading causes of death (Table B), with rates for males being at least twice as great as those for females for 3 of these leading causes. The largest ratio was for suicide (3.9). Other high ratios were observed for unintentional injuries and Parkinson disease (2.3 each); Chronic liver disease and cirrhosis (1.8); heart disease, COVID-19, and diabetes (1.6 each); kidney disease and Influenza and pneumonia (1.4 each); cancer and Septicemia (1.3 each); hypertension (1.2); and stroke and Chronic lower respiratory diseases (1.1 each). Age-adjusted rates were lower for males than for females for one leading cause, Alzheimer disease (0.7).

Age-adjusted death rates for the American Indian and Alaska Native population were higher than for the White population for 8 of the 15 leading causes of death (Tables B and 6). The largest ratio was for Chronic liver disease and cirrhosis (4.4). Other causes for which the ratio was high include diabetes (2.2); unintentional injuries (1.9); COVID-19 and suicide (1.5 each); kidney disease and Influenza and pneumonia (1.4 each); and Septicemia (1.3). Age-adjusted rates for the American Indian and Alaska Native population were lower than for the White population for 6 of the 15 leading causes of death. The smallest American Indian and Alaska Native-to-White ratio was for Parkinson disease (0.4); that is, the risk of dying from Parkinson disease was more than two times greater for the White population than for the American Indian and Alaska Native population. Other causes with a low American Indian and Alaska Native-to-White ratio were Alzheimer disease (0.5); Chronic lower respiratory diseases (0.7); cancer and stroke (0.8 each); and heart disease (0.9).

Age-adjusted death rates for the Black population were higher than for the White population for 10 of the 15 leading causes of death (Tables B and 6). The largest ratio was for kidney disease (2.2). Other causes for which the ratio was high include diabetes and hypertension (2.0 each); Septicemia (1.7); stroke (1.5); heart disease (1.3); unintentional injuries and COVID-19 (1.2 each); and cancer and Influenza and pneumonia (1.1 each). For five of the leading causes, age-adjusted rates were lower for the Black population than for the White population. The smallest Black-to-White ratios were for suicide and Parkinson disease (0.5 each). Other causes with low Black-to-White ratios were Chronic liver disease and cirrhosis (0.6); Chronic lower respiratory diseases (0.7); and Alzheimer disease (0.9).

Age-adjusted death rates for the White population were higher than for the Asian population for each of the 15 leading causes of death (Tables B and 6). The highest ratios were for Chronic lower respiratory diseases (4.7); followed by unintentional injuries and Chronic liver disease and cirrhosis (3.6 each); suicide (2.6); Septicemia (2.2); heart disease (2.1); Alzheimer disease (1.9); Parkinson disease (1.8); cancer and COVID-19 (1.7 each); Influenza and pneumonia (1.4); stroke and kidney disease (1.3 each); diabetes (1.2); and hypertension (1.1).

Age-adjusted death rates for the White population were higher than Native Hawaiian or Other Pacific Islander population for 7 of the 15 leading causes of death (Tables B and 6). The highest ratios were for Chronic lower respiratory disease and chronic liver disease and cirrhosis (2.0 each); followed by Alzheimer disease (1.5); unintentional injuries (1.4); suicide (1.2); and cancer and Influenza and pneumonia (1.1 each). Age-adjusted death rates for the White population were lower than for the Native Hawaiian or Other Pacific Islander population for 6 of the 15 leading causes of death. The smallest ratio was for diabetes (0.4); followed by kidney disease (0.6); stroke (0.7); hypertension (0.8); and COVID-19 and Septicemia (0.9 each).

Age-adjusted death rates for the White population were higher than for the Hispanic population for 10 of the 15 leading causes of death (Tables B and 6). The largest ratio was for Chronic lower respiratory diseases (2.8). Other causes with high ratios include suicide (2.2); Parkinson disease (1.6); heart disease (1.5); cancer and unintentional injuries (1.4 each);

Septicemia (1.3); Influenza and pneumonia (1.2); and stroke and Alzheimer disease (1.1 each). Age-adjusted rates were lower for the White population than for the Hispanic population for 3 of the 15 leading causes. The smallest White-to-Hispanic ratio was for diabetes (0.8), followed by COVID-19 and Chronic liver disease and cirrhosis (0.9 each).

Assault (homicide), the 16th leading cause of death in 2022, dropped from among the 15 leading causes of death in 2010. In 2022, the age-adjusted rate for homicide decreased 6.1%, from 8.2 in 2021 to 7.7 per 100,000 standard population in 2022 (Table 11). Homicide remains a major issue for some age groups. Homicide was among the 15 leading causes of death in 2022 for age groups 1–4 (3rd), 5–14 (3rd), 15–24 (2nd), 25–34 (3rd), 35–44 (6th), 45–54 (10th), and 55–64 (14th) (14) (For leading causes of infant death, see “Infant mortality”).

Although Human immunodeficiency virus (HIV) disease has not been among the 15 leading causes of death since 1997 (20), it is still considered a major public health problem for some age groups. The age-adjusted death rate (1.3 deaths per 100,000 U.S. standard population) for HIV disease remained unchanged statistically in 2022 from 2021 (Table 11). Historically, for all ages combined, HIV disease mortality reached its highest level in 1995 after a period of increase from 1987 through 1994. Subsequently, the rate for this disease decreased an average of 33.0% per year from 1995 through 1998, and 6.1% per year from 1999 through 2021 (9,13). In 2022, HIV disease was among the 15 leading causes of death for age groups 25–34 (13th), 35–44 (13th), and 45–54 (15th). The influence of HIV disease for younger age groups and by race and ethnicity shows among the 25–34 age group, where its rank is 9th for males, 7th for Black males, 11th for Black females, and 10th for Hispanic males.

Enterocolitis due to *Clostridium difficile* (*C. difficile*)—A predominantly antibiotic-associated inflammation of the intestines caused by *C. difficile*, a gram-positive, anaerobic, spore-forming bacillus—is often acquired in hospitals or other health care facilities with long-term patients or residents (21,22). The number of deaths from *C. difficile* climbed from 793 deaths in 1999 to a high of 8,085 deaths in 2011 (14). From 2011 to 2021, the number of deaths from this cause trended downward. In 2022, the number of deaths from *C. difficile* was 4,231 (126 more deaths than in 2021). The age-adjusted death rate in 2022 was unchanged from 2021, 1.0 deaths per 100,000 standard population. About 85.4% of deaths from *C. difficile* occurred among people age 65 and older (Table 7).

Leading causes of death in 2022 for the total population and for specific subpopulations are detailed further in a companion *National Vital Statistics Report* on leading causes by age, Hispanic origin and race, and sex (2).

Other selected causes

Dementia-related mortality

In 2022, 292,881 people died of dementia-related causes in the United States (Tables 7, 9, and 22). Deaths from dementia-related causes were presented for the first time in this report series in 2018 to provide a more comprehensive estimate of the

burden of mortality from Alzheimer disease and other dementias in the United States.

Dementia-related causes include conditions with similar physical signs and symptoms that, collectively, are considered to be a good indicator of dementia mortality (23). Dementia is characterized by memory impairment and cognitive decline (24–26). Causes of death attributable to dementia-related mortality include ICD–10 codes F01, Vascular dementia; F03, Unspecified dementia; G30, Alzheimer disease; and G31, Other degenerative diseases of nervous system, not elsewhere classified. Alzheimer disease, the sixth leading cause of death, is the most common cause of dementia, but other dementias, including Lewy body dementia, frontotemporal degeneration, vascular dementia, and mixed dementias, are often indistinguishable from Alzheimer disease in their symptoms and outcomes and may coexist with Alzheimer disease (22–24).

Certification and coding rule changes can impact data analysis of component causes of dementia. In 2022, Alzheimer disease accounted for 41.0% of all dementia deaths; Unspecified dementia for 31.9%; Other degenerative diseases of nervous system, not elsewhere classified for 19.9%; and Vascular dementia for 7.1%. For detailed information, see CDC WONDER (14). Changes in the percentage of deaths assigned to individual causes comprising dementia may be the result of many factors (25). Combining the types of dementia provides a more comprehensive and stable measure of dementia mortality.

The age-adjusted death rate for dementia-related causes decreased 2.8% in 2022 from 2021 for the total population (from 72.4 to 70.4). The rate for females decreased 4.2% (80.2 to 76.8). The rate for males was unchanged (59.7) (Tables 11 and 22).

Among race-ethnicity groups—Age-adjusted rates for American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) females decreased 11.8% (51.6 to 45.5). Rates for the total and female Black non-Hispanic (subsequently, Black) populations decreased 5.6% (71.8 to 67.8) and 7.9% (75.9 to 69.9), respectively. Rates for the total and female White non-Hispanic (subsequently, White) population decreased 2.8% (77.4 to 75.2) and 4.3% (86.3 to 82.6), respectively. Changes in age-adjusted death rates for dementia-related causes were not statistically significant for the other race-ethnicity-sex categories.

Drug-induced mortality

In 2022, a total of 112,109 people died of drug-induced causes in the United States (Tables 7, 9, and 23). The category of drug-induced causes includes deaths from drug overdose as well as from other medical conditions caused by use of legal or illegal drugs. In 2022, drug-overdose deaths accounted for 96.3% of all drug-induced deaths (Tables 7 and 9). The drug-induced category excludes deaths indirectly related to drug use, as well as newborn deaths due to the mother's drug use. (For a list of all drug-induced causes including those specifically classified as drug-overdose causes, see Technical Notes.)

The age-adjusted death rate for drug-induced causes did not change significantly for the total, male, and female populations from 2021 to 2022 (Tables 6, 11, and 23). For males in 2022, the

age-adjusted death rate for drug-induced causes was 2.3 times the rate for females.

Among race-ethnicity groups—Age-adjusted rates increased from 2021 and 2022 by 15.1% for the American Indian and Alaska Native population (20.3% for American Indian and Alaska Native males), 12.2% for the Asian population (14.5% for Asian males), 7.4% for the Black population (8.5% for Black males), and 6.8% for the Hispanic population (8.4% for Hispanic males). Rates for the White population decreased 3.4% (3.9% for White males and 2.8% for White females) (Tables 11 and 23). Changes in age-adjusted rates from 2021 to 2022 were not significant for other race-ethnicity-sex categories.

Alcohol-induced mortality

In 2022, a total of 51,191 people died of alcohol-induced causes in the United States (Tables 7, 9, and 24). This category includes deaths from dependent and nondependent use of alcohol, and deaths from accidental poisoning by alcohol. It excludes unintentional injuries, homicides, and other causes indirectly related to alcohol use, and deaths due to fetal alcohol syndrome. For a list of alcohol-induced causes, see Technical Notes.

The age-adjusted death rate for alcohol-induced causes decreased 6.3% for the total population, from 14.4 in 2021 to 13.5 in 2022 (Tables 6, 11, and 24). For males in 2022, the age-adjusted death rate for alcohol-induced causes was 2.5 times the rate for females. The rate decreased 6.7% for males and 6.0% for females from 2021 to 2022 (Tables 11 and 24).

Among race-ethnicity groups—Age-adjusted rates decreased in 2022 from 2021 by 14.5% for the American Indian and Alaska Native population (15.2% for American Indian and Alaska Native males and 13.5% for American Indian and Alaska Native females), 13.4% for the Black population (12.7% for Black males and 14.9% for Black females), 4.0% for Hispanic males, and 5.1% for the White population (6.0% for White males and 4.2% for White females) (Tables 11 and 24). The age-adjusted rate for alcohol-induced death did not change significantly in 2022 from 2021 for other race-ethnicity-sex categories.

Firearm-related mortality

In 2022, 48,204 people died from firearm-related injuries in the United States (Tables 7, 9, and 25). The age-adjusted death rate for firearm-related injuries for the total and male populations decreased significantly from 2021 to 2022 by 2.7% and 2.8%, respectively (Tables 6, 11, and 25). For males in 2022, the age-adjusted death rate for firearm-related injuries was 6.0 times the rate for females.

Among race-ethnicity groups—The age-adjusted death rate increased in 2022 from 2021 by 16.2% for the American Indian and Alaska Native population (17.4% for American Indian and Alaska Native males). The rates decreased 7.8% for the total, male, and female Black populations (Tables 11 and 25). The age-adjusted death rates for firearm-related injuries did not change significantly in 2022 from 2021 for other race-ethnicity-sex categories.

Effect on life expectancy of changes in mortality by age and cause of death

Changes in mortality by age and cause of death can have a major effect on life expectancy. In other words, year-to-year changes in life expectancy may be influenced by changes in age-specific rates for certain causes, particularly for younger age groups. Life expectancy at birth for the total population increased by 1.1 years (from 76.4 years in 2021 to 77.5 years in 2022) primarily because of decreases in mortality from COVID-19, heart disease, cancer, unintentional injuries, and homicide (Table 5). The increase in life expectancy for the total population was slightly offset by increases in mortality from Influenza and pneumonia, perinatal conditions, kidney disease, nutritional deficiencies, and congenital malformations. Life expectancy at birth for males increased by 1.3 years (from 73.5 years in 2021 to 74.8 years in 2022) due to decreases in mortality from COVID-19, heart disease, cancer, homicide, and unintentional injuries. The increase in life expectancy was offset somewhat by increases in mortality from perinatal conditions, Influenza and pneumonia, kidney disease, nutritional deficiencies, and congenital malformations. For the female population, life expectancy increased by 0.9 year (from 79.3 years in 2021 to 80.2 years in 2022) due to decreases in mortality from COVID-19, unintentional injuries, heart disease, diabetes, and cancer. This increase was offset by increases in mortality from Influenza and pneumonia, Chronic lower respiratory diseases, nutritional deficiencies, kidney disease, and perinatal conditions. (For a discussion of the major causes contributing to the change in life expectancy, see Technical Notes.)

Life expectancy for the American Indian and Alaska Native non-Hispanic (subsequently, American Indian and Alaska Native) population increased by 2.2 years to 67.8 years. This increase was primarily due to decreases in mortality from COVID-19, Chronic liver disease and cirrhosis, cancer, suicide, and diabetes. The increase in life expectancy was offset by increases in mortality due to unintentional injuries, homicide, perinatal conditions, Influenza and pneumonia, and congenital malformations. Life expectancy for the American Indian and Alaska Native male population increased 2.3 years to 64.5 years. This increase, primarily due to decreases in mortality from COVID-19, Chronic liver disease and cirrhosis, heart disease, cancer, and suicide, was offset somewhat by increases in mortality from unintentional injuries, perinatal conditions, homicide, legal intervention, and congenital malformation. Life expectancy for the American Indian and Alaska Native female population increased 2.1 years from 69.2 years in 2021 to 71.3 years in 2022. This increase in life expectancy, primarily due to decreases in mortality from COVID-19, Chronic liver disease and cirrhosis, Chronic lower respiratory diseases, diabetes, and Alzheimer disease, was offset somewhat by increases in mortality from Influenza and pneumonia, homicide, heart disease, cancer, and benign neoplasms.

Life expectancy for the Black non-Hispanic (subsequently, Black) population in 2022 increased 1.6 years to 72.8 years. This increase, due to decreases in mortality from COVID-19, heart disease, homicide, diabetes, and Chronic liver disease and

cirrhosis, was offset by increases in mortality from perinatal conditions, kidney disease, congenital malformations, nutritional deficiencies, and legal intervention. Life expectancy for the Black male population in 2022 increased 1.5 years to 69.1 years. This increase, due to decreases in mortality from COVID-19, homicide, heart disease, diabetes, and cancer, was offset by increases in mortality from perinatal conditions, congenital malformations, kidney disease, anemias, and legal intervention. Life expectancy for the Black female population increased 1.5 years in 2022 to 76.5 years due to decreases in mortality from COVID-19, heart disease, diabetes, homicide, and Chronic liver disease and cirrhosis. This increase was offset somewhat by increases in mortality from perinatal conditions, kidney disease, Influenza and pneumonia, aortic aneurysm, and unintentional injuries.

Life expectancy for the Asian non-Hispanic (subsequently, Asian) population increased 0.9 year to 84.4 years. The increase, primarily due to decreases in mortality from COVID-19, stroke, cancer, hypertension, and diabetes, was offset by increases in mortality due to kidney disease, heart disease, congenital malformations, unintentional injuries, and nutritional deficiencies. Life expectancy for the Asian male population increased 1.1 years in 2022 to 82.3 years due to decreases in mortality from COVID-19, stroke, perinatal conditions, diabetes, and cancer. This increase was offset somewhat by increases in mortality from unintentional injuries, kidney disease, congenital malformations, septicemia, and meningitis. Life expectancy for the Asian female population increased by 0.7 year in 2022 to 86.3 years due to decreases in mortality from COVID-19, cancer, stroke, perinatal conditions, and unintentional injuries. This increase was offset somewhat by increases in mortality from Influenza and pneumonia, kidney disease, nutritional deficiencies, pneumonitis, and *C. difficile*.

Life expectancy for the White non-Hispanic (subsequently, White) population increased 0.8 year in 2022 to 77.5 years (Table 4). This increase, due to decreases in mortality from COVID-19, unintentional injuries, heart disease, cancer, and Chronic liver disease and cirrhosis, was offset to some extent by increases in mortality from Influenza and pneumonia, perinatal conditions, nutritional deficiencies, kidney disease, and Chronic lower respiratory diseases. Life expectancy for the White male population increased 1.1 years in 2022 to 75.1 years due to decreases in mortality from COVID-19, unintentional injuries, heart disease, cancer, and suicide. This increase was offset somewhat by increases in mortality from perinatal conditions, Influenza and pneumonia, kidney disease, nutritional deficiencies, and Parkinson disease. Life expectancy for White females increased 0.6 year in 2022 to 80.1 years. This increase, due to decreases in mortality from COVID-19, unintentional injuries, diabetes, cancer, and medical complications, was offset somewhat by increases in mortality from Influenza and pneumonia, Chronic lower respiratory diseases, nutritional deficiencies, perinatal conditions, and kidney disease.

Life expectancy for the Hispanic population increased by 2.2 years to 80.0 years. This increase, due to decreases in mortality from COVID-19, heart disease, diabetes, cancer, and medical complications, was offset by increases in mortality

from unintentional injuries, Influenza and pneumonia, congenital malformations, perinatal conditions, and nutritional deficiencies. Life expectancy for the Hispanic male population in 2022 increased 2.4 years to 77.0 years due to decreases in mortality from COVID-19, heart disease, Chronic liver disease, diabetes, and homicide. This increase was offset somewhat by increases in mortality from unintentional injuries, perinatal conditions, congenital malformations, Influenza and pneumonia, and legal intervention. Life expectancy for the Hispanic female population increased by 1.7 years in 2022 to 82.8 years due to decreases in mortality from COVID-19, heart disease, cancer, diabetes, and stroke. This increase was offset somewhat by increases in mortality from Influenza and pneumonia, congenital malformations, chronic liver disease, kidney disease, and homicide.

The difference in life expectancy between the male and female populations decreased 0.4 year in 2022 to 5.4 years (Table 4). The narrowing in the male–female life expectancy gap was due primarily to greater decreases in mortality for the male population from COVID-19, heart disease, homicide, cancer, and Chronic lower respiratory diseases (data not shown).

Injury mortality by mechanism and intent

In 2022, a total of 307,785 deaths were classified as injury-related (Table 12). Injury data are presented using the external cause-of-injury mortality matrix for ICD–10, as jointly conceived by the International Collaborative Effort on Injury Statistics and the Injury Control and Emergency Health Services section of the American Public Health Association (27,28). The ICD codes for injuries have two essential dimensions: the mechanism of the injury and its manner or intent. The mechanism involves the circumstances of the injury (such as fall, motor-vehicle traffic, or poisoning). The manner or intent involves whether the injury was purposefully inflicted (where it can be determined) and, when intentional, whether the injury was self-inflicted (suicide) or inflicted upon another person (assault). In other report tables showing cause of death, the focus is on manner or intent, with subcategories showing selected mechanisms. The matrix has two distinct advantages for the analysis of injury mortality data: It contains a comprehensive list of mechanisms, and data can be displayed by mechanism with subcategories of intent, or vice versa. Four major mechanisms of injury in 2022—poisoning, firearm, fall, and motor-vehicle traffic—accounted for 82.3% of all injury deaths.

A total of 112,728 deaths occurred as the result of poisonings in 2022, accounting for 36.6% of all injury deaths (Table 12). The age-adjusted death rate for poisoning did not change significantly from 2021 to 2022. Most poisoning deaths were either unintentional (91.3%) or suicides (5.5%). However, 3.0% of poisoning deaths were of undetermined intent. The age-adjusted death rate for unintentional poisoning was unchanged statistically from 2021 to 2022.

Firearm injuries resulted in 48,204 deaths in 2022 (Table 12), accounting for 15.7% of all injury deaths. The age-adjusted death rate for firearm injuries (all intents) in 2022 decreased 2.7% from 2021. The two major component causes of firearm

injury deaths in 2022 were suicide (56.1%) and homicide (40.8%). The age-adjusted death rate for firearm suicide did not change significantly from 2021 to 2022. The age-adjusted rate for firearm homicide decreased 7.5%, from 6.7 in 2021 to 6.2 in 2022.

A total of 47,984 people died as the result of falls in 2022, accounting for 15.6% of all injury deaths (Table 12). The age-adjusted death rate for falls in 2022 decreased 1.7%, from 11.8 in 2021 to 11.6 in 2022. The overwhelming majority of fall-related deaths (97.2%) were unintentional.

Motor vehicle traffic-related injuries in 2022 resulted in 44,534 deaths, accounting for 14.5% of all injury deaths (Table 12). The age-adjusted death rate for these injuries decreased 3.0%, from 13.3 in 2021 to 12.9 in 2022.

Marital status

For those age 15 and older, the number of deaths in 2022 among people who were married was 1,170,668; widowed, 998,284; divorced, 576,726; and never married, 474,143 (Table 13); see Technical Notes. Those who were widowed had the highest age-adjusted death rate (1,880.3 per 100,000 U.S. standard population), followed by never-married people (1,623.5), divorced people (1,514.4), and married people (790.4). Never-married people had an age-adjusted death rate 45.4% higher than those ever-married (more than 2 times the rate of currently married people and 7.2% higher than divorced people). Widowed people had a rate more than twice that of married people, 24.2% higher than divorced people, and 15.8% higher than never-married people. Divorced people had a rate 1.9 times the rate of those who were married at the time of death.

For all age groups 15 and older, age-specific death rates for married people were much lower than those for never-married people. For those ages 15–24, divorced people had the highest death rate, whereas for those ages 25–34, 35–44, and 75 and older, widowed people had the highest death rate. Never-married people had the highest death rate among those ages 45–54, 55–64, and 65–74 (Table 13).

For each marital status group in 2022, males had higher age-adjusted death rates than females, ranging from 55.9% greater for those never married and widowed to 69.5% higher for those divorced at the time of death (Table 13).

Educational attainment

Age-specific and age-adjusted death rates are shown by educational attainment for age groups in the range of 25–64 years (Table 14). In 2022, a total of 358,700 decedents ages 25–64 had received a high school diploma or equivalent, compared with 271,100 who had completed some college or collegiate degree and 137,271 who had achieved less than a high school diploma or equivalent. In 2022, age-adjusted death rates for those with less than a high school diploma or GED (695.7 per 100,000 U.S. standard population) and for those with a high school diploma or GED (724.4) were more than 3 times the rate for those with some college or collegiate degree (220.3).

Injury at work

Deaths, crude death rates, and age-adjusted death rates for injury at work are shown for those age 15 and older (Tables 15 and 16). Age-adjusted death rates for injury at work were computed using age-specific death rates and the 2000 U.S. standard population for those age 15 and older; see “Computing rates” in Technical Notes. Information on deaths attributed to injuries at work is derived from a separate item on the death certificate that asks the medical certifier whether the death resulted from an injury sustained at work. This item is on the death certificate of all states.

State of residence

Mortality patterns varied considerably by state (Tables 17 and 20). The state with the highest age-adjusted death rate in 2022 was West Virginia (1,115.6 deaths per 100,000 U.S. standard population), with a rate 39.7% above the national rate (798.8). The state with the lowest age-adjusted death rate was Hawaii (615.9), with a rate 22.9% below the national rate. The age-adjusted death rate for West Virginia was 81.1% higher than the rate for Hawaii.

Variations in mortality by state were associated with differences in socioeconomic status, racial and ethnic composition, as well as with differences in risk of specific causes of death (29).

Infant mortality

In 2022, a total of 20,553 deaths occurred in children younger than age 1 year (Tables C, D, 19, and 20). This number represents 633 more infant deaths in 2022 than in 2021. The ratio of male-to-female IMRs was 1.2, the same as in 2021. IMR was 5.60 infant deaths per 1,000 live births, the neonatal mortality rate (deaths of infants ages 0–27 days per 1,000 live births) was 3.58, and the postneonatal mortality rate (deaths of infants ages 28 days–11 months per 1,000 live births) was 2.03 in 2022 (Figure 5 and Tables C and 18; see Technical Notes for information on alternative data sources).

The infant, neonatal, and postneonatal mortality rates increased significantly by 2.9%, 2.9%, and 4.1% from 2021 to 2022, respectively.

The 10 leading causes of infant death in 2022 accounted for 65.2% of all infant deaths in the United States (Table D). By rank, the 10 leading causes were:

1. Congenital malformations, deformations and chromosomal abnormalities (congenital malformations)
2. Disorders related to short gestation and low birth weight, not elsewhere classified (low birth weight)
3. Sudden infant death syndrome (SIDS)
4. Accidents (unintentional injuries)
5. Newborn affected by maternal complications of pregnancy (maternal complications)

Table C. Number of infant, neonatal, and postneonatal deaths and mortality rate, by sex: United States, 2021–2022

[Rates are infant (younger than 1 year), neonatal (younger than 28 days), and postneonatal (28 days–11 months) deaths per 1,000 live births in specified group]

Age and sex	2022		2021		Percent change ¹ from 2021 to 2022
	Number	Rate	Number	Rate	
Infant					
Total	20,553	5.60	19,920	5.44	2.9
Male	11,371	6.07	10,909	5.82	4.3
Female	9,182	5.12	9,011	5.03	1.8
Neonatal					
Total	13,125	3.58	12,768	3.48	2.9
Male	7,130	3.80	6,899	3.68	3.3
Female	5,995	3.34	5,869	3.28	1.8
Postneonatal					
Total	7,428	2.03	7,152	1.95	4.1
Male	4,241	2.26	4,010	2.14	5.6
Female	3,187	1.78	3,142	1.75	1.7

¹Based on a comparison of 2022 and 2021 mortality rates.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

6. Newborn affected by complications of placenta, cord and membranes (cord and placental complications)
7. Bacterial sepsis of newborn
8. Respiratory distress of newborn
9. Intrauterine hypoxia and birth asphyxia
10. Diseases of the circulatory system

The 10 leading causes of infant death remained unchanged in 2022 from 2021, although two causes changed ranks (30). Intrauterine hypoxia and birth asphyxia, the 10th leading cause of infant death in 2021, became the 9th leading cause in 2022,

while Diseases of the circulatory system, the 9th leading cause in 2021, became the 10th leading cause in 2022. Among the 10 leading causes, IMR increased 8.9% from 2021 for maternal complications and 13.8% for bacteria sepsis of newborn. Changes in rates among the other leading causes of infant death were not statistically significant (Table D).

IMRs by race for non-Hispanic origin that are based on the mortality file may be somewhat understated and are better measured using data from the linked file of live births and infant deaths (31); see Technical Notes. Infant mortality data presented in this report use the general mortality file, not the linked file of live births and infant deaths. IMRs for the Hispanic population are

Table D. Number of infant deaths, percentage of total infant deaths, and infant mortality rate for 2022, and percent change in infant mortality rates from 2021 to 2022 for the 10 leading causes of infant death in 2022: United States

[Rates are infant deaths per 100,000 live births]


Rank ¹	Cause of death (based on <i>International Classification of Diseases, 10th Revision</i>)	Number	Percent of total deaths ²	Rate	Percent change ³ from 2021 to 2022
...	All causes	20,553	100.0	560.4	3.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	3,970	19.3	108.2	0.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,884	14.0	78.6	-2.2
3	Sudden infant death syndrome (R95)	1,529	7.4	41.7	4.8
4	Accidents (unintentional injuries) (V01–X59)	1,354	6.6	36.9	3.7
5	Newborn affected by maternal complications of pregnancy (P01)	1,215	5.9	33.1	8.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	649	3.2	17.7	-3.3
7	Bacterial sepsis of newborn (P36)	636	3.1	17.3	13.8
8	Respiratory distress of newborn (P22)	455	2.2	12.4	9.7
9	Intrauterine hypoxia and birth asphyxia (P20–P21)	362	1.8	9.9	1.0
10	Diseases of the circulatory system (I00–I99)	356	1.7	9.7	-11.8
...	All other causes (residual)	7,143	34.8	194.8	...

... Category not applicable.

¹Based on number of deaths; see Technical Notes in this report.²Percentages may not add to 100 due to rounding.³Based on a comparison of the 2022 infant mortality rate with the 2021 infant mortality rate.

NOTE: Due to rounding, percent changes based on rates per 100,000 live births may differ from those computed using rates per 1,000 live births.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Figure 5. Infant, neonatal, and postneonatal mortality rates: United States, 1960–2022

not adjusted for misclassification; see Technical Notes. Because these rates are not adjusted, misclassification of Hispanic origin should be considered when interpreting rate disparities between Hispanic and non-Hispanic populations (16).

In 2022, the IMR for White non-Hispanic infants was 4.39 per 1,000 live births, an increase of 4.5% from 4.20 in 2021. In 2022, the IMR was 5.08 for Hispanic infants, 8.16 for American Indian and Alaska Native non-Hispanic infants, 2.85 for Asian non-Hispanic infants, 11.34 for Black non-Hispanic infants, and 8.79 for Native Hawaiian or Other Pacific Islander non-Hispanic infants (Table 18). The changes in IMRs in 2022 from 2021 for these groups were not statistically significant.

Maternal mortality

In 2022, a total of 817 women died of maternal causes in the United States—388 fewer deaths than in 2021 (Table 21) (30). The overall maternal mortality rate decreased 32.2%, from 32.9 deaths per 100,000 live births in 2021 to 22.3 in 2022. COVID-19 was listed on the death certificate as a contributing cause of death for 10.8% of maternal deaths in 2022 (14). The maternal

mortality rate in 2022 was highest for Black non-Hispanic (subsequently, Black) women (49.5), followed by White non-Hispanic (subsequently, White) (19.0), Hispanic (16.9), and Asian non-Hispanic (subsequently, Asian) (13.2) women. The maternal mortality rate decreased 39.6% for Hispanic women (from 28.0 in 2021 to 16.9 in 2022), 29.2% for Black women (from 69.9 in 2021 to 49.5 in 2022) and 28.6% for White women (from 26.6 to 19.0). The change in maternal mortality rates for Asian women was not significant. In 2022, the maternal mortality rates for American Indian and Alaska Native women and Native Hawaiian or Other Pacific Islander women did not meet National Center for Health Statistics standards of reliability because the numbers of deaths are too low.

Maternal deaths and death rates shown in this report are based on the new method for coding maternal deaths that was adopted by the National Center for Health Statistics starting with the 2018 data year (30) (Technical Notes). This method restricts application of the pregnancy checkbox to decedents ages 10–44 for coding cause of death to a maternal cause when the certificate has no mention of a maternal-related condition but has a positive checkbox entry (30). For women age 45 and older, the checkbox

is used in coding cause of death only if a positive checkbox entry is accompanied by a mention of a maternal-related condition as a cause of death. Maternal deaths include deaths of women while pregnant or within 42 days of being pregnant, from any cause related to or aggravated by the pregnancy but exclude deaths from external causes (that is, accidents, homicides, and suicides); for more information, see “Maternal Mortality in the United States: Changes in Coding, Publication, and Data Release, 2018” (32) and Technical Notes.

References

- Hoyert DL, Singh GK, Rosenberg HM. Sources of data on socio-economic differential mortality in the United States. *Jour Off Stat* 1995;11(3):233–60.
- Curtin SC, Tejada-Vera B, Bastian BA. Deaths: Leading causes for 2022. *Natl Vital Stat Rep*. 2024 Dec;73(10):1–117. DOI: <https://dx.doi.org/10.15620/cdc/164020>.
- Arias E, Xu JQ, Kochanek KD. United States life tables, 2022. *Natl Vital Stat Rep*. 2025 Apr;74(2):1–64. DOI: <https://dx.doi.org/10.15620/cdc/174575>.
- World Health Organization. International statistical classification of diseases and related health problems, 10th revision. 2008 ed. 2009.
- National Vital Statistics System. Volume 1. ICD–10, International statistical classification of diseases and related health problems. Tabular list. (Modified by NCHS for use in the classification and analysis of medical mortality data in the United States) NCHS Instruction Manual; part 2e, vol 1. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
- National Vital Statistics System. Volume 1. ICD–10, International statistical classification of diseases and related health problems. Alphabetical index. (Modified by NCHS for use in the classification and analysis of medical mortality data in the United States) NCHS Instruction Manual; part 2e, vol 3. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
- National Center for Health Statistics. 2003 revision of the U.S. Standard Certificate of Death. 2003. Available from: <https://www.cdc.gov/nchs/data/dvs/DEATH11-03final-acc.pdf>.
- Office of Management and Budget. Revisions to the standards for the classification of federal data on race and ethnicity. *Fed Regist* 62FR58782. 1997. Available from: <https://www.govinfo.gov/content/pkg/FR-1997-10-30/pdf/97-28653.pdf>.
- Murphy SL, Xu JQ, Kochanek KD, Arias E, Tejada-Vera B. Deaths: Final Data for 2018. *Natl Vital Stat Rep*. 2021 Jan;69(13):1–83. PMID: 33541516. Available from: <https://www.cdc.gov/nchs/data/nvsr/nvsr69/nvsr69-13-508.pdf>.
- Office of Management and Budget. Race and ethnic standards for federal statistics and administrative reporting. Statistical Policy Directive no 15. 1977. Available from: <https://wonder.cdc.gov/wonder/help/populations/bridged-race/directive15.html>.
- Heron M. Comparability of Race-specific Mortality Data Based on 1977 Versus 1997 Reporting Standards. *Natl Vital Stat Rep*. 2021 Apr;70(3):1–31. PMID: 34029180. DOI: <https://dx.doi.org/10.15620/cdc/103476>.
- National Center for Health Statistics. Vital statistics online data portal: Mortality multiple cause files. National Center for Health Statistics. Published annually. Available from: https://www.cdc.gov/nchs/data_access/VitalStatsOnline.htm.
- National Center for Health Statistics. National Vital Statistics System: Data release policy for vital statistics micro-data files. Available from: https://www.cdc.gov/nchs/nvss/dvs_data_release.htm.
- Centers for Disease Control and Prevention. CDC WONDER. Available from: <https://wonder.cdc.gov/>.
- Tolson GC, Barnes JM, Gay GA, Kowaleski JL. The 1989 revision of the U.S. Standard Certificates and Reports. *Vital Health Stat* 4. 1991 Jun;(28):1–34. PMID: 1713727. Available from: https://www.cdc.gov/nchs/data/series/sr_04/sr04_028.pdf.
- Arias E, Heron M, Hakes J. The validity of race and Hispanic-origin reporting on death certificates in the United States: An update. *Vital Health Stat* 2. 2016 Aug 1;(172):1–29. PMID: 28436642. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_172.pdf.
- Centers for Disease Control and Prevention. CDC WONDER. Single-race population estimates, United States, 2020–2022. July 1st resident population by state, age, sex, single-race, and Hispanic origin. Data source: Vintage 2022 estimates released by U.S. Census Bureau on June 22, 2023. Available from: <https://wonder.cdc.gov/single-race-population.html>.
- Arias E. United States life tables by Hispanic origin. National Center for Health Statistics. *Vital Health Stat* 2. 2010 Oct;(152):1–35. PMID: 21313788. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_152.pdf.
- Arias E, Xu JQ. Arias E, Xu J. United States Life Tables, 2019. *Natl Vital Stat Rep*. 2022 Mar;70(19):1–59. PMID: 35319436. DOI: <https://dx.doi.org/10.15620/cdc/113096>.
- Hoyert DL, Kochanek KD, Murphy SL. Deaths: Final data for 1997. *Natl Vital Stat Rep*. 1999 Jun 30;47(19):1–104. PMID: 10410536. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_19.pdf.
- Sunenshine RH, McDonald LC. Clostridium difficile-associated disease: New challenges from an established pathogen. *Cleve Clin J Med*. 2006 Feb;73(2):187–97. PMID: 16478043. Available from: <https://www.ccm.org/content/73/2/187.long>.
- Redelings MD, Sorvillo FJ, Mascola L. Increase in Clostridium difficile-related mortality rates, United States, 1999–2004. *Emerg Infect Dis*. 2007 Sep;13(9):1417–9. PMID: 18252127. Available from: https://wwwnc.cdc.gov/eid/article/13/9/06-1116_article.
- World Health Organization. Dementia. Available from: <https://www.who.int/news-room/fact-sheets/detail/dementia>.

24. National Institute on Aging. Alzheimer's disease fact sheet. Available from: <https://www.nia.nih.gov/alzheimers/publication/alzheimers-disease-fact-sheet>.
25. U.S. Department of Health and Human Services. National plan to address Alzheimer's disease: 2019 update. 2019. Available from: <https://aspe.hhs.gov/system/files/pdf/262601/NatlPlan2019.pdf>.
26. Kramarow EA, Tejada-Vera B. Dementia Mortality in the United States, 2000–2017. *Natl Vital Stat Rep*. 2019 Mar;68(2):1–29. PMID: 31112120. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr68/nvsr68_02-508.pdf.
27. National Center for Health Statistics. Proceedings of the international collaborative effort on injury statistics. 1995 Mar;1:1–330. Available from: https://stacks.cdc.gov/view/cdc/11212/cdc_11212_DS1.pdf?download-document-submit=Download.
28. Fingerhut LA, Cox CS, Warner M, participants of the International Collaborative Effort on Injury Statistics. International comparative analysis of injury mortality: Findings from the ICE on Injury Statistics. *Advance Data from Vital and Health Statistics*; no 303. National Center for Health Statistics. 1998. Available from: <https://www.cdc.gov/nchs/data/ad/ad303.pdf>.
29. Pamuk ER, Makuc DM, Heck KE, Reuben C, Lochner K. Socioeconomic status and health chartbook. Health, United States, 1998. National Center for Health Statistics. 1998. Available from: <https://www.cdc.gov/nchs/data/hus/hus98.pdf>.
30. Murphy SL, Kochanek KD, Xu JQ, Arias E. Deaths: Final data for 2021. *Natl Vital Stat Rep*. 2024 Oct;73(8):1–139. DOI: <https://dx.doi.org/10.15620/cdc/158787>.
31. National Center for Health Statistics. User guide to the 2022 period/2021 cohort linked birth/infant death public use file. 2024. Available from: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/DVS/period-cohort-linked/22PE21CO_linkedUG.pdf.
32. Hoyert DL, Miniño AM. Maternal Mortality in the United States: Changes in Coding, Publication, and Data Release, 2018. *Natl Vital Stat Rep*. 2020 Jan;69(2):1–18. PMID: 32510319. Available from: <https://www.cdc.gov/nchs/data/nvsr/nvsr69/nvsr69-02-508.pdf>.
33. World Health Organization. International statistical classification of diseases and related health problems, 10th revision. 1992.
34. National Center for Health Statistics. Comparability of cause-of-death between ICD revisions. 2015. Available from: https://www.cdc.gov/nchs/nvss/mortality/comparability_icd.htm.
35. National Center for Health Statistics. Updated comparability ratios (ICD–10 and ICD–9). 2004. Available from: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Datasets/Comparability/icd9_icd10/.
36. Anderson RN, Miniño AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD–9 and ICD–10: Preliminary estimates. *Natl Vital Stat Rep*. 2001 May 18;49(2):1–32. PMID: 11381674. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_02.pdf.
37. Faust MM, Dolman AB. Comparability of mortality statistics for the sixth and seventh revisions: United States, 1958. *Vital Statistics—Special Reports* 51(4):248–97. National Center for Health Statistics. 1965. Available from: https://www.cdc.gov/nchs/data/spec_rpt51_04.pdf.
38. Klebba AJ, Dolman AB. Comparability of mortality statistics for the seventh and eighth revisions of the International Classification of Diseases, United States. *Vital Health Stat* 2. 1975;(66):1–93. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_066.pdf.
39. Klebba AJ, Scott JH. Estimates of selected comparability ratios based on dual coding of 1976 death certificates by the eighth and ninth revisions of the International Classification of Diseases. *Monthly Vital Statistics Report*; vol 28 no 11. National Center for Health Statistics. 1980. Available from: https://www.cdc.gov/nchs/data/mvssr/supp/mv28_11s.pdf.
40. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2a: Instructions for classifying the underlying cause-of-death. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
41. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2b: Instructions for classifying the multiple causes of death. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
42. National Center for Health Statistics. Instruction manuals. Current manuals. Part 2c: ICD–10 ACME decision tables for classifying underlying causes of death. Published annually. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
43. National Center for Health Statistics. Instruction manuals. Historic manuals—Prior to ICD–10. Part 2g: Data entry instructions for the mortality medical indexing, classification, and references system (MICAR), 1996–1997. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
44. National Center for Health Statistics. Instruction manuals. Historic manuals—ICD–10. Part 2s: SuperMICAR data entry instructions. Available from: https://www.cdc.gov/nchs/nvss/instruction_manuals.htm.
45. National Center for Health Statistics. Multiple cause of death public use file Control Total Table 1. Available from: https://ftp.cdc.gov/pub/Health_Statistics/NCHS/Dataset_Documentation/DVS/mortality/Multiple-Cause-of-Death-Public-Use-File-Control-Total-Table-2022.pdf.
46. Chamblee RF, Evans MC. TRANSAX: The NCHS system for producing multiple cause-of-death statistics, 1968–78. *Vital Health Stat* 1 1986;(20):1–83. 1986. Available from: https://www.cdc.gov/nchs/data/series/sr_01/sr01_020acc.pdf.
47. Israel RA, Rosenberg HM, Curtin LR. Analytical potential for multiple cause-of-death data. *Am J Epidemiol* 1986 Aug;124(2):161–79. PMID: 3524199. Available from: <https://aje.oxfordjournals.org/content/124/2/161.full.pdf>.
48. National Center for Health Statistics. Instruction manual, part 9: ICD–10 cause-of-death lists for tabulating mortality statistics (updated October 2020 to include WHO updates to ICD–10 for data year 2020). 2020.

- Available from: <https://www.cdc.gov/nchs/data/dvs/Part9InstructionManual2020-508.pdf>.
49. Hoyert DL, Arias E, Smith BL, Murphy SL, Kochanek KD. Deaths: Final data for 1999. *Natl Vital Stat Rep*. 2001 Sep 21;49(8):1–113. PMID: 11591077. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr49/nvsr49_08.pdf.
 50. National Center for Health Statistics. Excess deaths associated with COVID-19. 2023 Available from: https://www.cdc.gov/nchs/nvss/vsrr/covid19/excess_deaths.htm.
 51. Ahmad FB, Cisewski JA, Xu J, Anderson RN. Provisional mortality data—United States, 2022. *MMWR Morb Mortal Wkly Rep* 2023 May 5;72(18):488–92. DOI: <http://dx.doi.org/10.15585/mmwr.mm7218a3>.
 52. National Center for Health Statistics, National Vital Statistics System. Computer edits for mortality data, including separate section for fetal deaths effective 2020. NCHS Instruction Manual; part 11. 2020. Available from: https://www.cdc.gov/nchs/data/dvs/manuals/2020_PT11_B.pdf.
 53. National Center for Health Statistics. Instruction manual, part 20: ICD–10 cause-of-death querying, 2013. 2013. Available from: https://www.cdc.gov/nchs/data/dvs/Instruction_Manual_revise20_2013.pdf.
 54. Arias E, Eschbach K, Schauman WS, Backlund EL, Sorlie PD. The Hispanic mortality advantage and ethnic misclassification on US death certificates. *Am J Public Health* 2010 Apr 1;100 Suppl 1(Suppl 1):S171–7. Available from: <https://ajph.aphapublications.org/doi/pdf/10.2105/AJPH.2008.135863>.
 55. Rosenberg HM, Maurer JD, Sorlie PD, Johnson NJ, MacDorman MF, Hoyert DL, et al. Quality of death rates by race and Hispanic origin: A summary of current research, 1999. *National Center for Health Statistics. Vital Health Stat 2*. 1999 Sept;(128):1–15. PMID: 10611854. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_128.pdf.
 56. Sorlie PD, Rogot E, Johnson NJ. Validity of demographic characteristics on the death certificate. *Epidemiology*. 1992 Mar;3(2):181–4.
 57. Arias E, Xu JQ, Curtin S, Bastian B, Tejada-Vera B. Mortality Profile of the Non-Hispanic American Indian or Alaska Native Population, 2019. *Natl Vital Stat Rep*. 2021 Nov;70(12):1–27. PMID: 34842523. DOI: <https://dx.doi.org/10.15620/cdc:110370>.
 58. Mulry M. Summary of accuracy and coverage evaluation for Census 2000. Research Report Series Statistics #2006–3. Washington, DC: U.S. Census Bureau. 2006. Available from: <https://www.census.gov/library/working-papers/2006/adrm/rrs2006-03.html>.
 59. Poe GS, Powell-Griner E, McLaughlin JK, Placek PJ, Thompson GB, Robinson K. Comparability of the death certificate and the 1986 National Mortality Followback Survey. *National Center for Health Statistics. Vital Health Stat 2*. 1993 Nov;(118):1–57. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_118.pdf.
 60. U.S. Census Bureau. DSSD 2010 census coverage measurement memorandum series 2010–G–01. 2012. Available from: <https://www2.census.gov/programs-surveys/decennial/2010/technical-documentation/methodology/g-series/g01.pdf>.
 61. Osterman MJK, Hamilton BE, Martin JA, Driscoll AK, Valenzuela CP. Births: Final Data for 2022. *Natl Vital Stat Rep*. 2024 Apr;73(2):1–56. PMID: 38625869. DOI: <https://dx.doi.org/10.15620/cdc:145588>.
 62. Sirken MG. Comparison of two methods of constructing abridged life tables by reference to a “standard” table. *National Center for Health Statistics. Vital Health Stat 2*. 1966 Feb;(4):1–19. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_004.pdf.
 63. Anderson RN. Method for constructing complete annual U.S. life tables. *National Center for Health Statistics. Vital Health Stat 2*. 1999 Dec;(129):1–36. Available from: https://www.cdc.gov/nchs/data/series/sr_02/sr02_129.pdf.
 64. National Center for Health Statistics. U.S. decennial life tables for 1989–91. *Methodology of the National and State Life Tables*; vol 1 no 2:1–13. 1998. Available from: https://www.cdc.gov/nchs/data/lifetables/life89_1_2.pdf.
 65. Wei R, Curtin LR, Arias E, Anderson RN. U.S. decennial life tables for 1999–2001: State life tables. *Natl Vital Stat Rep*. 2012 Sep 14;60(9):1–66. PMID: 24979971. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr57/nvsr57_04.pdf.
 66. Miniño AM, Murphy SL, Xu JQ, Kochanek KD. Deaths: Final data for 2008. *Natl Vital Stat Rep*. 2011 Dec 7;59(10):1–126. PMID: 22808755. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr59/nvsr59_10.pdf.
 67. Arias E. United States life tables, 2008. *Natl Vital Stat Rep*. 2012 Sep 24;61(3):1–63. PMID: 24974590. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr61/nvsr61_03.pdf.
 68. Kochanek KD, Maurer JD, Rosenberg HM. Causes of death contributing to changes in life expectancy: United States, 1984–89. *National Center for Health Statistics. Vital Health Stat*. 1994 May;20(23):1–37. Available from: https://www.cdc.gov/nchs/data/series/sr_20/sr20_023.pdf.
 69. Arriaga EE. Changing trends in mortality decline during the last decades. In: Ruzicka L, Wunsch G, Kane P, editors. *Differential mortality: Methodological issues and biosocial factors*. Oxford, England: Clarendon Press;1989.
 70. Arriaga EE. Measuring and explaining the change in life expectancies. *Demography* 1984 Feb;21(1):83–96.
 71. Miniño AM, Anderson RN, Fingerhut LA, Boudreault MA, Warner M. Deaths: injuries, 2002. *Natl Vital Stat Rep*. 2006 Jan 31;54(10):1–124. PMID: 16485447. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr54/nvsr54_10.pdf.
 72. Sorlie PD, Johnson NJ. Validity of education information on the death certificate. *Epidemiology*. 1996 Jul;7(4):437–9. PMID: 8793373.
 73. Arias E, Liebler CA, Garcia MA, Sáenz R. Data impacts of changes in U.S. Census Bureau procedures for race and ethnicity data. *SSM Popul Health*. 2025 Jan 9; 29:101742. PMID: 39897609. DOI: <https://doi.org/10.1016/j.ssmph.2024.101742>.
 74. U.S. Census Bureau. American Community Survey, 2022 1-year ACS public use microdata sample. Available from: https://www2.census.gov/programs-surveys/acs/tech-docs/pums/2022ACS_PUMS_User_Guide.pdf.

75. U.S. Census Bureau. International data base. Population by single years of age and sex, 2022. 2023. Available from: https://www.census.gov/data-tools/demo/idb/#/dashboard?YR_ANIM=2022&FIPS_SINGLE=&dashPages=DASH&COUNTRY_YEAR=2022&COUNTRY_YR_ANIM=2024.
76. Anderson RN, Rosenberg HM. Age standardization of death rates: Implementation of the year 2000 standard. *Natl Vital Stat Rep.* 1998 Oct 7;47(3):1–17. PMID: 9796247. Available from: https://www.cdc.gov/nchs/data/nvsr/nvsr47/nvsr47_03.pdf.
77. Brillinger DR. The natural variability of vital rates and associated statistics. *Biometrics.* 1986 Dec; 42(4):693–734. PMID: 3814721.
78. Fay MP, Feuer EJ. Confidence intervals for directly standardized rates: A method based on the gamma distribution. *Stat Med* 1997 Apr 15;16(7):791–801. PMID: 9131766.
79. National Center for Health Statistics. Vital statistics of the United States: Mortality, 1999. Technical appendix. 2004:1–93. Available from: <https://www.cdc.gov/nchs/data/statab/techap99.pdf>.
80. Schenker N, Gentleman JF. On judging the significance of differences by examining the overlap between confidence intervals. *Am Stat.* 2001 Aug;55(3):182–6. Available from: https://www.jstor.org/stable/2685796?seq=1#page_scan_tab_contents.
81. Arnold SF. *Mathematical statistics.* Englewood Cliffs, NJ: Prentice Hall; 1990.
- causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022 53
10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022. 63
11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022. 73
12. Number of deaths, death rate, and age-adjusted death rate for injury deaths, by mechanism and intent of death: United States, 2022. 83
13. Number of deaths, death rate, and age-adjusted death rate for age 15 and older, by marital status and sex: United States, 2022. 85
14. Number of deaths, death rate, and age-adjusted death rate for ages 25–64, by educational attainment and sex: United States, 2022. 86
15. Number of deaths and death rate by age and age-adjusted death rate for injury at work for age 15 and older, by Hispanic origin and race and sex: United States, 2022. 87
16. Number of deaths, death rate, and age-adjusted death rate for injury at work for age 15 and older, by Hispanic origin and race and sex: United States, 1997–2022. 88
17. Number of deaths, death rate, and age-adjusted death rate for major causes of death: United States, each state, Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, and Northern Marianas, 2022. 90
18. Infant, neonatal, and postneonatal mortality rates, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000–2022. 97
19. Number of infant deaths and infant mortality rate for 130 selected causes, by Hispanic origin and race: United States, 2022. 99
20. Number of infant deaths and infant mortality rate, by Hispanic origin and race for the United States, each state, Puerto Rico, U.S. Virgin Islands, Guam, American Samoa, and Northern Marianas, and by sex for the United States: 2022. 106
21. Number of maternal deaths and maternal mortality rate for selected causes, by Hispanic origin and race: United States, 2022. 108
22. Number of deaths, death rate, and age-adjusted death rate for dementia-related causes, by Hispanic origin and race and sex: United States, 1999–2022. 110
23. Number of deaths, death rate, and age-adjusted death rate for drug-induced causes, by Hispanic origin and race and sex: United States, 1999–2022. 112
24. Number of deaths, death rate, and age-adjusted death rate for alcohol-induced causes, by Hispanic origin and race and sex: United States, 1999–2022. 114
25. Number of deaths, death rate, and age-adjusted death rate for injury by firearms, by Hispanic origin and race and sex: United States, 1999–2022. 116

List of Detailed Tables

1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2022 22
2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2022. 24
3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2022. 28
4. Life expectancy at selected ages, by Hispanic origin and race and sex: United States, 2022 30
5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2022 31
6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022 33
7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022 43
8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022 48
9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related

Table 1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2022

[Excludes deaths of nonresidents of the United States. Data for specified race or Hispanic-origin groups other than non-Hispanic White and non-Hispanic Black should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys, although misclassification is very minor for the Hispanic and non-Hispanic Asian or Pacific Islander populations; see Technical Notes in this report]

Hispanic origin and race and year	Number			Crude death rate ¹			Age-adjusted death rate ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All origins and races³									
2022.....	3,279,857	1,719,250	1,560,607	984.1	1,040.2	928.9	798.8	954.5	666.1
2021.....	3,464,231	1,838,108	1,626,123	1,043.8	1,118.2	970.8	879.7	1,048.0	733.3
2020.....	3,383,729	1,769,884	1,613,845	1,027.0	1,090.8	965.1	835.4	998.3	695.1
2019.....	2,854,838	1,473,823	1,381,015	869.7	911.7	829.0	715.2	846.7	602.7
2018.....	2,839,205	1,458,469	1,380,736	867.8	905.2	831.6	723.6	855.5	611.3
2017.....	2,813,503	1,439,111	1,374,392	863.8	897.2	831.4	731.9	864.5	619.7
2016.....	2,744,248	1,400,232	1,344,016	849.3	880.2	819.3	728.8	861.0	617.5
2015.....	2,712,630	1,373,404	1,339,226	844.0	868.0	820.7	733.1	863.2	624.2
2014.....	2,626,418	1,328,241	1,298,177	823.7	846.4	801.7	724.6	855.1	616.7
2013.....	2,596,993	1,306,034	1,290,959	821.5	839.1	804.4	731.9	863.6	623.5
2012.....	2,543,279	1,273,722	1,269,557	810.2	824.5	796.4	732.8	865.1	624.7
2011.....	2,515,458	1,254,978	1,260,480	807.3	818.7	796.3	741.3	875.3	632.4
2010.....	2,468,435	1,232,432	1,236,003	799.5	812.0	787.4	747.0	887.1	634.9
2000.....	2,403,351	1,177,578	1,225,773	854.0	853.0	855.0	869.0	1,053.8	731.4
1990.....	2,148,463	1,113,417	1,035,046	863.8	918.4	812.0	938.7	1,202.8	750.9
1980.....	1,989,841	1,075,078	914,763	878.3	976.9	785.3	1,039.1	1,348.1	817.9
1970.....	1,921,031	1,078,478	842,553	945.3	1,090.3	807.8	1,222.6	1,542.1	971.4
1960.....	1,711,982	975,648	736,334	954.7	1,104.5	809.2	1,339.2	1,609.0	1,105.3
1950.....	1,452,454	827,749	624,705	963.8	1,106.1	823.5	1,446.0	1,674.2	1,236.0
1940.....	1,417,269	791,003	626,266	1,076.4	1,197.4	954.6	1,785.0	1,976.0	1,599.4
Hispanic⁴									
2022.....	275,684	155,240	120,444	433.0	481.0	383.7	614.7	748.8	498.4
2021.....	315,664	181,195	134,469	503.9	571.3	434.7	724.7	884.9	582.7
2020.....	305,708	175,585	130,123	498.6	567.8	428.2	723.6	903.8	570.1
2019.....	212,397	117,683	94,714	350.7	384.9	315.7	523.8	633.2	430.7
2018.....	204,719	113,045	91,674	341.9	373.9	309.3	524.1	633.1	431.7
2017.....	197,249	108,579	88,670	334.6	364.6	304.0	524.7	631.8	434.2
2016.....	188,254	103,532	84,722	327.6	356.8	297.7	525.8	631.8	436.4
2015.....	179,457	98,170	81,287	317.1	343.2	290.4	525.3	628.9	438.3
2014.....	169,387	92,474	76,913	305.8	330.1	281.0	523.3	626.8	437.5
2013.....	163,241	88,880	74,361	301.9	323.7	279.4	535.4	639.8	448.6
2012.....	156,419	85,238	71,181	295.0	316.5	272.7	539.1	643.9	452.5
2011.....	149,635	81,887	67,748	287.5	309.7	264.6	540.7	647.3	452.8
2010.....	144,490	79,622	64,868	286.2	310.8	260.9	558.6	677.7	463.4
2000.....	107,254	60,172	47,082	303.8	331.3	274.6	665.7	818.1	546.0
Non-Hispanic, single race⁵									
American Indian and Alaska Native:									
2022.....	23,613	12,721	10,892	975.4	1,062.4	890.2	947.9	1,084.3	816.1
2021.....	26,972	14,724	12,248	1,100.0	1,216.9	986.2	1,109.2	1,282.7	946.6
2020.....	24,725	13,431	11,294	1,016.5	1,123.4	913.2	1,036.2	1,205.9	881.5
2019.....	18,057	9,732	8,325	741.6	812.1	673.3	782.5	901.9	673.3
2018.....	17,790	9,678	8,112	735.9	813.5	660.8	790.8	918.7	673.1
Asian:									
2022.....	89,591	46,137	43,454	441.9	472.3	413.6	417.5	501.7	350.7
2021.....	92,432	48,386	44,046	469.5	511.7	430.5	461.7	554.9	386.3
2020.....	91,175	47,699	43,476	470.8	515.8	429.6	457.7	557.4	378.5
2019.....	70,532	35,914	34,618	373.1	398.7	349.8	372.8	442.4	317.2
2018.....	68,768	35,089	33,679	367.2	393.4	343.3	381.2	454.1	324.1
Black:									
2022.....	411,934	219,538	192,396	979.2	1,085.4	880.8	1,002.8	1,257.5	809.0
2021.....	449,764	238,599	211,165	1,074.5	1,185.8	971.4	1,118.0	1,374.0	917.2
2020.....	449,213	237,703	211,510	1,084.3	1,200.0	978.4	1,119.0	1,399.0	905.2
2019.....	346,677	181,363	165,314	842.5	921.8	769.9	884.0	1,092.8	724.9
2018.....	341,408	177,958	163,450	834.7	909.8	765.9	892.6	1,102.8	733.7

Table 1. Number of deaths, death rate, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, 2000, and 2010–2022—Con.

[Excludes deaths of nonresidents of the United States. Data for specified race or Hispanic-origin groups other than non-Hispanic White and non-Hispanic Black should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys, although misclassification is very minor for the Hispanic and non-Hispanic Asian or Pacific Islander populations; see Technical Notes in this report]

Hispanic origin and race and year	Number			Crude death rate ¹			Age-adjusted death rate ²		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Non-Hispanic, single race ⁵ —Con.									
Native Hawaiian or Other Pacific Islander:									
2022	4,592	2,461	2,131	722.1	765.3	677.9	782.0	859.5	705.4
2021	5,223	2,906	2,317	834.0	917.5	748.6	924.3	1,042.8	806.0
2020	4,439	2,489	1,950	723.5	804.6	641.1	821.3	947.9	699.8
2019	3,491	1,938	1,553	585.8	646.5	524.4	679.0	769.0	589.5
2018	3,277	1,786	1,491	558.9	605.4	511.8	675.7	758.1	597.3
White:									
2022	2,448,093	1,267,526	1,180,567	1,247.6	1,299.7	1,196.1	822.2	972.1	692.7
2021	2,548,809	1,337,385	1,211,424	1,294.9	1,368.8	1,222.1	893.9	1,055.6	751.4
2020	2,484,072	1,278,612	1,205,460	1,262.4	1,317.7	1,208.6	834.7	985.0	703.1
2019	2,183,844	1,115,767	1,068,077	1,106.8	1,146.6	1,068.1	739.9	868.8	627.4
2018	2,182,552	1,108,848	1,073,704	1,104.8	1,138.2	1,072.3	748.7	878.0	636.5

¹Rates are on an annual basis per 100,000 population in specified group; see Technical Notes.

²Age-adjusted rates are per 100,000 U.S. standard population. For method of computation, see Technical Notes.

³Includes origins and races not shown separately; see Technical Notes.

⁴Includes people of Hispanic origin of any race. The Hispanic-origin category is consistent with 1997 Office of Management and Budget (OMB) standards; see Technical Notes.

⁵Only one race was reported on the death certificate. Hispanic-origin and race categories are consistent with 1997 OMB standards; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2022

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race³ should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	All ages	Age group											Age not stated	Age-adjusted rate ²
		Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
Number														
Total	3,279,857	20,553	4,156	6,239	35,232	74,369	111,605	183,284	417,541	668,581	824,903	933,291	103	...
Male	1,719,250	11,371	2,353	3,587	25,637	52,511	73,486	114,847	254,278	384,985	427,507	368,601	87	...
Female	1,560,607	9,182	1,803	2,652	9,595	21,858	38,119	68,437	163,263	283,596	397,396	564,690	16	...
Single race ³	3,260,336	19,576	3,959	6,001	34,287	72,863	110,088	181,423	414,713	665,106	821,717	930,500	103	...
Male	1,708,400	10,857	2,245	3,452	24,988	51,467	72,538	113,703	252,590	383,065	425,888	367,520	87	...
Female	1,551,936	8,719	1,714	2,549	9,299	21,396	37,550	67,720	162,123	282,041	395,829	562,980	16	...
Two or more races ⁴	19,521	977	197	238	945	1,506	1,517	1,861	2,828	3,475	3,186	2,791	–	...
Male	10,850	514	108	135	649	1,044	948	1,144	1,688	1,920	1,619	1,081	–	...
Female	8,671	463	89	103	296	462	569	717	1,140	1,555	1,567	1,710	–	...
Hispanic, total ⁵	275,684	4,765	869	1,386	8,035	13,791	17,794	25,143	39,748	49,780	54,345	60,018	10	...
Male	155,240	2,612	466	787	6,040	10,542	12,875	16,955	25,576	28,932	27,163	23,283	9	...
Female	120,444	2,153	403	599	1,995	3,249	4,919	8,188	14,172	20,848	27,182	36,735	1	...
Hispanic, single race ³	273,099	4,552	827	1,356	7,892	13,568	17,563	24,896	39,386	49,394	53,974	59,681	10	...
Male	153,772	2,502	442	769	5,938	10,380	12,727	16,807	25,358	28,709	26,974	23,157	9	...
Female	119,327	2,050	385	587	1,954	3,188	4,836	8,089	14,028	20,685	27,000	36,524	1	...
American Indian and Alaska Native	1,705	41	11	8	90	143	158	180	290	330	246	208	–	...
Male	1,001	25	10	3	58	99	118	123	175	193	108	89	–	...
Female	704	16	1	5	32	44	40	57	115	137	138	119	–	...
Asian	886	39	12	12	43	54	53	72	109	156	182	154	–	...
Male	486	17	7	9	28	41	34	47	69	88	86	60	–	...
Female	400	22	5	3	15	13	19	25	40	68	96	94	–	...
Black	5,353	305	62	54	257	440	438	544	714	880	808	851	–	...
Male	3,137	165	28	34	192	332	322	364	436	515	422	327	–	...
Female	2,216	140	34	20	65	108	116	180	278	365	386	524	–	...
Native Hawaiian or Other Pacific Islander	315	10	6	4	22	30	23	33	55	54	42	36	–	...
Male	191	7	3	1	16	24	17	25	38	35	15	10	–	...
Female	124	3	3	3	6	6	6	8	17	19	27	26	–	...
White	264,840	4,157	736	1,278	7,480	12,901	16,891	24,067	38,218	47,974	52,696	58,432	10	...
Male	148,957	2,288	394	722	5,644	9,884	12,236	16,248	24,640	27,878	26,343	22,671	9	...
Female	115,883	1,869	342	556	1,836	3,017	4,655	7,819	13,578	20,096	26,353	35,761	1	...
Hispanic, two or more races ⁴	2,585	213	42	30	143	223	231	247	362	386	371	337	–	...
Male	1,468	110	24	18	102	162	148	148	218	223	189	126	–	...
Female	1,117	103	18	12	41	61	83	99	144	163	182	211	–	...

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race³ should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	All ages	Age group											Age not stated	Age-adjusted rate ²
		Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
Number—Con.														
Non-Hispanic, total	2,994,727	15,560	3,275	4,849	27,152	60,421	93,436	157,417	375,693	615,996	768,717	872,167	44	...
Male	1,557,750	8,627	1,881	2,796	19,568	41,863	60,349	97,367	227,139	354,078	399,232	344,814	36	...
Female	1,436,977	6,933	1,394	2,053	7,584	18,558	33,087	60,050	148,554	261,918	369,485	527,353	8	...
Non-Hispanic, single race ³	2,977,823	14,798	3,121	4,641	26,350	59,140	92,154	155,807	373,233	612,917	765,903	869,715	44	...
Male	1,548,383	8,224	1,798	2,679	19,021	40,982	59,552	96,374	225,671	352,385	397,802	343,859	36	...
Female	1,429,440	6,574	1,323	1,962	7,329	18,158	32,602	59,433	147,562	260,532	368,101	525,856	8	...
American Indian and Alaska Native	23,613	210	64	110	664	1,729	2,302	2,781	4,298	4,712	4,053	2,690	—	...
Male	12,721	120	32	61	447	1,097	1,461	1,663	2,442	2,491	1,923	984	—	...
Female	10,892	90	32	49	217	632	841	1,118	1,856	2,221	2,130	1,706	—	...
Asian	89,591	624	120	193	823	1,580	2,595	4,942	8,971	16,293	22,628	30,822	—	...
Male	46,137	356	70	110	569	1,097	1,661	3,071	5,562	9,376	11,746	12,519	—	...
Female	43,454	268	50	83	254	483	934	1,871	3,409	6,917	10,882	18,303	—	...
Black	411,934	5,800	1,116	1,502	9,366	17,383	23,234	35,376	73,375	96,857	80,186	67,734	5	...
Male	219,538	3,230	634	874	7,180	12,282	14,840	21,498	43,345	54,062	38,510	23,078	5	...
Female	192,396	2,570	482	628	2,186	5,101	8,394	13,878	30,030	42,795	41,676	44,656	—	...
Native Hawaiian or Other Pacific Islander	4,592	89	26	26	92	207	347	559	889	1,057	841	459	—	...
Male	2,461	49	18	14	71	141	219	315	501	544	401	188	—	...
Female	2,131	40	8	12	21	66	128	244	388	513	440	271	—	...
White	2,448,093	8,075	1,795	2,810	15,405	38,241	63,676	112,149	285,700	493,998	658,195	768,010	39	...
Male	1,267,526	4,469	1,044	1,620	10,754	26,365	41,371	69,827	173,821	285,912	345,222	307,090	31	...
Female	1,180,567	3,606	751	1,190	4,651	11,876	22,305	42,322	111,879	208,086	312,973	460,920	8	...
Non-Hispanic, two or more races ⁴	16,904	762	154	208	802	1,281	1,282	1,610	2,460	3,079	2,814	2,452	—	...
Male	9,367	403	83	117	547	881	797	993	1,468	1,693	1,430	955	—	...
Female	7,537	359	71	91	255	400	485	617	992	1,386	1,384	1,497	—	...
Not stated or not classifiable origin ⁶	9,446	228	12	4	45	157	375	724	2,100	2,805	1,841	1,106	49	...
Male	6,260	132	6	4	29	106	262	525	1,563	1,975	1,112	504	42	...
Female	3,186	96	6	—	16	51	113	199	537	830	729	602	7	...

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race³ should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	All ages	Age group											Age not stated	Age-adjusted rate ²
		Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
								Rate ⁷						
Total ⁸	984.1	558.0	28.0	15.3	79.5	163.4	255.4	453.3	992.1	1,978.7	4,708.2	14,389.6	...	798.8
Male	1,040.2	603.9	31.0	17.1	113.0	226.4	332.2	566.3	1,230.1	2,419.7	5,527.0	16,139.6	...	954.5
Female	928.9	510.0	24.8	13.3	44.3	98.0	176.7	339.6	762.4	1,586.3	4,061.0	13,438.5	...	666.1
Single race ³	1,008.7	569.7	28.5	15.6	81.0	165.5	257.9	456.9	999.0	1,990.6	4,736.1	14,482.1	...	804.5
Male	1,065.9	617.9	31.6	17.5	115.4	229.2	335.3	570.4	1,238.1	2,434.4	5,560.3	16,254.3	...	961.5
Female	952.4	519.3	25.3	13.6	45.0	99.2	178.4	342.5	767.9	1,595.6	4,084.7	13,519.9	...	670.6
Two or more races ⁵	193.8	395.5	20.2	9.7	46.8	102.3	150.1	257.7	495.5	922.6	1,868.5	4,597.1	...	346.7
Male	216.5	408.4	21.7	10.8	63.3	141.8	194.7	331.5	623.7	1,096.1	2,147.8	4,750.2	...	409.6
Female	171.4	382.1	18.7	8.6	29.8	62.8	108.6	190.2	379.9	771.8	1,647.2	4,505.3	...	291.4
Hispanic, total ¹⁵	433.0	477.4	22.2	13.1	74.4	142.1	196.3	327.7	702.5	1,522.8	3,682.6	11,107.1	...	614.7
Male	481.0	514.1	23.4	14.6	109.3	209.7	270.9	432.1	905.9	1,900.2	4,368.6	12,379.5	...	748.8
Female	383.7	439.4	21.0	11.6	37.8	69.4	114.0	218.4	499.9	1,193.7	3,183.0	10,427.8	...	498.4
Hispanic, single race ³	443.4	480.5	22.3	13.5	75.9	144.3	199.0	331.7	710.4	1,540.2	3,721.6	11,208.1	...	621.4
Male	492.3	518.9	23.4	15.0	111.6	213.1	274.8	437.5	915.8	1,920.8	4,413.5	12,496.2	...	757.1
Female	393.1	440.8	21.1	11.9	38.5	70.4	115.3	220.8	505.5	1,208.0	3,217.6	10,520.6	...	503.7
American Indian and Alaska Native	86.9	97.4	*	*	28.3	49.5	57.2	79.9	179.5	390.0	791.9	2,210.4	...	143.9
Male	98.1	116.8	*	*	35.6	65.1	78.0	101.6	207.8	459.7	766.1	2,594.8	...	165.7
Female	74.8	*	*	*	20.7	32.2	32.0	54.7	148.7	321.4	813.2	1,990.0	...	122.9
Asian	130.7	256.2	*	*	39.2	54.4	58.4	98.5	212.6	546.0	1,563.7	4,075.2	...	229.5
Male	142.8	*	*	*	49.5	82.2	73.5	129.3	277.5	674.3	1,795.4	4,662.0	...	275.1
Female	118.5	294.0	*	*	*	*	*	68.0	151.5	438.2	1,401.7	3,772.1	...	192.8
Black	160.8	409.0	22.1	8.3	46.1	87.4	96.2	159.2	287.8	625.8	1,364.8	4,272.9	...	257.3
Male	190.5	435.3	19.6	10.3	67.3	129.3	144.7	226.0	374.1	812.0	1,741.5	4,890.8	...	329.7
Female	131.7	381.9	24.6	6.3	23.9	43.8	49.8	99.7	211.4	472.9	1,103.7	3,960.7	...	198.9
Native Hawaiian or Other Pacific Islander	129.7	*	*	*	58.7	82.2	62.0	124.4	308.8	617.3	1,259.4	3,543.3	...	230.5
Male	152.3	*	*	*	*	124.0	*	180.6	426.2	849.5	*	*	...	253.8
Female	105.6	*	*	*	*	*	*	*	*	*	1,416.6	4,037.3	...	198.6
White	478.3	513.3	23.1	14.4	79.8	152.3	212.1	351.9	754.5	1,629.3	3,917.7	11,724.9	...	655.0
Male	530.0	554.9	24.3	15.9	117.7	225.0	292.0	463.0	972.2	2,031.8	4,649.1	13,064.0	...	798.1
Female	424.9	470.1	21.9	12.8	40.1	74.0	123.4	234.9	536.4	1,278.0	3,385.4	11,009.4	...	530.6
Hispanic, two or more races ⁴	124.5	419.8	21.3	6.0	35.4	72.6	95.1	146.5	317.9	622.1	1,458.5	4,279.4	...	258.4
Male	140.9	426.1	23.8	*	49.7	103.6	121.9	178.7	402.7	798.7	1,781.7	4,558.6	...	311.6
Female	107.9	413.2	*	*	20.7	40.4	68.3	115.4	241.0	477.6	1,227.3	4,128.4	...	214.1

Table 2. Number of deaths and death rate by age, and age-adjusted death rate, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for specified categories other than Black non-Hispanic, single race and White non-Hispanic, single race³ should be interpreted with caution because of inconsistencies between reporting these items on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and race and sex	All ages	Age group											Age not stated	Age-adjusted rate ²
		Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
Rate ⁷ —Con.														
Non-Hispanic, total	1,110.7	579.5	29.9	16.0	80.9	168.8	269.8	480.5	1,031.3	2,018.4	4,791.1	14,669.3	...	820.4
Male	1,171.2	627.5	33.6	18.0	114.0	230.5	347.5	595.3	1,272.6	2,460.9	5,612.7	16,453.0	...	977.4
Female	1,051.8	529.1	26.1	13.9	46.3	105.3	191.7	366.1	799.6	1,623.7	4,136.8	13,698.3	...	686.9
Non-Hispanic, single race ³	1,138.2	594.6	30.7	16.4	82.5	170.8	272.2	483.8	1,037.6	2,029.2	4,817.1	14,759.3	...	825.8
Male	1,200.0	645.1	34.6	18.4	116.4	233.1	350.3	598.9	1,279.9	2,474.5	5,643.9	16,565.5	...	984.0
Female	1,078.1	541.5	26.7	14.2	47.0	106.5	193.4	368.9	804.7	1,632.0	4,158.7	13,777.0	...	691.3
American Indian and Alaska Native	975.4	833.5	60.0	33.7	182.9	466.8	734.7	1,003.1	1,467.7	2,198.2	4,134.7	8,138.4	...	947.9
Male	1,062.4	932.8	58.9	36.6	242.2	581.5	930.0	1,209.6	1,741.4	2,515.4	4,365.5	7,885.2	...	1,084.3
Female	890.2	729.8	61.1	30.6	121.6	347.7	538.3	800.0	1,216.1	1,925.8	3,946.3	8,292.0	...	816.1
Asian	441.9	306.0	14.1	8.4	33.4	49.2	78.6	173.7	398.9	962.5	2,735.3	9,206.1	...	417.5
Male	472.3	338.1	16.0	9.3	45.7	69.1	104.8	228.0	531.6	1,249.6	3,256.5	10,082.0	...	501.7
Female	413.6	271.7	12.2	7.4	20.8	29.8	54.4	124.9	283.5	733.9	2,332.5	8,689.8	...	350.7
Black	979.2	1,066.7	52.8	26.6	153.8	264.9	412.6	701.8	1,468.2	2,791.3	5,429.7	13,311.4	...	1,002.8
Male	1,085.4	1,170.5	59.2	30.6	233.3	374.0	545.0	899.8	1,866.7	3,600.0	6,722.8	15,380.3	...	1,257.5
Female	880.8	959.6	46.2	22.6	72.5	155.7	288.6	523.4	1,122.4	2,174.3	4,610.2	12,446.1	...	809.0
Native Hawaiian or Other Pacific Islander	722.1	1,100.4	76.7	29.2	103.2	207.0	349.9	731.0	1,310.0	2,322.6	4,213.6	6,699.8	...	782.0
Male	765.3	1,166.7	*	*	155.4	274.5	428.7	809.4	1,490.9	2,496.6	4,306.7	6,934.7	...	859.5
Female	677.9	1,028.8	*	*	48.3	135.7	266.1	649.8	1,132.5	2,162.8	4,132.2	6,545.9	...	705.4
White	1,247.6	472.8	25.4	14.0	67.2	156.8	259.7	468.0	1,007.3	1,993.4	4,883.6	15,332.2	...	822.2
Male	1,299.7	509.9	28.8	15.7	91.3	211.4	331.3	573.3	1,233.6	2,409.1	5,695.4	17,191.0	...	972.1
Female	1,196.1	433.8	21.8	12.2	41.8	99.7	185.4	359.1	783.8	1,611.3	4,220.1	14,301.9	...	692.7
Non-Hispanic, two or more races ⁴	211.5	388.2	19.9	10.7	49.7	110.0	167.0	290.9	538.5	978.7	1,939.7	4,640.7	...	366.8
Male	235.9	402.8	20.9	11.7	66.7	151.9	218.1	378.7	678.0	1,149.8	2,207.8	4,776.7	...	433.0
Female	187.4	373.1	18.7	9.5	32.1	68.4	120.6	211.9	412.8	828.3	1,723.5	4,557.9	...	308.2

... Category not applicable.

— Quantity zero.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

¹Death rates for "Younger than 1" (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

²For method of computation, see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Two or more races were reported on the death certificate; see Technical Notes.

⁵Includes people of Hispanic origin of any race; see Technical Notes.

⁶Includes origin not stated or not classifiable; see Technical Notes.

⁷Data for age not stated included in "All ages" but not distributed among age groups.

⁸Includes deaths with origin not stated and origin not classifiable; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2022

[Rates are per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Populations used for computing death rates for Hispanic total are postcensal estimates based on the 2010 census estimated as of July 1, 2022; populations used for computing death rates for Mexican, Puerto Rican, Cuban, Central American, South American, Dominican, and Other and unknown Hispanic are estimates based on the 2022 1-year American Community Survey adjusted to control totals. The control totals are 2010-based postcensal estimates for the United States for July 1, 2022; see Technical Notes. Hispanic origin is reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and sex	Age group												Age not stated	Age- adjusted rate ²
	All ages	Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
		Number												
Hispanic	275,684	4,765	869	1,386	8,035	13,791	17,794	25,143	39,748	49,780	54,345	60,018	10	...
Male	155,240	2,612	466	787	6,040	10,542	12,875	16,955	25,576	28,932	27,163	23,283	9	...
Female	120,444	2,153	403	599	1,995	3,249	4,919	8,188	14,172	20,848	27,182	36,735	1	...
Central American	18,332	525	96	137	959	1,461	1,776	2,075	2,633	2,950	2,813	2,906	1	...
Male	10,625	290	49	77	775	1,203	1,382	1,506	1,674	1,577	1,215	876	1	...
Female	7,707	235	47	60	184	258	394	569	959	1,373	1,598	2,030	–	...
Cuban	20,303	100	19	27	132	285	396	744	1,917	2,971	5,141	8,570	1	...
Male	10,356	55	9	12	90	212	287	504	1,313	1,849	2,657	3,367	1	...
Female	9,947	45	10	15	42	73	109	240	604	1,122	2,484	5,203	–	...
Dominican	7,178	119	18	35	176	374	428	564	867	1,360	1,531	1,706	–	...
Male	3,721	65	9	23	125	276	293	347	520	762	752	549	–	...
Female	3,457	54	9	12	51	98	135	217	347	598	779	1,157	–	...
Mexican	148,306	2,725	476	812	4,785	7,853	10,384	14,945	22,734	27,693	27,718	28,177	4	...
Male	86,166	1,476	258	473	3,602	6,063	7,561	10,098	14,834	16,291	14,115	11,392	3	...
Female	62,140	1,249	218	339	1,183	1,790	2,823	4,847	7,900	11,402	13,603	16,785	1	...
Puerto Rican	33,536	418	91	115	605	1,365	1,952	2,932	4,928	6,443	7,445	7,241	1	...
Male	18,305	230	52	64	447	1,002	1,383	1,972	3,161	3,672	3,603	2,718	1	...
Female	15,231	188	39	51	158	363	569	960	1,767	2,771	3,842	4,523	–	...
South American	13,121	144	32	51	243	427	501	804	1,500	2,213	3,221	3,985	–	...
Male	6,627	77	18	26	173	330	340	488	881	1,241	1,581	1,472	–	...
Female	6,494	67	14	25	70	97	161	316	619	972	1,640	2,513	–	...
Other and unknown Hispanic	34,908	734	137	209	1,135	2,026	2,357	3,079	5,169	6,150	6,476	7,433	3	...
Male	19,440	419	71	112	828	1,456	1,629	2,040	3,193	3,540	3,240	2,909	3	...
Female	15,468	315	66	97	307	570	728	1,039	1,976	2,610	3,236	4,524	–	...

Table 3. Number of deaths and death rate by age, and age-adjusted death rate, by specified Hispanic origin and sex: United States, 2022—Con.

[Rates are per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Populations used for computing death rates for Hispanic total are postcensal estimates based on the 2010 census estimated as of July 1, 2022; populations used for computing death rates for Mexican, Puerto Rican, Cuban, Central American, South American, Dominican, and Other and unknown Hispanic are estimates based on the 2022 1-year American Community Survey adjusted to control totals. The control totals are 2010-based postcensal estimates for the United States for July 1, 2022; see Technical Notes. Hispanic origin is reported separately on the death certificate. People of Hispanic origin may be of any race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys; see Technical Notes]

Hispanic origin and sex	All ages	Age group											Age not stated	Age-adjusted rate ²
		Younger than 1 ¹	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older		
							Rate ³							
Hispanic	433.0	477.4	22.2	13.1	74.4	142.1	196.3	327.7	702.5	1,522.8	3,682.6	11,107.1	...	614.7
Male	481.0	514.1	23.4	14.6	109.3	209.7	270.9	432.1	905.9	1,900.2	4,368.6	12,379.5	...	748.8
Female	383.7	439.4	21.0	11.6	37.8	69.4	114.0	218.4	499.9	1,193.7	3,183.0	10,427.8	...	498.4
Central American	280.7	479.1	21.2	12.6	86.4	135.4	167.8	272.0	525.0	1,182.2	3,017.9	9,982.8	...	517.9
Male	316.4	539.5	21.3	14.0	132.9	213.5	237.8	373.1	681.5	1,493.6	3,793.3	10,109.6	...	632.0
Female	242.9	420.8	21.1	11.2	34.9	50.1	82.6	158.4	374.9	953.9	2,612.0	9,929.1	...	421.8
Cuban	833.6	430.8	*	10.3	49.1	81.8	118.4	221.2	574.2	1,424.0	3,827.0	12,658.8	...	587.4
Male	838.0	491.9	*	*	64.6	119.3	157.9	275.2	783.7	1,854.5	4,593.3	14,416.6	...	717.2
Female	829.0	374.1	*	*	32.4	42.8	71.4	156.7	363.1	1,030.0	3,247.5	11,733.0	...	478.2
Dominican	299.5	291.9	*	9.3	48.3	97.4	126.7	196.4	381.7	946.5	2,454.9	9,554.7	...	427.1
Male	327.5	268.0	*	12.3	68.7	144.7	181.3	275.0	540.0	1,280.9	3,252.6	10,828.4	...	547.3
Female	274.3	327.0	*	*	28.0	50.7	76.6	134.7	265.2	710.2	1,985.0	9,049.7	...	346.9
Mexican	396.4	496.7	20.4	12.3	70.4	136.1	197.0	337.6	753.0	1,623.5	3,865.6	12,105.9	...	649.5
Male	450.8	533.7	21.6	14.0	103.7	202.4	274.0	441.4	967.2	1,973.3	4,545.0	13,203.5	...	779.8
Female	339.6	459.2	19.2	10.5	35.6	64.5	112.4	226.6	531.8	1,295.4	3,346.5	11,459.3	...	531.2
Puerto Rican	567.9	483.7	24.9	11.9	67.2	157.0	239.0	419.7	827.4	1,762.6	4,082.7	11,644.3	...	688.1
Male	621.2	561.9	27.8	12.8	97.6	222.6	332.2	571.7	1,091.2	2,206.5	4,818.8	13,245.6	...	848.4
Female	514.8	413.2	21.8	11.0	35.8	86.6	142.1	271.4	577.6	1,391.6	3,571.1	10,855.6	...	555.7
South American	281.1	252.1	13.6	8.4	39.0	61.5	67.7	118.1	281.6	723.0	2,312.3	7,913.0	...	344.9
Male	293.2	272.6	*	8.4	54.8	96.2	92.9	150.3	346.2	960.5	2,941.1	8,496.4	...	419.1
Female	269.8	232.0	*	8.3	22.8	27.6	43.1	88.8	222.6	549.6	1,917.1	7,607.1	...	290.6
Other and unknown Hispanic	830.5	1,117.8	52.3	31.3	159.4	356.7	438.1	674.8	1,195.4	2,055.3	4,381.6	13,669.9	...	904.1
Male	913.4	1,380.1	55.1	31.9	226.3	481.3	576.7	875.4	1,501.8	2,581.3	5,095.9	14,083.8	...	1,083.6
Female	745.6	892.2	49.6	30.7	88.6	214.7	284.9	465.5	899.1	1,610.3	3,842.3	13,416.4	...	735.2

... Category not applicable.

— Quantity zero.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

¹Death rates for ages younger than 1 (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

²For method of computation, see Technical Notes.

³Data for age not stated included in "All ages" but not distributed among age groups.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 4. Life expectancy at selected ages, by Hispanic origin and race and sex: United States, 2022

[Life expectancies are preliminary estimates and may change slightly when updated data become available. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards]

Exact age	Non-Hispanic ^{2,4}																	
	Total ¹			Hispanic ^{2,3}			American Indian and Alaska Native			Asian			Black			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
0.....	77.5	74.8	80.2	80.0	77.0	82.8	67.8	64.5	71.3	84.4	82.3	86.3	72.8	69.1	76.5	77.5	75.1	80.1
1.....	76.9	74.3	79.6	79.4	76.4	82.2	67.4	64.2	70.8	83.7	81.6	85.6	72.6	68.9	76.3	76.9	74.4	79.4
5.....	73.0	70.4	75.7	75.4	72.5	78.3	63.6	60.4	67.0	79.8	77.7	81.6	68.8	65.1	72.4	72.9	70.5	75.5
10.....	68.0	65.4	70.7	70.5	67.5	73.3	58.7	55.5	62.1	74.8	72.7	76.6	63.8	60.1	67.5	68.0	65.6	70.5
15.....	63.1	60.5	65.8	65.5	62.6	68.4	53.8	50.6	57.2	69.8	67.8	71.7	58.9	55.2	62.6	63.0	60.6	65.6
20.....	58.3	55.7	60.9	60.7	57.8	63.5	49.2	46.1	52.6	64.9	62.9	66.7	54.3	50.7	57.7	58.2	55.8	60.7
25.....	53.5	51.1	56.0	56.0	53.2	58.6	44.8	41.8	48.0	60.1	58.0	61.8	49.8	46.4	53.0	53.4	51.1	55.8
30.....	48.9	46.6	51.3	51.3	48.7	53.8	40.7	38.0	43.6	55.2	53.2	56.9	45.3	42.2	48.3	48.8	46.6	51.0
35.....	44.3	42.1	46.5	46.7	44.2	49.0	36.9	34.3	39.6	50.4	48.4	52.0	40.9	38.0	43.7	44.2	42.1	46.3
40.....	39.8	37.7	41.9	42.1	39.8	44.2	33.4	31.0	35.8	45.6	43.7	47.1	36.6	33.8	39.3	39.7	37.7	41.7
45.....	35.3	33.4	37.3	37.6	35.4	39.5	29.8	27.8	31.9	40.8	38.9	42.2	32.5	29.8	34.9	35.2	33.3	37.1
50.....	30.9	29.1	32.8	33.1	31.0	34.9	26.7	24.9	28.4	36.1	34.3	37.4	28.4	25.9	30.6	30.8	29.1	32.6
55.....	26.7	25.0	28.4	28.7	26.8	30.3	23.5	21.9	25.0	31.4	29.8	32.7	24.4	22.1	26.4	26.6	25.0	28.2
60.....	22.7	21.1	24.2	24.5	22.7	25.9	20.6	19.3	21.8	26.9	25.4	28.0	20.8	18.6	22.6	22.6	21.1	24.0
65.....	18.9	17.5	20.2	20.5	19.0	21.7	17.8	16.8	18.7	22.5	21.2	23.4	17.4	15.5	19.0	18.8	17.5	20.0
70.....	15.3	14.2	16.3	16.7	15.4	17.7	14.9	14.2	15.5	18.4	17.2	19.1	14.4	12.8	15.6	15.2	14.1	16.2
75.....	12.0	11.0	12.8	13.2	12.1	13.9	12.4	11.9	12.6	14.5	13.6	15.0	11.5	10.2	12.4	11.8	10.9	12.6
80.....	8.9	8.1	9.5	10.0	9.1	10.4	9.9	9.6	10.0	10.8	10.1	11.1	8.9	7.8	9.5	8.8	8.0	9.4
85.....	6.4	5.8	6.8	7.2	6.5	7.4	7.9	7.7	7.8	7.7	7.2	7.8	6.6	5.8	7.0	6.3	5.7	6.7
90.....	4.4	3.9	4.6	5.0	4.4	5.0	6.2	6.1	6.0	5.1	4.8	5.1	4.8	4.2	5.0	4.3	3.8	4.5
95.....	3.0	2.7	3.1	3.4	3.0	3.3	4.8	4.9	4.6	3.4	3.2	3.2	3.4	3.1	3.5	2.9	2.6	3.0
100.....	2.1	1.9	2.2	2.4	2.2	2.3	3.9	4.0	3.6	2.3	2.2	2.1	2.5	2.3	2.5	2.0	1.8	2.1

¹Includes origins and races not shown separately; see Technical Notes in this report.²Based on death rates adjusted for Hispanic-origin and race misclassification on death certificates; see Technical Notes.³Includes persons of Hispanic origin of any race; see Technical Notes.⁴Only one race was reported on the death certificate; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2022

Hispanic origin and race and year	Both sexes	Male	Female
All origins and races ¹			
2022 ²	77.5	74.8	80.2
2021 ²	76.4	73.5	79.3
2020 ²	77.0	74.2	79.9
2019 ^{2,3}	78.8	76.3	81.4
2018 ²	78.7	76.2	81.2
2017 ²	78.6	76.1	81.1
2016 ²	78.7	76.2	81.1
2015 ²	78.7	76.3	81.1
2014 ²	78.9	76.5	81.3
2013 ²	78.8	76.4	81.2
2012 ²	78.8	76.4	81.2
2011 ²	78.7	76.3	81.1
2010 ²	78.7	76.2	81.0
2009 ²	78.5	76.0	80.9
2008 ²	78.2	75.6	80.6
2007.....	78.1	75.5	80.6
2006.....	77.8	75.2	80.3
2005.....	77.6	75.0	80.1
2004.....	77.6	75.0	80.1
2003.....	77.2	74.5	79.7
2002.....	77.0	74.4	79.6
2001.....	77.0	74.3	79.5
2000.....	76.8	74.1	79.3
1990.....	75.4	71.8	78.8
1980.....	73.7	70.0	77.4
1970.....	70.8	67.1	74.7
1960.....	69.7	66.6	73.1
1950.....	68.2	65.6	71.1
1940.....	62.9	60.8	65.2
Hispanic ^{4–6}			
2022 ²	80.0	77.0	82.8
2021 ²	77.8	74.6	81.1
2020 ²	77.9	74.6	81.3
2019 ^{2,3}	81.9	79.1	84.4
2018 ²	81.8	79.1	84.3
2017 ²	81.8	79.1	84.3
2016 ²	81.8	79.1	84.3
2015 ^{2,7}	81.9	79.3	84.3
2014 ²	82.1	79.4	84.5
2013 ²	81.9	79.2	84.2
2012 ²	81.9	79.3	84.3
2011 ²	81.8	79.2	84.2
2010 ²	81.7	78.8	84.3
2009 ²	81.1	78.4	83.5
2008 ²	80.8	78.0	83.3
2007.....	80.7	77.8	83.2
2006.....	80.3	77.5	82.9

Table 5. Life expectancy at birth, by Hispanic origin and race and sex: United States, 1940, 1950, 1960, 1970, 1980, 1990, and 2000–2022—Con.

Hispanic origin and race and year	Both sexes	Male	Female
Non-Hispanic, single race ^{6,8}			
American Indian and Alaska Native:			
2022 ²	67.8	64.5	71.3
2021 ²	65.6	62.2	69.2
2020 ²	67.1	63.8	70.7
2019 ^{2,3}	71.8	68.6	75.0
2018 ²	---	---	---
Asian:			
2022 ²	84.4	82.3	86.3
2021 ²	83.5	81.2	85.6
2020 ²	83.6	81.1	85.9
2019 ^{2,3}	85.6	83.5	87.4
2018 ²	---	---	---
Black:			
2022 ²	72.8	69.1	76.5
2021 ²	71.2	67.6	75.0
2020 ²	71.5	67.8	75.4
2019 ^{2,3}	74.8	71.3	78.1
2018 ²	74.7	71.3	78.0
White:			
2022 ²	77.5	75.1	80.1
2021 ²	76.7	74.0	79.5
2020 ²	77.4	74.8	80.1
2019 ^{2,3}	78.8	76.3	81.3
2018 ²	78.6	76.2	81.1

--- Data not available.

¹Includes origins and races not shown separately; see Technical Notes in this report.

²Life expectancies for 2008–2022 were calculated using newly revised methodology described in the Technical Notes.

³Life expectancies were updated using final data; therefore, data may differ from preliminary data previously published; see Technical Notes.

⁴Includes persons of Hispanic origin of any race. The Hispanic-origin category is consistent with 1997 OMB standards; see Technical Notes.

⁵Data by Hispanic origin available beginning with data year 2006.

⁶Based on death rate adjusted for race and Hispanic-origin misclassification; see Technical Notes.

⁷Life expectancies were revised using updated Medicare data; as a result, data may differ from those previously published; see Technical Notes.

⁸Only one race was reported on the death certificate. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards; see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group											Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
All causes													
2022.....	984.1	558.0	28.0	15.3	79.5	163.4	255.4	453.3	992.1	1,978.7	4,708.2	14,389.6	798.8
2021.....	1,043.8	558.8	25.0	14.3	88.9	180.8	287.9	531.0	1,117.1	2,151.3	5,119.4	15,743.3	879.7
2020.....	1,027.0	524.3	22.7	13.7	84.2	159.5	248.0	473.5	1,038.9	2,072.3	4,997.0	15,210.9	835.4
2019.....	869.7	553.0	23.3	13.4	69.7	128.8	199.2	392.4	883.3	1,764.6	4,308.3	13,228.6	715.2
2018.....	867.8	557.8	24.0	13.3	70.2	128.8	194.7	395.9	886.7	1,783.3	4,386.1	13,450.7	723.6
2017.....	863.8	567.0	24.3	13.6	74.0	132.8	195.2	401.5	885.8	1,790.9	4,472.6	13,573.6	731.9
2016.....	849.3	583.4	25.3	13.4	74.9	129.0	192.2	405.5	883.8	1,788.6	4,474.8	13,392.1	728.8
2015.....	844.0	589.6	24.9	13.2	69.5	116.7	180.1	404.0	875.3	1,796.8	4,579.2	13,673.9	733.1
2014.....	823.7	588.0	24.0	12.7	65.5	108.4	175.2	404.8	870.3	1,786.3	4,564.2	13,407.9	724.6
2013.....	821.5	594.7	25.5	13.0	64.8	106.1	172.0	406.1	860.0	1,802.1	4,648.1	13,660.4	731.9
2012.....	810.2	599.3	26.3	12.6	66.4	105.4	170.7	405.4	854.2	1,802.5	4,674.5	13,678.6	732.8
2011.....	807.3	600.1	26.3	13.2	67.7	104.7	172.0	409.8	849.4	1,846.2	4,753.0	13,779.3	741.3
2010.....	799.5	623.4	26.5	12.9	67.7	102.9	170.5	407.1	851.9	1,875.1	4,790.2	13,934.3	747.0
2009.....	794.5	659.7	27.4	13.8	69.8	104.4	180.0	418.1	856.7	1,888.7	4,820.2	13,660.1	749.6
2008.....	812.9	678.9	29.3	13.9	74.2	105.1	181.0	419.6	867.1	1,958.4	4,998.1	14,332.4	774.9
2007.....	804.6	702.5	29.4	15.2	78.8	107.2	186.0	420.3	866.7	1,976.0	4,987.1	14,160.9	775.3
2006.....	813.1	705.8	29.1	15.2	81.4	109.0	192.0	427.5	881.3	2,031.4	5,096.1	14,426.7	791.8
2005.....	828.4	710.2	29.9	16.3	80.7	106.8	194.9	431.9	898.5	2,109.7	5,251.8	14,982.4	815.0
2004.....	818.8	695.9	30.3	16.7	79.7	104.1	194.9	426.8	903.2	2,141.0	5,267.4	14,777.6	813.7
2003.....	843.9	704.9	31.8	16.9	81.1	105.2	202.6	433.1	937.3	2,235.0	5,451.3	15,401.4	843.5
2002.....	849.5	709.5	31.4	17.4	80.9	105.1	204.2	431.0	948.7	2,300.3	5,543.8	15,589.5	855.9
2001.....	848.0	687.0	33.4	17.2	80.2	105.6	203.5	426.7	972.5	2,344.2	5,573.7	15,432.6	858.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4	869.0
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6	875.6
Diseases of heart (I00–I09,I11,I13,I20–I51)													
2022.....	210.9	6.5	0.7	0.5	1.9	8.3	28.1	79.9	203.7	415.7	1,016.6	3,835.8	167.2
2021.....	209.6	7.4	0.8	0.5	2.2	9.1	29.4	84.9	208.7	416.6	1,044.2	4,078.7	173.8
2020.....	211.5	6.5	0.7	0.4	2.0	8.6	28.9	84.6	208.8	414.2	1,017.2	3,822.1	168.2
2019.....	200.8	7.1	0.8	0.4	2.0	7.6	25.2	76.2	190.4	388.8	991.2	3,798.3	161.5
2018.....	200.3	7.5	0.7	0.4	2.1	7.8	25.5	77.4	191.7	392.4	1,008.3	3,844.8	163.6
2017.....	198.8	7.7	0.8	0.4	2.1	8.1	25.4	77.1	190.7	392.9	1,028.4	3,882.9	165.0
2016.....	196.6	7.4	0.7	0.5	2.2	7.7	25.9	79.5	189.6	392.5	1,037.1	3,873.4	165.5
2015.....	197.2	7.3	0.9	0.5	2.3	8.0	25.6	79.3	188.1	389.5	1,071.6	3,986.5	168.5
2014.....	192.7	8.0	0.9	0.5	2.2	7.7	25.6	80.1	185.8	385.2	1,070.2	3,920.9	167.0
2013.....	193.3	7.8	1.1	0.4	2.1	7.6	25.6	80.3	184.6	390.3	1,095.1	4,013.9	169.8
2012.....	191.0	8.5	1.0	0.4	2.2	7.6	25.9	79.7	184.6	388.3	1,103.7	4,046.1	170.5
2011.....	191.5	7.7	1.0	0.5	2.3	7.9	26.2	80.7	183.2	399.0	1,134.7	4,111.6	173.7
2010.....	193.6	8.3	1.0	0.5	2.4	7.8	25.8	81.6	186.6	409.2	1,172.0	4,285.2	179.1
2009.....	195.4	9.6	0.9	0.5	2.4	7.8	26.7	82.3	190.0	422.8	1,210.8	4,316.9	182.8
2008.....	202.8	9.6	1.2	0.6	2.5	8.1	26.9	85.2	195.3	441.4	1,271.7	4,598.4	192.1
2007.....	204.5	10.2	1.1	0.6	2.5	8.1	27.7	85.2	197.8	454.8	1,308.6	4,668.1	196.1
2006.....	211.7	8.6	1.0	0.6	2.5	8.4	28.5	88.0	205.1	483.0	1,378.0	4,877.6	205.5
2005.....	220.7	8.9	0.9	0.6	2.6	8.3	29.2	89.7	212.8	512.3	1,458.5	5,188.3	216.8
2004.....	222.8	10.5	1.2	0.6	2.5	8.1	29.5	90.2	217.1	535.7	1,504.1	5,233.8	221.6
2003.....	236.1	11.0	1.2	0.6	2.7	8.3	30.8	92.4	232.3	579.8	1,607.7	5,570.7	236.3
2002.....	242.3	12.7	1.1	0.6	2.5	8.0	30.7	93.9	240.5	612.0	1,673.2	5,726.3	244.6
2001.....	245.7	11.9	1.5	0.7	2.5	8.0	29.6	92.4	248.9	632.6	1,723.0	5,784.1	249.5
2000.....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1	257.6
1999.....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0	266.5

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group										85 and older	Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84		
Malignant neoplasms (C00–C97)													
2022.....	182.5	1.1	1.8	2.0	3.2	8.0	25.6	82.5	249.8	529.4	979.3	1,573.5	142.3
2021.....	182.4	1.5	1.8	1.9	3.1	7.9	25.8	82.5	252.4	532.1	1,017.2	1,712.9	146.6
2020.....	182.8	1.4	2.0	1.9	3.1	7.8	25.5	85.7	260.0	539.1	990.0	1,538.1	144.1
2019.....	182.7	1.5	1.8	1.9	3.3	7.8	25.7	87.1	263.3	543.3	1,005.9	1,571.0	146.2
2018.....	183.2	1.3	2.0	2.1	3.2	8.1	25.8	89.6	269.6	554.4	1,031.5	1,577.7	149.1
2017.....	183.9	1.4	2.0	2.1	3.2	8.0	26.7	92.7	273.4	567.5	1,060.2	1,600.3	152.5
2016.....	185.1	1.7	2.4	2.1	3.3	8.5	26.9	96.5	280.6	578.3	1,081.7	1,620.3	155.8
2015.....	185.4	1.3	2.2	2.1	3.4	8.4	26.9	99.7	284.1	594.3	1,100.8	1,628.6	158.5
2014.....	185.6	1.3	2.0	2.1	3.6	8.3	27.8	103.2	287.6	603.1	1,125.9	1,632.9	161.2
2013.....	185.0	1.6	2.1	2.2	3.4	8.6	28.1	105.5	288.2	616.9	1,139.4	1,635.4	163.2
2012.....	185.6	1.6	2.4	2.2	3.6	8.7	28.0	108.5	293.2	632.2	1,161.7	1,658.9	166.5
2011.....	185.1	1.8	2.2	2.1	3.7	8.4	28.8	109.3	295.8	647.6	1,179.1	1,676.2	169.0
2010.....	186.2	1.6	2.1	2.2	3.7	8.8	28.8	111.6	300.1	666.1	1,202.2	1,729.5	172.8
2009.....	185.0	1.8	2.2	2.2	3.8	9.0	30.2	112.8	301.7	668.2	1,213.0	1,699.3	173.5
2008.....	186.0	1.7	2.4	2.2	3.8	8.8	30.1	113.4	304.7	688.4	1,230.9	1,724.6	176.4
2007.....	186.9	1.7	2.3	2.4	3.8	8.7	31.0	114.2	311.4	702.9	1,250.1	1,739.4	179.3
2006.....	187.6	1.9	2.4	2.2	3.8	9.3	32.2	116.3	317.7	716.3	1,259.2	1,748.3	181.8
2005.....	189.3	1.9	2.4	2.5	4.0	9.2	33.5	118.6	323.9	733.2	1,272.8	1,778.2	185.1
2004.....	189.2	1.8	2.5	2.5	4.1	9.3	33.6	119.0	330.8	746.8	1,278.6	1,767.4	186.8
2003.....	192.0	1.9	2.5	2.6	4.0	9.5	35.1	122.1	341.6	763.5	1,299.7	1,792.3	190.9
2002.....	193.7	1.9	2.6	2.6	4.2	9.8	36.0	124.1	349.7	787.2	1,308.8	1,812.4	194.3
2001.....	194.3	1.6	2.7	2.4	4.2	10.1	36.8	125.8	359.4	799.7	1,313.7	1,802.9	196.5
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4	199.6
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8	200.8
Accidents (unintentional injuries) (V01–X59,Y85–Y86)													
2022.....	68.1	36.8	8.7	4.0	33.1	72.7	84.6	77.6	80.8	68.5	127.8	417.4	64.0
2021.....	67.8	36.6	8.5	4.2	36.7	75.7	84.0	77.2	78.2	64.7	126.8	446.0	64.7
2020.....	61.0	32.0	7.4	3.8	35.5	68.0	73.7	68.9	68.2	57.2	113.3	383.6	57.6
2019.....	52.7	33.5	7.3	3.6	27.5	53.4	57.8	57.1	58.6	54.5	115.5	377.4	49.3
2018.....	51.1	30.4	7.7	3.5	28.0	53.9	54.9	55.4	56.0	52.3	111.3	368.6	48.0
2017.....	52.2	33.4	7.9	3.8	31.1	56.6	55.8	57.7	55.7	50.7	113.3	374.9	49.4
2016.....	49.9	30.7	7.9	4.0	31.9	53.7	51.8	54.6	52.7	49.1	110.7	365.7	47.4
2015.....	45.6	32.5	7.8	3.7	28.5	44.8	43.9	49.8	47.7	47.0	111.5	364.5	43.2
2014.....	42.6	29.4	7.6	3.6	26.8	39.8	39.6	47.4	44.9	45.1	108.7	349.1	40.5
2013.....	41.3	29.3	8.3	3.7	26.4	37.8	38.0	46.5	43.4	43.5	107.4	340.0	39.4
2012.....	40.7	29.6	8.4	3.8	27.1	37.5	37.1	46.1	41.0	44.0	107.8	336.9	39.1
2011.....	40.6	29.1	8.5	4.0	28.2	37.1	37.5	46.4	39.8	44.5	107.0	333.8	39.1
2010.....	39.1	28.1	8.6	4.0	28.3	35.5	36.0	43.7	38.4	43.3	106.1	328.4	38.0
2009.....	38.5	29.5	9.0	4.1	28.6	34.5	36.4	44.5	36.5	42.1	103.5	310.9	37.5
2008.....	40.1	31.8	9.1	4.6	32.5	36.3	38.1	45.8	37.4	43.9	105.7	318.3	39.2
2007.....	41.1	31.0	9.9	5.4	36.8	37.7	39.6	46.2	36.8	44.4	105.0	313.6	40.4
2006.....	40.8	28.4	10.1	5.6	37.9	38.0	40.5	45.5	35.8	43.8	104.7	299.2	40.2
2005.....	39.9	27.0	10.5	5.9	37.1	35.7	38.9	43.2	35.4	45.7	106.0	303.5	39.5
2004.....	38.3	26.2	10.4	6.5	36.8	33.2	37.6	40.7	32.9	43.5	103.6	295.8	38.1
2003.....	37.7	23.8	11.0	6.4	36.9	32.0	38.0	38.8	32.7	43.7	101.6	294.3	37.6
2002.....	37.1	23.9	10.6	6.6	37.7	31.9	37.4	36.7	31.3	44.0	101.1	289.6	37.1
2001.....	35.6	24.3	11.2	6.9	35.8	30.0	35.4	33.9	30.5	42.6	100.7	282.2	35.7
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5	34.9
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4	35.3
COVID-19 (U07.1) ⁴													
2022.....	56.0	3.8	0.7	0.3	1.0	3.6	8.8	23.9	57.6	124.5	292.2	818.2	44.5
2021.....	125.6	2.6	0.4	0.3	3.3	13.5	36.9	90.6	172.2	305.1	609.7	1,354.4	104.1
2020.....	106.5	0.9	*	0.1	1.2	4.9	14.4	42.0	99.3	234.3	589.8	1,645.0	85.0

See footnotes at end of table.

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group										Age- adjusted rate ³	
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84		85 and older
Cerebrovascular diseases (I60–I69)													
2022.....	49.6	2.4	0.3	0.2	0.4	1.3	4.9	13.8	33.7	84.0	265.0	1,043.9	39.5
2021.....	49.1	2.7	0.4	0.2	0.4	1.4	5.0	14.1	34.2	84.1	274.8	1,111.1	41.1
2020.....	48.6	2.8	0.4	0.2	0.4	1.3	4.8	14.1	33.4	81.0	262.9	1,017.9	38.8
2019.....	45.7	2.7	0.3	0.2	0.4	1.3	4.2	12.6	30.5	76.4	254.2	977.3	37.0
2018.....	45.2	2.5	0.3	0.2	0.3	1.2	4.1	12.3	30.3	76.8	256.0	984.3	37.1
2017.....	44.9	2.5	0.4	0.2	0.4	1.3	4.4	12.3	30.3	76.4	263.1	993.5	37.6
2016.....	44.0	3.1	0.3	0.2	0.3	1.3	4.6	12.5	29.7	76.0	265.5	972.9	37.3
2015.....	43.7	2.2	0.3	0.2	0.4	1.3	4.4	12.3	29.6	75.5	273.0	975.8	37.6
2014.....	41.7	2.4	0.2	0.2	0.4	1.3	4.3	12.3	29.3	74.5	265.7	929.7	36.5
2013.....	40.8	2.7	0.2	0.2	0.3	1.2	4.2	12.4	28.9	74.2	268.9	906.0	36.2
2012.....	40.9	2.6	0.3	0.2	0.4	1.3	4.3	12.8	28.7	75.7	272.2	931.2	36.9
2011.....	41.4	3.4	0.3	0.2	0.4	1.3	4.2	12.8	29.4	78.2	285.4	943.7	37.9
2010.....	41.9	3.3	0.3	0.2	0.4	1.3	4.6	13.1	29.3	81.7	288.3	993.8	39.1
2009.....	42.0	3.7	0.3	0.2	0.4	1.3	4.6	13.7	29.7	82.8	294.9	992.2	39.6
2008.....	44.1	3.4	0.4	0.2	0.4	1.3	4.8	13.7	30.6	87.3	313.3	1,071.0	42.1
2007.....	45.1	3.2	0.3	0.2	0.5	1.3	5.0	14.5	31.7	91.4	320.8	1,110.7	43.5
2006.....	46.0	3.5	0.3	0.2	0.5	1.3	5.1	14.6	32.9	94.9	333.9	1,131.7	44.8
2005.....	48.6	3.1	0.4	0.2	0.5	1.4	5.2	15.0	32.7	99.8	358.4	1,239.7	48.0
2004.....	51.3	3.2	0.3	0.2	0.5	1.4	5.4	14.8	34.0	106.6	385.6	1,331.9	51.2
2003.....	54.4	2.5	0.3	0.2	0.5	1.5	5.6	15.0	35.5	111.9	409.8	1,446.0	54.6
2002.....	56.6	3.0	0.3	0.2	0.4	1.4	5.4	15.1	37.1	119.6	430.0	1,520.1	57.2
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.0	38.3	122.9	443.3	1,532.0	58.4
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2	60.9
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8	61.6
Chronic lower respiratory diseases (J40–J47)													
2022.....	44.2	*	0.3	0.3	0.4	0.8	1.7	7.4	40.7	114.6	283.9	575.8	34.3
2021.....	42.9	*	0.2	0.2	0.4	0.8	1.7	7.8	41.2	112.7	285.9	600.6	34.7
2020.....	46.3	*	*	0.4	0.5	0.9	1.9	8.8	44.4	118.5	300.1	612.6	36.4
2019.....	47.8	*	0.2	0.4	0.4	0.8	1.6	8.8	44.2	124.6	318.0	654.3	38.2
2018.....	48.7	*	0.3	0.3	0.4	0.7	1.6	9.1	44.5	128.6	335.7	682.4	39.7
2017.....	49.2	*	0.2	0.3	0.4	0.7	1.7	9.4	44.4	133.8	347.6	700.6	40.9
2016.....	47.8	0.7	0.3	0.3	0.5	0.8	1.7	10.1	43.0	134.1	347.2	676.9	40.6
2015.....	48.2	0.7	0.3	0.4	0.5	0.7	1.7	10.1	42.7	136.6	357.9	705.1	41.6
2014.....	46.1	*	0.3	0.3	0.4	0.8	1.9	10.1	41.2	134.9	349.0	670.5	40.5
2013.....	47.2	0.6	0.4	0.4	0.4	0.7	1.9	10.6	40.5	141.2	367.0	699.3	42.1
2012.....	45.7	0.5	0.3	0.3	0.3	0.7	1.8	10.2	39.4	140.0	364.0	687.8	41.5
2011.....	45.9	0.8	0.3	0.3	0.4	0.6	1.8	10.4	39.5	144.3	374.9	697.9	42.5
2010.....	44.7	0.9	0.3	0.3	0.3	0.7	1.7	9.9	39.0	146.3	369.9	690.7	42.2
2009.....	44.8	0.7	0.4	0.3	0.4	0.7	1.8	10.4	40.0	147.5	376.4	684.9	42.7
2008.....	46.4	0.8	0.3	0.3	0.4	0.6	1.9	9.9	41.1	155.9	395.4	722.7	44.7
2007.....	42.5	1.0	0.4	0.3	0.3	0.7	1.9	9.5	38.6	145.5	367.1	652.0	41.4
2006.....	41.8	0.7	0.3	0.3	0.4	0.6	1.9	9.1	38.8	147.0	362.0	641.3	41.0
2005.....	44.3	0.8	0.4	0.3	0.3	0.7	2.0	9.4	41.6	158.4	385.0	691.9	43.9
2004.....	41.7	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.1	152.1	366.2	643.2	41.6
2003.....	43.6	0.8	0.4	0.3	0.5	0.7	2.2	8.7	43.1	161.7	382.2	670.2	43.7
2002.....	43.4	1.0	0.4	0.3	0.5	0.8	2.3	8.7	42.2	162.0	385.8	670.3	43.9
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.4	44.5	167.3	379.3	658.3	43.9
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6	44.2
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0	45.4

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group										85 and older	Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84		
Alzheimer disease (G30)													
2022.....	36.0	*	*	*	*	*	*	0.3	3.5	26.7	206.2	1,131.5	28.9
2021.....	36.0	*	*	*	*	*	*	0.3	3.2	26.4	214.3	1,243.6	31.0
2020.....	40.7	*	*	*	*	*	*	0.2	3.3	28.6	229.3	1,287.3	32.4
2019.....	37.0	*	*	*	*	*	*	0.3	3.0	24.9	210.2	1,191.3	29.8
2018.....	37.3	*	*	*	*	*	*	0.3	2.9	24.7	213.9	1,225.3	30.5
2017.....	37.3	*	*	*	*	*	*	0.2	2.8	24.5	219.7	1,244.7	31.0
2016.....	35.9	*	*	*	*	*	*	0.2	2.7	23.6	214.1	1,216.9	30.3
2015.....	34.4	*	*	*	*	*	*	0.2	2.4	22.4	211.9	1,174.2	29.4
2014.....	29.3	*	*	*	*	*	*	0.2	2.1	19.6	185.6	1,006.8	25.4
2013.....	26.8	*	*	*	*	*	*	0.2	2.2	18.1	171.6	929.5	23.5
2012.....	26.6	*	*	*	*	*	*	0.2	2.2	17.9	175.4	936.1	23.8
2011.....	27.3	*	*	*	*	*	*	0.2	2.2	19.2	183.9	967.1	24.7
2010.....	27.0	*	*	*	*	*	*	0.3	2.1	19.8	184.5	987.1	25.1
2009.....	25.8	*	*	*	*	*	*	0.2	2.0	19.4	179.1	945.3	24.2
2008.....	27.1	*	*	*	*	*	*	0.2	2.2	21.1	192.5	1,002.2	25.8
2007.....	24.8	*	*	*	*	*	*	0.2	2.2	20.2	175.8	928.7	23.8
2006.....	24.3	*	*	*	*	*	*	0.2	2.1	19.9	175.0	923.4	23.7
2005.....	24.2	*	*	*	*	*	*	0.2	2.1	20.2	177.0	935.5	24.0
2004.....	22.5	*	*	*	*	*	*	0.2	1.8	19.5	168.5	875.3	22.6
2003.....	21.9	*	*	*	*	*	*	0.2	2.0	20.7	164.1	846.8	22.1
2002.....	20.5	*	*	*	*	*	*	0.1	1.9	19.6	157.7	790.9	20.8
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.6	147.2	725.4	19.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7	18.1
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3	16.5
Diabetes mellitus (E10–E14)													
2022.....	30.4	*	*	0.1	0.7	2.6	6.6	18.2	41.4	79.8	150.6	287.3	24.1
2021.....	31.1	*	*	0.1	0.8	2.8	6.8	18.7	43.5	83.0	160.4	309.9	25.4
2020.....	31.0	*	*	0.1	0.7	2.5	6.9	18.7	42.5	83.6	154.7	293.4	24.8
2019.....	26.7	*	*	0.1	0.6	1.9	5.3	15.5	36.5	73.3	138.9	259.5	21.6
2018.....	26.0	*	*	0.1	0.6	1.8	5.5	15.4	35.3	72.1	137.5	260.4	21.4
2017.....	25.7	*	*	0.1	0.6	1.8	5.2	15.1	35.5	71.9	140.8	262.4	21.5
2016.....	24.8	*	*	0.1	0.5	1.8	5.1	14.6	34.4	69.9	137.9	263.6	21.0
2015.....	24.7	*	*	0.1	0.4	1.8	4.9	14.4	34.7	70.6	143.0	267.0	21.3
2014.....	24.0	*	*	0.1	0.4	1.6	4.9	13.9	33.3	69.0	141.8	268.6	20.9
2013.....	23.9	*	*	0.1	0.4	1.6	4.8	13.5	33.2	68.5	145.7	279.5	21.2
2012.....	23.6	*	*	0.1	0.4	1.5	4.6	13.0	32.5	69.7	145.8	285.7	21.2
2011.....	23.7	*	*	0.1	0.4	1.6	4.5	13.4	33.3	72.0	148.8	289.5	21.6
2010.....	22.4	*	*	0.1	0.4	1.5	4.4	12.5	32.0	67.6	144.1	285.5	20.8
2009.....	22.4	*	*	0.1	0.4	1.5	4.5	12.8	32.1	69.6	145.8	282.6	21.0
2008.....	23.2	*	*	0.1	0.5	1.4	4.4	12.6	33.3	74.7	153.2	298.9	22.0
2007.....	23.7	*	*	0.1	0.4	1.5	4.6	13.1	34.1	76.7	161.9	302.2	22.8
2006.....	24.3	*	*	0.1	0.4	1.7	4.8	13.1	35.8	80.6	166.2	310.4	23.6
2005.....	25.4	*	*	0.1	0.5	1.6	4.7	13.4	36.9	85.7	177.0	338.8	24.9
2004.....	25.0	*	*	0.1	0.4	1.5	4.6	13.4	36.8	86.2	176.6	328.2	24.7
2003.....	25.6	*	*	0.1	0.4	1.7	4.6	13.9	38.3	90.0	180.7	335.1	25.5
2002.....	25.5	*	*	0.1	0.4	1.6	4.8	13.7	37.5	90.9	182.4	337.0	25.6
2001.....	25.0	*	*	0.1	0.4	1.5	4.3	13.6	38.1	91.0	181.1	328.6	25.4
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7	25.0
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2	25.0

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10) and year	All ages ¹	Age group										85 and older	Age- adjusted rate ³
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84		
Nephritis, nephrotic syndrome and nephrosis (N00-N07, N17-N19,N25-N27)													
2022.....	17.4	1.0	*	*	0.1	0.8	2.4	6.6	15.8	38.3	97.0	264.2	13.8
2021.....	16.4	1.3	*	*	0.1	0.8	2.2	6.2	14.9	36.4	96.1	271.1	13.6
2020.....	15.9	1.2	*	*	0.2	0.7	2.0	5.8	14.7	35.2	89.3	248.1	12.7
2019.....	15.7	1.6	*	*	0.2	0.7	1.8	5.6	13.8	34.7	92.2	250.9	12.7
2018.....	15.7	2.0	*	*	0.1	0.6	1.8	5.4	13.6	35.6	94.3	257.9	12.9
2017.....	15.5	2.0	*	*	0.1	0.6	1.7	5.2	13.5	34.7	95.8	267.1	13.0
2016.....	15.5	1.6	*	*	0.1	0.6	1.8	5.0	13.6	34.6	98.1	270.1	13.1
2015.....	15.5	2.1	*	*	0.1	0.6	1.7	4.9	13.3	35.1	99.7	281.8	13.4
2014.....	15.1	2.3	*	*	0.2	0.5	1.7	4.7	12.6	34.3	98.6	282.4	13.2
2013.....	14.9	2.2	*	*	0.1	0.6	1.5	4.6	12.6	33.8	99.0	285.4	13.2
2012.....	14.5	2.1	*	*	0.2	0.5	1.6	4.7	12.3	33.3	99.9	280.0	13.1
2011.....	14.6	1.9	*	*	0.2	0.5	1.6	4.4	12.5	34.2	101.4	292.1	13.4
2010.....	16.3	2.7	*	0.1	0.2	0.6	1.8	4.9	13.9	39.3	115.7	333.8	15.3
2009.....	16.0	2.8	*	*	0.2	0.7	2.0	5.2	13.5	38.7	115.1	321.4	15.1
2008.....	15.9	3.5	*	*	0.2	0.6	1.8	5.0	14.1	39.9	113.3	325.6	15.1
2007.....	15.4	3.5	0.1	0.1	0.2	0.7	1.8	5.1	13.4	39.4	112.4	317.9	14.9
2006.....	15.2	4.0	*	*	0.2	0.7	1.8	5.2	13.7	38.8	111.0	316.2	14.8
2005.....	14.9	4.0	*	0.1	0.2	0.7	1.7	4.8	13.5	38.8	110.2	313.1	14.7
2004.....	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.5	38.1	108.2	306.4	14.5
2003.....	14.6	4.6	*	0.1	0.2	0.7	1.8	4.9	13.6	39.7	109.3	309.3	14.7
2002.....	14.2	4.4	*	0.1	0.2	0.7	1.7	4.7	12.9	39.0	108.9	303.4	14.4
2001.....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.1	40.0	104.0	293.8	14.1
2000.....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8	13.5
1999.....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9	13.0
Chronic liver disease and cirrhosis (K70,K73-K74)													
2022.....	16.4	*	*	*	0.1	3.9	12.6	23.3	39.2	39.0	36.7	30.2	13.8
2021.....	17.0	*	*	*	0.1	4.0	13.4	25.8	41.3	38.4	36.4	31.0	14.5
2020.....	15.7	*	*	*	0.1	3.5	11.7	23.5	38.1	36.5	34.5	27.0	13.3
2019.....	13.5	*	*	*	0.1	2.4	8.2	19.8	33.9	33.1	32.2	26.5	11.3
2018.....	13.1	*	*	*	0.1	2.2	7.5	19.6	33.0	32.5	32.5	25.5	11.1
2017.....	12.8	*	*	*	0.1	2.0	7.3	19.6	32.7	31.7	31.3	26.8	10.9
2016.....	12.5	*	*	*	0.1	2.1	7.0	19.5	32.4	30.7	31.9	24.5	10.7
2015.....	12.5	*	*	*	0.1	1.9	7.0	20.5	32.5	30.5	31.9	25.1	10.8
2014.....	12.0	*	*	*	0.1	1.7	6.4	19.9	31.9	29.6	30.4	23.4	10.4
2013.....	11.5	*	*	*	0.1	1.6	6.2	20.1	30.4	28.1	29.9	23.0	10.2
2012.....	11.1	*	*	*	0.1	1.4	6.1	20.1	29.1	27.6	29.3	21.4	9.9
2011.....	10.8	*	*	*	0.1	1.2	6.0	19.8	28.2	26.3	29.3	22.1	9.7
2010.....	10.3	*	*	*	0.1	1.2	5.9	19.2	26.8	26.3	27.7	21.8	9.4
2009.....	10.0	*	*	*	0.1	1.1	6.0	18.7	25.9	25.4	27.2	21.1	9.1
2008.....	9.9	*	*	*	0.1	1.1	6.1	18.5	25.0	26.3	28.0	21.9	9.2
2007.....	9.7	*	*	*	0.1	1.0	6.0	18.7	24.2	26.2	28.2	21.7	9.1
2006.....	9.2	*	*	*	0.1	0.8	5.9	17.8	22.6	25.6	28.9	21.1	8.8
2005.....	9.3	*	*	*	0.1	0.8	6.2	17.7	23.3	26.8	28.9	21.3	8.9
2004.....	9.2	*	*	*	*	0.8	6.4	18.0	22.4	27.4	28.7	21.1	9.0
2003.....	9.5	*	*	*	*	0.9	6.8	18.3	22.9	29.2	29.9	21.2	9.3
2002.....	9.5	*	*	*	0.1	1.0	7.1	18.0	22.8	29.3	31.3	22.5	9.4
2001.....	9.5	*	*	*	0.1	1.0	7.4	18.4	22.9	29.8	30.2	22.7	9.5
2000.....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1	9.5
1999.....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2	9.6

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes]

Cause of death (based on ICD-10) and year	All ages ¹	Age group											Age- adjusted rate ³
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)													
2022.....	14.8	...	...	1.2	13.6	19.0	18.7	19.2	18.7	16.0	20.3	23.0	14.2
2021.....	14.5	...	...	1.5	15.2	19.5	18.1	18.2	17.0	15.3	19.6	22.4	14.1
2020.....	14.0	...	...	1.5	14.2	18.4	17.4	18.0	16.9	14.5	18.4	20.9	13.5
2019.....	14.5	...	...	1.3	13.9	17.5	18.1	19.6	19.4	15.5	18.6	20.1	13.9
2018.....	14.8	...	...	1.5	14.5	17.6	18.2	20.0	20.2	16.3	18.7	19.1	14.2
2017.....	14.5	...	...	1.3	14.5	17.5	17.9	20.2	19.0	15.6	18.0	20.1	14.0
2016.....	13.9	...	...	1.1	13.2	16.5	17.4	19.7	18.7	15.4	18.2	19.0	13.5
2015.....	13.7	...	...	1.0	12.5	15.7	17.1	20.3	18.9	15.2	17.9	19.4	13.3
2014.....	13.4	...	...	1.0	11.6	15.1	16.6	20.2	18.8	15.6	17.5	19.3	13.0
2013.....	13.0	...	...	1.0	11.1	14.8	16.2	19.7	18.1	15.0	17.1	18.6	12.6
2012.....	12.9	...	...	0.8	11.1	14.7	16.7	20.0	18.0	14.0	16.8	17.8	12.6
2011.....	12.7	...	...	0.7	11.0	14.6	16.2	19.8	17.1	14.1	16.5	16.9	12.3
2010.....	12.4	...	...	0.7	10.5	14.0	16.0	19.6	17.5	13.7	15.7	17.6	12.1
2009.....	12.0	...	...	0.6	10.0	13.1	16.1	19.2	16.4	13.7	15.8	16.4	11.8
2008.....	11.8	...	...	0.5	9.9	13.2	15.9	18.6	16.0	13.6	16.1	16.4	11.6
2007.....	11.5	...	...	0.5	9.6	13.3	15.7	17.7	15.3	12.4	16.2	17.0	11.3
2006.....	11.2	...	...	0.5	9.8	12.7	15.2	17.2	14.4	12.4	15.8	17.3	11.0
2005.....	11.0	...	...	0.7	9.9	12.7	15.1	16.5	13.7	12.4	16.8	18.3	10.9
2004.....	11.1	...	...	0.7	10.3	12.9	15.2	16.6	13.7	12.2	16.3	17.6	11.0
2003.....	10.9	...	...	0.6	9.6	12.9	15.0	15.9	13.7	12.6	16.4	17.9	10.8
2002.....	11.0	...	...	0.6	9.8	12.8	15.3	15.8	13.5	13.4	17.7	18.9	10.9
2001.....	10.7	...	...	0.7	9.9	12.8	14.7	15.1	13.2	13.2	17.4	17.8	10.7
2000.....	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6	10.4
1999.....	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3	10.5
Influenza and pneumonia (J09-J18)													
2022.....	14.1	4.2	0.9	0.3	0.4	1.1	2.3	4.7	12.0	27.8	72.9	244.2	11.3
2021.....	12.6	3.5	0.3	0.1	0.3	0.9	1.9	4.4	11.3	25.3	69.9	231.8	10.5
2020.....	16.3	3.3	0.5	0.3	0.4	1.3	2.7	6.2	14.8	32.1	84.1	273.6	13.0
2019.....	15.2	4.1	0.8	0.3	0.4	1.0	2.3	5.1	12.5	27.3	77.2	294.7	12.3
2018.....	18.1	4.6	0.8	0.3	0.5	1.0	2.3	5.6	13.9	31.7	94.2	377.6	14.9
2017.....	17.1	4.0	0.7	0.3	0.4	0.9	1.9	4.8	12.0	29.6	93.8	375.3	14.3
2016.....	15.9	4.2	0.6	0.2	0.4	1.0	2.2	5.0	12.1	28.5	88.5	340.3	13.5
2015.....	17.8	4.4	0.6	0.2	0.4	0.9	1.7	4.7	11.3	29.5	101.6	421.4	15.2
2014.....	17.3	4.7	0.7	0.2	0.5	1.3	2.8	6.3	13.4	29.8	96.4	385.9	15.1
2013.....	18.0	4.5	0.6	0.3	0.4	1.0	2.2	5.1	12.2	29.5	103.7	441.0	15.9
2012.....	16.1	4.0	0.6	0.2	0.3	0.8	1.7	4.1	10.2	26.1	98.2	408.4	14.4
2011.....	17.3	5.2	0.7	0.3	0.5	1.2	2.1	5.0	11.0	28.9	104.0	439.2	15.7
2010.....	16.2	4.9	0.6	0.2	0.4	0.9	1.9	4.3	9.9	27.9	102.4	426.2	15.1
2009.....	17.5	6.3	0.9	0.6	1.0	2.0	3.2	6.5	11.7	29.5	107.0	433.8	16.5
2008.....	18.5	5.5	0.9	0.2	0.5	0.9	2.1	5.1	10.9	30.5	118.6	512.3	17.6
2007.....	17.5	5.4	0.7	0.3	0.4	0.8	1.8	4.3	9.5	28.2	113.5	506.7	16.8
2006.....	18.9	6.5	0.8	0.2	0.4	0.9	1.9	4.6	9.9	31.6	127.3	547.0	18.4
2005.....	21.3	6.6	0.7	0.3	0.4	0.9	2.1	5.1	11.2	35.1	142.0	644.9	21.0
2004.....	20.4	6.8	0.8	0.2	0.4	0.8	2.0	4.6	10.8	34.2	139.1	622.8	20.4
2003.....	22.5	8.1	1.0	0.4	0.5	1.0	2.2	5.2	11.2	36.9	150.8	703.0	22.6
2002.....	22.8	6.7	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.2	156.6	732.4	23.2
2001.....	21.8	7.5	0.7	0.2	0.5	0.9	2.2	4.6	10.8	36.2	148.3	700.1	22.2
2000.....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1	23.7
1999.....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8	23.5

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group										85 and older	Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84		
Essential hypertension and hypertensive renal disease (I10,I12,I15)													
2022.....	13.0	*	*	*	0.1	0.4	1.6	4.5	11.8	24.6	61.6	254.0	10.3
2021.....	12.9	*	*	*	0.1	0.4	1.5	4.8	12.1	24.9	63.2	270.8	10.7
2020.....	12.7	*	*	*	0.1	0.3	1.4	4.5	11.8	23.9	60.3	250.3	10.1
2019.....	11.1	*	*	*	*	0.3	1.2	3.6	9.8	20.9	54.3	227.1	8.9
2018.....	11.0	*	*	*	*	0.3	1.2	3.7	9.7	20.2	54.4	229.4	8.9
2017.....	10.8	*	*	*	*	0.3	1.3	3.9	9.5	19.7	55.8	231.4	9.0
2016.....	10.3	*	*	*	*	0.3	1.1	3.8	9.0	19.0	53.7	222.3	8.6
2015.....	10.0	*	*	*	*	0.3	1.2	3.4	8.8	18.1	54.1	222.7	8.5
2014.....	9.5	*	*	*	0.0	0.2	1.1	3.3	8.4	16.9	51.3	217.0	8.2
2013.....	9.7	*	*	*	0.1	0.3	1.0	3.5	8.0	17.3	53.7	231.6	8.5
2012.....	9.3	*	*	*	*	0.2	0.8	3.0	7.8	16.1	51.7	230.7	8.2
2011.....	8.9	*	*	*	*	0.2	1.0	3.1	7.0	16.6	51.4	222.7	8.1
2010.....	8.6	*	*	*	0.0	0.3	1.0	3.1	7.3	16.7	51.8	212.0	8.0
2009.....	8.4	*	*	*	0.1	0.3	1.0	3.1	7.1	16.3	51.0	208.0	7.8
2008.....	8.5	*	*	*	0.1	0.3	1.0	3.0	7.2	16.5	51.9	215.3	8.0
2007.....	8.0	*	*	*	0.1	0.2	0.9	2.8	6.4	15.9	49.2	209.1	7.6
2006.....	8.0	*	*	*	0.0	0.3	0.9	3.0	6.8	16.5	50.8	206.1	7.7
2005.....	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.5	55.5	228.0	8.3
2004.....	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	16.9	52.5	212.2	7.9
2003.....	7.6	*	*	*	0.1	0.2	0.8	2.5	6.3	16.8	51.6	199.4	7.6
2002.....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	15.9	48.1	189.6	7.1
2001.....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.4	47.6	175.6	6.9
2000.....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9	6.5
1999.....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1	6.2
Septicemia (A40–A41)													
2022.....	12.7	3.3	0.4	0.1	0.3	1.0	2.3	6.2	14.7	31.4	66.9	145.4	10.1
2021.....	12.4	3.2	0.2	0.1	0.2	0.9	2.6	6.0	15.1	31.1	67.4	153.3	10.2
2020.....	12.2	3.1	0.3	0.1	0.2	0.8	2.3	6.2	14.7	29.7	63.5	143.1	9.7
2019.....	11.7	3.5	0.3	0.2	0.3	0.8	1.9	5.3	13.4	28.0	64.5	150.2	9.5
2018.....	12.4	3.9	0.3	0.1	0.2	0.9	2.0	5.7	14.1	30.0	69.4	167.4	10.2
2017.....	12.6	3.7	0.3	0.1	0.3	0.9	2.1	5.8	13.9	30.9	72.4	173.3	10.6
2016.....	12.6	4.8	0.4	0.2	0.3	0.9	2.2	5.8	14.3	30.6	73.9	174.2	10.7
2015.....	12.7	4.5	0.3	0.2	0.3	0.9	2.0	5.9	14.1	31.4	75.3	185.8	11.0
2014.....	12.2	4.0	0.3	0.2	0.3	0.8	2.1	5.8	14.2	31.1	73.1	176.9	10.7
2013.....	12.1	3.9	0.3	0.1	0.3	0.8	2.0	5.6	13.6	30.5	76.4	179.6	10.7
2012.....	11.4	4.5	0.4	0.1	0.3	0.8	1.9	5.3	12.9	29.2	73.9	173.4	10.3
2011.....	11.5	4.5	0.4	0.2	0.3	0.8	2.0	5.5	13.0	29.5	74.4	179.7	10.5
2010.....	11.3	5.5	0.4	0.2	0.3	0.8	1.9	5.2	12.6	30.1	76.0	179.0	10.6
2009.....	11.6	5.5	0.4	0.2	0.3	0.9	2.2	5.4	13.1	31.4	79.2	182.4	11.0
2008.....	11.8	7.0	0.6	0.2	0.3	0.9	2.1	5.7	13.3	31.4	82.0	189.8	11.3
2007.....	11.6	6.8	0.5	0.2	0.4	0.7	2.1	5.5	12.8	32.2	79.5	190.8	11.2
2006.....	11.5	6.7	0.6	0.2	0.3	0.7	2.0	5.2	12.6	31.6	82.1	193.0	11.2
2005.....	11.6	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.8	32.2	81.3	203.4	11.4
2004.....	11.4	6.8	0.5	0.2	0.3	0.8	1.9	5.4	12.8	32.1	81.5	199.6	11.3
2003.....	11.7	7.0	0.5	0.2	0.4	0.8	2.1	5.3	13.0	32.3	84.8	213.7	11.8
2002.....	11.8	7.5	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.5	86.3	213.4	11.9
2001.....	11.3	7.8	0.7	0.2	0.3	0.7	1.8	5.0	12.4	32.6	82.2	210.3	11.5
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7	11.3
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7	11.3

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group											Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Parkinson disease (G20–G21)													
2022.....	12.0	*	*	*	*	*	*	0.2	2.1	19.0	102.1	225.4	9.5
2021.....	11.6	*	*	*	*	*	*	0.2	2.0	18.6	102.5	246.0	9.8
2020.....	12.2	*	*	*	*	*	*	0.2	2.1	19.0	105.3	237.4	9.9
2019.....	10.8	*	*	*	*	*	*	0.2	1.9	16.8	93.3	215.9	8.8
2018.....	10.3	*	*	*	*	*	*	0.2	1.8	16.2	91.0	214.6	8.7
2017.....	9.8	*	*	*	*	*	*	0.1	1.7	15.7	89.6	206.3	8.4
2016.....	9.2	*	*	*	*	*	*	0.2	1.6	14.3	85.5	198.6	8.0
2015.....	8.7	*	*	*	*	*	*	0.2	1.5	13.8	82.4	190.6	7.7
2014.....	8.2	*	*	*	*	*	*	0.2	1.4	13.0	79.2	182.0	7.4
2013.....	8.0	*	*	*	*	*	*	0.2	1.5	12.7	78.5	178.2	7.3
2012.....	7.6	*	*	*	*	*	*	0.1	1.4	12.3	76.2	172.3	7.0
2011.....	7.4	*	*	*	*	*	*	0.1	1.3	12.8	76.0	168.1	7.0
2010.....	7.1	*	*	*	*	*	*	0.2	1.3	11.8	74.8	165.9	6.8
2009.....	6.7	*	*	*	*	*	*	0.2	1.3	11.2	70.8	157.0	6.5
2008.....	6.7	*	*	*	*	*	*	0.2	1.2	12.3	71.2	157.4	6.6
2007.....	6.7	*	*	*	*	*	*	0.1	1.2	11.7	71.5	157.0	6.5
2006.....	6.6	*	*	*	*	*	*	0.2	1.2	12.0	69.5	157.6	6.5
2005.....	6.6	*	*	*	*	*	*	0.2	1.4	12.8	71.1	156.0	6.6
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	11.9	67.4	145.1	6.2
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.6	67.6	145.8	6.3
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.1	63.8	142.2	6.0
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.5	137.0	5.9
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9	5.7
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4	5.4
Dementia-related causes ⁶													
2022.....	87.9	*	*	0.1	*	*	0.1	0.9	9.4	67.4	476.2	2,809.6	70.4
2021.....	84.3	*	*	0.1	*	0.0	0.2	0.9	8.6	65.0	477.6	2,949.3	72.4
2020.....	92.3	*	*	*	*	0.0	0.1	0.8	8.7	68.1	495.1	2,946.9	73.3
2019.....	82.8	*	*	0.1	*	0.1	0.1	0.9	8.2	59.3	448.6	2,688.9	66.6
2018.....	81.6	*	*	*	*	0.1	0.1	0.8	7.8	57.9	447.0	2,700.3	66.6
2017.....	80.4	*	*	0.0	*	*	0.1	0.7	7.5	56.9	450.9	2,707.3	66.7
2016.....	77.2	*	*	0.1	*	0.1	0.1	0.8	7.3	54.7	441.6	2,626.4	64.9
2015.....	76.5	*	*	0.1	*	*	0.1	0.8	6.8	53.0	447.2	2,637.4	65.2
2014.....	75.2	*	0.1	0.1	*	*	0.1	0.8	6.9	52.7	450.6	2,611.3	64.9
2013.....	74.1	*	*	*	*	*	0.1	0.9	7.4	52.2	449.7	2,601.8	64.8
2012.....	71.2	*	0.2	*	*	*	0.1	0.9	6.8	50.3	445.2	2,532.7	63.3
2011.....	68.3	*	0.1	0.1	*	*	0.1	0.8	6.5	50.4	436.6	2,458.6	61.8
2010.....	63.6	*	*	*	*	*	0.1	0.9	6.3	48.3	412.1	2,352.4	58.8
2009.....	57.9	*	0.1	0.1	*	0.1	0.2	0.8	5.7	45.2	383.3	2,151.3	54.2
2008.....	58.9	*	0.2	0.1	*	*	0.1	0.9	6.0	46.4	396.7	2,213.2	55.9
2007.....	51.8	*	0.2	0.1	*	*	0.1	0.8	5.5	42.3	350.0	1,976.0	49.8
2006.....	50.8	*	0.1	0.1	*	*	0.1	0.8	5.5	42.4	346.2	1,967.0	49.5
2005.....	43.8	*	0.2	*	*	*	0.1	0.6	4.5	36.3	302.1	1,735.4	43.4
2004.....	39.0	*	0.2	0.1	*	*	0.1	0.6	3.9	33.1	275.4	1,556.9	39.1
2003.....	38.1	*	0.2	0.1	*	*	0.1	0.6	4.0	34.1	269.5	1,523.1	38.4
2002.....	35.5	*	0.2	0.1	*	*	0.1	0.4	3.6	31.9	254.9	1,428.9	36.1
2001.....	32.5	*	0.2	0.0	*	0.1	0.1	0.5	3.8	30.5	234.6	1,299.3	33.1
2000.....	29.7	*	0.2	0.1	0.1	0.1	0.1	0.5	3.6	29.4	218.5	1,180.9	30.5
1999.....	26.7	*	0.2	0.1	*	*	0.1	0.4	3.4	27.0	197.9	1,062.5	27.5

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group											Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Drug-induced causes ⁶													
2022.....	33.6	2.0	1.2	0.4	15.3	51.8	65.1	57.3	50.7	20.7	6.3	4.7	33.8
2021.....	33.5	2.1	0.8	0.3	17.5	54.2	64.0	56.2	48.0	18.5	6.0	5.3	33.6
2020.....	29.2	1.7	0.4	0.3	17.0	48.8	55.8	49.1	39.9	14.7	4.9	5.1	29.5
2019.....	22.7	1.3	0.3	0.2	11.5	36.9	42.3	38.9	32.7	12.6	5.2	4.6	22.8
2018.....	21.7	0.8	0.2	0.1	11.0	36.8	40.0	37.3	30.4	11.4	4.8	5.1	21.8
2017.....	22.7	0.9	0.2	0.2	13.0	39.8	40.6	39.8	30.0	10.5	4.5	5.3	22.8
2016.....	20.8	0.9	0.3	0.1	12.8	35.9	36.6	36.5	27.7	9.2	4.1	5.3	20.8
2015.....	17.2	0.7	0.4	0.1	10.0	28.0	29.6	31.9	23.3	8.1	4.4	5.6	17.2
2014.....	15.6	0.6	0.3	0.1	8.9	24.0	26.2	29.8	21.7	7.6	4.4	5.0	15.5
2013.....	14.7	0.8	0.3	0.1	8.6	21.7	24.1	29.0	20.6	7.1	4.4	5.3	14.6
2012.....	14.0	0.8	0.2	0.1	8.3	20.9	23.1	28.3	17.9	6.5	4.0	5.1	13.8
2011.....	14.0	0.6	0.2	0.1	8.9	20.9	23.4	28.2	17.1	6.0	4.0	4.9	13.9
2010.....	13.1	0.6	0.3	0.2	8.4	19.2	21.7	26.5	16.2	5.2	4.0	5.5	12.9
2009.....	12.8	0.8	0.2	0.1	8.0	17.8	21.5	26.9	14.9	5.4	4.5	5.1	12.6
2008.....	12.7	0.5	0.3	0.1	8.3	17.4	22.2	26.8	14.0	5.2	4.0	5.0	12.6
2007.....	12.7	0.8	0.3	0.2	8.5	17.5	22.6	26.8	13.4	4.6	3.9	5.2	12.6
2006.....	12.9	1.1	0.2	0.1	8.5	17.2	23.5	26.7	12.1	5.2	6.0	8.8	12.8
2005.....	11.3	0.9	0.2	0.1	7.3	14.6	21.5	23.6	10.6	4.7	5.4	8.3	11.3
2004.....	10.5	0.7	0.2	0.2	6.9	12.9	21.1	21.7	9.0	4.2	4.8	6.7	10.5
2003.....	9.9	0.6	0.2	0.1	6.3	12.3	20.7	20.0	8.0	4.1	4.2	6.3	9.9
2002.....	9.1	0.7	0.2	0.1	5.4	11.3	19.8	18.0	6.8	3.6	3.8	6.0	9.1
2001.....	7.6	0.5	0.2	0.1	4.5	9.5	17.0	14.7	5.4	3.0	3.5	5.2	7.6
2000.....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7	7.0
1999.....	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8	6.8
Alcohol-induced causes ⁶													
2022.....	15.4	*	*	*	0.5	6.6	17.0	26.8	39.7	28.8	15.7	7.6	13.5
2021.....	16.3	*	*	*	0.5	6.7	18.0	30.3	42.0	29.6	15.4	7.8	14.4
2020.....	14.9	*	*	*	0.4	6.0	16.0	27.5	39.3	27.1	14.4	6.5	13.1
2019.....	11.9	*	*	*	0.4	4.2	11.1	22.0	32.5	22.5	12.6	6.4	10.4
2018.....	11.4	*	*	*	0.3	3.7	10.0	21.6	31.5	22.2	12.4	6.1	9.9
2017.....	11.0	*	*	*	0.3	3.4	9.4	21.8	30.2	20.9	11.7	6.4	9.6
2016.....	10.8	*	*	*	0.4	3.6	9.2	21.4	29.7	20.3	11.8	6.3	9.5
2015.....	10.3	*	*	*	0.4	3.2	8.7	21.6	28.2	19.1	11.2	5.8	9.1
2014.....	9.6	*	*	*	0.3	2.8	8.0	20.4	26.8	17.6	10.5	5.6	8.5
2013.....	9.2	*	*	*	0.3	2.5	7.7	20.1	25.3	16.6	10.3	4.9	8.2
2012.....	8.8	*	*	*	0.4	2.4	7.4	20.0	24.1	15.8	10.3	5.0	8.0
2011.....	8.6	*	*	*	0.4	2.1	7.6	19.8	22.7	15.2	9.6	5.1	7.7
2010.....	8.3	*	*	*	0.3	2.2	7.5	19.1	21.9	15.8	9.6	5.3	7.6
2009.....	8.0	*	*	*	0.4	1.8	7.6	18.7	20.8	15.1	9.2	4.8	7.4
2008.....	8.0	*	*	*	0.4	2.0	7.6	18.6	20.7	15.3	9.4	5.2	7.4
2007.....	7.7	*	*	*	0.4	1.9	7.3	18.2	19.9	15.2	9.6	5.0	7.2
2006.....	7.4	*	*	*	0.3	1.6	7.5	17.5	19.2	14.5	9.7	5.3	7.0
2005.....	7.3	*	*	*	0.4	1.4	7.5	17.6	19.4	14.9	9.2	5.0	7.0
2004.....	7.2	*	*	*	0.3	1.6	7.7	17.3	18.6	15.5	9.2	4.6	7.0
2003.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.0	9.2	4.3	7.0
2002.....	7.0	*	*	*	0.3	1.5	8.1	16.9	18.3	15.4	9.3	4.6	6.9
2001.....	7.1	*	*	*	0.3	1.6	8.3	17.1	18.3	15.5	9.6	5.1	7.0
2000.....	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4	7.0
1999.....	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5	7.1

Table 6. Death rate by age, and age-adjusted death rate, for the 15 leading causes of death in 2022, dementia-related causes, drug-induced causes, alcohol-induced causes, and injury by firearms: United States, 1999–2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes]

Cause of death (based on ICD–10) and year	All ages ¹	Age group											Age- adjusted rate ³
		Younger than 1 ²	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Injury by firearms ⁶													
2022.....	14.5	*	0.9	1.5	21.1	22.9	18.1	14.8	13.4	11.9	16.5	18.5	14.2
2021.....	14.7	*	0.9	1.6	23.5	24.8	18.1	14.5	12.1	11.7	16.2	18.3	14.6
2020.....	13.7	*	0.8	1.5	22.2	22.7	16.7	13.3	11.8	10.9	15.3	16.2	13.6
2019.....	12.1	*	0.5	1.0	17.4	18.1	14.6	12.7	12.3	11.1	15.0	16.1	11.9
2018.....	12.1	*	0.6	1.1	17.2	17.7	14.6	12.8	12.7	12.0	15.4	14.7	11.9
2017.....	12.2	*	0.5	1.1	17.7	18.5	14.4	13.1	12.3	11.4	14.8	15.6	12.0
2016.....	12.0	*	0.6	0.9	17.2	18.2	14.5	12.8	11.9	11.4	14.7	14.3	11.8
2015.....	11.3	*	0.5	0.9	15.7	16.8	13.1	12.4	11.7	11.3	14.5	14.5	11.1
2014.....	10.5	*	0.4	0.9	14.0	14.7	12.1	12.2	11.4	11.5	13.9	15.0	10.3
2013.....	10.6	*	0.4	0.8	14.1	15.3	12.3	12.3	11.5	11.3	14.1	13.9	10.4
2012.....	10.7	*	0.4	0.8	14.7	15.3	12.4	12.4	11.6	10.8	14.1	13.6	10.5
2011.....	10.4	*	0.5	0.8	14.4	15.0	11.7	12.2	11.0	10.9	13.7	13.1	10.2
2010.....	10.3	*	0.4	0.7	14.2	15.0	11.7	12.0	11.1	10.7	12.7	13.2	10.1
2009.....	10.2	*	0.4	0.7	14.4	14.5	11.9	11.8	10.8	10.9	13.3	12.5	10.1
2008.....	10.4	*	0.5	0.7	15.4	15.4	11.8	11.5	10.8	10.7	13.2	12.5	10.3
2007.....	10.4	*	0.4	0.8	16.0	15.9	12.0	11.1	10.1	9.8	13.1	12.7	10.3
2006.....	10.4	*	0.4	0.9	16.7	15.7	11.6	11.2	9.7	9.9	12.9	12.5	10.3
2005.....	10.4	*	0.4	0.8	16.1	16.1	11.7	11.2	9.7	10.2	13.6	13.0	10.3
2004.....	10.1	*	0.3	0.7	15.6	15.3	11.4	11.0	9.8	10.1	13.3	12.7	10.0
2003.....	10.4	*	0.3	0.8	16.5	15.8	11.6	11.1	10.0	10.3	13.4	13.2	10.3
2002.....	10.5	*	0.4	0.8	16.6	15.6	12.2	10.8	10.2	10.8	14.4	13.2	10.5
2001.....	10.4	*	0.5	0.8	16.6	15.5	11.7	10.5	10.1	10.9	14.3	13.1	10.3
2000.....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2	10.2
1999.....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5	10.3

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Data for age not stated included in "All ages" but not distributed among age groups.

²Death rates for ages younger than 1 year (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

³For computation method, see Technical Notes.

⁴COVID-19 became an official cause of death in 2020; rates for years before 2020 are not applicable.

⁵Data include September 11, 2001, related deaths for which death certificates were filed as of October 24, 2002; see Technical Notes from "Deaths: Final Data for 2001," National Vital Statistics Reports, vol 53, no 3.

⁶For the list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group											Age not stated
		Younger than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
All causes	3,279,857	20,553	4,156	6,239	35,232	74,369	111,605	183,284	417,541	668,581	824,903	933,291	103
Salmonella infections. (A01-A02)	82	5	3	—	2	2	2	6	11	16	25	10	—
Shigellosis and amebiasis (A03,A06)	17	—	3	1	—	—	—	3	2	6	1	1	—
Certain other intestinal infections. (A04,A07-A09)	7,380	141	20	15	16	33	85	253	758	1,646	2,256	2,157	—
Tuberculosis. (A16-A19)	565	—	—	2	5	21	25	41	128	138	123	82	—
Respiratory tuberculosis. (A16)	391	—	—	1	5	11	21	24	85	93	85	66	—
Other tuberculosis (A17-A19)	174	—	—	1	—	10	4	17	43	45	38	16	—
Whooping cough (A37)	2	1	—	—	—	—	—	1	—	—	—	—	—
Scarlet fever and erysipelas (A38,A46)	3	—	—	—	—	—	1	—	—	1	1	—	—
Meningococcal infection (A39)	27	1	1	1	4	2	8	5	3	1	1	—	—
Septicemia (A40-A41)	42,261	123	60	59	113	447	1,024	2,488	6,167	10,623	11,723	9,433	1
Syphilis (A50-A53)	61	15	1	—	—	—	4	7	5	12	10	7	—
Acute poliomyelitis. (A80)	—	—	—	—	—	—	—	—	—	—	—	—	—
Arthropod-borne viral encephalitis. (A83-A84,A85.2)	6	—	—	—	—	—	—	—	2	3	1	—	—
Measles (B05)	—	—	—	—	—	—	—	—	—	—	—	—	—
Viral hepatitis. (B15-B19)	3,107	2	4	—	2	25	129	356	1,057	1,116	306	110	—
Human immunodeficiency virus (HIV) disease (B20-B24)	4,941	1	—	—	43	481	777	1,002	1,407	902	271	57	—
Malaria (B50-B54)	8	—	—	—	1	—	1	1	4	—	1	—	—
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99,U07.1)	198,029	287	238	225	524	1,821	4,207	10,426	26,038	44,837	54,181	55,244	1
Malignant neoplasms. (C00-C97)	608,371	40	266	835	1,421	3,641	11,177	33,363	105,133	178,860	171,573	102,057	5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	11,884	—	—	1	6	65	205	866	2,903	3,662	2,731	1,445	—
Malignant neoplasm of esophagus. (C15)	16,016	—	—	1	6	33	226	1,040	3,456	5,524	4,099	1,629	2
Malignant neoplasm of stomach. (C16)	10,853	—	1	—	17	124	457	943	2,078	2,926	2,669	1,637	1
Malignant neoplasms of colon, rectum and anus (C18-C21)	54,268	—	—	1	35	349	1,647	5,072	10,491	13,981	12,751	9,941	—
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	28,937	4	9	16	26	122	373	1,354	6,098	10,692	7,237	3,006	—
Malignant neoplasm of pancreas (C25)	48,323	1	—	2	4	70	490	2,403	8,783	15,445	14,239	6,886	—
Malignant neoplasm of larynx (C32)	3,933	—	—	—	1	2	25	202	965	1,423	931	384	—
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	131,982	—	—	1	12	118	771	4,484	24,394	44,583	40,755	16,863	1
Malignant melanoma of skin. (C43)	8,243	—	1	—	15	106	293	566	1,362	2,248	2,161	1,491	—
Malignant neoplasm of breast (C50)	42,672	—	—	—	7	383	1,856	4,083	8,165	10,747	9,864	7,567	—
Malignant neoplasm of cervix uteri. (C53)	4,051	—	—	—	4	152	562	791	932	835	507	268	—
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	12,764	—	—	—	—	46	211	723	2,568	4,541	3,257	1,417	1
Malignant neoplasm of ovary (C56)	13,214	—	—	2	33	104	287	1,061	2,717	3,835	3,549	1,626	—
Malignant neoplasm of prostate (C61)	33,363	—	—	—	2	2	21	357	2,777	8,385	11,604	10,215	—
Malignant neoplasms of kidney and renal pelvis (C64-C65)	14,488	2	12	29	22	57	198	919	2,633	4,214	4,012	2,390	—
Malignant neoplasm of bladder (C67)	17,334	—	—	1	1	20	100	330	1,640	3,910	5,870	5,462	—
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	17,937	9	56	302	243	446	823	1,692	3,881	5,365	3,708	1,412	—
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	56,959	14	98	206	383	594	996	2,180	6,551	14,392	18,901	12,644	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group											Age not stated
		Younger than 1	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Hodgkin disease (C81)	1,056	—	—	2	12	48	59	102	140	259	273	161	—
Non-Hodgkin lymphoma (C82–C85)	19,794	—	3	20	55	150	320	776	2,335	5,058	6,670	4,407	—
Leukemia (C91–C95)	23,324	14	95	184	313	384	526	873	2,437	5,620	7,561	5,317	—
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	12,616	—	—	—	1	10	87	425	1,617	3,420	4,340	2,716	—
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	169	—	—	—	2	2	4	4	22	35	57	43	—
All other and unspecified malignant neoplasms (C17,C23–C24, C26–C31,C37–C41,C44–C49,C51–C52,C57–C60, C62–C63,C66,C68–C69,C73–C80,C97)	81,150	10	89	273	604	848	1,636	4,297	12,739	22,152	22,728	15,774	—
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	16,034	32	37	51	74	106	256	503	1,437	3,376	5,216	4,946	—
Anemias (D50–D64)	6,021	5	13	31	88	153	218	289	612	1,110	1,512	1,990	—
Diabetes mellitus (E10–E14)	101,209	2	7	48	324	1,188	2,879	7,364	17,410	26,963	26,388	18,634	2
Nutritional deficiencies (E40–E64)	21,020	13	10	4	13	47	113	282	1,010	2,728	5,458	11,342	—
Malnutrition (E40–E46)	20,552	8	9	4	13	46	104	258	945	2,632	5,366	11,167	—
Other nutritional deficiencies (E50–E64)	468	5	1	—	—	1	9	24	65	96	92	175	—
Meningitis (G00,G03)	604	43	18	17	15	29	59	61	93	130	96	43	—
Parkinson disease (G20–G21)	39,915	1	—	—	—	6	12	86	879	6,424	17,886	14,621	—
Alzheimer disease (G30)	120,122	—	—	—	—	—	6	111	1,480	9,005	36,133	73,387	—
Major cardiovascular diseases (I00–I78)	936,436	340	156	335	1,078	4,765	15,656	40,956	107,984	182,806	242,054	340,276	30
Diseases of heart (I00–I09,I11,I13,I20–I51)	702,880	241	103	218	848	3,789	12,258	32,298	85,733	140,475	178,108	248,782	27
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	4,294	1	1	1	10	61	127	196	457	818	1,215	1,407	—
Hypertensive heart disease (I11)	72,070	—	—	2	58	539	2,146	4,701	10,137	13,199	14,820	26,465	3
Hypertensive heart and renal disease (I13)	16,091	—	—	1	2	45	158	456	1,050	2,279	4,288	7,812	—
Ischemic heart diseases (I20–I25)	371,506	11	5	8	102	990	4,956	17,223	50,365	83,121	97,826	116,879	20
Acute myocardial infarction (I21–I22)	103,905	4	1	3	47	394	1,877	6,109	16,744	25,801	27,512	25,409	4
Other acute ischemic heart diseases (I24)	4,823	2	1	1	3	21	83	308	823	1,146	1,182	1,253	—
Other forms of chronic ischemic heart disease (I20,I25)	262,778	5	3	4	52	575	2,996	10,806	32,798	56,174	69,132	90,217	16
Atherosclerotic cardiovascular disease, so described (I25.0)	81,672	—	1	1	21	351	1,767	6,079	16,731	22,134	17,820	16,755	12
All other forms of chronic ischemic heart disease (I20,I25.1–I25.9)	181,106	5	2	3	31	224	1,229	4,727	16,067	34,040	51,312	73,462	4
Other heart diseases (I26–I51)	238,919	229	97	206	676	2,154	4,871	9,722	23,724	41,058	59,959	96,219	4
Acute and subacute endocarditis (I33)	1,803	2	—	3	17	109	177	213	323	403	359	197	—
Diseases of pericardium and acute myocarditis (I30–I31,I40)	1,237	9	3	18	22	43	65	101	182	292	283	219	—
Heart failure (I50)	87,941	13	5	8	43	247	714	1,962	5,820	12,599	22,750	43,779	1
All other forms of heart disease (I26–I28, I34–I38,I42–I49,I51)	147,938	205	89	177	594	1,755	3,915	7,446	17,399	27,764	36,567	52,024	3
Essential hypertension and hypertensive renal disease (I10,I12,I15)	43,293	3	—	2	24	175	697	1,832	4,982	8,315	10,790	16,472	1
Cerebrovascular diseases (I60–I69)	165,393	87	49	100	157	599	2,150	5,563	14,173	28,384	46,421	67,708	2
Atherosclerosis (I70)	3,936	2	1	—	—	2	10	68	291	684	999	1,879	—
Other diseases of circulatory system (I71–I78)	20,934	7	3	15	49	200	541	1,195	2,805	4,948	5,736	5,435	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group											Age not stated
		Younger than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older	
Aortic aneurysm and dissection (I71)	9,999	1	—	7	29	128	357	747	1,435	2,225	2,762	2,308	—
Other diseases of arteries, arterioles and capillaries (I72-I78)	10,935	6	3	8	20	72	184	448	1,370	2,723	2,974	3,127	—
Other disorders of circulatory system (I80-I99)	5,216	16	1	3	60	191	480	613	926	1,028	1,009	888	1
Influenza and pneumonia (J09-J18)	47,052	154	129	131	168	490	985	1,905	5,067	9,408	12,774	15,840	1
Influenza (J09-J11)	5,944	12	53	76	61	112	217	310	771	1,147	1,475	1,710	—
Pneumonia (J12-J18)	41,108	142	76	55	107	378	768	1,595	4,296	8,261	11,299	14,130	1
Other acute lower respiratory infections (J20-J22,U04)	278	43	24	4	2	3	4	8	21	42	43	84	—
Acute bronchitis and bronchiolitis (J20-J21)	182	40	22	3	1	2	2	3	11	25	25	48	—
Other and unspecified acute lower respiratory infections (J22,U04)	96	3	2	1	1	1	2	5	10	17	18	36	—
Chronic lower respiratory diseases (J40-J47)	147,382	10	39	106	197	368	732	2,987	17,138	38,712	49,746	37,345	2
Bronchitis, chronic and unspecified (J40-J42)	358	4	7	2	4	3	3	13	44	76	82	120	—
Emphysema (J43)	7,905	1	—	1	1	2	34	179	983	2,205	2,716	1,783	—
Asthma (J45-J46)	3,602	2	26	95	181	322	366	480	580	574	487	489	—
Other chronic lower respiratory diseases (J44,J47)	135,517	3	6	8	11	41	329	2,315	15,531	35,857	46,461	34,953	2
Pneumoconioses and chemical effects (J60-J66,J68,U07.0)	562	—	—	2	1	3	5	13	39	135	196	168	—
Pneumonitis due to solids and liquids (J69)	20,052	8	5	7	49	132	278	599	1,763	3,647	5,892	7,672	—
Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)	48,082	172	103	99	143	319	641	1,584	5,054	11,010	15,433	13,522	2
Peptic ulcer (K25-K28)	4,105	—	1	2	8	34	116	285	624	995	1,069	971	—
Diseases of appendix (K35-K38)	487	3	1	4	4	11	20	36	81	91	115	121	—
Hernia (K40-K46)	2,712	10	9	5	6	9	49	126	316	528	699	955	—
Chronic liver disease and cirrhosis (K70,K73-K74)	54,803	1	2	2	58	1,786	5,501	9,401	16,484	13,178	6,428	1,959	3
Alcoholic liver disease (K70)	30,910	—	—	—	46	1,508	4,335	6,688	10,362	6,017	1,683	270	1
Other chronic liver disease and cirrhosis (K73-K74)	23,893	1	2	2	12	278	1,166	2,713	6,122	7,161	4,745	1,689	2
Cholelithiasis and other disorders of gallbladder (K80-K82)	4,483	1	—	1	18	21	48	131	396	904	1,360	1,603	—
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57,937	38	4	11	64	358	1,029	2,679	6,668	12,948	17,003	17,135	—
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	810	2	1	4	2	4	12	30	80	135	255	285	—
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	330	—	—	2	5	14	20	38	47	85	75	44	—
Renal failure (N17-N19)	56,756	36	3	5	57	340	995	2,608	6,535	12,718	16,661	16,798	—
Other disorders of kidney (N25,N27)	41	—	—	—	—	—	2	3	6	10	12	8	—
Infections of kidney (N10-N12,N13.6,N15.1)	1,205	4	3	3	5	15	37	78	142	285	356	276	1
Hyperplasia of prostate (N40)	839	—	—	—	—	—	—	8	39	103	244	445	—
Inflammatory diseases of female pelvic organs (N70-N76)	232	—	—	—	7	3	19	15	44	50	47	47	—
Pregnancy, childbirth and the puerperium (O00-O99)	1,263	...	...	5	164	591	489	12	1	1	—	—	—
Pregnancy with abortive outcome (O00-O07)	33	...	...	—	5	18	10	—	—	—	—	—	—
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	1,230	...	...	5	159	573	479	12	1	1	—	—	—
Certain conditions originating in the perinatal period (P00-P96)	10,203	10,068	62	28	15	7	5	3	5	5	1	4	—
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10,149	3,970	441	446	412	490	542	783	1,310	842	559	354	—

See footnotes at end of table.

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group											Age not stated
		Younger than 1	1–4	5–14	15–24	25–34	35–44	45–54	55–64	65–74	75–84	85 and older	
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	34,110	2,691	305	133	598	1,334	1,699	2,099	3,634	5,003	6,010	10,582	22
All other diseases (residual)	410,973	563	417	802	1,995	5,670	10,832	18,951	40,782	68,055	103,290	159,609	7
Accidents (unintentional injuries) (V01–X59,Y85–Y86)	227,039	1,354	1,288	1,652	14,669	33,058	36,972	31,394	34,017	23,150	22,392	27,074	19
Transport accidents (V01–V99,Y85)	49,178	80	386	945	7,215	8,797	7,586	6,545	7,273	5,482	3,435	1,431	3
Motor vehicle accidents (V02–V04,V09.0, V09.2,V12–V14,V19.0–V19.2,V19.4–V19.6, V20–V79,V80.3–V80.5,V81.0–V81.1,V82.0–V82.1, V83–V86,V87.0–V87.8,V88.0–V88.8, V89.0,V89.2)	46,027	79	366	900	6,997	8,406	7,084	6,080	6,567	4,949	3,219	1,377	3
Other land transport accidents (V01,V05–V06,V09.1, V09.3–V09.9,V10–V11,V15–V18,V19.3,V19.8–V19.9, V80.0–V80.2,V80.6–V80.9,V81.2–V81.9, V82.2–V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	1,195	1	13	19	106	184	207	157	240	174	74	20	–
Water, air and space, and other and unspecified transport accidents and their sequelae (V90–V99,Y85)	1,956	–	7	26	112	207	295	308	466	359	142	34	–
Nontransport accidents (W00–X59,Y86)	177,861	1,274	902	707	7,454	24,261	29,386	24,849	26,744	17,668	18,957	25,643	16
Falls (W00–W19)	46,630	6	21	21	149	383	682	1,200	3,247	6,698	13,396	20,825	2
Accidental discharge of firearms (W32–W34)	463	–	49	35	120	69	55	42	36	31	18	8	–
Accidental drowning and submersion (W65–W74)	4,168	39	458	222	454	500	502	421	562	552	321	135	2
Accidental exposure to smoke, fire and flames (X00–X09)	3,478	15	85	158	125	180	232	311	689	864	555	264	–
Accidental poisoning and exposure to noxious substances (X40–X49)	102,958	28	75	99	6,205	22,295	26,767	21,364	19,334	5,924	697	159	11
Other and unspecified nontransport accidents and their sequelae (W20–W31,W35–W64, W75–W99,X10–X39,X50–X59,Y86)	20,164	1,186	214	172	401	834	1,148	1,511	2,876	3,599	3,970	4,252	1
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	49,476	...	...	502	6,040	8,663	8,185	7,781	7,864	5,396	3,549	1,493	3
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	27,032	...	...	180	3,246	4,230	3,759	3,838	4,412	3,536	2,686	1,145	–
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60–X71,X75–X84,Y87.0)	22,444	...	...	322	2,794	4,433	4,426	3,943	3,452	1,860	863	348	3
Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	24,849	252	343	546	6,262	6,712	4,765	2,740	1,824	877	399	127	2
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	19,651	16	77	395	5,814	5,836	3,861	1,947	1,080	422	163	40	–
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3,*U01.5–*U01.9,*U02, X85–X92,X96–Y09,Y87.1)	5,198	236	266	151	448	876	904	793	744	455	236	87	2
Legal intervention (Y35,Y89.0)	810	–	–	1	120	248	209	134	70	23	5	–	–
Events of undetermined intent (Y10–Y34,Y87.2,Y89.9)	5,604	123	127	98	403	1,014	1,174	1,025	959	458	153	69	1
Discharge of firearms, undetermined intent (Y22–Y24)	415	–	14	18	103	94	62	45	41	20	14	4	–
Other and unspecified events of undetermined intent and their sequelae (Y10–Y21,Y25–Y34,Y87.2,Y89.9)	5,189	123	113	80	300	920	1,112	980	918	438	139	65	1
Operations of war and their sequelae (Y36,Y89.1)	7	–	–	–	–	–	2	1	–	1	3	–	–
Complications of medical and surgical care (Y40–Y84,Y88)	3,726	20	15	22	41	72	148	289	653	1,003	912	551	–

Table 7. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	All ages	Age group										85 and older	Age not stated
		Younger than 1	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84		
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ¹	4,231	1	—	—	5	10	46	135	419	989	1,383	1,243	—
COVID-19. (U07.1) ¹	186,552	141	101	131	447	1,640	3,841	9,678	24,252	42,062	51,188	53,070	1
Dementia-related causes ¹	292,881	10	16	23	9	16	52	371	3,946	22,770	83,437	182,229	2
Drug-induced deaths ¹	112,109	72	182	160	6,782	23,565	28,427	23,169	21,350	6,983	1,103	304	12
Drug overdose deaths ¹	107,941	71	182	160	6,696	23,029	27,583	22,352	20,252	6,434	930	241	11
Alcohol-induced deaths ¹	51,191	1	—	1	200	2,992	7,429	10,852	16,718	9,745	2,752	496	5
Firearm-related injuries ¹	48,204	16	140	628	9,363	10,436	7,904	5,981	5,625	4,030	2,884	1,197	—

— Quantity zero.

... Category not applicable.

¹Included in selected categories above. For list of ICD-10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group										
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
All causes	984.1	558.0	28.0	15.3	79.5	163.4	255.4	453.3	992.1	1,978.7	4,708.2	14,389.6
Salmonella infections. (A01-A02)	0.0	*	*	*	*	*	*	*	*	*	0.1	*
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections. (A04,A07-A09)	2.2	3.8	0.1	*	*	0.1	0.2	0.6	1.8	4.9	12.9	33.3
Tuberculosis. (A16-A19)	0.2	*	*	*	*	0.0	0.1	0.1	0.3	0.4	0.7	1.3
Respiratory tuberculosis. (A16)	0.1	*	*	*	*	*	0.0	0.1	0.2	0.3	0.5	1.0
Other tuberculosis (A17-A19)	0.1	*	*	*	*	*	*	*	0.1	0.1	0.2	*
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40-A41)	12.7	3.3	0.4	0.1	0.3	1.0	2.3	6.2	14.7	31.4	66.9	145.4
Syphilis (A50-A53)	0.0	*	*	*	*	*	*	*	*	*	*	*
Acute poliomyelitis. (A80)	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis. (A83-A84,A85.2)	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis. (B15-B19)	0.9	*	*	*	*	0.1	0.3	0.9	2.5	3.3	1.7	1.7
Human immunodeficiency virus (HIV) disease (B20-B24)	1.5	*	*	*	0.1	1.1	1.8	2.5	3.3	2.7	1.5	0.9
Malaria. (B50-B54)	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05,A20-A36,A42-A44,A48-A49, A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04, B06-B09,B25-B49,B55-B99,U07.1)	59.4	7.8	1.6	0.6	1.2	4.0	9.6	25.8	61.9	132.7	309.2	851.8
Malignant neoplasms. (C00-C97)	182.5	1.1	1.8	2.0	3.2	8.0	25.6	82.5	249.8	529.4	979.3	1,573.5
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)	3.6	*	*	*	*	0.1	0.5	2.1	6.9	10.8	15.6	22.3
Malignant neoplasm of esophagus. (C15)	4.8	*	*	*	*	0.1	0.5	2.6	8.2	16.3	23.4	25.1
Malignant neoplasm of stomach. (C16)	3.3	*	*	*	*	0.3	1.0	2.3	4.9	8.7	15.2	25.2
Malignant neoplasms of colon, rectum and anus (C18-C21)	16.3	*	*	*	0.1	0.8	3.8	12.5	24.9	41.4	72.8	153.3
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	8.7	*	*	*	0.1	0.3	0.9	3.3	14.5	31.6	41.3	46.3
Malignant neoplasm of pancreas (C25)	14.5	*	*	*	*	0.2	1.1	5.9	20.9	45.7	81.3	106.2
Malignant neoplasm of larynx (C32)	1.2	*	*	*	*	*	0.1	0.5	2.3	4.2	5.3	5.9
Malignant neoplasms of trachea, bronchus and lung (C33-C34)	39.6	*	*	*	*	0.3	1.8	11.1	58.0	131.9	232.6	260.0
Malignant melanoma of skin. (C43)	2.5	*	*	*	*	0.2	0.7	1.4	3.2	6.7	12.3	23.0
Malignant neoplasm of breast (C50)	12.8	*	*	*	*	0.8	4.2	10.1	19.4	31.8	56.3	116.7
Malignant neoplasm of cervix uteri. (C53)	1.2	*	*	*	*	0.3	1.3	2.0	2.2	2.5	2.9	4.1
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)	3.8	*	*	*	*	0.1	0.5	1.8	6.1	13.4	18.6	21.8
Malignant neoplasm of ovary (C56)	4.0	*	*	*	0.1	0.2	0.7	2.6	6.5	11.4	20.3	25.1
Malignant neoplasm of prostate (C61)	10.0	*	*	*	*	*	0.0	0.9	6.6	24.8	66.2	157.5
Malignant neoplasms of kidney and renal pelvis (C64-C65)	4.3	*	*	0.1	0.0	0.1	0.5	2.3	6.3	12.5	22.9	36.8
Malignant neoplasm of bladder (C67)	5.2	*	*	*	*	0.0	0.2	0.8	3.9	11.6	33.5	84.2
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)	5.4	*	0.4	0.7	0.5	1.0	1.9	4.2	9.2	15.9	21.2	21.8

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group										
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)	17.1	*	0.7	0.5	0.9	1.3	2.3	5.4	15.6	42.6	107.9	194.9
Hodgkin disease (C81)	0.3	*	*	*	*	0.1	0.1	0.3	0.3	0.8	1.6	2.5
Non-Hodgkin lymphoma (C82-C85)	5.9	*	*	0.0	0.1	0.3	0.7	1.9	5.5	15.0	38.1	67.9
Leukemia (C91-C95)	7.0	*	0.6	0.4	0.7	0.8	1.2	2.2	5.8	16.6	43.2	82.0
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.8	*	*	*	*	*	0.2	1.1	3.8	10.1	24.8	41.9
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.7
All other and unspecified malignant neoplasms (C17,C23-C24, C26-C31,C37-C41,C44-C49,C51-C52,C57-C60, C62-C63,C66,C68-C69,C73-C80,C97)	24.3	*	0.6	0.7	1.4	1.9	3.7	10.6	30.3	65.6	129.7	243.2
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4.8	0.9	0.2	0.1	0.2	0.2	0.6	1.2	3.4	10.0	29.8	76.3
Anemias (D50-D64)	1.8	*	*	0.1	0.2	0.3	0.5	0.7	1.5	3.3	8.6	30.7
Diabetes mellitus (E10-E14)	30.4	*	*	0.1	0.7	2.6	6.6	18.2	41.4	79.8	150.6	287.3
Nutritional deficiencies (E40-E64)	6.3	*	*	*	*	0.1	0.3	0.7	2.4	8.1	31.2	174.9
Malnutrition (E40-E46)	6.2	*	*	*	*	0.1	0.2	0.6	2.2	7.8	30.6	172.2
Other nutritional deficiencies (E50-E64)	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.5	2.7
Meningitis (G00,G03)	0.2	1.2	*	*	*	0.1	0.1	0.2	0.2	0.4	0.5	0.7
Parkinson disease (G20-G21)	12.0	*	*	*	*	*	*	0.2	2.1	19.0	102.1	225.4
Alzheimer disease (G30)	36.0	*	*	*	*	*	*	0.3	3.5	26.7	206.2	1,131.5
Major cardiovascular diseases (I00-I78)	281.0	9.2	1.1	0.8	2.4	10.5	35.8	101.3	256.6	541.0	1,381.5	5,246.4
Diseases of heart (I00-I09,I11,I13,I20-I51)	210.9	6.5	0.7	0.5	1.9	8.3	28.1	79.9	203.7	415.7	1,016.6	3,835.8
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)	1.3	*	*	*	*	0.1	0.3	0.5	1.1	2.4	6.9	21.7
Hypertensive heart disease (I11)	21.6	*	*	*	0.1	1.2	4.9	11.6	24.1	39.1	84.6	408.0
Hypertensive heart and renal disease (I13)	4.8	*	*	*	*	0.1	0.4	1.1	2.5	6.7	24.5	120.4
Ischemic heart diseases (I20-I25)	111.5	*	*	*	0.2	2.2	11.3	42.6	119.7	246.0	558.4	1,802.1
Acute myocardial infarction (I21-I22)	31.2	*	*	*	0.1	0.9	4.3	15.1	39.8	76.4	157.0	391.8
Other acute ischemic heart diseases (I24)	1.4	*	*	*	*	0.0	0.2	0.8	2.0	3.4	6.7	19.3
Other forms of chronic ischemic heart disease (I20,I25)	78.8	*	*	*	0.1	1.3	6.9	26.7	77.9	166.3	394.6	1,391.0
Atherosclerotic cardiovascular disease, so described (I25.0)	24.5	*	*	*	0.0	0.8	4.0	15.0	39.8	65.5	101.7	258.3
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)	54.3	*	*	*	0.1	0.5	2.8	11.7	38.2	100.7	292.9	1,132.6
Other heart diseases (I26-I51)	71.7	6.2	0.7	0.5	1.5	4.7	11.1	24.0	56.4	121.5	342.2	1,483.5
Acute and subacute endocarditis (I33)	0.5	*	*	*	*	0.2	0.4	0.5	0.8	1.2	2.0	3.0
Diseases of pericardium and acute myocarditis (I30-I31,I40)	0.4	*	*	*	0.0	0.1	0.1	0.2	0.4	0.9	1.6	3.4
Heart failure (I50)	26.4	*	*	*	0.1	0.5	1.6	4.9	13.8	37.3	129.8	675.0
All other forms of heart disease (I26-I28, I34-I38,I42-I49,I51)	44.4	5.6	0.6	0.4	1.3	3.9	9.0	18.4	41.3	82.2	208.7	802.1
Essential hypertension and hypertensive renal disease (I10,I12,I15)	13.0	*	*	*	0.1	0.4	1.6	4.5	11.8	24.6	61.6	254.0
Cerebrovascular diseases (I60-I69)	49.6	2.4	0.3	0.2	0.4	1.3	4.9	13.8	33.7	84.0	265.0	1,043.9

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group										
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Atherosclerosis. (I70)	1.2	*	*	*	*	*	*	0.2	0.7	2.0	5.7	29.0
Other diseases of circulatory system (I71-I78)	6.3	*	*	*	0.1	0.4	1.2	3.0	6.7	14.6	32.7	83.8
Aortic aneurysm and dissection (I71)	3.0	*	*	*	0.1	0.3	0.8	1.8	3.4	6.6	15.8	35.6
Other diseases of arteries, arterioles and capillaries (I72-I78)	3.3	*	*	*	0.0	0.2	0.4	1.1	3.3	8.1	17.0	48.2
Other disorders of circulatory system (I80-I99)	1.6	*	*	*	0.1	0.4	1.1	1.5	2.2	3.0	5.8	13.7
Influenza and pneumonia (J09-J18)	14.1	4.2	0.9	0.3	0.4	1.1	2.3	4.7	12.0	27.8	72.9	244.2
Influenza (J09-J11)	1.8	*	0.4	0.2	0.1	0.2	0.5	0.8	1.8	3.4	8.4	26.4
Pneumonia (J12-J18)	12.3	3.9	0.5	0.1	0.2	0.8	1.8	3.9	10.2	24.4	64.5	217.9
Other acute lower respiratory infections (J20-J22,U04)	0.1	1.2	0.2	*	*	*	*	*	0.0	0.1	0.2	1.3
Acute bronchitis and bronchiolitis (J20-J21)	0.1	1.1	0.1	*	*	*	*	*	*	0.1	0.1	0.7
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	*	*	*	*	*	*	*	*	*	*	0.6
Chronic lower respiratory diseases (J40-J47)	44.2	*	0.3	0.3	0.4	0.8	1.7	7.4	40.7	114.6	283.9	575.8
Bronchitis, chronic and unspecified (J40-J42)	0.1	*	*	*	*	*	*	*	0.1	0.2	0.5	1.9
Emphysema (J43)	2.4	*	*	*	*	*	0.1	0.4	2.3	6.5	15.5	27.5
Asthma (J45-J46)	1.1	*	0.2	0.2	0.4	0.7	0.8	1.2	1.4	1.7	2.8	7.5
Other chronic lower respiratory diseases (J44,J47)	40.7	*	*	*	*	0.1	0.8	5.7	36.9	106.1	265.2	538.9
Pneumoconioses and chemical effects (J60-J66,J68,U07.0)	0.2	*	*	*	*	*	*	*	0.1	0.4	1.1	2.6
Pneumonitis due to solids and liquids (J69)	6.0	*	*	*	0.1	0.3	0.6	1.5	4.2	10.8	33.6	118.3
Other diseases of respiratory system. (J00-J06,J30-J39,J67,J70-J98)	14.4	4.7	0.7	0.2	0.3	0.7	1.5	3.9	12.0	32.6	88.1	208.5
Peptic ulcer (K25-K28)	1.2	*	*	*	*	0.1	0.3	0.7	1.5	2.9	6.1	15.0
Diseases of appendix (K35-K38)	0.1	*	*	*	*	*	0.0	0.1	0.2	0.3	0.7	1.9
Hernia (K40-K46)	0.8	*	*	*	*	*	0.1	0.3	0.8	1.6	4.0	14.7
Chronic liver disease and cirrhosis (K70,K73-K74)	16.4	*	*	*	0.1	3.9	12.6	23.3	39.2	39.0	36.7	30.2
Alcoholic liver disease (K70)	9.3	*	*	*	0.1	3.3	9.9	16.5	24.6	17.8	9.6	4.2
Other chronic liver disease and cirrhosis (K73-K74)	7.2	*	*	*	*	0.6	2.7	6.7	14.5	21.2	27.1	26.0
Cholelithiasis and other disorders of gallbladder (K80-K82)	1.3	*	*	*	*	0.0	0.1	0.3	0.9	2.7	7.8	24.7
Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17.4	1.0	*	*	0.1	0.8	2.4	6.6	15.8	38.3	97.0	264.2
Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.5	4.4
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)	0.1	*	*	*	*	*	0.0	0.1	0.1	0.3	0.4	0.7
Renal failure (N17-N19)	17.0	1.0	*	*	0.1	0.7	2.3	6.5	15.5	37.6	95.1	259.0
Other disorders of kidney (N25,N27)	0.0	*	*	*	*	*	*	*	*	*	*	*
Infections of kidney (N10-N12,N13.6,N15.1)	0.4	*	*	*	*	*	0.1	0.2	0.3	0.8	2.0	4.3
Hyperplasia of prostate (N40)	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	6.9
Inflammatory diseases of female pelvic organs (N70-N76)	0.1	*	*	*	*	*	*	*	0.1	0.1	0.3	0.7
Pregnancy, childbirth and the puerperium (O00-O99)	0.4	...	...	*	0.4	1.3	1.1	*	*	*	*	*
Pregnancy with abortive outcome (O00-O07)	0.0	...	...	*	*	*	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (O10-O99)	0.4	...	...	*	0.4	1.3	1.1	*	*	*	*	*
Certain conditions originating in the perinatal period. (P00-P96)	3.1	273.4	0.4	0.1	*	*	*	*	*	*	*	*

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group										
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3.0	107.8	3.0	1.1	0.9	1.1	1.2	1.9	3.1	2.5	3.2	5.5
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)	10.2	73.1	2.1	0.3	1.3	2.9	3.9	5.2	8.6	14.8	34.3	163.2
All other diseases (residual)	123.3	15.3	2.8	2.0	4.5	12.5	24.8	46.9	96.9	201.4	589.5	2,460.9
Accidents (unintentional injuries). (V01-X59,Y85-Y86)	68.1	36.8	8.7	4.0	33.1	72.7	84.6	77.6	80.8	68.5	127.8	417.4
Transport accidents (V01-V99,Y85)	14.8	2.2	2.6	2.3	16.3	19.3	17.4	16.2	17.3	16.2	19.6	22.1
Motor vehicle accidents (V02-V04,V09.0, V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6, V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1, V83-V86,V87.0-V87.8,V88.0-V88.8, V89.0,V89.2)	13.8	2.1	2.5	2.2	15.8	18.5	16.2	15.0	15.6	14.6	18.4	21.2
Other land transport accidents (V01,V05-V06,V09.1, V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9, V80.0-V80.2,V80.6-V80.9,V81.2-V81.9, V82.2-V82.9,V87.9, V88.9,V89.1,V89.3,V89.9)	0.4	*	*	*	0.2	0.4	0.5	0.4	0.6	0.5	0.4	0.3
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.6	*	*	0.1	0.3	0.5	0.7	0.8	1.1	1.1	0.8	0.5
Nontransport accidents (W00-X59,Y86)	53.4	34.6	6.1	1.7	16.8	53.3	67.3	61.5	63.5	52.3	108.2	395.4
Falls (W00-W19)	14.0	*	0.1	0.1	0.3	0.8	1.6	3.0	7.7	19.8	76.5	321.1
Accidental discharge of firearms (W32-W34)	0.1	*	0.3	0.1	0.3	0.2	0.1	0.1	0.1	0.1	*	*
Accidental drowning and submersion (W65-W74)	1.3	1.1	3.1	0.5	1.0	1.1	1.1	1.0	1.3	1.6	1.8	2.1
Accidental exposure to smoke, fire and flames (X00-X09)	1.0	*	0.6	0.4	0.3	0.4	0.5	0.8	1.6	2.6	3.2	4.1
Accidental poisoning and exposure to noxious substances (X40-X49)	30.9	0.8	0.5	0.2	14.0	49.0	61.3	52.8	45.9	17.5	4.0	2.5
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64, W75-W99,X10-X39,X50-X59,Y86)	6.1	32.2	1.4	0.4	0.9	1.8	2.6	3.7	6.8	10.7	22.7	65.6
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14.8	...	...	1.2	13.6	19.0	18.7	19.2	18.7	16.0	20.3	23.0
Intentional self-harm (suicide) by discharge of firearms (X72-X74)	8.1	...	...	0.4	7.3	9.3	8.6	9.5	10.5	10.5	15.3	17.7
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)	6.7	...	...	0.8	6.3	9.7	10.1	9.8	8.2	5.5	4.9	5.4
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7.5	6.8	2.3	1.3	14.1	14.8	10.9	6.8	4.3	2.6	2.3	2.0
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)	5.9	*	0.5	1.0	13.1	12.8	8.8	4.8	2.6	1.2	0.9	0.6
Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02, X85-X92,X96-Y09,Y87.1)	1.6	6.4	1.8	0.4	1.0	1.9	2.1	2.0	1.8	1.3	1.3	1.3
Legal intervention (Y35,Y89.0)	0.2	*	*	*	0.3	0.5	0.5	0.3	0.2	0.1	*	*
Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)	1.7	3.3	0.9	0.2	0.9	2.2	2.7	2.5	2.3	1.4	0.9	1.1
Discharge of firearms, undetermined intent (Y22-Y24)	0.1	*	*	*	0.2	0.2	0.1	0.1	0.1	0.1	*	*
Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)	1.6	3.3	0.8	0.2	0.7	2.0	2.5	2.4	2.2	1.3	0.8	1.0

See footnotes at end of table.

Table 8. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by age: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	All ages ¹	Age group										
		Younger than 1 ²	1-4	5-14	15-24	25-34	35-44	45-54	55-64	65-74	75-84	85 and older
Operations of war and their sequelae. (Y36,Y89.1)	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care. (Y40-Y84,Y88)	1.1	0.5	*	0.1	0.1	0.2	0.3	0.7	1.6	3.0	5.2	8.5
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ³	1.3	*	*	*	*	*	0.1	0.3	1.0	2.9	7.9	19.2
COVID-19. (U07.1) ³	56.0	3.8	0.7	0.3	1.0	3.6	8.8	23.9	57.6	124.5	292.2	818.2
Dementia-related causes ³	87.9	*	*	0.1	*	*	0.1	0.9	9.4	67.4	476.2	2,809.6
Drug-induced deaths ³	33.6	2.0	1.2	0.4	15.3	51.8	65.1	57.3	50.7	20.7	6.3	4.7
Drug overdose deaths ³	32.4	1.9	1.2	0.4	15.1	50.6	63.1	55.3	48.1	19.0	5.3	3.7
Alcohol-induced deaths ³	15.4	*	*	*	0.5	6.6	17.0	26.8	39.7	28.8	15.7	7.6
Firearm-related injuries ³	14.5	*	0.9	1.5	21.1	22.9	18.1	14.8	13.4	11.9	16.5	18.5

0.0 Quantity more than zero but less than 0.05.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Data for age not stated included in "all ages" but not distributed among age groups.

²Death rates for younger than 1 (based on population estimates) differ from infant mortality rates (based on live births); see Technical Notes.

³Included in selected categories above. For list of ICD-10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	3,279,857	1,719,250	1,560,607	275,684	155,240	120,444	23,613	12,721	10,892	89,591	46,137	43,454	411,934	219,538	192,396	4,592	2,461	2,131	2,448,093	1,267,526	1,180,567
Salmonella infections (A01-A02)	82	49	33	13	10	3	—	—	—	6	5	1	12	6	6	1	—	1	50	28	22
Shigellosis and amebiasis (A03,A06)	17	8	9	7	5	2	—	—	—	—	—	—	2	—	2	—	—	—	8	3	5
Certain other intestinal infections (A04,A07-A09)	7,380	3,030	4,350	625	284	341	57	21	36	182	73	109	765	338	427	12	6	6	5,674	2,273	3,401
Tuberculosis (A16-A19)	565	354	211	101	75	26	13	6	7	147	101	46	94	57	37	3	1	2	206	113	93
Respiratory tuberculosis (A16)	391	258	133	72	55	17	10	5	5	118	87	31	63	40	23	3	1	2	124	69	55
Other tuberculosis (A17-A19)	174	96	78	29	20	9	3	1	2	29	14	15	31	17	14	—	—	—	82	44	38
Whooping cough (A37)	2	1	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	2	1	1
Scarlet fever and erysipelas (A38,A46)	3	1	2	1	1	—	—	—	—	—	—	—	—	—	—	—	—	—	2	—	2
Meningococcal infection (A39)	27	16	11	9	6	3	—	—	—	1	—	1	10	4	6	—	—	—	6	5	1
Septicemia (A40-A41)	42,261	20,977	21,284	3,259	1,652	1,607	316	148	168	968	508	460	7,134	3,403	3,731	70	34	36	30,197	15,053	15,144
Syphilis (A50-A53)	61	37	24	11	10	1	3	2	1	4	3	1	22	9	13	1	1	—	18	11	7
Acute poliomyelitis (A80)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Arthropod-borne viral encephalitis (A83-A84, A85.2)	6	5	1	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	6	5	1
Measles (B05)	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—	—
Viral hepatitis (B15-B19)	3,107	1,944	1,163	485	303	182	53	24	29	191	111	80	417	268	149	12	10	2	1,899	1,198	701
Human immunodeficiency virus (HIV) disease (B20-B24)	4,941	3,802	1,139	751	614	137	41	30	11	50	48	2	2,401	1,674	727	5	3	2	1,613	1,371	242
Malaria (B50-B54)	8	7	1	—	—	—	—	—	—	—	—	—	4	3	1	—	—	—	4	4	—
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05, A20-A36,A42-A44, A48-A49,A54-A79, A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09, B25-B49,B55-B99,U07.1)	198,029	108,820	89,209	21,584	12,533	9,051	1,815	918	897	5,863	3,334	2,529	23,358	11,579	11,779	307	170	137	143,767	79,535	64,232
Malignant neoplasms (C00-C97)	608,371	319,336	289,035	47,021	24,146	22,875	3,137	1,497	1,640	19,864	10,023	9,841	70,698	35,442	35,256	849	400	449	462,834	245,676	217,158

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non–Hispanic, single race ³																					
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	11,884	8,434	3,450	706	510	196	67	41	26	489	336	153	1,171	853	318	26	18	8	9,324	6,593	2,731	
Malignant neoplasm of esophagus (C15)	16,016	12,699	3,317	848	676	172	73	53	20	351	272	79	1,200	842	358	18	16	2	13,424	10,760	2,664	
Malignant neoplasm of stomach (C16)	10,853	6,437	4,416	2,061	1,148	913	86	45	41	864	503	361	1,863	1,067	796	25	16	9	5,877	3,607	2,270	
Malignant neoplasms of colon, rectum and anus (C18–C21)	54,268	29,029	25,239	4,958	2,778	2,180	345	186	159	2,011	1,096	915	7,025	3,723	3,302	77	41	36	39,487	21,000	18,487	
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	28,937	18,884	10,053	4,047	2,580	1,467	234	126	108	1,667	1,070	597	3,535	2,376	1,159	67	47	20	19,152	12,512	6,640	
Malignant neoplasm of pancreas (C25)	48,323	25,186	23,137	3,895	1,958	1,937	241	102	139	1,634	773	861	5,867	2,841	3,026	64	27	37	36,314	19,310	17,004	
Malignant neoplasm of larynx (C32)	3,933	3,125	808	238	205	33	22	17	5	60	52	8	609	480	129	–	–	–	2,980	2,356	624	
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	131,982	69,413	62,569	5,822	3,282	2,540	672	299	373	3,920	2,150	1,770	14,046	7,737	6,309	167	79	88	106,506	55,422	51,084	
Malignant melanoma of skin (C43)	8,243	5,357	2,886	270	153	117	12	7	5	63	36	27	125	60	65	1	–	1	7,731	5,075	2,656	
Malignant neoplasm of breast (C50)	42,672	461	42,211	3,483	27	3,456	201	1	200	1,429	11	1,418	6,413	95	6,318	67	1	66	30,793	321	30,472	
Malignant neoplasm of cervix uteri (C53)	4,051	...	4,051	593	...	593	30	...	30	162	...	162	649	...	649	20	...	20	2,567	...	2,567	
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	12,764	...	12,764	1,153	...	1,153	56	...	56	444	...	444	2,671	...	2,671	32	...	32	8,326	...	8,326	
Malignant neoplasm of ovary (C56)	13,214	...	13,214	1,217	...	1,217	60	...	60	533	...	533	1,312	...	1,312	21	...	21	9,991	...	9,991	
Malignant neoplasm of prostate (C61)	33,363	33,363	...	2,475	2,475	...	133	133	...	714	714	...	5,543	5,543	...	33	33	...	24,274	24,274	...	
Malignant neoplasms of kidney and renal pelvis (C64–C65)	14,488	9,519	4,969	1,364	921	443	121	79	42	385	234	151	1,422	910	512	16	8	8	11,087	7,304	3,783	
Malignant neoplasm of bladder (C67)	17,334	12,460	4,874	819	586	233	57	35	22	357	252	105	1,313	833	480	16	12	4	14,693	10,688	4,005	

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	17,937	10,082	7,855	1,503	817	686	60	39	21	483	271	212	1,237	651	586	28	11	17	14,506	8,215	6,291
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	56,959	32,511	24,448	4,832	2,670	2,162	203	100	103	1,761	999	762	5,912	3,156	2,756	69	37	32	43,852	25,354	18,498
Hodgkin disease (C81)	1,056	643	413	150	94	56	3	2	1	24	15	9	99	66	33	–	–	–	770	463	307
Non-Hodgkin lymphoma (C82–C85)	19,794	11,300	8,494	1,753	965	788	65	33	32	740	430	310	1,439	823	616	22	10	12	15,667	8,976	6,691
Leukemia (C91–C95)	23,324	13,487	9,837	1,897	1,051	846	72	34	38	676	396	280	2,031	1,084	947	33	18	15	18,476	10,820	7,656
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	12,616	6,992	5,624	1,020	552	468	62	31	31	312	153	159	2,325	1,174	1,151	14	9	5	8,810	5,028	3,782
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	169	89	80	12	8	4	1	–	1	9	5	4	18	9	9	–	–	–	129	67	62
All other and unspecified malignant neoplasms (C17,C23–C24, C26–C31,C37–C41, C44–C49,C51–C52, C57–C60,C62–C63,C66, C68–C69,C73–C80,C97)	81,150	42,376	38,774	6,737	3,360	3,377	464	234	230	2,537	1,254	1,283	8,785	4,275	4,510	102	54	48	61,950	32,885	29,065
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	16,034	8,467	7,567	1,019	524	495	67	32	35	447	215	232	1,541	721	820	19	7	12	12,857	6,919	5,938
Anemias (D50–D64)	6,021	2,768	3,253	417	198	219	35	15	20	164	78	86	1,251	585	666	10	2	8	4,107	1,872	2,235
Diabetes mellitus (E10–E14)	101,209	57,557	43,652	12,508	7,033	5,475	1,219	659	560	3,709	1,982	1,727	17,954	9,371	8,583	303	160	143	64,701	37,886	26,815
Nutritional deficiencies (E40–E64)	21,020	7,760	13,260	1,190	468	722	121	44	77	469	168	301	2,053	834	1,219	13	3	10	17,082	6,202	10,880
Malnutrition (E40–E46)	20,552	7,555	12,997	1,161	456	705	114	40	74	464	167	297	2,008	816	1,192	13	3	10	16,703	6,032	10,671
Other nutritional deficiencies (E50–E64)	468	205	263	29	12	17	7	4	3	5	1	4	45	18	27	–	–	–	379	170	209

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Meningitis (G00,G03)	604	348	256	77	47	30	5	3	2	25	21	4	130	74	56	2	–	2	359	199	160
Parkinson disease (G20–G21)	39,915	24,503	15,412	2,508	1,430	1,078	113	63	50	1,232	742	490	1,919	1,093	826	19	13	6	33,969	21,065	12,904
Alzheimer disease (G30)	120,122	37,475	82,647	9,552	2,904	6,648	337	110	227	3,448	1,069	2,379	9,284	2,749	6,535	98	28	70	96,899	30,434	66,465
Major cardiovascular diseases (I00–I78)	936,436	492,088	444,348	67,463	36,990	30,473	4,838	2,629	2,209	27,175	14,066	13,109	122,434	64,109	58,325	1,418	776	642	706,141	369,268	336,873
Diseases of heart (I00–I09,I11,I13,I20–I51)	702,880	386,766	316,114	47,715	27,442	20,273	3,753	2,152	1,601	18,091	9,980	8,111	88,964	48,430	40,534	1,004	575	429	537,969	294,757	243,212
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	4,294	1,522	2,772	279	112	167	31	15	16	147	55	92	365	125	240	15	6	9	3,423	1,196	2,227
Hypertensive heart disease (I11)	72,070	36,143	35,927	5,020	2,738	2,282	502	278	224	1,728	820	908	12,084	6,633	5,451	91	54	37	51,974	25,215	26,759
Hypertensive heart and renal disease (I13)	16,091	7,521	8,570	1,000	492	508	140	66	74	376	173	203	2,555	1,282	1,273	23	9	14	11,873	5,420	6,453
Ischemic heart diseases (I20–I25)	371,506	223,952	147,554	27,594	16,840	10,754	1,973	1,211	762	10,956	6,538	4,418	43,103	24,839	18,264	551	335	216	284,345	172,181	112,164
Acute myocardial infarction (I21–I22)	103,905	62,571	41,334	7,805	4,664	3,141	572	346	226	3,133	1,855	1,278	11,888	6,695	5,193	164	90	74	79,750	48,545	31,205
Other acute ischemic heart diseases (I24)	4,823	2,756	2,067	257	165	92	35	20	15	111	66	45	785	477	308	4	3	1	3,594	2,003	1,591
Other forms of chronic ischemic heart disease (I20,I25)	262,778	158,625	104,153	19,532	12,011	7,521	1,366	845	521	7,712	4,617	3,095	30,430	17,667	12,763	383	242	141	201,001	121,633	79,368
Atherosclerotic cardiovascular disease, so described (I25.0)	81,672	52,180	29,492	7,027	4,794	2,233	540	340	200	2,336	1,468	868	12,755	8,093	4,662	118	81	37	57,732	36,571	21,161
All other forms of chronic ischemic heart disease. (I20,I25.1–I25.9)	181,106	106,445	74,661	12,505	7,217	5,288	826	505	321	5,376	3,149	2,227	17,675	9,574	8,101	265	161	104	143,269	85,062	58,207
Other heart diseases (I26–I51)	238,919	117,628	121,291	13,822	7,260	6,562	1,107	582	525	4,884	2,394	2,490	30,857	15,551	15,306	324	171	153	186,354	90,745	95,609
Acute and subacute endocarditis. (I33)	1,803	1,098	705	162	100	62	19	8	11	35	25	10	260	159	101	3	2	1	1,308	793	515

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of pericardium and acute myocarditis (I30–I31,I40)	1,237	634	603	125	64	61	12	4	8	36	25	11	204	93	111	2	1	1	847	439	408
Heart failure..... (I50)	87,941	41,657	46,284	4,912	2,434	2,478	354	187	167	1,718	835	883	10,476	5,101	5,375	94	52	42	69,972	32,820	37,152
All other forms of heart disease (I26–I28, I34–I38,I42–I49,I51)	147,938	74,239	73,699	8,623	4,662	3,961	722	383	339	3,095	1,509	1,586	19,917	10,198	9,719	225	116	109	114,227	56,693	57,534
Essential hypertension and hypertensive renal disease (I10,I12,I15)	43,293	20,237	23,056	3,806	1,902	1,904	219	102	117	1,903	868	1,035	7,673	3,750	3,923	68	37	31	29,296	13,393	15,903
Cerebrovascular diseases (I60–I69)	165,393	71,819	93,574	14,224	6,673	7,551	747	308	439	6,482	2,852	3,630	22,656	10,293	12,363	298	137	161	119,929	51,042	68,887
Atherosclerosis..... (I70)	3,936	1,889	2,047	288	144	144	13	9	4	132	54	78	373	186	187	4	2	2	3,095	1,477	1,618
Other diseases of circulatory system (I71–I78)	20,934	11,377	9,557	1,430	829	601	106	58	48	567	312	255	2,768	1,450	1,318	44	25	19	15,852	8,599	7,253
Aortic aneurysm and dissection (I71)	9,999	5,962	4,037	597	411	186	37	22	15	343	206	137	1,146	661	485	26	15	11	7,757	4,584	3,173
Other diseases of arteries, arterioles and capillaries (I72–I78)	10,935	5,415	5,520	833	418	415	69	36	33	224	106	118	1,622	789	833	18	10	8	8,095	4,015	4,080
Other disorders of circulatory system (I80–I99)	5,216	2,652	2,564	470	267	203	36	21	15	81	40	41	1,032	497	535	3	2	1	3,533	1,796	1,737
Influenza and pneumonia (J09–J18)	47,052	24,060	22,992	4,166	2,177	1,989	392	198	194	1,766	908	858	5,335	2,772	2,563	63	26	37	34,951	17,775	17,176
Influenza..... (J09–J11)	5,944	2,717	3,227	580	271	309	74	37	37	139	58	81	546	256	290	18	6	12	4,546	2,064	2,482
Pneumonia (J12–J18)	41,108	21,343	19,765	3,586	1,906	1,680	318	161	157	1,627	850	777	4,789	2,516	2,273	45	20	25	30,405	15,711	14,694
Other acute lower respiratory infections (J20–J22,U04)	278	128	150	34	18	16	2	1	1	9	5	4	38	21	17	1	1	–	191	80	111
Acute bronchitis and bronchiolitis (J20–J21)	182	89	93	27	15	12	1	1	–	7	5	2	32	18	14	–	–	–	114	49	65
Other and unspecified acute lower respiratory infections (J22,U04)	96	39	57	7	3	4	1	–	1	2	–	2	6	3	3	1	1	–	77	31	46
Chronic lower respiratory diseases (J40–J47)	147,382	69,004	78,378	5,615	2,796	2,819	761	347	414	1,785	1,058	727	10,990	5,454	5,536	109	58	51	127,003	58,720	68,283
Bronchitis, chronic and unspecified. (J40–J42)	358	139	219	20	8	12	3	1	2	12	4	8	33	10	23	–	–	–	288	115	173
Emphysema (J43)	7,905	4,165	3,740	253	155	98	44	23	21	134	96	38	514	298	216	5	4	1	6,888	3,547	3,341
Asthma (J45–J46)	3,602	1,415	2,187	383	162	221	35	12	23	121	53	68	1,012	472	540	9	4	5	1,972	684	1,288

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																					
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	
Other chronic lower respiratory diseases (J44,J47)	135,517	63,285	72,232	4,959	2,471	2,488	679	311	368	1,518	905	613	9,431	4,674	4,757	95	50	45	117,855	54,374	63,481	
Pneumoconioses and chemical effects (J60–J66, J68,U07.0)	562	521	41	20	19	1	7	7	–	2	2	–	17	14	3	–	–	–	512	475	37	
Pneumonitis due to solids and liquids. (J69)	20,052	11,751	8,301	1,342	733	609	91	49	42	700	395	305	2,322	1,314	1,008	18	13	5	15,435	9,153	6,282	
Other diseases of respiratory system (J00–J06, J30–J39,J67,J70–J98)	48,082	24,822	23,260	4,049	2,110	1,939	299	148	151	1,323	694	629	4,746	2,307	2,439	36	19	17	37,344	19,384	17,960	
Peptic ulcer (K25–K28)	4,105	2,191	1,914	358	211	147	32	17	15	163	82	81	431	255	176	5	3	2	3,079	1,597	1,482	
Diseases of appendix (K35–K38)	487	267	220	46	31	15	4	3	1	15	5	10	81	50	31	1	1	–	337	175	162	
Hernia (K40–K46)	2,712	1,223	1,489	245	114	131	24	9	15	41	18	23	209	108	101	1	1	–	2,172	960	1,212	
Chronic liver disease and cirrhosis. (K70,K73–K74)	54,803	34,340	20,463	8,933	6,067	2,866	1,579	851	728	910	585	325	4,017	2,454	1,563	46	31	15	38,779	24,008	14,771	
Alcoholic liver disease (K70)	30,910	21,059	9,851	5,126	4,018	1,108	1,202	685	517	415	314	101	2,218	1,423	795	26	20	6	21,553	14,357	7,196	
Other chronic liver disease and cirrhosis. (K73–K74)	23,893	13,281	10,612	3,807	2,049	1,758	377	166	211	495	271	224	1,799	1,031	768	20	11	9	17,226	9,651	7,575	
Cholelithiasis and other disorders of gallbladder (K80–K82)	4,483	2,287	2,196	475	252	223	43	19	24	163	79	84	449	191	258	6	2	4	3,326	1,730	1,596	
Nephritis, nephrotic syndrome and nephrosis (N00–N07, N17–N19,N25–N27)	57,937	30,178	27,759	5,299	2,808	2,491	434	227	207	1,985	1,041	944	10,921	5,349	5,572	121	60	61	38,813	20,470	18,343	
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01,N04)	810	429	381	65	34	31	4	2	2	28	19	9	126	63	63	2	1	1	581	307	274	
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified. (N02–N03, N05–N07,N26)	330	196	134	20	10	10	5	4	1	24	18	6	44	17	27	1	–	1	229	142	87	
Renal failure (N17–N19)	56,756	29,532	27,224	5,212	2,764	2,448	424	221	203	1,933	1,004	929	10,744	5,266	5,478	118	59	59	37,972	20,003	17,969	

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other disorders of kidney. (N25,N27)	41	21	20	2	—	2	1	—	1	—	—	—	7	3	4	—	—	—	31	18	13
Infections of kidney. (N10–N12,N13.6,N15.1)	1,205	437	768	129	39	90	14	5	9	20	6	14	115	50	65	1	—	1	918	334	584
Hyperplasia of prostate. (N40)	839	839	...	75	75	...	3	3	...	25	25	...	73	73	...	—	—	...	659	659	...
Inflammatory diseases of female pelvic organs (N70–N76)	232	...	232	29	...	29	3	...	3	10	...	10	40	...	40	—	...	—	148	...	148
Pregnancy, childbirth and the puerperium. (O00–O99)	1,263	...	1,263	227	...	227	26	...	26	46	...	46	395	...	395	5	...	5	551	...	551
Pregnancy with abortive outcome. (O00–O07)	33	...	33	7	...	7	2	...	2	1	...	1	11	...	11	1	...	1	11	...	11
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	1,230	...	1,230	220	...	220	24	...	24	45	...	45	384	...	384	4	...	4	540	...	540
Certain conditions originating in the perinatal period. (P00–P96)	10,203	5,695	4,508	2,402	1,346	1,056	82	48	34	365	207	158	2,976	1,687	1,289	41	22	19	3,826	2,098	1,728
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	10,149	5,276	4,873	1,865	959	906	82	42	40	251	137	114	1,530	797	733	24	8	16	6,147	3,208	2,939
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	34,110	17,133	16,977	2,954	1,810	1,144	307	197	110	762	383	379	5,299	2,865	2,434	49	26	23	24,235	11,538	12,697
All other diseases. . . . (residual)	410,973	179,851	231,122	28,956	13,645	15,311	2,873	1,363	1,510	9,208	3,911	5,297	45,588	20,363	25,225	440	215	225	321,231	138,958	182,273
Accidents (unintentional injuries). (V01–X59, Y85–Y86)	227,039	151,629	75,410	28,699	21,954	6,745	3,146	2,071	1,075	4,046	2,658	1,388	34,644	24,776	9,868	314	227	87	152,647	97,463	55,184
Transport accidents (V01–V99,Y85)	49,178	35,769	13,409	8,871	6,689	2,182	892	602	290	1,054	694	360	8,214	5,996	2,218	95	72	23	29,383	21,247	8,136

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motor vehicle accidents (V02-V04,V09.0, V09.2,V12-V14, V19.0-V19.2, V19.4-V19.6,V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1,V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0,V89.2)	46,027	33,274	12,753	8,424	6,330	2,094	844	569	275	979	632	347	7,822	5,671	2,151	88	68	20	27,241	19,565	7,676
Other land transport accidents. (V01, V05-V06,V09.1, V09.3-V09.9, V10-V11,V15-V18, V19.3,V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1,V89.3,V89.9)	1,195	951	244	245	202	43	24	16	8	34	29	5	169	144	25	2	1	1	703	547	156
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	1,956	1,544	412	202	157	45	24	17	7	41	33	8	223	181	42	5	3	2	1,439	1,135	304
Nontransport accidents (W00-X59,Y86)	177,861	115,860	62,001	19,828	15,265	4,563	2,254	1,469	785	2,992	1,964	1,028	26,430	18,780	7,650	219	155	64	123,264	76,216	47,048
Falls (W00-W19)	46,630	23,819	22,811	2,884	1,757	1,127	246	139	107	1,348	744	604	2,161	1,238	923	47	28	19	39,679	19,755	19,924
Accidental discharge of firearms (W32-W34)	463	409	54	59	51	8	13	11	2	2	1	1	125	111	14	3	3	—	253	225	28
Accidental drowning and submersion (W65-W74)	4,168	3,092	1,076	700	572	128	61	40	21	169	123	46	642	502	140	15	12	3	2,488	1,775	713

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental exposure to smoke, fire and flames (X00–X09)	3,478	2,079	1,399	241	157	84	35	23	12	52	25	27	676	397	279	1	–	1	2,428	1,445	983
Accidental poisoning and exposure to noxious substances (X40–X49)	102,958	73,988	28,970	14,137	11,433	2,704	1,665	1,093	572	1,002	822	180	19,979	14,792	5,187	117	93	24	63,851	44,176	19,675
Other and unspecified nontransport accidents and their sequelae (W20–W31, W35–W64,W75–W99, X10–X39,X50–X59,Y86)	20,164	12,473	7,691	1,807	1,295	512	234	163	71	419	249	170	2,847	1,740	1,107	36	19	17	14,565	8,840	5,725
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	49,476	39,273	10,203	5,122	4,128	994	650	475	175	1,459	1,025	434	3,826	3,069	757	95	75	20	37,481	29,861	7,620
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	27,032	23,538	3,494	2,074	1,847	227	256	222	34	372	326	46	2,237	1,936	301	24	20	4	21,732	18,895	2,837
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60–X71,X75–X84,Y87.0)	22,444	15,735	6,709	3,048	2,281	767	394	253	141	1,087	699	388	1,589	1,133	456	71	55	16	15,749	10,966	4,783
Assault (homicide) (*U01–*U02,X85–Y09,Y87.1)	24,849	19,977	4,872	4,482	3,684	798	386	302	84	310	201	109	13,236	11,352	1,884	52	44	8	5,980	4,097	1,883
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	19,651	16,428	3,223	3,500	2,977	523	224	185	39	196	137	59	11,575	10,127	1,448	37	34	3	3,830	2,735	1,095
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3, *U01.5–*U01.9,*U02, X85–X92,X96–Y09,Y87.1)	5,198	3,549	1,649	982	707	275	162	117	45	114	64	50	1,661	1,225	436	15	10	5	2,150	1,362	788
Legal intervention (Y35,Y89.0)	810	762	48	191	183	8	33	33	–	13	11	2	198	183	15	1	1	–	364	342	22
Events of undetermined intent (Y10–Y34,Y87.2,Y89.9)	5,604	3,666	1,938	552	404	148	103	70	33	107	72	35	1,340	931	409	7	6	1	3,390	2,122	1,268

See footnotes at end of table.

Table 9. Number of deaths from 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Discharge of firearms, undetermined intent (Y22–Y24)	415	312	103	60	48	12	9	6	3	3	2	1	124	102	22	–	–	–	214	151	63
Other and unspecified events of undetermined intent and their sequelae..... (Y10–Y21,Y25–Y34, Y87.2,Y89.9)	5,189	3,354	1,835	492	356	136	94	64	30	104	70	34	1,216	829	387	7	6	1	3,176	1,971	1,205
Operations of war and their sequelae..... (Y36,Y89.1)	7	7	–	1	1	–	–	–	–	–	–	–	–	–	–	–	–	–	6	6	–
Complications of medical and surgical care (Y40–Y84,Y88)	3,726	1,928	1,798	347	176	171	31	14	17	81	42	39	603	287	316	11	6	5	2,634	1,396	1,238
Enterocolitis due to <i>Clostridium</i> <i>difficile</i> (A04.7) ⁴	4,231	1,808	2,423	347	159	188	30	7	23	106	45	61	429	203	226	7	5	2	3,277	1,372	1,905
COVID-19..... (U07.1) ⁴	186,552	102,660	83,892	20,499	11,923	8,576	1,690	861	829	5,450	3,120	2,330	21,896	10,778	11,118	285	157	128	135,484	75,114	60,370
Dementia-related causes ⁴	292,881	97,470	195,411	19,213	6,209	13,004	897	300	597	7,591	2,463	5,128	23,599	7,687	15,912	202	70	132	240,132	80,270	159,862
Drug-induced deaths ⁴	112,109	78,614	33,495	14,573	11,563	3,010	1,641	1,040	601	1,184	909	275	21,455	15,749	5,706	137	104	33	70,734	47,595	23,139
Drug overdose deaths ⁴	107,941	75,814	32,127	14,131	11,228	2,903	1,543	980	563	1,142	875	267	20,725	15,222	5,503	125	96	29	67,974	45,811	22,163
Alcohol-induced deaths ⁴	51,191	36,426	14,765	7,467	6,028	1,439	1,904	1,180	724	660	539	121	4,339	3,025	1,314	44	34	10	36,099	25,139	10,960
Firearm-related injuries ⁴	48,204	41,302	6,902	5,853	5,081	772	530	452	78	583	474	109	14,203	12,410	1,793	65	58	7	26,323	22,285	4,038

– Quantity zero.

... Category not applicable.

¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes people of Hispanic origin of any race; see Technical Notes.

³Only one race reported on death certificate; see Technical Notes.

⁴Included in selected categories above. For list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	984.1	1,040.2	928.9	433.0	481.0	383.7	975.4	1,062.4	890.2	441.9	472.3	413.6	979.2	1,085.4	880.8	722.1	765.3	677.9	1,247.6	1,299.7	1,196.1
Salmonella infections (A01–A02)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	0.0
Shigellosis and amebiasis. (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07–A09)	2.2	1.8	2.6	1.0	0.9	1.1	2.4	1.8	2.9	0.9	0.7	1.0	1.8	1.7	2.0	*	*	*	2.9	2.3	3.4
Tuberculosis. (A16–A19)	0.2	0.2	0.1	0.2	0.2	0.1	*	*	*	0.7	1.0	0.4	0.2	0.3	0.2	*	*	*	0.1	0.1	0.1
Respiratory tuberculosis (A16)	0.1	0.2	0.1	0.1	0.2	*	*	*	*	0.6	0.9	0.3	0.1	0.2	0.1	*	*	*	0.1	0.1	0.1
Other tuberculosis (A17–A19)	0.1	0.1	0.0	0.0	0.1	*	*	*	*	0.1	*	*	0.1	*	*	*	*	*	0.0	0.0	0.0
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas. (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection. (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Septicemia. (A40–A41)	12.7	12.7	12.7	5.1	5.1	5.1	13.1	12.4	13.7	4.8	5.2	4.4	17.0	16.8	17.1	11.0	10.6	11.5	15.4	15.4	15.3
Syphilis (A50–A53)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	*	*	*
Acute poliomyelitis. (A80)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83–A84, A85.2)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis. (B15–B19)	0.9	1.2	0.7	0.8	0.9	0.6	2.2	2.0	2.4	0.9	1.1	0.8	1.0	1.3	0.7	*	*	*	1.0	1.2	0.7
Human immunodeficiency virus (HIV) disease (B20–B24)	1.5	2.3	0.7	1.2	1.9	0.4	1.7	2.5	*	0.2	0.5	*	5.7	8.3	3.3	*	*	*	0.8	1.4	0.2
Malaria. (B50–B54)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae. (A00,A05, A20–A36,A42–A44, A48–A49,A54–A79, A81–A82,A85.0–A85.1, A85.8,A86–B04,B06–B09, B25–B49,B55–B99,U07.1)	59.4	65.8	53.1	33.9	38.8	28.8	75.0	76.7	73.3	28.9	34.1	24.1	55.5	57.2	53.9	48.3	52.9	43.6	73.3	81.6	65.1
Malignant neoplasms (C00–C97)	182.5	193.2	172.0	73.9	74.8	72.9	129.6	125.0	134.0	98.0	102.6	93.7	168.0	175.2	161.4	133.5	124.4	142.8	235.9	251.9	220.0

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	3.6	5.1	2.1	1.1	1.6	0.6	2.8	3.4	2.1	2.4	3.4	1.5	2.8	4.2	1.5	4.1	*	*	4.8	6.8	2.8
Malignant neoplasm of esophagus (C15)	4.8	7.7	2.0	1.3	2.1	0.5	3.0	4.4	1.6	1.7	2.8	0.8	2.9	4.2	1.6	*	*	*	6.8	11.0	2.7
Malignant neoplasm of stomach (C16)	3.3	3.9	2.6	3.2	3.6	2.9	3.6	3.8	3.4	4.3	5.1	3.4	4.4	5.3	3.6	3.9	*	*	3.0	3.7	2.3
Malignant neoplasms of colon, rectum and anus (C18–C21)	16.3	17.6	15.0	7.8	8.6	6.9	14.3	15.5	13.0	9.9	11.2	8.7	16.7	18.4	15.1	12.1	12.7	11.5	20.1	21.5	18.7
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	8.7	11.4	6.0	6.4	8.0	4.7	9.7	10.5	8.8	8.2	11.0	5.7	8.4	11.7	5.3	10.5	14.6	6.4	9.8	12.8	6.7
Malignant neoplasm of pancreas (C25)	14.5	15.2	13.8	6.1	6.1	6.2	10.0	8.5	11.4	8.1	7.9	8.2	13.9	14.0	13.9	10.1	8.4	11.8	18.5	19.8	17.2
Malignant neoplasm of larynx (C32)	1.2	1.9	0.5	0.4	0.6	0.1	0.9	*	*	0.3	0.5	*	1.4	2.4	0.6	*	*	*	1.5	2.4	0.6
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	39.6	42.0	37.2	9.1	10.2	8.1	27.8	25.0	30.5	19.3	22.0	16.8	33.4	38.3	28.9	26.3	24.6	28.0	54.3	56.8	51.8
Malignant melanoma of skin (C43)	2.5	3.2	1.7	0.4	0.5	0.4	*	*	*	0.3	0.4	0.3	0.3	0.3	0.3	*	*	*	3.9	5.2	2.7
Malignant neoplasm of breast (C50)	12.8	0.3	25.1	5.5	0.1	11.0	8.3	*	16.3	7.0	*	13.5	15.2	0.5	28.9	10.5	*	21.0	15.7	0.3	30.9
Malignant neoplasm of cervix uteri (C53)	1.2	...	2.4	0.9	...	1.9	1.2	...	2.5	0.8	...	1.5	1.5	...	3.0	3.1	...	6.4	1.3	...	2.6
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	3.8	...	7.6	1.8	...	3.7	2.3	...	4.6	2.2	...	4.2	6.3	...	12.2	5.0	...	10.2	4.2	...	8.4
Malignant neoplasm of ovary (C56)	4.0	...	7.9	1.9	...	3.9	2.5	...	4.9	2.6	...	5.1	3.1	...	6.0	3.3	...	6.7	5.1	...	10.1
Malignant neoplasm of prostate (C61)	10.0	20.2	...	3.9	7.7	...	5.5	11.1	...	3.5	7.3	...	13.2	27.4	...	5.2	10.3	...	12.4	24.9	...
Malignant neoplasms of kidney and renal pelvis (C64–C65)	4.3	5.8	3.0	2.1	2.9	1.4	5.0	6.6	3.4	1.9	2.4	1.4	3.4	4.5	2.3	*	*	*	5.7	7.5	3.8
Malignant neoplasm of bladder (C67)	5.2	7.5	2.9	1.3	1.8	0.7	2.4	2.9	1.8	1.8	2.6	1.0	3.1	4.1	2.2	*	*	*	7.5	11.0	4.1

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	5.4	6.1	4.7	2.4	2.5	2.2	2.5	3.3	1.7	2.4	2.8	2.0	2.9	3.2	2.7	4.4	*	*	7.4	8.4	6.4
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	17.1	19.7	14.6	7.6	8.3	6.9	8.4	8.4	8.4	8.7	10.2	7.3	14.1	15.6	12.6	10.9	11.5	10.2	22.3	26.0	18.7
Hodgkin disease (C81)	0.3	0.4	0.2	0.2	0.3	0.2	*	*	*	0.1	*	*	0.2	0.3	0.2	*	*	*	0.4	0.5	0.3
Non-Hodgkin lymphom (C82–C85)	5.9	6.8	5.1	2.8	3.0	2.5	2.7	2.8	2.6	3.6	4.4	3.0	3.4	4.1	2.8	3.5	*	*	8.0	9.2	6.8
Leukemia (C91–C95)	7.0	8.2	5.9	3.0	3.3	2.7	3.0	2.8	3.1	3.3	4.1	2.7	4.8	5.4	4.3	5.2	*	*	9.4	11.1	7.8
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	3.8	4.2	3.3	1.6	1.7	1.5	2.6	2.6	2.5	1.5	1.6	1.5	5.5	5.8	5.3	*	*	*	4.5	5.2	3.8
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.1	0.1	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.1	0.1	0.1
All other and unspecified malignant neoplasms (C17,C23–C24, C26–C31,C37–C41, C44–C49,C51–C52, C57–C60,C62–C63,C66, C68–C69,C73–C80,C97)	24.3	25.6	23.1	10.6	10.4	10.8	19.2	19.5	18.8	12.5	12.8	12.2	20.9	21.1	20.6	16.0	16.8	15.3	31.6	33.7	29.4
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00–D48)	4.8	5.1	4.5	1.6	1.6	1.6	2.8	2.7	2.9	2.2	2.2	2.2	3.7	3.6	3.8	*	*	*	6.6	7.1	6.0
Anemias (D50–D64)	1.8	1.7	1.9	0.7	0.6	0.7	1.4	*	1.6	0.8	0.8	0.8	3.0	2.9	3.0	*	*	*	2.1	1.9	2.3
Diabetes mellitus (E10–E14)	30.4	34.8	26.0	19.6	21.8	17.4	50.4	55.0	45.8	18.3	20.3	16.4	42.7	46.3	39.3	47.6	49.8	45.5	33.0	38.8	27.2
Nutritional deficiencies (E40–E64)	6.3	4.7	7.9	1.9	1.4	2.3	5.0	3.7	6.3	2.3	1.7	2.9	4.9	4.1	5.6	*	*	*	8.7	6.4	11.0
Malnutrition (E40–E46)	6.2	4.6	7.7	1.8	1.4	2.2	4.7	3.3	6.0	2.3	1.7	2.8	4.8	4.0	5.5	*	*	*	8.5	6.2	10.8
Other nutritional deficiencies (E50–E64)	0.1	0.1	0.2	0.0	*	*	*	*	*	*	*	*	0.1	*	0.1	*	*	*	0.2	0.2	0.2

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Meningitis (G00,G03)	0.2	0.2	0.2	0.1	0.1	0.1	*	*	*	0.1	0.2	*	0.3	0.4	0.3	*	*	*	0.2	0.2	0.2
Parkinson disease (G20–G21)	12.0	14.8	9.2	3.9	4.4	3.4	4.7	5.3	4.1	6.1	7.6	4.7	4.6	5.4	3.8	*	*	*	17.3	21.6	13.1
Alzheimer disease (G30)	36.0	22.7	49.2	15.0	9.0	21.2	13.9	9.2	18.6	17.0	10.9	22.6	22.1	13.6	29.9	15.4	8.7	22.3	49.4	31.2	67.3
Major cardiovascular diseases (I00–I78)	281.0	297.7	264.5	106.0	114.6	97.1	199.8	219.6	180.5	134.0	144.0	124.8	291.0	317.0	267.0	223.0	241.3	204.2	359.9	378.7	341.3
Diseases of heart (I00–I09,I11,I13,I20–I51)	210.9	234.0	188.2	74.9	85.0	64.6	155.0	179.7	130.8	89.2	102.2	77.2	211.5	239.4	185.6	157.9	178.8	136.5	274.2	302.2	246.4
Acute rheumatic fever and chronic rheumatic heart diseases (I00–I09)	1.3	0.9	1.6	0.4	0.3	0.5	1.3	*	*	0.7	0.6	0.9	0.9	0.6	1.1	*	*	*	1.7	1.2	2.3
Hypertensive heart disease (I11)	21.6	21.9	21.4	7.9	8.5	7.3	20.7	23.2	18.3	8.5	8.4	8.6	28.7	32.8	25.0	14.3	16.8	11.8	26.5	25.9	27.1
Hypertensive heart and renal disease (I13)	4.8	4.6	5.1	1.6	1.5	1.6	5.8	5.5	6.0	1.9	1.8	1.9	6.1	6.3	5.8	3.6	*	*	6.1	5.6	6.5
Ischemic heart diseases (I20–I25)	111.5	135.5	87.8	43.3	52.2	34.3	81.5	101.1	62.3	54.0	66.9	42.0	102.5	122.8	83.6	86.6	104.2	68.7	144.9	176.6	113.6
Acute myocardial infarction (I21–I22)	31.2	37.9	24.6	12.3	14.4	10.0	23.6	28.9	18.5	15.5	19.0	12.2	28.3	33.1	23.8	25.8	28.0	23.5	40.6	49.8	31.6
Other acute ischemic heart diseases (I24)	1.4	1.7	1.2	0.4	0.5	0.3	1.4	1.7	*	0.5	0.7	0.4	1.9	2.4	1.4	*	*	*	1.8	2.1	1.6
Other forms of chronic ischemic heart disease (I20,I25)	78.8	96.0	62.0	30.7	37.2	24.0	56.4	70.6	42.6	38.0	47.3	29.5	72.3	87.3	58.4	60.2	75.3	44.9	102.4	124.7	80.4
Atherosclerotic cardiovascular disease, so described (I25.0)	24.5	31.6	17.6	11.0	14.9	7.1	22.3	28.4	16.3	11.5	15.0	8.3	30.3	40.0	21.3	18.6	25.2	11.8	29.4	37.5	21.4
All other forms of chronic ischemic heart disease (I20,I25.1–I25.9)	54.3	64.4	44.4	19.6	22.4	16.8	34.1	42.2	26.2	26.5	32.2	21.2	42.0	47.3	37.1	41.7	50.1	33.1	73.0	87.2	59.0
Other heart diseases (I26–I51)	71.7	71.2	72.2	21.7	22.5	20.9	45.7	48.6	42.9	24.1	24.5	23.7	73.3	76.9	70.1	50.9	53.2	48.7	95.0	93.1	96.9
Acute and subacute endocarditis (I33)	0.5	0.7	0.4	0.3	0.3	0.2	*	*	*	0.2	0.3	*	0.6	0.8	0.5	*	*	*	0.7	0.8	0.5

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Diseases of pericardium and acute myocarditis (I30–I31,I40)	0.4	0.4	0.4	0.2	0.2	0.2	*	*	*	0.2	0.3	*	0.5	0.5	0.5	*	*	*	0.4	0.5	0.4
Heart failure..... (I50)	26.4	25.2	27.5	7.7	7.5	7.9	14.6	15.6	13.6	8.5	8.5	8.4	24.9	25.2	24.6	14.8	16.2	13.4	35.7	33.7	37.6
All other forms of heart disease (I26–I28, I34–I38,I42–I49,I51)	44.4	44.9	43.9	13.5	14.4	12.6	29.8	32.0	27.7	15.3	15.4	15.1	47.3	50.4	44.5	35.4	36.1	34.7	58.2	58.1	58.3
Essential hypertension and hypertensive renal disease (I10,I12,I15)	13.0	12.2	13.7	6.0	5.9	6.1	9.0	8.5	9.6	9.4	8.9	9.9	18.2	18.5	18.0	10.7	11.5	9.9	14.9	13.7	16.1
Cerebrovascular diseases (I60–I69)	49.6	43.5	55.7	22.3	20.7	24.1	30.9	25.7	35.9	32.0	29.2	34.5	53.9	50.9	56.6	46.9	42.6	51.2	61.1	52.3	69.8
Atherosclerosis..... (I70)	1.2	1.1	1.2	0.5	0.4	0.5	*	*	*	0.7	0.6	0.7	0.9	0.9	0.9	*	*	*	1.6	1.5	1.6
Other diseases of circulatory system (I71–I78)	6.3	6.9	5.7	2.2	2.6	1.9	4.4	4.8	3.9	2.8	3.2	2.4	6.6	7.2	6.0	6.9	7.8	*	8.1	8.8	7.3
Aortic aneurysm and dissection (I71)	3.0	3.6	2.4	0.9	1.3	0.6	1.5	1.8	*	1.7	2.1	1.3	2.7	3.3	2.2	4.1	*	*	4.0	4.7	3.2
Other diseases of arteries, arterioles and capillaries (I72–I78)	3.3	3.3	3.3	1.3	1.3	1.3	2.9	3.0	2.7	1.1	1.1	1.1	3.9	3.9	3.8	*	*	*	4.1	4.1	4.1
Other disorders of circulatory system (I80–I99)	1.6	1.6	1.5	0.7	0.8	0.6	1.5	1.8	*	0.4	0.4	0.4	2.5	2.5	2.4	*	*	*	1.8	1.8	1.8
Influenza and pneumonia (J09–J18)	14.1	14.6	13.7	6.5	6.7	6.3	16.2	16.5	15.9	8.7	9.3	8.2	12.7	13.7	11.7	9.9	8.1	11.8	17.8	18.2	17.4
Influenza..... (J09–J11)	1.8	1.6	1.9	0.9	0.8	1.0	3.1	3.1	3.0	0.7	0.6	0.8	1.3	1.3	1.3	*	*	*	2.3	2.1	2.5
Pneumonia..... (J12–J18)	12.3	12.9	11.8	5.6	5.9	5.4	13.1	13.4	12.8	8.0	8.7	7.4	11.4	12.4	10.4	7.1	6.2	8.0	15.5	16.1	14.9
Other acute lower respiratory infections (J20–J22,U04)	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	0.1	0.1	*	*	*	*	0.1	0.1	0.1
Acute bronchitis and bronchiolitis . . . (J20–J21)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	*	*	*	*	*	0.1	0.1	0.1
Other and unspecified acute lower respiratory infections (J22,U04)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	0.0
Chronic lower respiratory diseases..... (J40–J47)	44.2	41.7	46.7	8.8	8.7	9.0	31.4	29.0	33.8	8.8	10.8	6.9	26.1	27.0	25.3	17.1	18.0	16.2	64.7	60.2	69.2
Bronchitis, chronic and unspecified..... (J40–J42)	0.1	0.1	0.1	0.0	*	*	*	*	*	*	*	*	0.1	*	0.1	*	*	*	0.1	0.1	0.2
Emphysema (J43)	2.4	2.5	2.2	0.4	0.5	0.3	1.8	1.9	1.7	0.7	1.0	0.4	1.2	1.5	1.0	*	*	*	3.5	3.6	3.4
Asthma..... (J45–J46)	1.1	0.9	1.3	0.6	0.5	0.7	1.4	*	1.9	0.6	0.5	0.6	2.4	2.3	2.5	*	*	*	1.0	0.7	1.3

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other chronic lower respiratory diseases (J44,J47)	40.7	38.3	43.0	7.8	7.7	7.9	28.0	26.0	30.1	7.5	9.3	5.8	22.4	23.1	21.8	14.9	15.5	14.3	60.1	55.8	64.3
Pneumoconioses and chemical effects (J60–J66, J68,U07.0)	0.2	0.3	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.3	0.5	0.0
Pneumonitis due to solids and liquids..... (J69)	6.0	7.1	4.9	2.1	2.3	1.9	3.8	4.1	3.4	3.5	4.0	2.9	5.5	6.5	4.6	*	*	*	7.9	9.4	6.4
Other diseases of respiratory system..... (J00–J06, J30–J39,J67,J70–J98)	14.4	15.0	13.8	6.4	6.5	6.2	12.4	12.4	12.3	6.5	7.1	6.0	11.3	11.4	11.2	5.7	*	*	19.0	19.9	18.2
Peptic ulcer (K25–K28)	1.2	1.3	1.1	0.6	0.7	0.5	1.3	*	*	0.8	0.8	0.8	1.0	1.3	0.8	*	*	*	1.6	1.6	1.5
Diseases of appendix (K35–K38)	0.1	0.2	0.1	0.1	0.1	*	*	*	*	*	*	*	0.2	0.2	0.1	*	*	*	0.2	0.2	0.2
Hernia (K40–K46)	0.8	0.7	0.9	0.4	0.4	0.4	1.0	*	*	0.2	*	0.2	0.5	0.5	0.5	*	*	*	1.1	1.0	1.2
Chronic liver disease and cirrhosis..... (K70,K73–K74)	16.4	20.8	12.2	14.0	18.8	9.1	65.2	71.1	59.5	4.5	6.0	3.1	9.5	12.1	7.2	7.2	9.6	*	19.8	24.6	15.0
Alcoholic liver disease..... (K70)	9.3	12.7	5.9	8.1	12.4	3.5	49.6	57.2	42.3	2.0	3.2	1.0	5.3	7.0	3.6	4.1	6.2	*	11.0	14.7	7.3
Other chronic liver disease and cirrhosis..... (K73–K74)	7.2	8.0	6.3	6.0	6.3	5.6	15.6	13.9	17.2	2.4	2.8	2.1	4.3	5.1	3.5	3.1	*	*	8.8	9.9	7.7
Cholelithiasis and other disorders of gallbladder..... (K80–K82)	1.3	1.4	1.3	0.7	0.8	0.7	1.8	*	2.0	0.8	0.8	0.8	1.1	0.9	1.2	*	*	*	1.7	1.8	1.6
Nephritis, nephrotic syndrome and nephrosis (N00–N07, N17–N19,N25–N27)	17.4	18.3	16.5	8.3	8.7	7.9	17.9	19.0	16.9	9.8	10.7	9.0	26.0	26.4	25.5	19.0	18.7	19.4	19.8	21.0	18.6
Acute and rapidly progressive nephritic and nephrotic syndrome (N00–N01,N04)	0.2	0.3	0.2	0.1	0.1	0.1	*	*	*	0.1	*	*	0.3	0.3	0.3	*	*	*	0.3	0.3	0.3
Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified. (N02–N03, N05–N07,N26)	0.1	0.1	0.1	0.0	*	*	*	*	*	0.1	*	*	0.1	*	0.1	*	*	*	0.1	0.1	0.1
Renal failure (N17–N19)	17.0	17.9	16.2	8.2	8.6	7.8	17.5	18.5	16.6	9.5	10.3	8.8	25.5	26.0	25.1	18.6	18.3	18.8	19.4	20.5	18.2

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD–10)*; see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Other disorders of kidney. (N25,N27)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	*	*
Infections of kidney. (N10–N12,N13.6,N15.1)	0.4	0.3	0.5	0.2	0.1	0.3	*	*	*	0.1	*	*	0.3	0.2	0.3	*	*	*	0.5	0.3	0.6
Hyperplasia of prostate. (N40)	0.3	0.5	...	0.1	0.2	...	*	*	...	0.1	0.3	...	0.2	0.4	...	*	*	...	0.3	0.7	...
Inflammatory diseases of female pelvic organs (N70–N76)	0.1	...	0.1	0.0	...	0.1	*	...	*	*	...	*	0.1	...	0.2	*	...	*	0.1	...	0.1
Pregnancy, childbirth and the puerperium. (O00–O99)	0.4	...	0.8	0.4	...	0.7	1.1	...	2.1	0.2	...	0.4	0.9	...	1.8	*	...	*	0.3	...	0.6
Pregnancy with abortive outcome. (O00–O07)	0.0	...	0.0	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*	*	...	*
Other complications of pregnancy, childbirth and the puerperium (O10–O99)	0.4	...	0.7	0.3	...	0.7	1.0	...	2.0	0.2	...	0.4	0.9	...	1.8	*	...	*	0.3	...	0.5
Certain conditions originating in the perinatal period. (P00–P96)	3.1	3.4	2.7	3.8	4.2	3.4	3.4	4.0	2.8	1.8	2.1	1.5	7.1	8.3	5.9	6.4	6.8	*	1.9	2.2	1.8
Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99)	3.0	3.2	2.9	2.9	3.0	2.9	3.4	3.5	3.3	1.2	1.4	1.1	3.6	3.9	3.4	3.8	*	*	3.1	3.3	3.0
Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00–R99)	10.2	10.4	10.1	4.6	5.6	3.6	12.7	16.5	9.0	3.8	3.9	3.6	12.6	14.2	11.1	7.7	8.1	7.3	12.4	11.8	12.9
All other diseases. (residual)	123.3	108.8	137.6	45.5	42.3	48.8	118.7	113.8	123.4	45.4	40.0	50.4	108.4	100.7	115.5	69.2	66.9	71.6	163.7	142.5	184.7
Accidents (unintentional injuries). (V01–X59, Y85–Y86)	68.1	91.7	44.9	45.1	68.0	21.5	129.9	173.0	87.9	20.0	27.2	13.2	82.3	122.5	45.2	49.4	70.6	27.7	77.8	99.9	55.9
Transport accidents (V01–V99,Y85)	14.8	21.6	8.0	13.9	20.7	7.0	36.8	50.3	23.7	5.2	7.1	3.4	19.5	29.6	10.2	14.9	22.4	7.3	15.0	21.8	8.2

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Motor vehicle accidents (V02-V04,V09.0, V09.2,V12-V14, V19.0-V19.2, V19.4-V19.6,V20-V79, V80.3-V80.5, V81.0-V81.1, V82.0-V82.1,V83-V86, V87.0-V87.8, V88.0-V88.8, V89.0,V89.2)	13.8	20.1	7.6	13.2	19.6	6.7	34.9	47.5	22.5	4.8	6.5	3.3	18.6	28.0	9.8	13.8	21.1	6.4	13.9	20.1	7.8
Other land transport accidents..... (V01, V05-V06,V09.1, V09.3-V09.9, V10-V11,V15-V18, V19.3,V19.8-V19.9, V80.0-V80.2, V80.6-V80.9, V81.2-V81.9, V82.2-V82.9, V87.9, V88.9, V89.1,V89.3,V89.9)	0.4	0.6	0.1	0.4	0.6	0.1	1.0	*	*	0.2	0.3	*	0.4	0.7	0.1	*	*	*	0.4	0.6	0.2
Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)	0.6	0.9	0.2	0.3	0.5	0.1	1.0	*	*	0.2	0.3	*	0.5	0.9	0.2	*	*	*	0.7	1.2	0.3
Nontransport accidents (W00-X59,Y86)	53.4	70.1	36.9	31.1	47.3	14.5	93.1	122.7	64.2	14.8	20.1	9.8	62.8	92.8	35.0	34.4	48.2	20.4	62.8	78.2	47.7
Falls (W00-W19)	14.0	14.4	13.6	4.5	5.4	3.6	10.2	11.6	8.7	6.6	7.6	5.7	5.1	6.1	4.2	7.4	8.7	*	20.2	20.3	20.2
Accidental discharge of firearms (W32-W34)	0.1	0.2	0.0	0.1	0.2	*	*	*	*	*	*	*	0.3	0.5	*	*	*	*	0.1	0.2	0.0
Accidental drowning and submersion (W65-W74)	1.3	1.9	0.6	1.1	1.8	0.4	2.5	3.3	1.7	0.8	1.3	0.4	1.5	2.5	0.6	*	*	*	1.3	1.8	0.7

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision (ICD-10)*; see Technical Notes in this report]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Accidental exposure to smoke, fire and flames. (X00–X09)	1.0	1.3	0.8	0.4	0.5	0.3	1.4	1.9	*	0.3	0.3	0.3	1.6	2.0	1.3	*	*	*	1.2	1.5	1.0
Accidental poisoning and exposure to noxious substances (X40–X49)	30.9	44.8	17.2	22.2	35.4	8.6	68.8	91.3	46.7	4.9	8.4	1.7	47.5	73.1	23.7	18.4	28.9	7.6	32.5	45.3	19.9
Other and unspecified nontransport accidents and their sequelae (W20–W31, W35–W64,W75–W99, X10–X39,X50–X59,Y86)	6.1	7.5	4.6	2.8	4.0	1.6	9.7	13.6	5.8	2.1	2.5	1.6	6.8	8.6	5.1	5.7	*	*	7.4	9.1	5.8
Intentional self-harm (suicide) (*U03,X60–X84,Y87.0)	14.8	23.8	6.1	8.0	12.8	3.2	26.8	39.7	14.3	7.2	10.5	4.1	9.1	15.2	3.5	14.9	23.3	6.4	19.1	30.6	7.7
Intentional self-harm (suicide) by discharge of firearms (X72–X74)	8.1	14.2	2.1	3.3	5.7	0.7	10.6	18.5	2.8	1.8	3.3	0.4	5.3	9.6	1.4	3.8	6.2	*	11.1	19.4	2.9
Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03, X60–X71,X75–X84,Y87.0)	6.7	9.5	4.0	4.8	7.1	2.4	16.3	21.1	11.5	5.4	7.2	3.7	3.8	5.6	2.1	11.2	17.1	*	8.0	11.2	4.8
Assault (homicide). (*U01–*U02,X85–Y09,Y87.1)	7.5	12.1	2.9	7.0	11.4	2.5	15.9	25.2	6.9	1.5	2.1	1.0	31.5	56.1	8.6	8.2	13.7	*	3.0	4.2	1.9
Assault (homicide) by discharge of firearms (*U01.4,X93–X95)	5.9	9.9	1.9	5.5	9.2	1.7	9.3	15.5	3.2	1.0	1.4	0.6	27.5	50.1	6.6	5.8	10.6	*	2.0	2.8	1.1
Assault (homicide) by other and unspecified means and their sequelae (*U01.0–*U01.3, *U01.5–*U01.9,*U02, X85–X92,X96–Y09,Y87.1)	1.6	2.1	1.0	1.5	2.2	0.9	6.7	9.8	3.7	0.6	0.7	0.5	3.9	6.1	2.0	*	*	*	1.1	1.4	0.8
Legal intervention (Y35,Y89.0)	0.2	0.5	0.0	0.3	0.6	*	1.4	2.8	*	*	*	*	0.5	0.9	*	*	*	*	0.2	0.4	0.0
Events of undetermined intent (Y10–Y34,Y87.2,Y89.9)	1.7	2.2	1.2	0.9	1.3	0.5	4.3	5.8	2.7	0.5	0.7	0.3	3.2	4.6	1.9	*	*	*	1.7	2.2	1.3

See footnotes at end of table.

Table 10. Death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Rates are on an annual basis per 100,000 population in specified group; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes in this report]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Discharge of firearms, undetermined intent (Y22–Y24)	0.1	0.2	0.1	0.1	0.1	*	*	*	*	*	*	*	0.3	0.5	0.1	*	*	*	0.1	0.2	0.1
Other and unspecified events of undetermined intent and their sequelae..... (Y10–Y21,Y25–Y34, Y87.2,Y89.9)	1.6	2.0	1.1	0.8	1.1	0.4	3.9	5.3	2.5	0.5	0.7	0.3	2.9	4.1	1.8	*	*	*	1.6	2.0	1.2
Operations of war and their sequelae..... (Y36,Y89.1)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Complications of medical and surgical care (Y40–Y84,Y88)	1.1	1.2	1.1	0.5	0.5	0.5	1.3	*	*	0.4	0.4	0.4	1.4	1.4	1.4	*	*	*	1.3	1.4	1.3
Enterocolitis due to <i>Clostridium difficile</i> (A04.7) ⁴	1.3	1.1	1.4	0.5	0.5	0.6	1.2	*	1.9	0.5	0.5	0.6	1.0	1.0	1.0	*	*	*	1.7	1.4	1.9
COVID-19..... (U07.1) ⁴	56.0	62.1	49.9	32.2	36.9	27.3	69.8	71.9	67.8	26.9	31.9	22.2	52.0	53.3	50.9	44.8	48.8	40.7	69.0	77.0	61.2
Dementia-related causes ⁴	87.9	59.0	116.3	30.2	19.2	41.4	37.1	25.1	48.8	37.4	25.2	48.8	56.1	38.0	72.8	31.8	21.8	42.0	122.4	82.3	162.0
Drug-induced deaths ⁴	33.6	47.6	19.9	22.9	35.8	9.6	67.8	86.9	49.1	5.8	9.3	2.6	51.0	77.9	26.1	21.5	32.3	10.5	36.0	48.8	23.4
Drug overdose deaths ⁴	32.4	45.9	19.1	22.2	34.8	9.2	63.7	81.8	46.0	5.6	9.0	2.5	49.3	75.3	25.2	19.7	29.9	9.2	34.6	47.0	22.5
Alcohol-induced deaths ⁴	15.4	22.0	8.8	11.7	18.7	4.6	78.6	98.5	59.2	3.3	5.5	1.2	10.3	15.0	6.0	6.9	10.6	*	18.4	25.8	11.1
Firearm-related injuries ⁴	14.5	25.0	4.1	9.2	15.7	2.5	21.9	37.7	6.4	2.9	4.9	1.0	33.8	61.4	8.2	10.2	18.0	*	13.4	22.9	4.1

0.0 Quantity more than zero but less than 0.05.

* Estimate does not meet National Center for Health Statistics standards of reliability; see Technical Notes.

... Category not applicable.

¹Includes deaths with origin not stated, origin not classifiable, and two or more races reported; see Technical Notes.

²Includes people of Hispanic origin of any race; see Technical Notes.

³Only one race was reported on the death certificate; see Technical Notes.

⁴Included in selected categories above. For list of ICD–10 codes included, see Technical Notes.

SOURCE: National Center for Health Statistics, National Vital Statistics System, mortality data file.

Table 11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022

[Age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD–10); see Technical Notes]

Cause of death (based on ICD–10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
All causes	798.8	954.5	666.1	614.7	748.8	498.4	947.9	1,084.3	816.1	417.5	501.7	350.7	1,002.8	1,257.5	809.0	782.0	859.5	705.4	822.2	972.1	692.7
Salmonella infections (A01–A02)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	0.0
Shigellosis and amebiasis (A03,A06)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Certain other intestinal infections (A04,A07–A09)	1.8	1.7	1.8	1.4	1.5	1.4	2.3	1.7	2.6	0.9	0.8	0.9	1.9	2.1	1.8	*	*	*	1.8	1.7	1.9
Tuberculosis (A16–A19)	0.1	0.2	0.1	0.2	0.4	0.1	*	*	*	0.7	1.1	0.4	0.2	0.3	0.1	*	*	*	0.1	0.1	0.1
Respiratory tuberculosis (A16)	0.1	0.1	0.0	0.2	0.3	*	*	*	*	0.5	0.9	0.2	0.1	0.2	0.1	*	*	*	0.0	0.0	0.0
Other tuberculosis (A17–A19)	0.0	0.0	0.0	0.1	0.1	*	*	*	*	0.1	*	*	0.1	*	*	*	*	*	0.0	0.0	0.0
Whooping cough (A37)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Scarlet fever and erysipelas (A38,A46)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Meningococcal infection (A39)	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Septicemia (A40–A41)	10.1	11.4	9.1	7.4	8.3	6.6	12.5	13.0	12.1	4.5	5.6	3.7	17.3	20.2	15.5	11.2	11.1	11.2	9.9	11.1	9.0
Syphilis (A50–A53)	0.0	0.0	0.0	*	*	*	*	*	*	*	*	*	0.0	*	*	*	*	*	*	*	*
Acute poliomyelitis (A80)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Arthropod-borne viral encephalitis (A83–A84, A85.2)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Measles (B05)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Viral hepatitis (B15–B19)	0.7	0.9	0.5	0.9	1.2	0.7	1.9	1.8	2.0	0.9	1.1	0.6	0.9	1.2	0.6	*	*	*	0.7	0.8	0.5
Human immunodeficiency virus (HIV) disease (B20–B24)	1.3	2.1	0.6	1.3	2.2	0.5	1.7	2.5	*	0.2	0.5	*	5.4	8.1	3.1	*	*	*	0.7	1.1	0.2
Malaria (B50–B54)	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*
Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05, A20–A36,A42–A44, A48–A49,A54–A79, A81–A82,A85.0–A85.1, A85.8,A86–B04,B06–B09, B25–B49,B55–B99,U07.1)	47.3	60.0	37.7	49.5	65.5	37.4	71.0	77.9	65.3	27.2	36.7	20.3	57.4	71.0	49.0	52.8	63.4	44.3	46.7	59.2	37.2
Malignant neoplasms (C00–C97)	142.3	167.3	124.2	104.3	121.9	92.2	118.2	122.7	115.0	90.1	105.0	79.5	165.0	202.9	142.2	141.0	139.8	143.8	149.0	174.6	130.0

See footnotes at end of table.

Table 11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																				
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White		
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female
Malignant neoplasms of lip, oral cavity and pharynx (C00–C14)	2.7	4.2	1.5	1.5	2.4	0.8	2.4	3.1	1.9	2.1	3.3	1.2	2.6	4.5	1.3	4.0	*	*	3.0	4.6	1.6
Malignant neoplasm of esophagus (C15)	3.7	6.4	1.4	1.8	3.1	0.7	2.6	4.1	1.4	1.6	2.7	0.6	2.6	4.4	1.4	*	*	*	4.3	7.4	1.6
Malignant neoplasm of stomach (C16)	2.6	3.4	2.0	4.4	5.4	3.6	3.3	3.8	2.9	3.9	5.2	2.9	4.4	6.1	3.3	4.1	*	*	1.9	2.6	1.4
Malignant neoplasms of colon, rectum and anus (C18–C21)	13.0	15.3	11.0	10.7	13.1	8.8	13.1	15.2	11.2	9.1	11.2	7.4	16.5	20.6	13.6	12.6	14.3	11.4	13.2	15.4	11.3
Malignant neoplasms of liver and intrahepatic bile ducts (C22)	6.6	9.3	4.2	8.8	12.1	6.0	8.3	9.7	7.1	7.5	10.8	4.8	7.7	12.0	4.6	10.6	15.2	6.2	6.0	8.4	3.9
Malignant neoplasm of pancreas (C25)	11.1	12.9	9.7	8.8	9.5	8.0	9.1	8.5	9.5	7.4	8.0	6.9	13.5	15.5	12.1	10.6	8.8	12.1	11.5	13.4	9.9
Malignant neoplasm of larynx (C32)	0.9	1.6	0.3	0.5	1.0	0.1	0.8	*	*	0.3	0.5	*	1.4	2.6	0.5	*	*	*	0.9	1.6	0.4
Malignant neoplasms of trachea, bronchus and lung (C33–C34)	30.1	35.2	26.0	13.6	17.6	10.7	24.7	24.2	25.2	17.8	22.6	14.2	32.3	43.3	24.8	29.0	29.1	29.1	33.2	37.9	29.5
Malignant melanoma of skin (C43)	2.0	2.8	1.3	0.6	0.7	0.5	*	*	*	0.3	0.4	0.2	0.3	0.4	0.3	*	*	*	2.6	3.7	1.7
Malignant neoplasm of breast (C50)	10.3	0.2	18.8	7.4	0.1	13.5	7.7	*	14.4	6.4	*	11.5	15.2	0.5	26.1	10.6	*	20.4	10.3	0.2	19.0
Malignant neoplasm of cervix uteri (C53)	1.1	...	2.1	1.2	...	2.2	1.3	...	2.6	0.7	...	1.3	1.5	...	2.8	3.1	...	6.3	1.0	...	2.0
Malignant neoplasms of corpus uteri and uterus, part unspecified (C54–C55)	2.9	...	5.4	2.4	...	4.5	2.0	...	3.8	2.0	...	3.5	5.9	...	10.2	5.3	...	10.2	2.6	...	4.9
Malignant neoplasm of ovary (C56)	3.1	...	5.8	2.6	...	4.7	2.2	...	4.1	2.4	...	4.3	3.0	...	5.2	3.3	...	6.5	3.3	...	6.1
Malignant neoplasm of prostate (C61)	7.7	18.6	...	6.3	15.4	...	5.2	12.0	...	3.4	8.3	...	13.7	36.5	...	6.3	13.8	...	7.5	17.8	...
Malignant neoplasms of kidney and renal pelvis (C64–C65)	3.4	5.0	2.1	3.0	4.5	1.8	4.7	6.8	3.1	1.7	2.4	1.2	3.3	5.0	2.1	*	*	*	3.6	5.2	2.2
Malignant neoplasm of bladder (C67)	4.1	6.9	2.0	2.1	3.6	1.0	2.2	3.0	1.5	1.7	2.8	0.9	3.3	5.4	2.0	*	*	*	4.6	7.8	2.2

See footnotes at end of table.

Table 11. Age-adjusted death rate for 113 selected causes, Enterocolitis due to *Clostridium difficile*, COVID-19, dementia-related causes, drug-induced causes, alcohol-induced causes, and firearm-related injuries, by Hispanic origin and race and sex: United States, 2022—Con.

[Age-adjusted rates are per 100,000 U.S. standard population; see Technical Notes in this report. Hispanic-origin and race categories are consistent with 1997 Office of Management and Budget standards. Data for some Hispanic-origin and race categories should be interpreted with caution because of inconsistencies in reporting these items on death certificates and surveys; see Technical Notes. An asterisk (*) preceding a cause-of-death code indicates that the code is not included in the *International Classification of Diseases, 10th Revision* (ICD-10); see Technical Notes]

Cause of death (based on ICD-10)	Non-Hispanic, single race ³																					
	Total ¹			Hispanic ²			American Indian and Alaska Native			Asian			Black			Native Hawaiian or Other Pacific Islander			White			
	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	Both sexes	Male	Female	
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70–C72)	4.3	5.2	3.5	3.0	3.4	2.6	2.3	3.1	1.6	2.2	2.7	1.7	2.8	3.4	2.4	4.5	*	*	5.1	6.2	4.1	
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81–C96)	13.6	17.6	10.4	10.8	13.4	8.9	7.9	8.2	7.7	8.1	10.7	6.3	14.3	18.6	11.4	11.8	12.4	11.0	14.2	18.5	10.8	
Hodgkin disease (C81)	0.3	0.3	0.2	0.3	0.5	0.2	*	*	*	0.1	*	*	0.2	0.3	0.1	*	*	*	0.3	0.4	0.2	
Non-Hodgkin lymphoma (C82–C85)	4.7	6.1	3.6	4.1	5.0	3.4	2.5	2.7	2.4	3.4	4.6	2.5	3.5	4.7	2.6	3.7	*	*	5.0	6.5	3.8	
Leukemia (C91–C95)	5.6	7.4	4.3	4.0	5.0	3.3	2.8	2.8	2.8	3.1	4.2	2.3	4.9	6.5	3.9	5.6	*	*	6.1	8.0	4.6	
Multiple myeloma and immunoproliferative neoplasms (C88,C90)	2.9	3.7	2.3	2.4	2.9	2.0	2.4	2.6	2.3	1.4	1.6	1.3	5.6	7.0	4.7	*	*	*	2.8	3.6	2.1	
Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)	0.0	0.1	0.0	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	*	0.0	0.0	0.0
All other and unspecified malignant neoplasms (C17,C23–C24, C26–C31,C37–C41, C44–C49,C51–C52, C57–C60,C62–C63,C66, C68–C69,C73–C80,C97)	19.2	22.6	16.6	14.9	16.4	13.7	17.6	18.9	16.3	11.6	13.2	10.3	20.7	24.2	18.3	16.5	18.1	15.0	20.1	23.8	17.3	
In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior. (D00–D48)	3.8	4.7	3.2	2.4	2.9	2.1	2.7	2.7	2.6	2.1	2.4	1.9	3.8	4.6	3.5	*	*	*	4.1	5.2	3.4	
Anemias (D50–D64)	1.5	1.6	1.4	1.0	1.1	0.9	1.4	*	1.4	0.8	0.9	0.7	3.0	3.3	2.9	*	*	*	1.3	1.4	1.2	
Diabetes mellitus (E10–E14)	24.1	30.5	18.8	28.3	34.7	22.9	47.7	55.1	40.8	17.2	21.5	13.9	42.9	52.8	35.5	49.9	53.8	46.0	21.3	27.6	16.0	
Nutritional deficiencies (E40–E64)	5.0	4.6	5.3	3.2	3.1	3.2	5.1	4.3	5.7	2.2	2.0	2.4	5.6	5.9	5.2	*	*	*	5.4	4.8	5.8	
Malnutrition (E40–E46)	4.9	4.4	5.2	3.1	3.1	3.2	4.8	3.9	5.5	2.2	2.0	2.3	5.4	5.8	5.1	*	*	*	5.3	4.7	5.6	
Other nutritional deficiencies. (E50–E64)	0.1	0.1	0.1	0.1	*	*	*	*	*	*	*	*	0.1	*	0.1	*	*	*	0.1	0.1	0.1	

See footnotes at end of table.