

# Irodov Problem 1.13

Rajarshi Dasgupta

July 25, 2025

**1.13** Point A moves uniformly with velocity  $v$  so that the vector  $\vec{v}$  is continually ‘aimed’ at point B which in its turn moves rectilinearly and uniformly with velocity  $u < v$ . At the initial moment of time  $\vec{v} \perp \vec{u}$  and the points are separated by a distance  $l$ . How soon will the points converge?

I want to discuss the solution of the above problem from ‘Problems in General Physics’ by I.E. Irodov [?]. Let  $T$  be the time taken for points A and B to converge. So, we need to find an expression for  $T$  in terms of the quantities  $u$ ,  $v$ , and  $l$ .

## 1 School kid’s solution

Let  $r$  be the distance between the points A and B, and  $\theta$  be the angle between the velocities  $\vec{v}$  and  $\vec{u}$ . We consider the relative velocity of A with respect to B along AB and orthogonal to AB to obtain the following relation.

$$\dot{r} = -v + u \cos \theta \quad (1)$$

$$r\dot{\theta} = -u \sin \theta \quad (2)$$

First, we will eliminate  $\dot{\theta}$  by differentiating equation 1 with respect to time and then substituting the expression for  $\dot{\theta}$  from equation 2.

$$\ddot{r} = -u\dot{\theta} \sin \theta$$

$$\implies r\ddot{r} = u^2 \sin^2 \theta$$




Figure 1: (a) Velocities of points A and B separated by a distance of  $l$  at the initial moment of time  $t = 0$  (b) Velocities of points A and B separated by a distance of  $r$  at time  $t \in (0, T)$ .

Now, we will eliminate  $\theta$  from the above expression by using the trigonometric identity  $\cos^2 \theta + \sin^2 \theta = 1$ , and equation 1.

$$\begin{aligned} r\ddot{r} &= u^2 \sin^2 \theta \\ &= u^2 (1 - \cos^2 \theta) \\ &= u^2 \left\{ 1 - \left[ \frac{\dot{r} + v}{u} \right]^2 \right\} \end{aligned}$$

Simplifying the above expression we obtain the following second order differential equation.

$$\begin{aligned} r\ddot{r} + \dot{r}^2 + 2\dot{r}v + v^2 - u^2 &= 0 \\ \implies \frac{d}{dt}(r\dot{r} + 2rv) + v^2 - u^2 &= 0 \end{aligned}$$

Now, integrating with respect to time from  $t = 0$  to  $T$  we get the following expression.

$$\begin{aligned} \int_{t=0}^{t=T} \frac{d}{dt}(r\dot{r} + 2rv) dt + \int_{t=0}^{t=T} (v^2 - u^2) dt &= 0 \\ \implies r(T)\dot{r}(T) + 2r(T)v - r(0)\dot{r}(0) - 2r(0)v + (v^2 - u^2)T &= 0 \end{aligned}$$

We will use the fact that

$$\begin{aligned} r(0) &= l \\ \dot{r}(0) &= -v + u \cos \pi/2 && \text{(from equation 1)} \\ &= -v \\ r(T) &= 0 && \text{(since A and B converge at time T)} \end{aligned}$$

to get the final expression for  $T$ .

$$T = \frac{lv}{v^2 - u^2} \quad (3)$$

## 2 Physics major's solution

To be written please wait

## 3 Numerical solution

To be written please wait

## Discussion

To be written please wait

## References

- [1] I. E. Irodov, *Problems in General Physics*, Mir Publishers, 1988.