

TOTAL MARKS: 100

TOTAL TIME: 3 HOURS

- (1) Question 1 is compulsory.
- (2) Attempt any **four** from the remaining questions.
- (3) Assume data wherever required.
- (4) Figures to the right indicate full marks.

1 (a) Choose the correct answer:

(4 marks)

(i) The general solution of the equation $x^2p^2 + 3xyp + 2y^2 = 0$ is _____

- (a) $(y^2x - c)(xy - c) = 0$
- (b) $(x - y - c)(x^2 + y^2 - c) = 0$
- (c) $(xy - c)(x^2y - c) = 0$
- (d) $(y - x - c)(x^2 + y^2 + c) = 0$

ii) The given differential equation is solvable for y, if it is possible to express y in terms of _____

- (a) y and p
- (b) x and p
- (c) x and y
- (d) y and x

iii) The singular solution of Clairaut's equation is _____

- (a) $y = xg(x) + f[g(x)]$
- (b) $y = cx + f(c)$
- (c) $cy + f(c)$
- (d) $y g^2(x) + f[g(x)]$

iv) The singular solution of the equation $y = px - \log p$ is _____

- (a) $y^2 = 4ax$
- (b) $x = 1 - \log x$
- (c) $y = 1 - \log \left(\frac{1}{x} \right)$
- (d) $x^2 = y \log x$

1 (b) Solve $p^2 - 2p \sinh x - 1 = 0$

(4 marks)

1 (c) Solve $y=2px+\tan^{-1}(xp^2)$

(6 marks)

1 (d) Obtain the general solution and singular solution of Clairaut's equation is $(y-px)(p-1)=p$

(6 marks)

2 (a) Choose the correct answer:

(4 marks)

(i) The complementary function of $[D^4+4]x=0$ is _____

(a) $x = e^{-4}[c_1 \cos t + c_2 \sin t] + e^1[c_3 \cot t + c_4 \sin t]$

(b) $x = [c_1 \cos t + c_2 \sin t] + [c_3 \cos t + c_4 \sin t]$

(c) $x = [c_1 + c_2 t]e^{-t}$

(d) $x = [c_1 + c_2 t]e^t$

Find the particular integral of $(D^3-3D^2+4)y=e^{2x}$ is _____

(a) $\frac{x^2 e^{2x}}{6}$

(b) $\frac{x^2 e^{3x}}{6}$

(c) $\frac{x^2 e^x}{6}$

(d) $\frac{x^2 e^{4x}}{6}$

iii) Roots of $\frac{d^2 y}{dx^2} + 4\frac{dy}{dx} + 5y = 0$ are _____

(a) $2 \pm i$

(b) $3 \pm i$

(c) $2 \pm 2i$

(d) $-2 \pm i$

iv) Find the particular integral of $(D^3 + 4D)y=\sin 2x$ is _____

$$(a) \frac{x \sin x}{8}$$

$$(b) \frac{-x \sin x}{8}$$

$$(c) \frac{-x \sin 2x}{8}$$

$$(d) \frac{x \sin 2x}{8}$$

2 (b)

(4 marks)

$$\text{Solve } \frac{d^2 y}{dx^3} + 6 \frac{d^2 y}{dx^2} + 11 \frac{dy}{dx} + 6y = e^x + 1$$

2 (c)

(6 marks)

$$\text{Solve } \frac{d^2 y}{dx^2} - 4y = \cosh(2x - 1) + 3^x$$

2 (d)

(6 marks)

$$\text{Solve } \frac{dy}{dx} + y = ze^x, \frac{dz}{dx} + z = y + e^x$$

3 (a) Choose the correct answer:

(4 marks)

(i) The Wronskian x and $x e^x$ is _____

- (a) e^x
- (b) e^{2x}
- (c) e^{-2x}
- (d) e^{-x}

(ii) The complementary function of $x^2 y'' - xy' - 3y = x^2 \log x$ is _____

(a) $c_1 \cos(\log x) + c_2 \sin(\log x)$

(b) $c_1 x^{-1} + c_2 x$

(c) $c_1 x + c_2 x^3$

(d) $c_1 \cos x + c_2 \sin x$

iii) To transform $(1+x)^2 y'' + (1+x)y' + y = 2 \sin \log(1+x)$ into a linear differential

equation with constant coefficient _____

- (a) $(1+x)=e^t$
- (b) $(1+x)=e^{-t}$
- (c) $(1+x)^2=e^t$
- (d) $(1-x)^2=e^t$

iv) The equation $a_0(ax+b)^2y''+a_1(ax+b)y'+a_2y=\phi(x)$ is _____

- (a) Simultaneous equation
- (b) Cauchy's linear equation
- (c) Legendre linear equation
- (d) Euler's equation

3 (b) Using the variation of parameters method to solve the equation $y''+2y'+y=e^{-x} \log x$. (6 marks)

3 (c) (6 marks)

$$\text{Solve } x^2 \frac{d^2 y}{dx^2} - (2m-1)x \frac{dy}{dx} + (m^2 + n^2)y = n^2 x^m \log x$$

3 (d) Obtain the Frobenius method solve the equation (6 marks)

$$x \frac{d^2 y}{dx^2} + \frac{dy}{dx} - y = 0$$

4 (a)i) Partial differential equation by eliminating a and b from the relation (4 marks)

$Z=(x-a)^2+(y-b)^2$ is _____

- (a) $p^2q^2=4z$
- (b) $pq=4z$
- (c) $r=4z$
- (d) $t=4$

ii) The Lagrange's linear partial differential equation $Pp+Qq=R$ the subsidiary equation is _____

- (a) $\frac{dx}{R} = \frac{dy}{P} = \frac{dz}{Q}$
- (b) $\frac{dx}{P} = \frac{dy}{Q} = \frac{dz}{R}$
- (c) $\frac{dx}{Q} = \frac{dy}{R} = \frac{dz}{P}$
- (d) $\frac{dx}{P} + \frac{dy}{Q} + \frac{dz}{R}$

iii) By the method of separation of variable we seek a solution in the form is

(a) $x = x + y$

(b) $z = x^2 + y^2$

(c) $x = z + y$

(d) $x = x(x)y(y)$

iv) The solution of

$$\frac{\partial^2 z}{\partial x^2} = \sin(xy) \text{ is}$$

(a) $z = -x^2 \sin(xy) + y f(x) + \phi(x)$

(b) $\frac{-\sin(xy)}{y} + x f(y) + \phi(y)$

(c) $z = \frac{-\sin xy}{x^2} + y f(x) + \phi(x)$

(d) None of these

4 (b) From the partial differential equation of all sphere of radius 3 units having their centre in the xy-plane. (4 marks)

4 (c) (6 marks)

$$\text{Solve } x(y^2 + z)p - y(x^2 + z) = z(x^2 - y^2)$$

4 (d) Use the method of separation of variable to solve (6 marks)

$$y^3 \frac{\partial z}{\partial x} + x^2 \frac{\partial z}{\partial y} = 0$$

5 (a) Choose the correct answer: (4 marks)

(i) The value of

$$\int_0^1 \int^{x^2} 0e^{y/x} dy dx \text{ is } \underline{\hspace{2cm}}$$

(a) 0

(b) 1

(c) 3

(d) 1/2

ii) The value of $\beta(1/2)$ is _____

(a) $2\sqrt{\pi}$

(b) $\pi/2$

(c) $\sqrt{\pi}$

(d) $\sqrt{2\pi}$

(iii) The integral

$$\int_0^a \int_y^a \frac{x}{x^2 + y^2} dx dy$$

after changing the order of integration is _____

(a) $\int_0^a \int_0^x dy dx$

(b) $\int_0^a \int_0^x \frac{x}{x^2 + y^2} dx dy$

(c) $\int_0^x \int_0^x \frac{x}{x^2 + y^2} dx dy$

(d) $\int_0^x \int_0^a \frac{x}{x^2 + y^2} dx dy$

iv) The value of $\beta(3, 1/2)$ is _____

(a) $\frac{15}{16}$

(b) $\frac{16}{15}$

(c) $\frac{16}{5}$

(d) $\frac{16}{3}$

5 (b) Change the order of integration in

(4 marks)

$$\int_0^{4a} \int_{\frac{x^2}{4x}}^{\sqrt{ax}} dy dx$$

and hence evaluate the same.

5 (c)

(6 marks)

Evaluate $\int_{-c}^c \int_{-b}^b \int_{-a}^a (x^2 + y^2 + z^2) dx dy dz$

5 (d)

(6 marks)

Prove that $\int_0^1 \frac{x^2}{\sqrt{1-x^4}} dx - \int_0^1 \frac{1}{\sqrt{1+x}} dx = \frac{\pi}{4\sqrt{2}}$

6 (a) Let S be the closed boundary surface of a region of volume V then for a vector field of defined in V and S $\int_S f \cdot \mathbf{n} ds$ is _____

(4 marks)

- (a) $\int_V \text{curl of } \mathbf{dv}$
- (b) $\int_V \mathbf{f} \cdot \mathbf{dv}$
- (c) $\int_V \text{grad } \mathbf{dv}$
- (d) None of these

If $\int_C \mathbf{f} \cdot d\mathbf{r}$ where $\mathbf{f} = 3xy\hat{i} - y^2\hat{j}$

and C is the part of the parabola $y=2x^2$ from the region (0, 0) to the point (1, 2) is

- (a) 7/6
- (b) -7/6
- (c) $3x+3y$
- (d) -35

iii) In the Green's theorem in the plane

$\oint_C Mdx + Ndy = \text{_____}$

- (a) $\iint_R \left[\frac{\partial M}{\partial y} + \frac{\partial N}{\partial x} \right] dx dy$
- (b) $\iint_R \left[\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right] dx dy$
- (c) $\iint_R \left[\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right] dx dy$
- (d) $\iint_R \left[\frac{\partial N}{\partial x} + \frac{\partial M}{\partial y} \right] dx dy$

iv) A necessary and sufficient condition that the line integral

$$\int_c \vec{F} \cdot d\vec{r}$$

for any closed curve C is _____

$$(a) \operatorname{div} \vec{F} = 0$$

$$(b) \operatorname{div} \vec{F} \neq 0$$

$$(c) \operatorname{curl} \vec{F} = 0$$

$$(d) \operatorname{grad} \vec{F} = 0$$

6 (b) Using the divergence theorem, evaluate

(4 marks)

$$\int_c f \cdot n ds \text{ where } f = 4xz\hat{i} - y^2\hat{j} + yz\hat{k}$$

and S is the surface of the cube bounded by $x=0, x=1, y=0, y=1, z=0, z=1$.

6 (c) Use the Green's theorem, evaluate

(6 marks)

$$\iint_c (2x^2 - y^2)dx + (x^2 + y^2)dy$$

where C is the triangle formed by the lines $x=0, y=0$ and $x+y=1$.

6 (d) Verify the Stoke's theorem for

(6 marks)

$$f = -y^3\hat{i} + x^3\hat{j}$$

where S is the circle disc $x^2+y^2 \leq 1, z=0$.

7 (a) Choose the correct answer:

(4 marks)

$$(i) L\{\sinh at\} = \underline{\hspace{2cm}}$$

$$(a) \frac{s}{s^2 - a^2}$$

$$(b) \frac{s}{s^2 + a^2}$$

$$(c) \frac{a}{s^2 - a^2}$$

$$(d) \frac{a}{s^2 + a^2}$$

- ii) if $L\{f(t)\}=F(s)$ then $L\{e^{at}f(t)\}$ is _____ (a) $F(s+a)$
 (b) $F(s-a)$
 (c) $F(s)$
 (d) None of these

$$iii) L \left\{ \frac{e^t \sin t}{t} \right\}$$

$$(a) \frac{\pi}{2} + \tan^{-1}(s-1)$$

$$(b) \frac{\pi}{2} + \tan^{-1} s$$

$$(c) \frac{\pi}{2} - \cot^{-1} s$$

$$(d) \cot^{-1}(s-1)$$

- iv) Transform of unit step function $L\{u(t-a)\}$ is, _____

$$(a) \frac{e^{as}}{s}$$

$$(b) \frac{e^{-s}}{s}$$

$$(c) \frac{e^{2s}}{s}$$

$$(d) \frac{e^{-as}}{s}$$

7 (b)

(4 marks)

$$\text{Evaluate } L \left\{ 3^t + \frac{\cos 2t - \cos 3t}{t} + t \sin t \right\}$$

- 7 (c) Find the Laplace transform of the triangular wave, given by

(6 marks)

$$f(t) = \begin{cases} t & 0 \\ 2C-1 & C < 1 < 2C \end{cases} \quad \text{and } f(t+2C) = f(t)$$

7 (d)

(6 marks)

$$\text{Express } f(t) = \begin{cases} \cos t & \text{if } 0 < t < \pi \\ \cos 2t & \text{if } \pi < t < 2\pi \\ \cos 3t & \text{if } t > 2\pi \end{cases}$$

in terms of unit step function and hence find $L\{f(t)\}$

8 (a) Choose the correct answer:

(4 marks)

$$L^{-1} \left\{ \cot^{-1} \left(\frac{s}{a} \right) \right\} = \text{_____}$$

(a) $\frac{\sin t}{t}$

(b) $\frac{\sin at}{t}$

(c) $\frac{\sinh at}{t}$

(d) $\frac{\sinh t}{t}$

$$(ii) L^{-1} \left\{ \frac{1}{4s^2 - 36} \right\} = \text{_____}$$

(a) $\frac{\cosh 6t}{4}$

(b) $\frac{\sin 3t}{12}$

(c) $\frac{\sinh 3t}{12}$

(d) $\frac{\cosh 3t}{6}$

$$iii) L^{-1} \left\{ \frac{1}{s(s^2 + a^2)} \right\} = \text{_____}$$

(a) $\frac{1 - \cos at}{a^2}$

(b) $\frac{1 + \cos at}{a^2}$

(c) $\frac{1 - \sin at}{a^2}$

(d) $\frac{1 + \sin 3t}{6}$

$$(iv) L^{-1} \left\{ \frac{s^2 - 3s + 4}{s^4} \right\} = \underline{\hspace{2cm}}$$

$$(a) 1 - 3t + 2t^3$$

$$(b) 1 + \frac{t^2}{3}$$

$$(c) t + \frac{3}{2} + 1$$

$$(d) t - \frac{3}{2}t^2 + \frac{2}{3}t^3$$

8 (b)

(4 marks)

$$Find L^{-1} \left\{ \frac{3s + 7}{s^2 - 2s - 3} \right\}$$

8 (c) Using Convolution theorem evaluate

(6 marks)

$$L^{-1} \left\{ \frac{1}{(s + 1)(s^2 + 4)} \right\}$$

8 (d)

(6 marks)

$$Solve \frac{d^2y}{dt^2} + 5\frac{dy}{dt} + 6y = 5e^{2t} \text{ given that } y(0) = 2, \frac{dy(0)}{dt} = 1$$

by using Laplace transform method.