The "IF" Command in Excel

TOTAL POINTS 2

1. Remember: IF(logical_test, [value_if_true], [value_if_false])

1 point

	A	B
1	dog	
2	cat	
3	rabbit	

Using the table above, what would be the result of the following formula?

=IF(A1="dog", "bark", "no bark")

bark

1. Remember: IF(logical_test, [value_if_true], [value_if_false])

1 point

	A	B
1	dog	
2	cat	
3	rabbit	

Using the table above, what would be the result of the following formula?

=IF(A1="dog", "bark", "no bark")

bark

2. Select all statements below that are true.

1 point

- [ ] =IF(1=2, "incorrect", "correct")

  returns: incorrect

- [x] =IF(TRUE, "yes", "no")

  returns: yes

- [x] =IF("apples"="oranges", "same", "different")

  returns: different

- [ ] =IF(A1>A1, "low", "high")

  returns: low

The “IF” Command in Excel Using Numerical Data

TOTAL POINTS 2

1. Given the formula

1 point

=IF(A1>1000, 1000, A1)

Which of the following is true?

- If the value of A1 is -20, the formula will return 1000.
- If the value of A1 is 2000, the formula will return 2000.
- No matter the value of A1, the lowest value this formula will return is 0.
- No matter the value of A1, the highest value this formula will return is 1000.

2.

1 point



Assume the formula shown above was entered into cell D1, then copied down to D2 and D3. The resulting values in cells D1, D2, and D3 would be:

- small, small, small
- small, small, large
- small, large, large
- large, large, large

The "Nested IF" Command in Excel

TOTAL POINTS 2

1. The formula

   1 point

   =IF(FALSE, 1, IF(TRUE, 2, 3))

   will return:

   - ○ 1
   - ◉ 2
   - ○ 3
   - ○ Error

2. Given the formula

   1 point

   =IF(A1<10, “low”, IF(A1<20, “middle”, “high”))

   If the value of A1 is 10, what will the formula return?

   middle

The "VLOOKUP" Function in Excel

TOTAL POINTS 2

1. 1 point

	A	B	C	D	E	F	G	H
1	Apples	2	8		Apples			
2	Bananas	5	7		=VLOOKUP(E1, $A$1:$C$3,3, FALSE)			
3	Candy	3	5					

What will the VLOOKUP formula return?

8

2. 1 point

6	Apples	1		Apples
7	Bananas	6		=VLOOKUP(D6, $A$6:$B$9,2, FALSE)
8	Candy	13		
9	Apples	2		

What will the VLOOKUP formula return?

1

Another "VLOOKUP" Example

TOTAL POINTS 2

1.

1 point

	A	B	C	D	E	F	G	H	I	J	K
1	Size	Label		Size	Label						
2	0	S		8	=VLOOKUP(D2,$A$2:$B$5,2,TRUE)						
3	5	M									
4	10	L									
5	15	XL									

VLOOKUP(lookup_value, table_array, col_index_num, [range_lookup])

What will the VLOOKUP formula return? Note the last parameter.

- ○ S
- ◉ M
- ○ L
- ○ XL
- ○ #N/A

2.

1 point

	A	B	C	D	E	F	G	H	I	J	K
1	Size	Label		Size	Label						
2	0	S		8	=VLOOKUP(D2,$A$2:$B$5,2,FALSE)						
3	5	M									
4	10	L									
5	15	XL									

VLOOKUP(lookup_value, table_array, col_index_num, [range_lookup])

What will the VLOOKUP formula return? Note the last parameter.

- ○ S
- ○ M
- ○ L
- ○ XL
- ◉ #N/A

The "HLOOKUP" Function in Excel

TOTAL POINTS 2

1.

1 point

	A	B	C	D	E	F
1	Apples	Bananas	Candy			
2	2	3	4			
3	5	6	7			
4	8	9	10			
5						
6	Candy					
7	=HLOOKUP(A6,$A$1:$C$4,3,FALSE)					
8	HLOOKUP(lookup_value, table_array, row_index_num, [range_lookup])					

What will the HLOOKUP formula return?

7

2.

1 point

	A	B	C	D	E	F	G	H	I
1	Size	0	5	10	15				
2	Label	S	M	L	XL				
3									
4	=HLOOKUP(10,$A$1:$E$2,2,FALSE)								
5	HLOOKUP(lookup_value, table_array, row_index_num, [range_lookup])								

What will the HLOOKUP formula return?

- ○ S
- ○ M
- ◉ L
- ○ XL
- ○ #N/A

Professor 'Know-it-all' Needs Help!

TOTAL POINTS 2

1. Professor 'Know-it-all' wants to implement another system to call upon a group of students at random. She has created 4 groups: 1 point

   Group 1: Students 1-5 Group 2: Students 6-10 Group 3: Students 11-15 Group 4: Students 16-20

   Her VLOOKUP function does not correctly return a random group. What is wrong with her function below?



   - ( ) The lookup_value parameter should be "RANDBETWEEN(1,16)".
   - ( ) The table_array parameter should be "A3:B6".
   - ( ) The col_index_num parameter should be "1".
   - (•) The [range_lookup] parameter should be "TRUE".

2. You inform Professor K. that a much simpler solution to her problem above would be to just use the RANDBETWEEN function to randomly return a value of 1, 2, 3, or 4 directly. Which of the following would correctly implement this solution? 1 point

   - (•) =RANDBETWEEN(1,4)
   - ( ) =RANDBETWEEN(1,16)
   - ( ) =RANDBETWEEN(1,20)
   - ( ) =RANDBETWEEN(0,5)