Reading Data into Excel

TOTAL POINTS 2

1.

1 / 1 point

```
ColumnA,ColumnB,ColumnC,ColumnD
1,2,3,4
5,6,7,8
9,10,11,12
```

The format shown in the example above is:

- (●) Comma delimited
- ( ) Space delimited
- ( ) Tab delimited
- ( ) Fixed width

2.

1 / 1 point

```
ColumnA ColumnB ColumnC ColumnD
100     200     300     400
50000   60000   70000   80000
9000000 10000000        11000000        12000000
```

The format shown in the example above is:

- ( ) Comma delimited
- ( ) Space delimited
- (●) Tab delimited
- ( ) Semicolon delimited

Basic Data Manipulation in Excel

TOTAL POINTS 2

1. To move to the edge of the current data region, press:

1 / 1 point

- ○ Page Up/Down key
- ◉ CTRL + arrow key
- ○ ALT + arrow key
- ○ Home/End key

2.

1 / 1 point

	A	B	C
1	Serial Number	Part Number	Revision
2	9254	123	A
3	9621	123	B
4	9067	234	A
5	9435	234	B
6	9929	234	C
7	9682	234	D

The data above is sorted:

- ◉ First by Part Number, then by Revision.
- ○ First by Revision, then by Part Number.
- ○ First by Serial Number, then by Revision.
- ○ First by Revision, then by Serial Number.

Arithmetic Manipulation in Excel

TOTAL POINTS 2

1. Typing which of the following into a cell would result in a cell value of 3?

1 / 1 point

- ○ 1,2
- ○ 1+2
- ○ =1,2
- ◉ =1+2

2.

1 / 1 point

	A	B	C
1	1	2	3
2	4	5	6
3	7	8	9

Using the data in the example above, what is the value of the following formula?

=A3+C1

10

Basic Functions in Excel

TOTAL POINTS 2

1.

1 / 1 point

	A	B	C
1	1	2	3
2	4	5	6
3	7	8	9

In the example above, typing which of the following into a cell would result in a cell value of 3? Select all that apply.

- [x] =1+2

Correct
You must select all of the correct answers, and none of the incorrect answers.

- [x] =SUM(1,2)

Correct
You must select all of the correct answers, and none of the incorrect answers.

- [x] =SUM(A1,B1)

Correct
You must select all of the correct answers, and none of the incorrect answers.

- [x] =SUM(A1:B1)

Correct
You must select all of the correct answers, and none of the incorrect answers.

2.

1 / 1 point

	A	B	C
1	USE	2	
2	IGNORE	5	
3	USE	8	
4			
5	=SUMIF(A1:A3,"USE",B1:B3)		

SUMIF(range, criteria, [sum_range])

In the example above, what would be the calculated value of the SUMIF statement in cell A5?

10

Functions Using Absolute and Relative References

TOTAL POINTS 2

1. Which symbol is used in a formula to “fix” a row or a column reference?

1 / 1 point

- ○ #
- ◉ $
- ○ %
- ○ &

✓ **Correct**

2. What would be the calculated value of the following formula?

1 / 1 point

=MAX(1,4,2,8,5)

8

Functions Explained

TOTAL POINTS 2

1.

1 / 1 point

	A	B	C	D	E
1		1	2	3	4
2	1				
3	2				
4	3				
5	4				

	A	B	C	D	E
1		1	2	3	4
2	1	1	2	3	4
3	2	2	4	6	8
4	3	3	6	9	12
5	4	4	8	12	16

Let's say you want each blank cell to equal its row header multiplied by its column header, as shown above. Which of the following cell B2 formulas would give the correct result when pasted into the remaining empty cells?

- ( ) =A2*B1
- (•) =$A2*B$1
- ( ) =A$2*$B1
- ( ) =A2*B1

2. If you want to always reference cell C5 in a formula, even when that formula is copied down rows or across columns, the reference to cell C5 should be written as:

1 / 1 point

- ( ) C5
- ( ) $C5
- ( ) C$5
- (•) C5