

Non-Hermitian Hamiltonians and Similarity Transformations

Francisco M. Fernández¹

Received: 8 March 2015 / Accepted: 10 June 2015
© Springer Science+Business Media New York 2015

Abstract We show that similarity (or equivalent) transformations enable one to construct non-Hermitian operators with real spectrum. In this way we can also prove and generalize the results obtained by other authors by means of a gauge-like transformation and its generalization. Such similarity transformations also reveal the connection with pseudo-Hermiticity in a simple and straightforward way.

Keywords Similarity transformation · Equivalent operators · Isospectral operators · Pseudosymmetric operators

1 Introduction

In the last years there has been great interest in the mathematical properties of non-Hermitian Hamiltonians, which was mainly aroused by the conjecture that the non-Hermitian Hamiltonians with real spectra studied so far [1–3] exhibited PT symmetry [4]. There is a vast literature on non-Hermitian Hamiltonians, some of which is reviewed elsewhere [5] (see also [6–14] and references therein). Later Mostafazadeh [15–17] showed that every Hamiltonian with a real spectrum is pseudo-Hermitian and that all the PT-symmetric Hamiltonians studied in the literature exhibited such property. On the other hand, the so-called space-time symmetry did not prove to be so robust in producing non-Hermitian operators with real spectra [18–21].

✉ Francisco M. Fernández
fernande@quimica.unlp.edu.ar

¹ INIFTA (UNLP, CCT La Plata-CONICET), División Química Teórica,
Bldv. 113 S/N, Sucursal 4, Casilla de Correo 16, 1900 La Plata, Argentina

Some time ago Ahmed [22] derived a family of one-dimensional non-Hermitian Hamiltonians with real spectrum by means of a *gauge-like* transformation. He argued that the eigenfunctions of the resulting PT-symmetric Hamiltonian did not satisfy the PT-orthogonality condition. Recently, Rath and Mallick [23] put forward a generalization of the gauge-like transformation that involves both the coordinate and momentum operators and leads to a non-Hermitian Hamiltonian that appears to be isospectral with the harmonic oscillator.

The purpose of this paper is to discuss the gauge-like transformation in a more general and rigorous setting. In Section 2 we outline the main ideas of the similarity (or equivalent) transformation between a non-Hermitian and a Hermitian Hamiltonian. In Section 3 we discuss the gauge-like transformation introduced by Ahmed and in Section 4 the somewhat more general transformation proposed by Rath and Mallick. In Section 5 we show how to generalize the latter. Finally, in Section 6 we summarize the main results and draw conclusions.

2 Similarity or Equivalent Transformation

Let H be a Hermitian operator with a discrete spectrum

$$H\psi_n = E_n\psi_n, \quad (1)$$

and a complete set of eigenvectors

$$\sum_n |\psi_n\rangle\langle\psi_n| = I, \quad \langle\psi_m|\psi_n\rangle = \delta_{mn}, \quad (2)$$

where I is the identity operator. Its spectral decomposition reads

$$H = \sum_n E_n |\psi_n\rangle\langle\psi_n|. \quad (3)$$

For every linear invertible operator U the similarity transformation

$$\tilde{H} = UHU^{-1}, \quad (4)$$

yields a new operator \tilde{H} that is not Hermitian unless $U^{-1} = U^\dagger$. We say that H and \tilde{H} are equivalent or similar. The transformed vectors

$$|\varphi_n\rangle = U|\psi_n\rangle, \quad (5)$$

are eigenvectors of \tilde{H}

$$\tilde{H}|\varphi_n\rangle = UHU^{-1}U|\psi_n\rangle = E_n|\varphi_n\rangle, \quad (6)$$

whereas

$$|\Phi_n\rangle = (U^{-1})^\dagger|\psi_n\rangle, \quad (7)$$

are eigenvectors of the adjoint operator \tilde{H}^\dagger

$$\tilde{H}^\dagger|\Phi_n\rangle = (U^{-1})^\dagger HU^\dagger(U^{-1})^\dagger|\psi_n\rangle = E_n|\Phi_n\rangle. \quad (8)$$

Both sets of vectors form a biorthonormal basis

$$\langle\Phi_m|\varphi_n\rangle = \langle\psi_m|\psi_n\rangle = \delta_{mn}, \quad (9)$$

that enables us to write

$$\tilde{H} = \sum_n E_n U|\psi_n\rangle\langle\psi_n|U^{-1} = \sum_n E_n |\varphi_n\rangle\langle\Phi_n|. \quad (10)$$

The basis set $\{|\varphi_n\rangle\}$ is orthonormal with the metric given by $(U^{-1})^\dagger U^{-1}$:

$$\langle\psi_m|\psi_n\rangle = \langle\varphi_m|(U^{-1})^\dagger U^{-1}|\varphi_n\rangle = \delta_{mn}. \quad (11)$$

On the other hand, the standard inner product

$$\langle\varphi_m|\varphi_n\rangle = \langle\psi_m|U^\dagger U|\psi_n\rangle, \quad (12)$$

is not necessarily finite.

It follows from (4) that

$$\tilde{H}^\dagger = (U^{-1})^\dagger H U^\dagger = (U^{-1})^\dagger U^{-1} \tilde{H} U U^\dagger = \eta \tilde{H} \eta^{-1} \quad (13)$$

where $\eta = (U^{-1})^\dagger U^{-1}$ is Hermitian and positive definite. We say that \tilde{H} is η -pseudo-Hermitian [15–17] and (11) becomes

$$\langle\varphi_m|\eta|\varphi_n\rangle = \delta_{mn}. \quad (14)$$

If A and B are two linear operators then

$$[\tilde{A}, \tilde{B}] = U[A, B]U^{-1}. \quad (15)$$

In particular, the commutator $[x, p] = iI$ between the coordinate x and momentum p is conserved

$$[\tilde{x}, \tilde{p}] = iI. \quad (16)$$

Summarizing A non-Hermitian operator \tilde{H} that is similar or equivalent to an Hermitian one H is pseudo Hermitian. In addition to it, both operators are isospectral. When the similarity transformation is unitary ($U^{-1} = U^\dagger$) it conserves the norm ($\langle\varphi_m|\varphi_n\rangle = \delta_{mn}$), $\eta = I$ and \tilde{H} is obviously Hermitian.

The results developed above are not new since they are contained in Mostafazadeh's papers [15–17]. We simply derived them here from the point of view of a similarity transformation in order to connect them with the papers of Ahmed [22] and Rath and Mallick [23] in a clearer way.

3 Gauge-Like Transformation

The gauge-like transformation for one-dimensional operators

$$H = \frac{1}{2}p^2 + V(x), \quad (17)$$

discussed by Ahmed [22] is a particular case of the similarity transformation outlined in Section 2. If we choose

$$U = e^{u(x)}, \quad (18)$$

then [24]

$$\tilde{p} = U p U^{-1} = p + [u, p] = p + iu', \quad \tilde{x} = x, \quad (19)$$

and

$$\tilde{H} = \frac{1}{2}(p + iu')^2 + V(x). \quad (20)$$

Therefore, H and \tilde{H} are isospectral as discussed in Section 2.

The transformation of the non-Hermitian operator

$$H_\beta = \frac{1}{2}[p + i\beta v(x)]^2 + V(x), \quad (21)$$

yields

$$\tilde{H}_\beta = \frac{1}{2}[p + i\beta v(x) + iu'(x)]^2 + V(x). \quad (22)$$

If $v(x)$ is real and

$$u'(x) = -2\beta v(x), \quad (23)$$

then

$$\tilde{H}_\beta = H_\beta^\dagger. \quad (24)$$

Since $u(x)$ is real then U is Hermitian and positive definite; therefore H_β is U -pseudo-Hermitian.

In particular, Ahmed chose $v(x) = x$ and $V(x) = (\alpha^2 + \beta^2)x^2/2$ so that

$$H_\beta = \frac{1}{2}(p + i\beta x)^2 + \frac{1}{2}(\alpha^2 + \beta^2)x^2, \quad (25)$$

and $u(x) = u_1(x) = -\beta x^2$ leads to (24). Note that if $u_2(x) = -\beta x^2/2$ then

$$e^{u_2} H_\beta e^{-u_2} = \frac{1}{2}p^2 + \frac{1}{2}(\alpha^2 + \beta^2)x^2 = H_{SHO} \quad (26)$$

from which we conclude that H_β and the simple harmonic oscillator H_{SHO} are isospectral. In this case the eigenfunctions $\varphi_n(x)$ of the former operator are square integrable provided $\alpha \neq 0$ [22]. These results are particular cases of those derived in Section 2 (note that $e^{u_2} (e^{u_2})^\dagger = e^{u_1}$).

Ahmed [22] also discussed the particular case $\beta = i\gamma$, γ real, that leads to the Hermitian operator

$$H_\gamma = \frac{1}{2}(p - \gamma x)^2 + \frac{1}{2}(\alpha^2 - \gamma^2)x^2, \quad (27)$$

and draw two curious conclusions. He stated that “Remarkably, the usual connection between the nodal structure with the quantum number n does not hold any more. Even the ground state may have nodes for some values of γ .” Since $|\varphi_n(x)| = |\psi_n(x)|$ it is obvious that both functions have the same number of nodes; in particular, the ground state $\varphi_0(x)$ is nodeless in the interval $(-\infty, \infty)$ as expected. He also said that “Eigenvalues (18) possess an interesting feature of becoming complex (conjugate) at the cost of eigenfunction (19) being delocalized as it would not vanish at $x = \pm\infty$. This interesting phase-transition of eigenvalues from real to complex takes place when $\gamma > \gamma_{critical} (= \alpha)$.” It is obvious that this *interesting phase transition* is due to the force constant chosen for H_{SHO} and has nothing to do with the transformation of one oscillator into the other. To see this point more clearly just choose

$$H_\gamma = \frac{1}{2}(p - \gamma x)^2 + \frac{1}{2}kx^2, \quad (28)$$

and the phase transition does not take place for any value of γ if $k > 0$.

4 Transformation of Coordinate and Momentum

Recently, Rath and Mallick [23] proposed the following generalization of the gauge-like transformation:

$$x \rightarrow \tilde{x} = \frac{1}{\sqrt{1 + \alpha\beta}}(x + i\alpha p), \quad p \rightarrow \tilde{p} = \frac{1}{\sqrt{1 + \alpha\beta}}(p + i\beta x), \quad (29)$$

that converts

$$H_{HO} = \frac{1}{2}(p^2 + x^2) \quad (30)$$

into the non-Hermitian operator

$$H = \frac{1}{2(1 + \alpha\beta)}[(p + i\beta x)^2 + (x + i\alpha p)^2]. \quad (31)$$

They did not place any restriction whatsoever on α and β but it is obvious that at least $\alpha\beta \neq -1$. By means of a non-rigorous procedure based on second quantization, an adjustable frequency and a truncated perturbation expansion the authors conjectured that the eigenvalues of H appeared to be exactly those of H_{HO} .

This conclusion follows straightforwardly from the similarity transformation

$$H = U H_{HO} U^{-1}, \quad (32)$$

where U is given by

$$UxU^{-1} = \tilde{x}, \quad UpU^{-1} = \tilde{p} \quad (33)$$

According to the results of Section 2 both operators are isospectral with eigenvalues

$$E_n = n + \frac{1}{2}, \quad n = 0, 1, \dots, \quad (34)$$

and H is η -pseudo-Hermitian. Clearly, no further calculation is required.

It only remains to determine whether the eigenfunctions of H are square integrable. To this end we resort to the construction of the eigenvectors of H_{HO} in second-quantization form [24]:

$$a|\psi_0\rangle = 0, \quad |\psi_n\rangle = \frac{1}{\sqrt{n!}}(a^\dagger)^n|\psi_0\rangle, \quad (35)$$

where

$$a = \frac{1}{\sqrt{2}}(x + ip), \quad a^\dagger = \frac{1}{\sqrt{2}}(x - ip). \quad (36)$$

It follows from (29), (33) and (35) that

$$\tilde{a}|\varphi_0\rangle = 0, \quad |\varphi_n\rangle = \frac{1}{\sqrt{n!}}(\tilde{a}^\dagger)^n|\varphi_0\rangle. \quad (37)$$

Since

$$\tilde{a} = \frac{1}{\sqrt{2(1 + \alpha\beta)}}[(1 - \beta)x + i(1 + \alpha)p], \quad (38)$$

then the ground state $\varphi_0(x)$ is a solution of the first-order differential equation

$$\varphi_0'(x) = -\frac{1 - \beta}{1 + \alpha}x\varphi_0(x), \quad (39)$$

that leads to

$$\varphi_0(x) = \left[\frac{1 - \beta}{\pi(1 + \alpha)} \right]^{1/4} \exp \left[-\frac{1 - \beta}{2(1 + \alpha)}x^2 \right]. \quad (40)$$

We appreciate that $\varphi_0(x)$ is square integrable (and, consequently, also all the other eigenfunctions $\varphi_n(x)$) provided that $(1 + \alpha)(1 - \beta) > 0$. The square-integrability of the eigenfunctions was not discussed by Rath and Mallick [23] in spite of the fact that the conditions just given appear explicitly in the zero and pole of their chosen frequency ω for Case II.

The operator that carries out the transformation (29) is of the form [24]

$$U = \exp(ax^2 + bp^2), \quad (41)$$

where

$$\begin{aligned} i\alpha &= \frac{(e^{2\sqrt{-ab}} - 1)\sqrt{-ab}}{a(e^{2\sqrt{-ab}} + 1)} \\ i\beta &= \frac{(1 - e^{2\sqrt{-ab}})\sqrt{-ab}}{b(e^{2\sqrt{-ab}} + 1)}, \end{aligned} \quad (42)$$

that leads to $\alpha/\beta = -b/a$.

5 A More General Coordinate-Momentum Transformation

A more general similarity transformation is given by [24]

$$\begin{aligned} \tilde{x} &= UxU^{-1} = U_{11}x + U_{12}p \\ \tilde{p} &= UpU^{-1} = U_{21}x + U_{22}p, \end{aligned} \quad (43)$$

where

$$U_{11}U_{22} - U_{21}U_{12} = 1, \quad (44)$$

follows from the condition $[\tilde{x}, \tilde{p}] = iI$. Since the matrix elements U_{ij} may be complex numbers the transformation depends on eight parameters that should satisfy two equations; therefore, there are only six independent parameters and the transformation is given by an exponential operator of the form [24]

$$U = \exp \left[\frac{a}{2}x^2 + \frac{c}{2}(xp + px) + \frac{b}{2}p^2 \right], \quad (45)$$

where a , b and c are complex numbers.

The application of this similarity transformation to the harmonic oscillator H_{HO} (30) yields the operator

$$\begin{aligned} \tilde{H} &= UH_{HO}U^{-1} \\ &= \frac{1}{2} \left[(U_{22}^2 + U_{12}^2)p^2 + (U_{11}^2 + U_{21}^2)x^2 + (U_{21}U_{22} + U_{11}U_{12})(xp + px) \right]. \end{aligned} \quad (46)$$

By means of well known operator formulas [24] it is not difficult to prove that

$$\begin{aligned} U_{11} &= \cosh(\theta) - \frac{c}{\theta} \sinh(\theta) \\ U_{12} &= -\frac{b}{\theta} \sinh(\theta) \\ U_{21} &= \frac{a}{\theta} \sinh(\theta) \\ U_{22} &= \cosh(\theta) + \frac{c}{\theta} \sinh(\theta) \\ \theta &= \sqrt{c^2 - ab}. \end{aligned} \quad (47)$$

In general, any operator of the form (46) with matrix elements U_{ij} that satisfy the condition (44) is equivalent (and therefore isospectral) to the harmonic oscillator (30). It is always η -pseudo-Hermitian and under certain conditions it may also be Hermitian or PT-symmetric.

For example, if $U_{22}^2 + U_{12}^2$ and $U_{11}^2 + U_{21}^2$ are both real and $(U_{21}U_{22} + U_{11}U_{12})$ purely imaginary, then \tilde{H} is PT-symmetric. The choice $U_{11} = U_{22} = 1$, $U_{12} = 0$, and $U_{21} = i\beta$ yields one of the examples given by Ahmed [22]. On the other hand, when $U_{11} = U_{22} = 1/\sqrt{1 + \alpha\beta}$, $U_{12} = i\alpha/\sqrt{1 + \alpha\beta}$, and $U_{21} = i\beta/\sqrt{1 + \alpha\beta}$ we obtain the model proposed by Rath and Mallick [23]. Obviously, when the coefficients of p^2 , x^2 and $xp + px$ are real \tilde{H} is Hermitian.

Arguing as in Section 4 we conclude that the eigenfunctions $\varphi_n(x)$ of \tilde{H} are square integrable provided that

$$\Re \frac{U_{11} + iU_{21}}{U_{22} - iU_{12}} > 0 \quad (48)$$

6 Conclusions

The purpose of this paper is to show that the results of Ahmed [22] and Rath and Mallick [23] can be straightforwardly derived and proved by suitable similarity transformations. In the former case there is no need of discussing the reality of the operator and its eigenfunctions or the orthogonality conditions. In fact, the proposition enunciated by the author does not explain the situation. Once we prove that a non-Hermitian operator is similar to an Hermitian one the reality of the spectrum of the former is certainly proved. Of course, caution must be exercised with respect to the square-integrability of its eigenfunctions.

With respect to the latter paper [23] the similarity transformation is a much more rigorous and straightforward way of proving that the non-Hermitian operator is isospectral with the harmonic oscillator. The results of both papers are merely particular cases of the general expressions derived by Mostafazadeh [15–17] and also of the equations derived in Section 2.

Finally, (46) with the restriction (44) enables us to construct a family of non-Hermitian operators with real spectrum. If necessary we can enlarge the number of cases by choosing $H_{HO} = p^2 + kx^2$, $k > 0$, instead of the operator (30) thus having one more independent parameter at our disposal.

Compliance with Ethical Standards Conflict of Interest: The authors declare that they have no conflict of interest. B. Bagchi and C. Quesne (Pseudo-Hermiticity, weak pseudo-Hermiticity and η -orthogonality condition. Phys. Lett. A 301, 173–176 (2002)) discussed the correct application of the η -orthogonality condition to the pseudo-Hermitian gauge-transformed Hamiltonian.

Acknowledgments The author would like to thank the referees for useful comments and suggestions that contributed to the improvement of this paper.

References

1. Alvarez, G.: Bender-Wu branch points in the cubic oscillator. J. Phys. A **28**, 4589–4598 (1995)
2. Delabaere, E., Pham, F.: Eigenvalues of complex Hamiltonians with PT-symmetry. II. Phys. Lett. A **250**, 29–32 (1998)
3. Fernández, F.M., Guardiola, R., Ros, J., Znojil, M.: Strong-coupling expansions for the PT-symmetric oscillators $V(x) = a(ix) + b(ix)^2 + c.(ix)^3$. J. Phys. A **31**, 10105–10112 (1998)
4. Bender, C.M., Boettcher, S.: Real Spectra in Non-Hermitian Hamiltonians Having PT Symmetry. Phys. Rev. Lett. **80**, 5243–5246 (1998)
5. Bender, C.M.: Making sense of non-Hermitian Hamiltonians. Rep. Prog. Phys. **70**, 947–1018 (2007)
6. Fernández, F.M., Guardiola, R., Ros, J., Znojil, M.: A family of complex potentials with real spectrum. J. Phys. A **32**, 3105–3116 (1999)

7. Krejčířík, D., Bíla, H., Znojil, M.: Closed formula for the metric in the Hilbert space of a PT-symmetric model. *J. Phys. A* **39**, 10143–10153 (2006)
8. Znojil, M.: Coupled-channel version of the PT-symmetric square well. *J. Phys. A* **39**, 441–455 (2006)
9. Znojil, M.: Solvable non-Hermitian discrete square well with closed-form physical inner product. *J. Phys. A* **47**, 435302 (2014)
10. Znojil, M.: PT-symmetric model with an interplay between kinematical and dynamical nonlocalities. *J. Phys. A* **48**, 195303 (2015)
11. Lévai, G., Znojil, M.: Conditions for complex spectra in a class of PT-symmetric potentials. *Mod. Phys. Lett. A* **16**, 1973–1981 (2001)
12. Znojil, M., Lévai, G.: Spontaneous beakdown of PT-symmetry in the solvable square-well model. *Mod. Phys. Lett. A* **16**, 2273–2280 (2001)
13. Znojil, M.: PT-symmetric harmonic oscillators. *Phys. Lett. A* **259**, 220–223 (1999)
14. Znojil, M.: PT-symmetric square well. *Phys. Lett. A* **285**, 7–10 (2001)
15. Mostafazadeh, A.: Pseudo-Hermiticity versus PT symmetry: The necessary condition for the reality of the spectrum of a non-Hermitian Hamiltonian. *J. Math. Phys.* **43**, 205–214 (2002)
16. Mostafazadeh, A.: Pseudo-Hermiticity versus PT-symmetry. II. A complete characterization of non-Hermitian Hamiltonians with a real spectrum. *J. Math. Phys.* **43**, 2814–2816 (2002)
17. Mostafazadeh, A.: Pseudo-Hermiticity versus PT-symmetry III: Equivalence of pseudo-Hermiticity and the presence of antilinear symmetries. *J. Math. Phys.* **43**, 3944–3951 (2002)
18. Fernández, F.M., García, J.: Non-Hermitian Hamiltonians with unitary and antiunitary symmetries. *Ann. Phys.* **342**, 195–204 (2014). arXiv:1309.0808 [quant-ph]
19. Fernández, F.M., García, J.: PT-symmetry broken by point-group symmetry. *J. Math. Phys.* **55**, 042107 (2014). arXiv:1308.6179v2 [quant-ph]
20. Amore, P., Fernández, F.M., García, J.: Is space-time symmetry a suitable generalization of parity-time symmetry? *Ann. Phys.* **350**, 533–548 (2014). arXiv:1405.5234 [quant-ph]
21. Amore, P., Fernández, F.M., García, J.: Non-Hermitian oscillators with Td symmetry. *Ann. Phys.* **353**, 238–251 (2014). arXiv:1409.2672 [quant-ph]
22. Ahmed, Z.: Pseudo-Hermiticity of Hamiltonians under gauge-like transformation: real spectrum of non-Hermitian Hamiltonians. *Phys. Lett. A* **294**, 287–291 (2002)
23. Rath, B., Mallick, P.: Zero energy correction method for non-Hermitian Harmonic oscillator with simultaneous transformation of co-ordinate and momentum. arXiv:1501.06161 [quant-ph]
24. Fernández, F.M., Castro, E.A.: Algebraic Methods in Quantum Chemistry and Physics, Mathematical Chemistry Series, CRC, Boca Raton, New York, London, Tokyo (1996)