


Average Weight

71.2


Height (m)	Favorite Color	Gender	Weight (kg)	Residual
1.6	Blue	Male	88	16.8
1.6	Green	Female	76	
1.5	Blue	Female	56	
1.8	Red	Male	73	
1.5	Green	Male	77	
1.4	Blue	Female	57	

Height (m)	Favorite Color	Gender	Weight (kg)	Residual
1.6	Blue	Male	88	16.8
1.6	Green	Female	76	4.8
1.5	Blue	Female	56	-15.2
1.8	Red	Male	73	1.8
1.5	Green	Male	77	5.8
1.4	Blue	Female	57	-14.2



So we replace these residuals with their average.

$$\frac{(1.8 + 5.8)}{2} = 3.8$$



Now the **Predicted Weight** = $71.2 + (0.1 \times 16.8) = 72.9$


Height (m)	Favorite Color	Gender	Weight (kg)
1.6	Blue	Male	88



That gives us...

$$71.2 + (0.1 \times 16.8) + (0.1 \times 15.1)$$

= 74.4




To review, the three ideas behind **AdaBoost** are...

1) **AdaBoost** combines a lot of "weak learners" to make classifications. The weak learners are almost always **stumps**.



2) Some **stumps** get more say in the classification than others.

3) Each **stump** is made by taking the previous **stump's** mistakes into account.



Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight
Yes	Yes	205	Yes	1/8
No	Yes	180	Yes	1/8
Yes	No	210	Yes	1/8
Yes	Yes	167	Yes	1/8
No	Yes	156	No	1/8
No	Yes	125	No	1/8
Yes	No	168	No	1/8
Yes	Yes	172	No	1/8

The **Total Error** for a stump is the sum of the weights associated with the *incorrectly* classified samples.




We use the **Total Error** to determine **Amount of Say** this stump has in the final classification with the following formula:

$$\text{Amount of Say} = \frac{1}{2} \log\left(\frac{1 - \text{Total Error}}{\text{Total Error}}\right)$$

Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight
Yes	Yes	205	Yes	1/8
No	Yes	180	Yes	1/8
Yes	No	210	Yes	1/8
Yes	Yes	167	Yes	1/8
No	Yes	156	No	1/8
No	Yes	125	No	1/8
Yes	No	168	No	1/8
Yes	Yes	172	No	1/8



Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight
Yes	Yes	205	Yes	1/8
No	Yes	180	Yes	1/8
Yes	No	210	Yes	1/8
Yes	Yes	167	Yes	1/8
No	Yes	156	No	1/8
No	Yes	125	No	1/8
Yes	No	168	No	1/8
Yes	Yes	172	No	1/8



The sum of the weights
Total Error = for the incorrectly
classified samples.

Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight
Yes	Yes	205	Yes	1/8
No	Yes	180	Yes	1/8
Yes	No	210	Yes	1/8
Yes	Yes	167	Yes	1/8
No	Yes	156	No	1/8
No	Yes	125	No	1/8
Yes	No	168	No	1/8
Yes	Yes	172	No	1/8

New Sample Weight = sample weight $\times e^{\text{amount of say}}$

This is the formula we will
use to *increase* the
Sample Weight for the
sample that was
incorrectly classified.



Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight
Yes	Yes	205	Yes	1/8
No	Yes	180	Yes	1/8
Yes	No	210	Yes	1/8
Yes	Yes	167	Yes	1/8
No	Yes	156	No	1/8
No	Yes	125	No	1/8
Yes	No	168	No	1/8
Yes	Yes	172	No	1/8

New Sample Weight = sample weight $\times e^{-\text{amount of say}}$

This is the formula we will use to *decrease* the **Sample Weights**.

Chest Pain	Blocked Arteries	Patient Weight	Heart Disease	Sample Weight	New Weight	Norm. Weight
Yes	Yes	205	Yes	1/8	0.05	0.07
No	Yes	180	Yes	1/8	0.05	0.07
Yes	No	210	Yes	1/8	0.05	0.07
Yes	Yes	167	Yes	1/8	0.33	0.49
No	Yes	156	No	1/8	0.05	0.07
No	Yes	125	No	1/8	0.05	0.07
Yes	No	168	No	1/8	0.05	0.07
Yes	Yes	172	No	1/8	0.05	0.07

Now, when we add up the **New Sample Weights**, we get 1 (plus or minus a little rounding error).