Assignment 18.3:

Calculate F Test for given 10, 20, 30, 40, 50 and 5,10,15, 20, 25.

For 10, 20, 30, 40, 50:

Ans:

F Test is generally defined as ratio of the variances of the given two set of values. First calculate standard deviation and variation of the given set of values.

The formula used to calculate SD is, Standard Deviation Formula

$$\sigma = \sqrt{\frac{1}{n-1}\sum_{i=1}^{n}(x_i-\overline{x})^2}$$

The standard deviation is represented by the symbol σ and variance is square of the standard deviation.

F Test Formula

$$F = \frac{\text{estimate of } \sigma^2 \text{ from means}}{\text{estimate of } \sigma^2 \text{ from individuals}}$$

Calculate Variance of first set

```
Total Inputs (N) = (10,20,30,40,50)

Total Inputs (N) = 5

Mean (x_m) = (x1+x1+x2...xn)/N

Mean (x_m) = 150/5

Mean (x_m) = 30

SD = sqrt (1/(N-1)*((x_1-x_m)^2+(x_2-x_m)^2+...+(x_n-x_m)^2))
= sqrt (1/(5-1)((10-30)^2+(20-30)^2+(30-30)^2+(40-30)^2+(50-30)^2))
= sqrt (1/4((-20)^2+(-10)^2+(0)^2+(10)^2+(20)^2))
= sqrt (1/4((400)+(100)+(0)+(100)+(400)))
= sqrt(250)
= 15.8114
Variance=SD<sup>2</sup>
Variance=15.8114<sup>2</sup>
Variance=250
```

Calculate Variance of second set

```
For 5, 10,15,20,25:
Total Inputs (N) = (5,10,15,20,25)
Total Inputs (N) = 5
Mean (x_m) = (x_1+x_2+x_3...x_N) / N
Mean (x_m) = 75/5
Means (x_m) = 15
SD = sqrt(1/(N-1)*((x_1-x_m)^2+(x_2-x_m)^2+..+(x_n-x_m)^2))
    = qrt(1/(5-1)((5-15)^2+(10-15)^2+(15-15)^2+(20-15)^2+(25-15)^2))
    = \operatorname{sqrt}(1/4((-10)^2+(-5)^2+(0)^2+(5)^2+(10)^2))
    = \operatorname{sqrt}(1/4((100)+(25)+(0)+(25)+(100)))
    = sqrt(62.5)
     = 7.9057
Variance=SD<sup>2</sup>
Variance=7.9057<sup>2</sup>
Variance=62.5
To calculate F Test
F Test = (variance of 10, 20,30,40,50) / (variance of 5, 10, 15, 20, 25)
= 250/62.5
= 4.
```

The F Test value is 4.