



CONTENTS

1	Introduction	3
1.1	Atoms	4
1.2	Mass and Acceleration	6
1.3	Mass and Gravity	7
1.4	Mass and Weight	8
2	Atomic and Molecular Mass	11
2.1	Molar Mass	15
2.2	Heavy atoms aren't stable	15
3	Work and Energy	17
3.1	Heat	18
3.2	Electricity	18
3.3	Chemical Energy	18
3.4	Kinetic Energy	19
3.5	Gravitational Potential Energy	19
3.6	Conservation of Energy	20
3.7	Efficiency	20
4	Units and Conversions	23
4.1	Conversion Factors	25
4.2	Conversion Factors and Ratios	26
4.3	When Conversion Factors Don't Work	27
5	Simple Machines	29

5.1	Levers	30
5.2	Ramps	31
5.3	Gears	33
5.4	Hydraulics	34
6	Buoyancy	37
6.1	The Mechanism of Buoyancy	38
7	Heat	41
7.1	Specific Heat Capacity	41
7.2	Getting to Equilibrium	43
7.3	Specific Heat Capacity Details	44
A	Answers to Exercises	47
	Index	53



CHAPTER 1

Introduction

This book will start you on the long and difficult trek to becoming a modern problem solver. Along the path, you will learn how to use the tools of math, computers, and science.

Why should you bother? There are big problems in this world that will require expert problem solvers. Those people will make the world a better place while enjoying interesting and lucrative careers. We are talking about engineers, scientists, doctors, computer programmers, architects, actuaries, and mathematicians. Right now, those occupations represent about 6% of all the jobs in the United States. Soon, that number is expected to rise above 10%. On average, people in that 10% of the population are expected to have salaries twice that of their non-technical counterparts.

Solving problems is difficult. At some point on this journey, you will see people who are better at solving problems than you are. You, like every other person who has gone on this journey, will think “I have worked so hard on this, but that person is better at it than I am. I should quit.” Don’t.

First, solving problems is like a muscle. The more you do, the better you get at it. It is OK to say “I am not good at this yet.” That just means you need more practice.

Second, you don't need to be the best in the world. 10 million people your age can be better at solving problems than you, *and you can still be in the top 10% of the world*. If you complete this journey, there will be problems for you to solve and a job where your problem-solving skills will be appreciated.

So where do we start?

1.1 Atoms


The famous physicist Richard Feynman once asked this question: “If, in some cataclysm, all of scientific knowledge were to be destroyed, and only one sentence was passed on to the next generation of creatures, what statement would contain the most information in the fewest words?”

His answer was “All things are made of atoms—little particles that move around in perpetual motion, attracting each other when they are a little distance apart, but repelling upon being squeezed into one another.”

That seems like a good place to start.

All things (including the air around you) are made of atoms. Atoms are very tiny – there are more atoms in a drop of water than there are drops of water in all the oceans.

Every atom has a nucleus that contains protons and neutrons. There is also a cloud of electrons flying around the nucleus. However, the mass of the atom comes mainly from the protons and neutrons, which are exponentially heavier than electrons.



Watch **Elements and atoms** from Khan Academy at https://youtu.be/IFKnq9QM6_A.

We classify atoms by the numbers of protons they have. An atom with one proton is a hydrogen atom, an atom with two protons is a helium atom, and so forth (refer to periodic

table on pg..). We say that hydrogen and helium are

elements because the classification of elements is based on proton number. And we give each element an atomic symbol. Hydrogen gets H. Helium gets He. Oxygen gets O. Carbon gets C, etc.

Often two hydrogen atoms will attach to an oxygen atom. The result is a water molecule. Why do they cluster together? because they share electrons in their clouds.


A molecule is described by the elements it contains. Water is H_2O because it has two hydrogen atoms and one oxygen atom.

There are many kinds of molecules. You know a few:


- Table salt is crystals made of NaCl molecules: a sodium atom attached to a chlorine atom.
- Baking soda, or sodium bicarbonate, is NaHCO_3 .
- Vinegar is a solution including acetic acid (CH_3COOH).
- O_2 is the oxygen molecules that you breathe out of the air (Air, a blend of gases, is mostly N_2).

Sometimes two hydrogen atoms form a molecule (H_2). Sometimes two oxygen atoms form a molecule (O_2). If you mix these together and light a match, they will rearrange themselves into water molecules. This is called a *chemical reaction*. In any chemical reaction, the atoms are rearranged into new molecules.

Some chemical reactions (like the burning of hydrogen gas described above) are *exothermic* – that is, they give off energy. Burning hydrogen gas happens quickly and gives off a lot of energy. If you have enough, it will make quite an explosion.



Other chemical reactions are *endothermic* – that is they consume energy. Photosynthesis, the process by which plants consume energy from the sun to make sugar from CO_2 and H_2O requires an endothermic chemical reaction.



1.2 Mass and Acceleration

Each atom has a mass, so everything that is made up of atoms has a mass, which is pretty much everything. We measure mass in grams. A paper clip is about 1 gram of steel. An adult human can weigh 70,000 grams, so for larger things we often talk about kilograms. A kilogram is 1000 grams.

The first interesting thing about mass is that objects with more mass require more force to accelerate. For example, pushing a bicycle so that it accelerates from a standstill to jogging speed in 2 seconds requires a lot less force than pushing a train so that it accelerates at the same rate.

You will probably find it useful to watch Khan Academy's summary of Newton's second law of motion: <https://youtu.be/ou9YMWlJgkE>

Newton's Second Law of Motion

The force necessary to accelerate an object of mass m is given by:

$$F = ma$$

That is the force is equal to the mass times the acceleration.

What are the units here? We already know that mass is measured in kilograms. We can measure velocity in meters per second, but that is different from acceleration. Acceleration is the rate of change in velocity. So if we want to go from 0 to 5 meters per second (that's jogging speed) in two seconds. That is a change in velocity of 2.5 meters per second every second. We would say this acceleration is 2.5m/s^2 .

What about measuring force? Newton decided to name the unit after himself: The force necessary to accelerate one kilogram at 1m/s^2 is known as a *newton*.

Exercise 1 Acceleration

While driving a bulldozer, you come across a train car (with no brakes and no locomotive) on a track in the middle of a city. The train car has a label telling you that it weighs 2,400 kg. There is a bomb welded to the interior of the train car, and the timer tells you that you can safely push the train car for 120 seconds. To get the train car to where it can explode safely, you need to accelerate it to 20 meters per second. Fortunately, the track is level and the train car's wheels have almost no rolling resistance. With what force, in newtons, do you need to push the train for those 120 seconds?

Working Space

Answer on Page 47

1.3 Mass and Gravity

The second interesting thing about mass is that masses are attracted to each other by the force we call *gravity*. The force of attraction between two objects is proportional to the product of their masses. As objects get farther away, the force decreases. That is why you are more attracted to the earth than you are to distant stars, which have much more mass than the earth.

Newton's Law of Universal Gravitation

Two masses (m_1 and m_2) that are a distance of r from each other, are attracted toward each other with a force of magnitude:

$$F = G \frac{m_1 m_2}{r^2}$$

where G is the universal gravitational constant. If you measure the mass in kilograms and the distance in meters, G is about 6.674×10^{-11} . That will get you the force of the attraction in newtons.

Exercise 2 Gravity

The earth's mass is about 6×10^{24} kilograms.

Your spacecraft's mass is 6,800 kilograms. Your spacecraft is also about 100,000 km from the center of the earth. (For reference, the moon is about 400,000 km from the center of the earth.)

What is the force of gravity that is pulling your spacecraft and the earth toward each other?


Working Space

Answer on Page 47

1.4 Mass and Weight

Gravity pulls on things proportional to their mass, so we often ignore the difference between mass and weight.

The weight of an object is the force due to the object's mass and gravity. When we say, "This potato weighs 1 pound," we actually mean "This potato weighs 1 pound on earth." That same potato would weigh about one-fifth of a pound on the moon.



But that potato has a mass of 0.45 kg anywhere in the universe.



CHAPTER 2

Atomic and Molecular Mass

A proton and a neutron have about the same mass. An electron, on the other hand, has much less mass: One neutron weighs about the same amount as 2000 electrons. Thus, the mass of any object comes mostly from the protons and neutrons in the nucleus of its atoms.

We know how many protons an atom has by what element it is, but how do we know the number neutrons?

If you buy a balloon filled with helium, it will have two different kinds of helium atoms: Most of the helium atoms will have 2 neutrons, but a few will have only 1 neutron. We say that these are two different *isotopes* of helium. We call them helium-4 (or ${}^4\text{He}$) and helium-3 (or ${}^3\text{He}$). Isotopes are named for the sum of protons and neutrons the atom has: helium-3 has 2 protons and 1 neutron.

Watch Khan Academy's **Atomic mass, number, and isotopes** at <https://www.khanacademy.org/science/chemistry/atomic-structure-and-properties/introduction-to-the-atom/v/atomic-number-mass-number-and-isotopes>

A hydrogen atom nearly always has just 1 proton and no neutrons. A helium atom nearly

always has 2 protons and 2 neutrons. So, if you have a 100 hydrogen atoms and 100 helium atoms, the helium will have about 4 times more mass than the hydrogen. We say "Hydrogen is about 1 atomic mass unit(amu), and helium-4 is about 4 atomic mass units."

What, precisely, is an atomic mass unit? It is defined as 1/12 of the mass of a carbon-12 atom. Scientists have measured the mass of helium-4, and it is about 4.0026 atomic mass units. (By the way, an atomic mass unit is also called a *dalton*.)

There is a square for each element. In the middle, you see the atomic symbol and the name of the element. In the upper right corner is the atomic number – the number of protons in the atom.

In the upper left corner is the atomic mass in atomic mass units.

Atomic Weight		
Atomic Number	6	12.011
Symbol	C	
Name	Carbon	

Look at the atomic mass of boron. About 80% of all boron atoms have six neutrons. The other 20% have only 5 neutrons. So most boron atoms have a mass of about 11 atomic mass units, but some have a mass of about 10 atomic mass units. The atomic mass of boron is equivalent to the average mass of a boron atom: 10.811.

Exercise 3 Mass of a Water Molecule

Using the periodic table, what is the average mass of one water molecule in atomic mass units?

Working Space

Answer on Page 48

2.1 Molar Mass

An atomic mass unit is a very, very, very small unit; we would much rather work in grams. It turns out that $6.02214076 \times 10^{23}$ atoms equal 1 mole (a standard measure for chemistry). Scientists use this number so much that they gave it a name: *the Avogadro constant* or *Avogadro's number*.

Watch Khan Academy's discussion of the mole at <https://www.khanacademy.org/science/ap-chemistry-beta/x2eef969c74e0d802:atomic-structure-and-properties/x2eef969c74e0d802:moles-and-molar-mass/v/the-mole-and-avogadro-s-number>

If you have 12 doughnuts, that's a dozen doughnuts. If you have $6.02214076 \times 10^{23}$ doughnuts, you have a *mole* of doughnuts. (Note: it isn't practical to measure doughnuts this way: A mole of doughnuts would be about the size of the earth. We use moles for small things like molecules.)

Let's say you want to know how much a mole of NaCl weighs. From the periodic table, you see that Na has an atomic mass of 22.98976 atomic mass units. And Cl has 35.453 atomic mass units. One atom of NaCl has a mass of $22.98976 + 35.453 = 58.44276$ atomic mass units. Then a mole of NaCl has a mass of 58.44276 grams. Handy, right?

Exercise 4 Burning Methane

Natural gas is mostly methane (CH_4). When one molecule of methane burns, two oxygen molecules (O_2) are consumed. One molecule of H_2O and one molecule of CO_2 are produced.

If I need 200 grams of water, how many grams of methane do I need to burn?

(This is how the hero in "The Martian" made water for his garden.)

Working Space

Answer on Page 48

2.2 Heavy atoms aren't stable

When you look at the periodic table, there are a surprisingly large number of elements. You might be told to "Drink milk so that you can get the calcium you need." However, no

one has told you “You should eat kale so that you get enough copernicium in your diet.”

Copernicium, with 112 protons and 173 neutrons, has only been observed in a lab. It is highly radioactive and unstable(meaning it decays): a copernicium atom usually lives for less than a minute before decaying.

The largest stable element is lead, which has 82 protons and between 122 and 126 neutrons. Elements with lower atomic numbers than lead, have at least one stable isotope. Elements with higher atomic numbers than lead don't.

Bismuth, with an atomic number of 83, is *almost* stable. In fact, most bismuth atoms will live for billions of years before decaying.



CHAPTER 3

Work and Energy

In this chapter, we are going to talk about how engineers define work and energy. We have already talked about force. Force is measured in newtons, and one newton is equal to the force necessary to accelerate one kilogram at a rate of 1m/s^2 .

When you lean on a wall, you are exerting a force on the wall, but you aren't doing any work. On the other hand, if you push a car for a mile, you are clearly doing work. Work, to an engineer, is the force you apply to something, as well as the distance that it moves, in the direction of the applied force. We measure work in *joules*. A joule is one newton of force over one meter.

[!!!DECIDE HOW TO IMPLEMENT FORCE VECTORS]



For example, if you push a car uphill with a force of 10 newtons for 12 meters, you have done 120 joules of work.

Work is how energy is transferred from one thing to another. When you push the car, you also burn sugars(energy of the body) in your blood. That energy is then transferred to the car: after it has been pushed uphill.

Thus, we measure the energy something consumes or generates in units of work: joules, kilowatt-hours, horsepower-hours, foot-pounds, BTUs(British Thermal Unit), and calories.

Let's go over a few different forms that energy can take.

Watch Khan Academy's **Changes in energy** at <https://www.khanacademy.org/science/ms-physics/x1baed5db7c1bb50b:energy/x1baed5db7c1bb50b:changes-in-energy/a/changes-in-energy>

3.1 Heat

When you heat something, you are transferring energy to it. The BTU is a common unit for heat: One BTU is the amount of heat required to raise the temperature of one pound of water, by one degree. One BTU is about 1,055 joules. In fact, when you buy and sell natural gas as fuel, it is priced by the BTU.

3.2 Electricity

Electricity is the movement of electrons. When you push electrons through a space that resists their passage (like a light bulb), energy is transferred from the power source (a battery) into the source of the resistance.

Let's say your lightbulb consumes 60 watts of electricity, and you leave it on for 24 hours. We would say that you have consumed 1.44 kilowatt hours or 3,600,000 joules.

Watch Khan Academy's **Introduction to charge** at <https://www.khanacademy.org/science/in-in-class10th-physics/in-in-electricity/in-in-electric-current-circuit/v/intro-to-charge>

3.3 Chemical Energy

As mentioned early, some chemical reactions consume energy and some produce energy. Thus, energy can be stored in the structure of a molecule. When a plant uses photosynthesis to rearrange water and carbon dioxide into a sugar molecule, it converts the energy in the sunlight(solar energy) into chemical energy. Remember photosynthesis is a process

that releases energy. Therefore, the sugar molecule has more chemical energy than the carbon dioxide and water molecules that were used in its creation.

In our diet, we measure this energy in *kilocalories*. A calorie is the energy necessary to raise one gram of water one degree Celsius: it is about 4.19 joules. This is a very small unit: an apple has about 100,000 calories(100 kilocalories), so people working with food started measuring everything in kilocalories.

Here is where things get confusing: People who work with food got tired of saying “kilo-calories”, so they just started using “Calorie” to mean 1,000 calories. This has created terrible confusion over the years. So if the C is capitalized, “Calorie” probably means kilocalorie.

3.4 Kinetic Energy

A mass in motion has energy. For example, if you are in a moving car and you slam on the breaks, the energy from the motion of the car will be converted into heat in the breaks and under the tires.

How much energy does the car have?

Formula for Kinetic Energy

$$E = \frac{1}{2}mv^2$$

where E is the energy in joules, m is the mass in kilograms, and v is the speed in meters per second.

3.5 Gravitational Potential Energy

Watch Khan Academy’s **Potential energy** at <https://youtu.be/oGzwVYPxKjg>

When you lift something heavy onto a shelf, you are giving it *potential energy*. The amount of energy that you transferred to it is proportional to its weight and the height that you lifted it.

On the surface of the earth, gravity will accelerate a heavy object downward at a rate of 9.8m/s^2 .

Formula for Gravitational Potential Energy

On earth, then, gravitational potential energy is given by

$$E = (9.8)mh$$

where E is the energy in joules, m is the mass of the object you lifted, and h is the height that you lifted it.

There are other kinds of potential energy. For example, when you draw a bow, you have given that bow potential energy. When you release it, the potential energy is transferred to the arrow, which expresses it as kinetic energy.

3.6 Conservation of Energy

The first law of thermodynamics says “Energy is neither created nor destroyed.”

Energy can change forms: Your cells consume chemical energy to give gravitational potential energy to a car you push up a hill. However, the total amount of energy in a closed system stays constant.

Exercise 5 The Energy of Falling

A 5 kg cannonball falls off the top of a 3 meter ladder. Just before it hits the floor, all of its gravitational potential energy has been converted into kinetic energy. How fast is the cannonball going when it hits the floor?

Working Space

Answer on Page 48

3.7 Efficiency

Watch Khan Academy’s **Laws of thermodynamics** at <https://www.khanacademy.org/science/ap-biology/cellular-energetics/cellular-energy/a/the-laws-of-thermodynamics>

Although energy is always conserved as it moves through different forms, scientists aren't always that good at controlling it.

For example, a car engine consumes the chemical energy in gasoline. Only about 20% of the energy consumed is used to turn the wheels. Most of the energy is actually lost as heat. If you run a car for a while, the engine gets very hot and the exhaust going out the tailpipe turns hot.

A human is about 25% efficient. Most of the loss is in the heat produced during the chemical reactions that turns food into motion.

In general, if you are trying to increase efficiency in any system, the solution is usually easy to identify because heat is produced. Reduce heat, Increase efficiency.

Light bulbs are an interesting case. To get the light of a 60 watt incandescent bulb, you can use an 8 watt LED or a 16 watt fluorescent light. Thus, we say that the LED light is much more efficient: If you run both, the incandescent bulb will consume 1.44 kilowatt-hours. The LED will consume only 0.192 kilowatt-hours.

Besides light, the incandescent bulb is producing a lot of heat. If it is inside your house, what happens to the heat? It warms your house.

In the winter, when you want light and heat, the incandescent bulb is 100% efficient!

In the summer, if you are running the air conditioner, the incandescent bulb is worse than just "inefficient at making light" – it is actually counteracting the air conditioner!



CHAPTER 4

Units and Conversions

At this point, you are working with a lot of units: grams for weight, joules for energy, newtons for force, meters for distance, seconds for time, etc. For each type of measurement, there are several different units; for example, distance can be measured in feet, miles, and light-years.

Some Equalencies

Distance	
1 mile	1.6093 kilometers
1 foot	0.3048 meters
1 inch	2.54 centimeters
1 light-year	9.461×10^{12} kilometers
Volume	
1 milliliter	1 cubic centimeter
1 quart	0.9461 liters
1 gallon	3.7854 liters
1 fluid ounce	29.6 milliliters
Mass	
1 pound	0.4535924 kilograms
1 ounce	0.4535924 grams
1 metric ton	1000 kilograms
Force	
1 newton	1 kilogram meter per sec ²
Pressure	
1 pascal	1 newton per square meter
1 bar	0.98692 atmosphere
1 pound per square inch	6897 pascals
Energy	
1 joule	1 newton meter
1 calorie	4.184 joules
1 kilowatt-hour	3.6×10^6 joules

(You don't need to memorize these! Just remember that this page is here.)

In the metric system, prefixes are often used to express a multiple. Here are the common prefixes:

Common Prefixes for Metric Units

giga	$\times 10^9$
mega	$\times 10^6$
kilo	$\times 10^3$
milli	$\div 10^3$
micro	$\div 10^6$
nano	$\div 10^9$

(These are worth memorizing. Here's a mnemonic: "King Henry Doesn't Usually Drink Chocolate Milk.")

4.1 Conversion Factors

Here is a really handy trick to remembering how to do conversions between units.

Often, you will be given a table like the one above, and someone will ask you “How many miles are in 0.23 light-years?” You know that 1 mile = 1.6093 kilometers and that 1 light-year is 9.461×10^{12} kilometers. How do you do the conversion?

The trick is to treat the two parts of the equality as a fraction that equals 1. That is, you think:

$$\frac{1 \text{ miles}}{1.6093 \text{ km}} = \frac{1.6093 \text{ km}}{1 \text{ miles}} = 1$$

and

$$\frac{1 \text{ light-years}}{9.461 \times 10^{12} \text{ km}} = \frac{9.461 \times 10^{12} \text{ km}}{1 \text{ light-years}} = 1$$

We call these fractions *conversion factors*.

Now, your problem is

$$0.23 \text{ light-years} \times \text{Some conversion factors} = ? \text{ miles}$$

Note that when you multiply fractions together, things in the numerators can cancel with things in the denominator:

$$\left(\frac{31\pi}{47}\right) \left(\frac{11}{37\pi}\right) = \left(\frac{31\cancel{\pi}}{47}\right) \left(\frac{11}{37\cancel{\pi}}\right) = \left(\frac{31}{47}\right) \left(\frac{11}{37}\right)$$

When working with conversion factors, you will do the same with the units:

$$\begin{aligned} 0.23 \text{ light-years} \left(\frac{9.461 \times 10^{12} \text{ km}}{1 \text{ light-years}}\right) \left(\frac{1 \text{ miles}}{1.6093 \text{ km}}\right) &= \\ 0.23 \cancel{\text{light-years}} \left(\times \frac{9.461 \times 10^{12} \cancel{\text{km}}}{1 \cancel{\text{light-years}}}\right) \left(\frac{1 \text{ miles}}{1.6093 \cancel{\text{km}}}\right) &= \frac{(0.23)(9.461 \times 10^{12})}{1.6093} \text{ miles} \end{aligned}$$

Exercise 6 Simple Conversion Factors

How many calories are in 4.5 kilowatt-hours?

Working Space

Answer on Page 48

4.2 Conversion Factors and Ratios

Conversion factors also work on ratios. For example, if you are told that a bug is moving 0.5 feet every 120 milliseconds. What is that in meters per second?

The problem then is

$$\frac{0.5 \text{ feet}}{120 \text{ milliseconds}} = \frac{? \text{ m}}{\text{second}}$$

So you will need conversion factors to replace the “feet” with “meters” and to replace “milliseconds” with “seconds”:

$$\left(\frac{0.5 \cancel{\text{ feet}}}{120 \cancel{\text{ milliseconds}}} \right) \left(\frac{0.3048 \text{ meters}}{1 \cancel{\text{ feet}}} \right) \left(\frac{1000 \cancel{\text{ milliseconds}}}{1 \text{ second}} \right) = \frac{(0.5)(0.3048)(1000)}{120} \text{ m/second}$$

Exercise 7 Conversion Factors

The hole in the bottom of the boat lets in 0.1 gallons every 2 minutes. How many milliliters per second is that?

Working Space

Answer on Page 49

4.3 When Conversion Factors Don't Work

Conversion factors only work when the units being converted are proportional to each other. Gallons and liters, for example, are proportional to each other: If you have n gallons, you have $n \times 3.7854$ liters.

Degrees celsius and degrees fahrenheit are *not* proportional to each other. If your food is n degrees celsius, it is $n \times \frac{9}{5} + 32$ degrees fahrenheit. You can't use conversion factors to convert celsius to fahrenheit.

Watch Khan Academy's video on this at <https://www.khanacademy.org/test-prep/sat/x0a8c2e5f:untitled-652/x0a8c2e5f:problem-solving-and-data-analysis-lessons-by-skill/a/gtp--sat-math--article--units--lesson>




CHAPTER 5

Simple Machines

As mentioned earlier, physicists define work to be the force applied times the distance it is applied over. So, if you pushed your car 100 meters with 17 newtons of force, you have done 1700 joules of work.

Humans have always had to move really heavy things, so many centuries ago we developed simple machines to decrease the amount of force necessary to execute those tasks. These include things like:

- Levers
- Pulleys
- Ramps
- Gears
- Hydraulics
- Screws



While these machines can decrease the force needed, they don't change the amount of work that must be done. So if the force is decreased to a third, the distance that you must apply the force is increased by a factor of three.

"Mechanical gain" is what we call the increase in force.

5.1 Levers

A lever rotates on a fulcrum. To decrease the necessary force, the load is placed nearer to the fulcrum than where the force is applied.

In particular, physicists talk about the *torque* created by a force. When you push on a lever, the torque is the product of the force you exert and the distance from the point of rotation.

Torque is typically measured in newton-meters.

To balance two torques, the products must be the same. So, assuming that the forces are applied in the proper direction,

$$R_L F_L = R_A F_A$$


where R_L and R_A are the distance from the fulcrum to the where the load's force and the applied force (respectively) are applied, and F_L and F_A are the amounts of the forces.

Exercise 8 Lever


Paul, who weighs 70 kilograms, sits on a see-saw 4 meters from the fulcrum. Jan, who weighs 50 kilograms, wants to balance. How far should Jan sit from the fulcrum?

*Working Space**Answer on Page 49*

Watch Khan Academy's video on levers: <https://www.khanacademy.org/science/physics/discoveries/simple-machines-explorations/a/lever>

**5.2 Ramps**

Ramps, or incline planes, let you roll or slide objects up to a higher level. Steeper ramps give you less mechanical gain. For example, it is much easier to roll a ball up a wheelchair



ramp than on a skateboard ramp.

Assuming the ramp has a constant steepness, the mechanical gain is equal to the ratio of the length of the ramp divided by the amount that it rises.

If you assume there is no friction, the force that you push a weight up the ramp will be:

$$F_A = \frac{V}{L} F_G$$

Where F_A is the force you need to push. L is the length of the ramp, V is the amount of vertical gain and F_G is the force of gravity on the mass.

(We haven't talked about the sine function yet, but in case you already know about it: Note that

$$\frac{V}{L} = \sin \theta$$

where θ is the angle between the ramp and level.)

Exercise 9 Ramp


A barrel of oil weighs 136 kilograms. You can push with a force of up to 300 newtons. You have to get the barrel onto a platform that is 2 meters. What is the shortest board that you can use as a ramp?

Working Space

Answer on Page 49

5.3 Gears

Gears (which might have a chain connecting them like on a bicycle) have teeth and come in pairs. You apply torque to one gear, and it applies torque to another. The torque is increased or decreased based on the ratio between the teeth on the gears.



If N_A is the number of teeth on the gear you are turning with a torque of T_A , and N_L is the number of teeth on the gear it is turning, the resulting torque is:

$$T_L = \frac{N_A}{N_L} T_A$$

Exercise 10 Gears

The bicycle is an interesting case because we are not trying to get mechanical gain. We want to spin the pedals slower with more force.

You like to pedal your bike at 70 revolutions per minute. The chainring that is connected to your pedals has 53 teeth. The circumference of your tire is 2.2 meters. You wish to ride a 583 meters per minute.

How many teeth should the rear sprocket have?


Working Space

Answer on Page 49

Watch Khan Academy's introduction to simple machines here: <https://www.khanacademy.org/science/physics/discoveries/simple-machines-explorations/a/simple-machines-and-how-to-use->

5.4 Hydraulics

In a hydraulic system, like the braking system of a car, you exert force on a piston filled with fluid. The fluid carries that pressure into another cylinder. The pressure of the fluid pushes the piston in that cylinder out.



The pressure in the hose can be measured in pounds per square inch (PSI) or newtons per square meter (Pascals or Pa). We will use Pascals.

To figure out how much pressure you create, you divide the force by the area of the piston head you are pushing.

To figure out how much force that creates on the other end, you multiply the pressure times the area of the piston head that is pushing the load.

Exercise 11 **Hydraulics**

Your car has disc brakes. When you put 2,500,000 pascals of pressure on the brake fluid, the car stops quickly. As the car designer, you would like that to require 12 newtons of force from the driver's foot. What should the radius of the master cylinder (the one the driver is pushing on) be?

Working Space

Answer on Page 50



CHAPTER 6


Buoyancy

When you put a boat into water, it will sink into the water until the mass of the water it displaces is equal to the mass of the boat. We think of this in terms of forces. Gravity pulls the mass of the boat down. The *buoyant force* pushes the boat up. A boat dropped into the water will bob up and down a bit before reaching an equilibrium where the two forces are equal.

Watch Khan Academy's introduction to buoyance at <https://www.khanacademy.org/science/in-in-class9th-physics-india/in-in-gravity/in-in-pressure-in-liquids-archimedes-principle/a/archimedes-principle-buoyancy-fluids-physics-khan-academy>

The buoyant force pushes things up – against the force of gravity. The force is equal to the weight of the fluid being replaced. So, for example, a cubic meter of freshwater has a mass of about 1000kg. If you submerge anything with a volume of one meter in freshwater on earth, the buoyant force will be about 9800 newtons.

For some things, like a block of styrofoam, this buoyant force will be sufficient to carry it to the surface. Once it reaches the surface, it will continue to rise (displacing less water) until the mass of the water it displaces is equal to its mass. And then we say “It floats!”



For some things, like a block of lead, the buoyant force is not sufficient to lift it to the surface, and thus we say “It sinks!”

This is why a helium balloon floats through the air. The air that it displaces weighs more than the balloon and the helium itself. (It is easy to forget that air has a mass, but it does.)

Exercise 12 Buoyancy

You have an aluminum box that has a heavy base, so it will always float upright. The box and its contents weigh 10 kg. Its base is 0.3 m x 0.4 m. It is 1m tall. When you drop it into freshwater ($1000\text{kg}/\text{m}^3$), how far will it sink before it reaches equilibrium.

Working Space

Answer on Page 50

6.1 The Mechanism of Buoyancy

As you dive down in the ocean, you will experience greater and greater pressure from the water. And if you take a balloon with you, you will gradually see it get smaller as the water pressure compresses the air in the balloon.

Let's say you are 3 meters below the surface of the water. What is the pressure in Pascals (newtons per square meter)? You can think of the water as a column of water crushing

down upon you. The pressure over a square meter is the weight of 3 cubic meters of water pressing down.


$$p = (3)(1000)(9.8) = 29,400 \text{ Pa}$$

This is called *hydrostatic pressure*. The general rule for hydrostatic pressure in Pascals p is

$$p = dgh$$

Where d is the density of the fluid in kg per cubic meter, g is the acceleration due to gravity in m/s^2 , and h is the height of the column of fluid above you.

So, where does buoyant force come from? Basically, the pressure pushing up on the deepest part of the object is higher than the pressure pushing down on the shallowest part of the object. That is where buoyancy comes from.



Exercise 13 **Hydrostatic Pressure**

You dive into a tank of olive oil on Mars. How much more hydrostatic pressure does your body experience at 5 meters deep than it did at the surface?

The density of olive oil is about 900 kg per square meter. The acceleration due to gravity on Mars is 3.721 m/s^2 .

Working Space

Answer on Page 50

Notice that although the pressure is increasing as you go deeper, the buoyant force will *not increase* because the buoyant force is always equal to the weight of the fluid that is displaced, regardless if that is 1 meter or 100 meters underwater.

Also, saltwater is denser than freshwater. That is why people float better in the sea than they do in a river.

And, lipids, like fats and oils, are less dense than water. That is why people with a lot of body fat tend to float better than people with less body fat. And why oil floats in a glass of water.



CHAPTER 7

Heat

Let's say you put a 1 kg aluminum pan that is 80°C into 3 liters of water that is 20°C . Energy, in the form of heat, will be transferred from the pan to the water until they are at the same temperature. (We call this "thermal equilibrium.")

What will the temperature of the water be?

7.1 Specific Heat Capacity

If you are heating something, the amount of energy you need to transfer to it depends on three things: the mass of the thing you are heating, the amount of temperature change you want, and the *specific heat capacity* of that substance.

Energy in Heat Transfer

The energy moved in a heat transfer is given by

$$E = mc\Delta_T$$

where m is the mass, Δ_T is the change in temperature, and c is the specific heat capacity of the substance.


(Note that this assumes no phase change. For example, this formula works nicely on warming liquid water, but it gets more complicated if you warm the water past its boiling point.)

Can we guess the specific heat capacity of a substance? It is very, very difficult to guess the specific heat of a substance, so we determine it by experimentation.

For example, someone determined that it took about 0.9 joules to raise the temperature of solid aluminum one degree Celsius. So we say "The specific heat capacity of aluminum is 0.9 J/g °C."

The specific heat capacity of liquid water is about 4.2 J/g °C.

To answer the question, then, the amount of energy given off by the pan must equal the amount of energy absorbed by the water. And they need to be the same temperature at the end. Let T be the final temperature of both.



Watch Khan Academy's discussion of heat capacity at <https://www.khanacademy.org/science/ap-chemistry-beta/x2eef969c74e0d802:thermodynamics/x2eef969c74e0d802:heat-capacity-and-calorimetry/v/heat-capacity>

Three liters of water weighs 3,000 grams, so the change in energy in the water will be:

$$E_W = mc\Delta_T = (3000)(4.2)(T - 20) = 12600T - 252000 \text{ joules}$$

The pan weighs 1000 grams, so the change in energy in the pan will be::

$$E_P = mc\Delta_T = (1000)(0.9)(T - 80) = 900T - 72000 \text{ joules}$$

Total energy stays the same so $E_W + E_P = 0$. So you need to solve

$$(12600T - 252000) + (900T - 72000) = 0$$

And find that the temperature at equilibrium will be

$$T = 24^\circ\text{C}$$

Exercise 14 Thermal Equilibrium

Just as you put the aluminium pan in the water as described above, someone also puts a 1.2 kg block of copper cooled to 10°C . The specific heat of solid copper is about $0.4 \text{ J/g } ^\circ\text{C}$.

What is the new temperature at equilibrium?


Working Space

Answer on Page 50

7.2 Getting to Equilibrium

When two objects with different temperatures are touching, the speed at which they exchange heat is proportional to the differences in their temperatures. Thus, as their temperatures get closer together, the heat exchange slows down.

In our example, the pan and the water will get close to equilibrium quickly, but they may never actually reach equilibrium.



Exercise 15 Cooling Your Coffee

You have been given a ridiculously hot cup of coffee and a small pitcher of chilled milk.

You need to start chugging your coffee in three minutes, and you want it as cool as possible at that time. When should you add the milk to the coffee?

Working Space

Answer on Page 51

7.3 Specific Heat Capacity Details

For any given substance, the specific heat capacity often changes a lot when the substance changes state. For example, ice is $2.1 \text{ J/g } ^\circ\text{C}$, whereas liquid water is $4.2 \text{ J/g } ^\circ\text{C}$.

Watch Khan Academy's discussion of the specific heat of water: <https://www.khanacademy.org/science/biology/water-acids-and-bases/water-as-a-solid-liquid-and-gas/v/specific-heat-of-water>

Even within a given state, the specific heat capacity varies a bit based on the temperature and pressure. If you are trying to do these sorts of calculations with great accuracy, you will want to find the specific heat capacity that matches your situation. For example, I might look for the specific heat capacity for water at 22°C at 1 atmosphere of pressure(

atm).



APPENDIX A

Answers to Exercises

Answer to Exercise 1 (on page 7)

To get the train to 20 meters per second in 120 seconds, you must accelerate it with a constant rate of $\frac{1}{6} \text{ m/s}^2$. You remember that $F = ma$, so $F = 2400 \times \frac{1}{6}$. Thus, you will push the train with a force of 400 newtons for the 120 seconds before the bomb goes off.

Answer to Exercise 2 (on page 8)

$$F = G \frac{m_1 m_2}{r^2} = (6.674 \times 10^{-11}) \frac{(6.8^3)(6 \times 10^{24})}{(10^5)^2} = 6.1 \times 10^6$$

About 6 million newtons.

Answer to Exercise 3 (on page 14)

The average hydrogen atom has a mass of 1.00794 atomic mass units.

The average oxygen atom has a mass of 15.9994.

$$2 \times 1.00794 + 15.9994 = 18.01528 \text{ atomic mass units.}$$

Answer to Exercise 4 (on page 15)

From the last exercise, you know that 1 mole of water weighs 18.01528 grams. So 200 grams of water is about 11.1 moles. So you need to burn 11.1 moles of methane.

What does one mole of methane weigh? Using the periodic table: $12.0107 + 4 \times 1.00794 = 16.04246$ grams.

$$16.0424 \times 11.10 = 178.1 \text{ grams of methane.}$$

Answer to Exercise 5 (on page 20)

At the top of the ladder, the cannonball has $(9.8)(5)(3) = 147$ joules of potential energy.

At the bottom, the kinetic energy $\frac{1}{2}(5)v^2$ must be equal to 147 joules. So $v^2 = \frac{294}{5}$. Thus it is going about 7.7 meters per second.

(Yes, a tiny amount of energy is lost to air resistance. For a dense object moving at these relatively slow speeds, this energy is negligible.)

Answer to Exercise 6 (on page 26)

$$4.5 \text{ kWh} \left(\frac{3.6 \times 10^6 \text{ joules}}{1 \text{ kWh}} \right) \left(\frac{1 \text{ calories}}{4.184 \text{ joules}} \right) = \frac{(4.5)(3.6 \times 10^6)}{4.184} = 1.08 \times 10^6 \text{ calories}$$

Answer to Exercise ?? (on page 26)

$$\frac{0.1 \text{ gallons}}{2 \text{ minutes}} \left(\frac{3.7854 \text{ liters}}{1 \text{ gallons}} \right) \left(\frac{1000 \text{ milliliters}}{1 \text{ liters}} \right) \left(\frac{1 \text{ minutes}}{60 \text{ seconds}} \right) =$$

$$\frac{(0.1)(3.7854)(1000)}{(2)(60)} \text{ ml/second} = 3.1545 \text{ ml/second}$$

Answer to Exercise 8 (on page 31)

Paul is exerting $(70)(9.8)$ newtons of force at 4 meters from the fulcrum, so he is creating a torque of 2,744 newton-meters of torque on the see-saw. Jan is creating $(50)(9.5) = 490$ newtons of force.

If r is the distance from the fulcrum to Jan's seat, to balance $490r = 2744$, so $r = 5.6$ meters.

Answer to Exercise 9 (on page 33)

To lift the barrel would require $136 \times 9.8 = 1,332.8$ newtons of force.

Letting L be the length of the ramp:

$$300 = \frac{2}{L} 1332.8$$

So $L = 8.885$ meters.

Answer to Exercise 9 (on page 34)

$$583 = (70)(2.2) \frac{53}{n}$$

Thus $n = 14$ teeth.

Answer to Exercise 11 (on page 35)

We are looking for r , the radius of the piston head in meters. The area of the piston head is πr^2 .

The pressure in pascals of the brake fluid is given by $12/(\pi r^2)$.

$$2,500,000 = \frac{12}{\pi r^2}$$

$$\text{So } r = \sqrt{\frac{12}{\pi \times 2.5 \times 10^6}} = 0.001236077446474 \text{ meters.}$$

Answer to Exercise 12 (on page 38)

Equilibrium will be achieved when the box has displaced 10 kg of water. That is, when it has displaced 0.01 cubic meters.

The area of the base of the box is 0.12 square meters. So if the box sinks x meters into the water it will displace $0.12x$ cubic meters.

Thus at equilibrium $x = \frac{0.01}{0.12} \approx 0.083$ m. So, the box will sink 8.3 cm into the water before reaching equilibrium.

Answer to Exercise 13 (on page 40)

$$p = dgh = (900)(3.721)(5) = 16,744.5 \text{ Pa}$$

Answer to Exercise 14 (on page 43)

$$E_C = (1200)(0.4)(T - 10) = 480T - 4800$$

Total energy stays constant:

$$0 = (12600T - 252000) + (900T - 72000) + (480T - 4800)$$

Solving for T gets you $T = 23.52^\circ \text{C}$.

Answer to Exercise 15 (on page 44)

During the 3 minutes, you want the coffee to give off as much of its heat as possible, so you want to maximize the difference between the temperature of the coffee and the temperature of the room around it.

You wait until the last moment to put the milk in.



INDEX

atom, 4
atomic mass, 14
atomic mass unit, 12
Avogadro's number, 15

BTU, 18

calories, 19
career, 3
chemical energy, 18
chemical reaction, 5
conversion factors, 25

efficiency, 21
electricity, 18
electrons, 4
elements, 5
endothermic, 6
energy
 conservation of, 20
exothermic, 5

heat, 18

isotopes, 11

Joule, 17

kinetic energy, 19

metric system
 prefixes, 24
mole, 15

molecules, 5

neutron, 11
neutrons, 4

periodic table of elements, 13
potential energy
 gravitational, 19

proton, 11
protons, 4

quitting, 3

specific heat capacity, 41

thermal equilibrium, 41

units table, 24

work, 18