

Common Diseases & Contraindicated Drugs

7 common conditions • 5 specific drug names each • academic study format

Important: Contraindication depends on the exact condition, severity, indication, dose and current labeling. Some entries are more accurately classified as avoid/major caution in specific circumstances.

1. Hypertension (High BP)

Specific drug name	Class	Status	Main reason	Warning
Ibuprofen	NSAID	AVOID / CAUTION	Fluid retention and reduced antihypertensive effect	May increase sodium/water retention and raise BP or worsen BP control.
Pseudoephedrine	Decongestant	AVOID	Sympathetic stimulation	Can increase blood pressure and heart rate.
Phenylephrine	Decongestant	AVOID	Vasoconstriction	Can cause vasoconstriction and raise blood pressure.
Ergotamine	Ergot alkaloid	CONTRAINDICATED in uncontrolled hypertension	Marked vasoconstriction	May significantly increase vascular resistance and BP.
Cocaine	Sympathomimetic stimulant	CONTRAINDICATED	Severe sympathetic stimulation	Can cause marked hypertension, tachycardia and cardiovascular complications.

2. Diabetes Mellitus

Specific drug name	Class	Status	Main reason	Warning
Prednisolone	Corticosteroid	AVOID / CAUTION	Hyperglycemia	Can increase hepatic glucose production and reduce insulin sensitivity.
Hydrochlorothiazide	Thiazide diuretic	CAUTION	Impaired glucose tolerance	May increase blood glucose, particularly at higher doses.
Olanzapine	Atypical antipsychotic	CAUTION / AVOID when possible	Hyperglycemia	Can worsen glucose regulation and increase blood glucose.
Niacin	Vitamin B3	CAUTION	Reduced insulin sensitivity	High doses can increase blood glucose.
Salbutamol (albuterol)	β2-agonist bronchodilator	CAUTION	Hyperglycemia	Systemic β2 stimulation can increase glucose levels.

3. Acute Myocardial Infarction (Heart Attack)

Specific drug name	Class	Status	Main reason	Warning
Ibuprofen	NSAID	AVOID / CAUTION	Increased cardiovascular risk	NSAIDs can increase thrombotic cardiovascular risk and may worsen outcomes after MI.
Sildenafil	PDE-5 inhibitor	CONTRAINDICATED with nitrates	Severe hypotension	Potentiates nitrate-mediated vasodilation and can cause a dangerous fall in BP.
Vardenafil	PDE-5 inhibitor	CONTRAINDICATED with nitrates	Severe hypotension	Combination with nitrates can cause a dangerous fall in BP.
Tadalafil	PDE-5 inhibitor	CONTRAINDICATED with nitrates	Severe hypotension	Potentiates nitrate-induced vasodilation.
Ergotamine	Ergot alkaloid	AVOID	Coronary vasoconstriction	Can cause vasoconstriction and worsen myocardial ischemia.

4. Stroke

Specific drug name	Class	Status	Main reason	Warning
Alteplase	Thrombolytic	CONTRAINDICATED after recent intracranial hemorrhage	Intracranial bleeding	Thrombolysis can cause or worsen life-threatening intracranial bleeding.
Warfarin	Anticoagulant	CONTRAINDICATED in active intracranial bleeding	Increased bleeding	Can worsen active intracranial hemorrhage.

Specific drug name	Class	Status	Main reason	Warning
Apixaban	DOAC anticoagulant	CONTRAINDICATED in active pathological bleeding	Increased bleeding	Should not be used during active pathological bleeding.
Rivaroxaban	DOAC anticoagulant	CONTRAINDICATED in active pathological bleeding	Increased bleeding	Can worsen active bleeding.
Dabigatran	DOAC anticoagulant	CONTRAINDICATED in active pathological bleeding	Increased bleeding	Can worsen active bleeding.

5. Heart Failure

Specific drug name	Class	Status	Main reason	Warning
Ibuprofen	NSAID	AVOID / MAJOR CAUTION	Fluid retention	Can cause sodium/water retention and worsen renal function and heart failure.
Verapamil	Non-DHP calcium-channel blocker	AVOID in HFrEF	Negative inotropy	Can reduce myocardial contractility and worsen systolic heart failure.
Diltiazem	Non-DHP calcium-channel blocker	AVOID in HFrEF	Negative inotropy	Can depress cardiac contractility in systolic heart failure.
Pioglitazone	Thiazolidinedione	CONTRAINDICATED in severe heart failure	Fluid retention	Can cause edema and worsen heart failure.
Rosiglitazone	Thiazolidinedione	AVOID / CONTRAINDICATED in severe HF	Fluid retention	Can cause fluid retention and exacerbate heart failure.

6. Asthma

Specific drug name	Class	Status	Main reason	Warning
Propranolol	Non-selective β -blocker	CONTRAINDICATED / AVOID	Bronchospasm	β 2 blockade can cause bronchoconstriction and worsen asthma.
Timolol	Non-selective β -blocker	CONTRAINDICATED / AVOID	Bronchospasm	Even ophthalmic timolol can produce systemic β -blockade and bronchospasm.
Aspirin	NSAID / antiplatelet	CONTRAINDICATED in aspirin-sensitive asthma	Bronchospasm	Can trigger bronchospasm in NSAID-exacerbated respiratory disease.
Ketorolac	NSAID	AVOID in NSAID-sensitive asthma	Bronchospasm	Can precipitate bronchospasm in susceptible patients.
Diclofenac	NSAID	AVOID in NSAID-sensitive asthma	Bronchospasm	Can trigger bronchospasm in susceptible patients.

7. Peptic Ulcer / Active GI Bleeding

Specific drug name	Class	Status	Main reason	Warning
Ibuprofen	NSAID	CONTRAINDICATED / AVOID	GI ulceration and bleeding	Can damage gastric mucosa and worsen ulceration and bleeding.
Diclofenac	NSAID	CONTRAINDICATED / AVOID	GI bleeding	Can increase gastrointestinal mucosal injury and bleeding.
Ketorolac	NSAID	CONTRAINDICATED / AVOID	GI ulceration and bleeding	Has significant gastrointestinal bleeding risk.
Aspirin	Antiplatelet / NSAID	CONTRAINDICATED in active major bleeding	Bleeding risk	Inhibits platelet aggregation and can worsen bleeding.
Naproxen	NSAID	CONTRAINDICATED / AVOID	GI ulceration and bleeding	Can damage gastric mucosa and increase gastrointestinal bleeding risk.