

Geomagnetically Induced Currents



G. S. Lakhina¹, R. Hajra² and B. T. Tsurutani³

¹Indian Institute of Geomagnetism, New Panvel (W), Navi Mumbai, India

²Indian Institute of Technology Indore, Simrol, Indore, India

³Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA

Definition

Geomagnetically induced current (GIC)	Intense, low-frequency, quasi-direct currents flowing in the conductor systems owing to rapid changes of the geomagnetic fields
Space weather	Dynamic variations at the Sun and in interplanetary space disturbing the near-Earth space environment creating conditions that are hostile to space-borne and ground-based technological systems and can even impose danger to human life

Introduction

Geomagnetically induced currents (GICs) are intense, low-frequency (~ 0.001 – 1 Hz) currents flowing in Earth-ground conductor systems owing to rapid changes of the geomagnetic fields ($dB/dt > 1$ nT s⁻¹) (Campbell 1980; Akasofu and Aspin 1982). According to Faraday's law of induction, a fluctuating geomagnetic field induces geoelectric fields which drive GICs in the ground and electrical power systems. The induced currents are essentially quasi-DC as their frequencies are much less than the frequency of the electrical grids operating at 50 or 60 Hz.

Intense GICs are known to cause extensive and expensive damages to a variety of technological systems such as electric power systems, telecommunication cables, and natural gas

pipelines. Strong GICs enter the neutral grounding of transformers (operating typically at ~ 50 Hz frequency) causing transformer core magnetic flux increase and consequent half-cycle saturation of the core. These in turn distort the AC waveform of the power signal, which can lead to transformer overheating, system-relay interference, reactive power loss, or even system damage.

Space Weather Events and Geomagnetically Induced Currents

Space weather generally refers to the physical conditions in the Sun-Earth system that can affect the performance of spacecraft and ground-based technological systems, including electric power distribution systems. The solar sources of the space weather are the solar activity transients such as solar flares and coronal mass ejections (CMEs), solar wind high-speed (~ 550 – 800 km s⁻¹) streams (HSSs), and corotating interaction regions (CIRs) formed by the HSS interaction with slow (~ 300 – 400 km s⁻¹) solar-wind streams. During all of the above phenomena, significant variability occurs in the solar-wind speed, plasma density, and strength and direction of the magnetic field. The dynamics of geospace, extending from the Earth's plasma environment called the ionosphere to the protecting magnetic bubble called the magnetosphere, is largely affected by this solar-wind variability.

Magnetic reconnection between the interplanetary magnetic field (IMF) and the dayside geomagnetic field is considered to be the main mechanism of energy transfer from solar wind to the Earth's magnetosphere (Dungey 1961). The process of magnetic reconnection works most efficiently when IMF is directed southward, opposite to the dayside magnetopause magnetic field direction (Tsurutani and Gonzalez 1997). Dayside magnetic reconnection leads to injection and storage of solar wind kinetic energy (energetic particles) in the magnetotail. This causes stretching of the magnetotail and formation of a near-Earth neutral line. Onset of magnetic

reconnection at this neutral line gives rise to an explosive release of energy from the magnetotail toward the inner magnetosphere. This near-midnight plasma injection excites a substorm (Tsurutani and Meng 1972), and auroras in the high-latitude nightside ionospheric region. If the southward IMF is intense with hours duration, such as during the magnetic cloud portion of an interplanetary coronal mass ejection (ICME) (Burlaga et al. 1981), the plasma injection is deeper and the particles get energized to $\sim 10\text{--}300$ keV. This leads to intensification of the ring current which produces a diamagnetic decrease in the Earth's magnetic field measured at near-equatorial magnetic stations. This latter effect is the *main phase* of the geomagnetic storm. After the storm main phase is over, the decay of the ring current starts the *recovery phase* of the magnetic storm. The energetic particles are lost by several physical processes, namely, charge-exchange, Coulomb collisions, wave-particle interactions, and convection out the front side of the magnetopause (Kozyra and Liemohn 2003).

The intensity of the magnetic storm is measured by the disturbance storm time (Dst) index, an hourly geomagnetic activity index, or by the SYM-H index with a temporal resolution of 1 min. The intensity of a substorm is measured by the auroral electrojet (AE) index. The intensities of both substorms and magnetic storms are expressed in nT. During intense magnetic storms, near-equatorial magnetic field disturbances of a few 100s nT are recorded for a few hours to days. This ground magnetic effect is due to the magnetospheric ring current which exists from $L = 2$ to $L = 8$ (L is the distance in Earth's radius R_E that a dipole magnetic field crosses the magnetic equator). During super magnetic storms, the magnetic disturbances could be higher, exceeding 1000 nT at times. Substorms cause high-latitude ground-magnetic disturbances from ~ 100 to 1000s nT lasting for a few minutes to hours. These magnetic effects are due to a combination of the auroral electrojet which is located in the ionosphere at a height of ~ 100 km and field-aligned currents coming from the magnetosphere to the ionosphere. The ring current effects of substorms are relatively minor compared to that of magnetic storms. During magnetic storm main phases, many substorms may also occur.

A second means of energy transfer from the solar wind to the magnetosphere is sudden solar wind ram pressure changes. One mechanism is the pressure pulse due to high plasma densities sunward of fast magnetosonic shocks (Kennel et al. 1985). These cause sudden impulses (SI^+ s), strong changes of the equatorial magnetic field of $10\text{--}100$ nT with $\sim 1\text{--}10$ min durations (Araki 1977), the triggering of nightside substorms (Heppner 1955), and in extreme cases, the formation of new radiation belts (Blake et al. 1992).

Geomagnetic field disturbances associated with magnetic storms and substorms are known as the key factor for the generation of GICs. Strongest GIC events are recorded in the

high ($>50^\circ$) geomagnetic latitude zone in association with large amplitude magnetic field fluctuations during intense and frequent auroral activities (Pirjola 2000). However, recent studies show evidence of mid-to-low latitude GICs in association with SI^+ events caused by solar wind ram pressure impingement on the dayside magnetopause of the Earth (Gaunt 2014; Carter et al. 2015).

Generation of GIC Events

A GIC event is generated by the high-frequency changes in geomagnetic fields due to space weather events. Geomagnetic storms, substorms, supersubstorms, magnetic pulsations, and sudden impulses (SI^+) are the major drivers for the generation of GICs (Pulkkinen et al. 2005; Viljanen et al. 2006; Ngwira et al. 2014; Tsurutani et al. 2020). In fact, a GIC event is essentially the ground manifestation of the complex space-weather chain starting from the solar eruptions, extending to the interplanetary space, and then to the surface of the Earth and below. Therefore, the GIC signals carry information about the processes involved in the Sun-Earth connection and space weather. Most models calculate GIC either from the measurements of geomagnetic field (B) and its time derivative dB/dt using magnetometers or from the magnetotelluric measurements of both B field and electric field E (Gaunt 2016; Pulkkinen et al. 2017).

Systematic studies on GICs and their effects on the electric power systems were started first in the high-latitude countries, e.g., Finland (Viljanen and Pirjola 1994) and Canada (Boteler 2001), where GICs might be expected to occur. Later, the GIC studies have been undertaken in the low- and mid-latitude regions such as Scotland, China, Japan, South Africa, Brazil, New Zealand, and Australia. An excellent overview of GIC impacts on electrical power systems is given by Gaunt (2016). Direct current (DC) blocking devices, such as neutral-point reactors and series capacitors, are being designed and installed in high voltage transmission grids to mitigate the flow of GICs through transformers (Kappenman et al. 1991; Thomson et al. 2010). The real-time monitoring of GICs provides a useful means to understand the power-system status, but it does not guarantee that the system will survive a future severe GIC event (Erinmez et al. 2002). At present, no reliable forecastable model for GICs exists (Gaunt 2016).

The severity of the GICs flowing through ground-based technological systems depends on several factors such as severity of the magnetic disturbances in the magnetosphere-ionosphere system, location and configuration of the system (i.e., power transformers, transmission lines, etc.), and conductivity of the Earth's surface and interior, coastal effects, etc. (Cilverd et al. 2018; Liu et al. 2018). The highest GIC levels occur for the fastest rate of change of the magnetic field. Therefore, dB/dt is considered as an important parameter to

quantify the relative level of GIC threat. For example, the Québec power station blackout during the 13–14 March 1989 magnetic superstorm occurred when the peak dB/dt was ~ 480 nT/min. Recent study of the March 1989 magnetic storm by Boteler (2019) shows that the peak dB/dt occurred during a shock-induced substorm, much before this superstorm attained its peak $Dst = -589$ nT value. It is interesting to note that dB/dt values exceeding 2000 nT/min have been observed, and much higher values ~ 5000 nT/min might have occurred during historic magnetic storms (Kappenman 2006). Recent studies show that compression of the magnetosphere caused by interplanetary shock impacts, discussed earlier, increases the magnetopause currents leading to large dB/dt variations in the mid- to low- latitudes as well as in the equatorial electrojets (Carter et al. 2015; Oliveira et al. 2018). Furthermore, in many magnetic storms, the GICs due to the SI^+ can be much larger than those recorded at the same station during the storm main phase (Zhang et al. 2015). Thus, even the power stations situated near the equatorial electrojet are susceptible to GIC risks. For maximum possible dB/dt values due to SI^+ s, we refer the reader to Tsurutani and Lakhina (2014).

Future Outlook

Impacts of GICs on electric power systems, natural gas pipelines, and telecommunications have direct socioeconomic impacts. This can be understood from some historical examples of large GIC events. The largest recorded magnetic storm at Earth with a peak $Dst = -1760$ nT was caused by a CME associated with an extremely intense solar flare on 1 September 1859 (Tsurutani et al. 2003). GICs on telegraph lines caused arcing that led to fires and electrical shocks (Loomis 1861). Two recent examples of GIC-related hazards are the collapse of the entire Hydro Québec power system during a superstorm (with peak $Dst = -589$ nT) on 13–14 March 1989 (Allen et al. 1989), and the transformer collapse near Malmö in Sweden during the 29–30 October 2003 superstorm (with peak $Dst \sim -400$ nT) (Pulkkinen et al. 2005; Wik et al. 2008). Due to growing dependence of modern society on electric technology (and greater vulnerability), large GIC impacts are now realized to be quite damaging to society as they can disrupt functioning of the power grids and communication cables, and cause erosion of oil and gas pipelines (Royal Academy of Engineering report 2013; Pulkkinen et al. 2017). Currently GICs are considered as the greatest space-weather hazard globally.

There has been tremendous progress in understanding the space weather processes and generation of GICs during the past two decades. An exhaustive review on the current status and future challenges of space-weather physics and solar-terrestrial physics (STP) is given by Tsurutani et al. (2020).

The ultimate challenge for the space community is to predict the occurrence of extreme magnetic storms and their associated extreme GICs. Although extreme events are rare, yet they are the ones that can cause the maximum damage. Presently, there is a limited understanding of the upper limit for the geoelectric fields that drive GICs. Additionally, the prediction of GICs with a lead time of 1–3 days is very limited or unreliable (Gaunt 2016; Pulkkinen et al. 2017) as reliable estimates for the strength of magnetic storms a couple of days in advance are not available. The major sources of uncertainty are prediction of the IMF southward components associated with the ICME and in the sheath region behind the interplanetary shock. Solar wind measurements taken upstream of the Earth at the Lagrange 1 (L1) point by NASA's Advanced Composition Explorer (ACE) and Deep Space Climate ObserVatoRy (DSCOVR) are being used to provide reliable estimates on the solar-wind drivers. This information is used in the physics-based and empirical models to calculate the currents in the magnetosphere and ionosphere to make real-time prediction of GICs with a lead time of ~ 30 min or so. For extreme events, such short-time notice is not sufficient to put any mitigation plan in place against hazardous extreme GICs. Since GICs depend on the ground conductivity and characteristics of regional power grids, it is a big challenge for the modelers to develop global GIC models that can predict the magnitude and duration of GIC events in different parts of the world.

Cross-References

- Electromagnetic Induction
- Electromagnetic Pulsations and Magnetic Storms
- Extreme Events
- Magnetotellurics
- Space Weather

Acknowledgments GSL thanks Indian National Science Academy, New Delhi, for support under the INSA-Honorary Scientist Scheme. The work of RH is funded by the Science & Engineering Research Board (SERB), a statutory body of the Department of Science & Technology (DST), Government of India through Ramanujan Fellowship. Portions of the research has been performed at the Jet Propulsion Laboratory, California Institute of Technology under contract with NASA.

Bibliography

- Akasofu SI, Aspnes JD (1982) Auroral effects on power transmission line systems. *Nature* 295:136–137. <https://doi.org/10.1038/295136a0>
- Allen J, Frank L, Sauer H, Reiff P (1989) Effects of the March 1989 solar activity. *EOS Trans* 70:1479–1488. <https://doi.org/10.1029/89EO00409>

- Araki T (1977) Global structure of geomagnetic sudden commencements. *Planet Space Sci* 25(4):373–384. [https://doi.org/10.1016/0032-0633\(77\)90053-8](https://doi.org/10.1016/0032-0633(77)90053-8)
- Blake JB, Kolassinski WA, Fillius RA, Mullen EG (1992) Injection of electrons and protons with energies of tens of MeV into $L < 3$ on March 24, 1991. *Geophys Res Lett* 19:821–824. <https://doi.org/10.1029/92GL00624>
- Boteler DH (2001) Assessment of geomagnetic hazard to power systems in Canada. *Nat Hazards* 23:101–120. <https://doi.org/10.1023/A:1011194414259>
- Boteler DH (2019) A 21st century view of the March 1989 magnetic storm. *Space Weather* 17:1427–1441. <https://doi.org/10.1029/2019SW002278>
- Burlaga LF, Sittler E, Mariani F, Schwenn R (1981) Magnetic loop behind an interplanetary shock: voyager, Helios, and IMP 8 observations. *J Geophys Res* 86:6673–6684. <https://doi.org/10.1029/JA086iA08p06673>
- Campbell WH (1980) Observation of electric currents in the Alaska oil pipeline resulting from auroral electrojet current sources. *Geophys J Int* 61:437–449
- Carter BA, Yizengaw E, Pradipta R, Halford AJ, Norman R, Zhang K (2015) Interplanetary shocks and the resulting geomagnetically induced currents at the equator. *Geophys Res Lett* 42:6554–6559. <https://doi.org/10.1002/2015GL065060>
- Cilverd MA, Rodger CJ, Brundell JB, Dalzell M, Martin I, MacManus DH et al (2018) Long-lasting geomagnetically induced currents and harmonic distortion observed in New Zealand during the 7–8 September 2017 disturbed period. *Space Weather* 16:704–717. <https://doi.org/10.1029/2018SW001822>
- Dungey JW (1961) Interplanetary magnetic field and the auroral zones. *Phys Rev Lett* 6:47–48. <https://doi.org/10.1103/PhysRevLett.6.47>
- Erinmez IA, Kappenman JG, Radasky WA (2002) Management of the geomagnetically induced current risks on the national grid company's electric power transmission system. *J Atmos Solar Terr Phys* 64:743–756
- Gaunt CT (2014) Reducing uncertainty – responses for electricity utilities to severe solar storms. *J. Space Weather Space Clim* 4:A01. <https://doi.org/10.1051/swsc/2013058>
- Gaunt CT (2016) Why space weather is relevant to electrical power systems. *Space Weather* 14:2–9. <https://doi.org/10.1002/2015SW001306>
- Heppner JP (1955) Note on the occurrence of world-wide SSCs during the onset of negative bays at college, Alaska. *J Geophys Res* 60:29–32. <https://doi.org/10.1029/JZ060i001p00029>
- Kappenman JG (2006) Great geomagnetic storms and extreme impulsive geomagnetic field disturbance events – an analysis of observational evidence including the great storm of May 1921. *Adv Space Res* 38:188–199. <https://doi.org/10.1016/j.asr.2005.08.055>
- Kappenman JG, Norr SR, Sweezy GA, Carlson DL, Albertson VD, Harder JE, Damsky BL (1991) GIC mitigation: a neutral blocking/bypass device to prevent the flow of GIC in power systems. *IEEE Trans Power Delivery* 6(3):1271–1281. <https://doi.org/10.1109/61.85876>
- Kennel CF, Edmiston JP, Hada T (1985) A quarter century of collisionless shock research. In: Stone RG, Tsurutani BT (eds) *Collisionless shocks in the heliosphere: a tutorial review*, Geophysical monograph series, vol 34. AGU, Washington, pp 1–36
- Kozyra JU, Liemohn MW (2003) Ring current energy input and decay. *Space Sci Rev* 109:105–131. <https://doi.org/10.1023/B:SPAC.0000007516.10433.ad>
- Liu CM, Wang X, Wang H, Zhao H (2018) Quantitative influence of coast effect on geomagnetically induced currents in power grids: a case study. *J Space Weather Space Clim* 8:A60. <https://doi.org/10.1051/swsc/2018046>
- Loomis E (1861) On the great auroral exhibition of Aug. 28th to Sept. 4, 1859, and on auroras generally. *Am J Sci* 82:318–335
- Ngwira CM, Pulkkinen A, Kuznetsova MM, Gloer A (2014) Modeling extreme “Carrington-type” space weather events using three-dimensional MHD code simulations. *J Geophys Res* 119:4456–4474. <https://doi.org/10.1002/2013JA019661>
- Oliveira DM, Arel D, Raeder J, Zesta E, Ngwira CM, Carter BA et al (2018) Geomagnetically induced currents caused by interplanetary shocks with different impact angles and speeds. *Space Weather* 16:636–647. <https://doi.org/10.1029/2018SW001880>
- Pirjola R (2000) Geomagnetically induced currents during magnetic storms. *IEEE Trans Plasma Sci* 28:1867–1873. <https://doi.org/10.1109/27.902215>
- Pulkkinen A, Lindahl S, Viljanen A, Pirjola R (2005) Geomagnetic storm of 29–31 October 2003: geomagnetically induced currents and their relation to problems in the Swedish high-voltage power transmission system. *Space Weather* 3:S08C03. <https://doi.org/10.1029/2004SW000123>
- Pulkkinen A, Bernabeu E, Thomson A, Viljanen A, Pirjola R, Boteler D, Eichner J, Cilliers PJ, Welling D, Savani NP, Weigel RS, Love JJ, Balch C, Ngwira CM, Crowley G, Schultz A, Kataoka R, Anderson B, Fugate D, Simpson JJ, Mac Alester M (2017) Geomagnetically induced currents: science, engineering, and applications readiness. *Space Weather* 15:828–856. <https://doi.org/10.1002/2016SW001501>
- Royal Academy of Engineering report (2013) *Extreme space weather impacts on engineered systems and infrastructure*. Royal Academy of Engineering, London
- Thomson AWP, Gaunt CT, Cilliers P, Wild JA, Opperman B, McKinnell LA, Kotze P, Ngwira CM, Lotz SI (2010) Present day challenges in understanding the geomagnetic hazard to national power grids. *Adv Space Res* 45(9):1182–1190. <https://doi.org/10.1016/j.asr.2009.11.023>
- Tsurutani BT, Gonzalez WD (1997) The interplanetary causes of magnetic storms: a review. In: Tsurutani BT et al (eds) *Magnetic storms*, Geophysical monograph series, vol 98. AGU, Washington, DC, pp 77–89. <https://doi.org/10.1029/GM098p0077>
- Tsurutani BT, Lakhina GS (2014) An extreme coronal mass ejection and consequences for the magnetosphere and Earth. *Geophys Res Lett* 41. <https://doi.org/10.1002/2013GL058825>
- Tsurutani BT, Meng CI (1972) Interplanetary magnetic-field variation and substorm activity. *J Geophys Res* 77(16):2964–2970
- Tsurutani BT, Gonzalez WD, Lakhina GS, Alex S (2003) The extreme magnetic storm of 1–2 September 1859. *J Geophys Res* 108:1268. <https://doi.org/10.1029/2002JA009504>
- Tsurutani BT, Lakhina GS, Hajra R (2020) The physics of space weather/solar-terrestrial physics (STP): what we know now and what the current and future challenges are. *Nonlinear Processes Geophys* 27:75–119. <https://doi.org/10.5194/npg-27-75-2020>
- Viljanen A, Pirjola R (1994) Geomagnetically induced currents in the Finnish high-voltage power system. *Surv Geophys* 15:383–408. <https://doi.org/10.1007/BF00665999>
- Viljanen A, Tanskanen EI, Pulkkinen A (2006) Relation between substorm characteristics and rapid temporal variations of the ground magnetic field. *Ann Geophys* 24:725–733
- Wik M, Viljanen A, Pirjola R, Pulkkinen A, Wintoft P, Lundstedt H (2008) Calculation of geomagnetically induced currents in the 400kV power grid in southern Sweden. *Space Weather* 6:S07005. <https://doi.org/10.1029/2007SW000343>
- Zhang JJ, Wang C, Sun TR, Liu CM, Wang KR (2015) GIC due to storm sudden commencement in low-latitude high-voltage power network in China: observation and simulation. *Space Weather* 13:643–655. <https://doi.org/10.1002/2015SW001263>