

Week 3 Quiz

Quiz, 10 questions

8/10 points (80%)

✓ **Congratulations! You passed!**

Next Item



1 / 1
points

1.

For which of the following situations would a 0/1 loss function make the most sense?

- ☐ Your prediction for the number of bikes sold this year by a local bike shop.
- ☐ Your prediction as to whether it will rain tomorrow.
- ☐ Your estimate of the market price of your house.
- ☒ Your answer choice on a Coursera multiple choice quiz.



Correct

This question refers to the following learning objective (s):

- Understand the concept of loss functions and how they relate to Bayesian decision making.



1 / 1
points

2.

Fill in the blank: Under a **quadratic loss function**, the summary statistic that minimizes the posterior expected loss is the _____ of the posterior.



Week 3 Quiz

Quiz, 10 questions

☐ Median

☐ Mode

☒ Mean

8/10 points (80%)

Correct

Correct Answer. The mean is the summary statistic that minimizes the posterior expected loss under the quadratic loss function.

This question refers to the following learning objective (s):

- Understand the concept of loss functions and how they relate to Bayesian decision making



1 / 1
points

3.

You are employed in the Human Resources department of a company and are asked to predict the number of employees that will quit their jobs in the coming year. Based on a Bayesian analysis of historical data, you determine the posterior predictive distribution of the number of quitters to follow a Poisson($\lambda = 10$) distribution. Given a **0/1** loss function, what is the prediction that minimizes posterior expected loss?

☐ a. 9

☐ b. 10

☐ c. 11

☒ d. Either a or b

Correct

Correct Answer. Both 9 and 10 are modes of Poisson ($\lambda = 10$). Since the loss function is 0/1, the mode of the posterior distribution minimizes posterior expected loss.

Week 3 Quiz

Quiz, 10 questions

This question refers to the following learning objective(s):

- Make optimal decisions given a posterior distribution and a loss function.

8/10 points (80%)



1 / 1
points

4.

Suppose that you are trying to decide whether a coin is biased towards heads ($p = 0.75$) or tails ($p = 0.25$). If you decide incorrectly, you incur a loss of 10. Flipping another coin incurs a cost of 1. If your current posterior probability of a head-biased coin is 0.6, should you make the decision now or flip another coin and then decide?



Flip another coin, since the minimum posterior expected loss of making the decision now is 4, while the minimum posterior expected loss of making the decision after seeing another coin flip is between 3 and 4.

Correct

This question refers to the following learning objective(s):

- Decide between hypotheses given a loss function:
Make optimal decisions given a posterior distribution and a loss function.



Flip another coin, since the minimum posterior expected loss of making the decision now is 4, while the minimum posterior expected loss of making the decision after seeing another coin flip is between 2 and 3.



Make the decision now, since the minimum posterior expected loss of making the decision now is 4, while the minimum posterior expected loss of making the decision after seeing another coin flip is between 4 and 5.



Week 3 Quiz

Quiz, 10 questions



Make the decision now, since the minimum posterior expected loss of making the decision now is 4, while the minimum posterior expected loss of making the decision after seeing another coin flip is between 5 and 6.

8/10 points (80%)



1 / 1
points

5.

You are testing a hypothesis H_1 against an alternative hypothesis H_2 using Bayes Factors. You calculate $BF[H_1 : H_2]$ to be 42.7. According to guidelines first given by Jeffreys (and presented in the lecture), what conclusion can be drawn from the data?



The data provides strong evidence against H_1 .



The data provides decisive evidence against H_1 .



The data provides strong evidence against H_2 .



Correct

This question refers to the following learning objective (s):

- Compare multiple hypotheses using Bayes Factors.



The data provides decisive evidence against H_2 .



1 / 1
points

6.

Suppose that you are trying to estimate the true proportion p of male births in the United States. Starting with a strong prior (Beta(500,500)) on the proportion, you randomly sample 10,000 birth certificates, observing 5029 males and 4971 females. What is the posterior probability that p is less than or equal to 0.5? (Hint: use function(s) in R to answer this question)



☐ 0.27

Week 3 Quiz ☐ 0.28

8/10 points (80%)

Quiz, 10 questions

☒ 0.29



Correct

This question refers to the following learning objective (s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation
- Make inferences about a proportion using a conjugate Beta prior.

☐ 0.30



1 / 1
points

7.

True or False: A Bayesian hypothesis test for a mean $\mu = 0$ using a concentrated prior on μ will yield nearly identical results to a hypothesis test with a high-variance prior μ ?

☐ True

☒ False



Correct

When testing hypotheses using Bayes Factors, prior specification is highly important. If the prior puts most of its mass on implausible values, the null model will be disproportionately favored.

This question refers to the following learning objective (s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation
- Compare multiple hypotheses using Bayes Factors.

Week 3 Quiz

Quiz, 10 questions



1 / 1
points

8/10 points (80%)

8.

Suppose you are trying to infer the difference in the proportion of yellow M&Ms between two bags of M&Ms. Each bag is estimated to contain roughly 30 M&Ms. To make inferences about the proportion, you draw 10 M&Ms from each bag **without** replacement and record the number of yellow M&Ms in each bag.

Is there a problem with the experimental design? If so, what is it?

- ☐ Yes, the probability of drawing a yellow M&M is not independent between groups.
- ☐ There is no problem with the experimental design.
- ☒ Yes, the probability of drawing a yellow M&M is not independent within groups.



Correct

This question refers to the following learning objective (s):

- Identify assumptions relating to a statistical inference.



0 / 1
points

9.

Suppose that when testing $H_0 : p = 0.5$ versus $H_1 : p \neq 0.5$ using Bayes Factors, we get the posterior probability $P(H_0 \mid \text{data}) = 0.25$. Conditional on H_1 , the posterior mean of p is 0.6. Under **quadratic** loss, what is the point estimate for p that minimizes expected posterior loss?

- ☐ 0.5
- ☒ 0.55



Week 3 Quiz

Quiz, 10 questions

This should not be selected

Under quadratic loss, the mean minimizes expected posterior loss. Use the posterior probabilities of each hypothesis to weight their respective means. Hint - if $H_0 : p = 0.5$ what is the posterior mean of p under H_0 ?

8/10 points (80%)

This question refers to the following learning objective (s):

- Create point estimates and credible intervals by averaging over multiple hypotheses
- Make optimal decisions given a posterior distribution and a loss function.

☐ 0.575

☐ 0.6



0 / 1
points

10.

True or False: The use of the reference prior $Beta(1/2, 1/2)$ has little bearing on the results of a hypothesis test comparing $p = 0.5$ versus $p \neq 0.5$, provided that the sample size is sufficiently large.



True

This should not be selected

This prior in fact has a significant bearing on the results of the hypothesis test. This is because it places most of its mass on values far from 0.5, so when calculating the Bayes Factor, the marginal likelihood of the data given the prior will be small if the proportion of successes is near 0.5, particularly if the sample size is large (see problems 6 and 7 for an example of this phenomenon.)

This question refers to the following learning objective (s):

- Conceptualize Lindley's paradox and how the Bayes Factor depends on prior elicitation.

Week 3 Quiz

8/10 points (80%)

Quiz, 10 questions



False

