1

Progressions (7) 11.9.5

EE23BTECH11051-Rajnil Malviya

Question:-

If a function Satisfying f(x + y) = f(x) f(y) for all $x, y \in N$ such that f(1) = 3 and $\sum_{x=1}^{n} f(x) = 120$, find the value of n.

Solution:- x = 1 and y = 1, we get

$$f(2) = f(1+1) \tag{1}$$

$$= f(1)f(1)$$
 (2)

$$= [f(1)]^2 (3)$$

$$f(3) = f(2+1) \tag{4}$$

$$= f(2)f(1)$$
 (5)

$$= [f(1)]^3 (6)$$

Using induction, we get;

$$f(x) = [f(1)]^x (7)$$

$$r = f(1) \tag{8}$$

$$=3 (9)$$

We can observe two non repeated poles z = 3 and z = 1

$$R_1 = \lim_{z \to 3} (z - 3) \frac{3z^{n+1}}{(z - 3)(z - 1)}$$
 (14)

$$=\frac{3^{n+2}}{2}$$
 (15)

$$R_2 = \lim_{z \to 1} (z - 1) \frac{3z^{n+1}}{(z - 3)(z - 1)}$$
 (16)

$$=\frac{-3}{2}\tag{17}$$

$$y(n) = R_1 + R_2 (18)$$

$$=\frac{3^{n+2}}{2}+\frac{-3}{2}\tag{19}$$

$$\implies 120 = \frac{3^{n+2} - 3}{2} \tag{20}$$

$$\implies n = 3$$
 (21)

Ans . n take values from n = 0 to n = 3, so there are total four terms .

Symbol	Value	Description
<i>x</i> (0)	3	first term
r	3	common ratio
y(n)	120	sum of all n terms
x(n)	$x(0) r^n u(n)$	$n+1^{th}$ term

TABLE I

Applying z-transformation on x(n);

$$\implies X(z) = \frac{3}{1 - 3z^{-1}} \quad |z| > |3| \tag{10}$$

$$Y(z) = X(z) U(z)$$
(11)

$$\implies Y(z) = \left(\frac{3}{1 - 3z^{-1}}\right) \left(\frac{1}{1 - z^{-1}}\right) \quad |z| > |r| \quad (12)$$

Using contour integration;

$$y(n) = \frac{1}{2\pi j} \oint_C \frac{3}{(1 - 3z^{-1})(1 - z^{-1})} dz$$
 (13)



