

H.W → 4

Q1) Expand  $\tan^{-1}(x+h)$  in powers of  $h$  and hence find the value of  $\tan^{-1}(1.003)$  upto 5 places of decimal.

Ans →  $f(x+h) = \tan^{-1}(x+h)$

let  $x+h = y$

∴  $f(y) = \tan^{-1}(y)$

let  $a = x$

∴  $f(y) = f(x) + (y-x)f'(x) + \frac{(y-x)^2}{2!}f''(x) + \frac{(y-x)^3}{3!}f'''(x) + \dots$

$$f(y) = f(x) + (y-x)f'(x) + \frac{(y-x)^2}{2!}f''(x) + \frac{(y-x)^3}{3!}f'''(x) + \dots$$

∴  $1.003 = y$

$1 + 0.003 = y$

$x = 1$

$h = 0.003$

$$\frac{1}{1+x^2}$$
$$= \frac{-2x}{(1+x^2)^2}$$

∴  $\tan^{-1}(1.003) = \tan^{-1}(1) + (0.003) \left( \frac{1}{1+1} \right) + (0.003)^2 \left( \frac{-2}{(1+1)^2} \right)$

$$= \frac{\pi}{4} + \frac{0.003}{2} + \frac{0.000006(-2)}{4}$$

~~0.6178~~  $0.78689$

Q.2> Using Taylor's theorem evaluate upto 4 places of decimals.

(i)  $\sqrt{1.02}$

Ans  $\rightarrow y = 1.02$

$$f(x+h) = \sqrt{x+h}$$

$$x = 1$$

$$h = 0.02$$

$$\text{let } x+h = y, \quad y = 1.02$$

$$\therefore f(y) = f(x) + (y-x)f'(x) + \frac{(y-x)^2}{2!} f''(x)$$

$$\therefore f(x) = \sqrt{x}$$

$$f'(x) = \frac{1}{2\sqrt{x}}$$

$$f''(x) = \frac{-1}{4x^{3/2}}$$

% 1 4 8 %

$$\therefore f(1.02) = f(1) + (0.02)f'(1) + \frac{(0.02)^2}{2!} f''(1)$$

$$= 1 + 0.02 \times \frac{1}{2} + \frac{(0.02)^2}{2} \times \left(-\frac{1}{4}\right)$$

$$= 1 + 0.01 - \frac{0.0004}{8}$$

$$= 1 + 0.01 - 0.00005$$

$$= 1.00995$$

ii)  $\sqrt{10}$

Ans  $\rightarrow f(x+h) = \sqrt{9+1}$

$x=9$

$h=1$

$y = x+h = 10$

So  $f(y) = \sqrt{y}$   
 $a=9$

So  $f(y) = f(a) + (y-a) f'(a) + \frac{(y-a)^2}{2} f''(a)$

~~$f(y) = f(y) + 9 f'(1) + \frac{81}{2} f''(1)$~~

$f(10) = f(9) + 1 \times f'(9) + \frac{1}{2} f''(9)$

$f(10) = 3 + \frac{1}{2\sqrt{9}} - \frac{1}{2} \times \frac{1}{4 \times 3^3}$

$f(10) = 3 + \frac{1}{6} - \frac{1}{216}$

$= 3 + 0.1667 - 0.00463$

$= 3.16207$



$$f(0) = \log(\cos 0) = \log(1) = 0$$

$$f'(0) = \frac{1}{\cos 0} \times (-\sin 0) = 0$$

$$f''(0) = -\sec^2 0 = -1$$

$$f'''(0) = -2 \sec 0 \times \csc 0 \quad \times$$

Q.4) Expand  $(2x^3 + 7x^2 + x - 1)$  in powers of  $(x-2)$ .

Ans  $\rightarrow a=2$

$$f(x) = 2x^3 + 7x^2 + x - 1$$

$$f(2) = 2(8) + 7(4) + 2 - 1 = 16 + 28 + 1 = 45$$

$$f'(2) = 6(2)^2 + 14(2) + 1 = 24 + 28 + 1 = 53$$

$$f''(2) = 12(2) + 14 = 24 + 14 = 38$$

$$f'''(2) = 12$$

$$f^{(4)}(2) = 0$$

$$\therefore f(x) = f(2) + \frac{(x-2)}{1!} f'(2) + \frac{(x-2)^2}{2!} f''(2) + \frac{(x-2)^3}{3!} f'''(2)$$

$$f(x) = 45 + 53(x-2) + \frac{(x-2)^2}{2} \times 38 + \frac{(x-2)^3}{6} \times 12$$

$$f(x) = 45 + 53x - 106 + 14(x-2)^2 + 2(x-2)^3$$