





All 19 free parameters fitted jointly; 161 valid RTs >1 s excluded. Likelihood conditions on 0 ≤ fixation-time response ≤ planned onset+1 s.
Each model prediction is normalized per exact onset, then pooled by retained parent trials (21,786). Late fixation-labelled rows are onset-censored.
RTD data/model are separately normalized over [−1,1] s; display only −0.1–0.6 s. No repeat filter is reapplied.



Dots: observed choices. Lines: onset-specific complete-mixture P(right | valid RT ∈ [0,1] s), averaged over fitted trials.
Posterior-mean predictions; proactive, reactive and contaminants estimated together. Lines join measured stimulus conditions.

ASD0047 | Joint-fit abort and convergence diagnostics






Parameter	Posterior mean	95% interval
V_A	3.133	[2.931, 3.342]
θ_A	6.693	[6.244, 7.166]
t_{nd} (ms)	-778.051	[-872.404, -685.271]
λ	1.974	[1.812, 2.144]
T_0 (ms)	64.335	[53.770, 75.900]
θ_E	2.652	[2.385, 2.936]
ℓ	0.857	[0.842, 0.871]
α	0.863	[0.640, 1.135]
w	0.547	[0.540, 0.555]
del_go (ms)	123.737	[106.755, 141.862]

Parameter	Posterior mean	95% interval
μ_{20} (ms)	97.262	[93.082, 101.572]
μ_{40} (ms)	107.224	[104.839, 109.690]
μ_{60} (ms)	104.521	[102.119, 106.944]
σ_{20} (ms)	10.629	[7.840, 14.199]
σ_{40} (ms)	12.700	[11.206, 14.294]
σ_{60} (ms)	15.545	[14.332, 16.834]
P_{lapse}	0.079	[0.067, 0.091]
$P_{lapse,\,right}$	0.292	[0.217, 0.378]
β_{lapse} (s^{-1})	1.671	[1.421, 1.956]

All times in parameter tables are ms. Abort histogram area is the observed abort fraction, not one.
Parameters update together; no proactive/lapse estimates are frozen. Intervals are variational credible intervals.
Proactive delay across starts: start 0: -778.1 [-872.4, -685.3] ms; start 1: -573.4 [-666.5, -484.0] ms
Competitive starts can reveal weak identification; the selected corner is not a claim of a unique solution.

ASD0047 | SHANK3 cohort 1 WT | Single-stage posterior
Proactive + contaminant parameters

19 jointly fitted parameters; 10,000 full-rank variational draws; selected seed 0



Diagonal titles: parameter name and posterior mean; solid mean, dashed central 95% limits.
Lower triangle: 68% and 95% joint contours. Every draw retained; padded axes do not depict full prior support.
Times in ms; only proactive t_nd can be negative. d=1 is fixed and has no posterior axis.