



Demoivre's theorem

Prof. Dr. A.N.M. Rezaul Karim

B.Sc (Honors), M.Sc in Mathematics (CU)
DCSA (BOU), PGD in ICT (BUET), Ph.D. (IU)

Professor

Department of Computer Science & Engineering
International Islamic University Chittagong





$$\frac{x}{r} = \cos \theta \quad \text{and} \quad \frac{y}{r} = \sin \theta$$

$$\Rightarrow x = r \cos \theta \quad \Rightarrow y = r \sin \theta$$

$$\therefore (x + iy) = (r \cos \theta + i r \sin \theta)$$

$$\therefore (x + iy) = r(\cos \theta + i \sin \theta)$$





Let another complex number $\mathbf{p_1(x_1 + iy_1)}$

$$\therefore \mathbf{x_1 + iy_1 = r_1(\cos \theta_1 + i \sin \theta_1)}$$

$$\therefore \mathbf{x_2 + iy_2 = r_2(\cos \theta_2 + i \sin \theta_2)}$$

$$\therefore \mathbf{x_3 + iy_3 = r_3(\cos \theta_3 + i \sin \theta_3)}$$

$$\therefore \mathbf{x_4 + iy_4 = r_4(\cos \theta_4 + i \sin \theta_4)}$$

$$\therefore \mathbf{x_n + iy_n = r_n(\cos \theta_n + i \sin \theta_n)}$$

State and Prove Demoivre's theorem

Statement: whatever be the value of n , positive or negative, integral or fractional, $\cos n\theta + i \sin n\theta$ is the value or one of the values of $(\cos \theta + i \sin \theta)^n$

Proof:

Case 1: when n is a positive integer,

By actual multiplication $(x_1 + iy_1)(x_2 + iy_2) = r_1(\cos \theta_1 + i \sin \theta_1)r_2(\cos \theta_2 + i \sin \theta_2)$

$$\therefore r_1(\cos \theta_1 + i \sin \theta_1)r_2(\cos \theta_2 + i \sin \theta_2)$$

$$\therefore r_1 r_2 (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2)$$

$$= r_1 r_2 \{ \cos \theta_1 \cos \theta_2 + i \cos \theta_1 \sin \theta_2 + i \sin \theta_1 \cos \theta_2 + i^2 \sin \theta_1 \sin \theta_2 \}$$

$$= r_1 r_2 \{ \cos \theta_1 \cos \theta_2 + i \cos \theta_1 \sin \theta_2 + i \sin \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 \} \quad [i^2 = -1]$$

$$= r_1 r_2 \{ \cos \theta_1 \cos \theta_2 - \sin \theta_1 \sin \theta_2 + i \sin \theta_1 \cos \theta_2 + i \cos \theta_1 \sin \theta_2 \}$$

$$[\cos A \cos B - \sin A \sin B = \cos(A + B) \text{ \& } \sin A \cos B + \cos A \sin B = \sin(A + B)]$$

$$= r_1 r_2 \{ \cos(\theta_1 + \theta_2) + i(\sin \theta_1 \cos \theta_2 + \cos \theta_1 \sin \theta_2) \}$$

$$= r_1 r_2 \{ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \}$$

$$\therefore r_1 r_2 (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) = r_1 r_2 \{ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \}$$

$$\therefore (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2) = \{ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \} \text{(i)}$$

$$\text{Now, } (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2)(\cos \theta_3 + i \sin \theta_3)$$

$$= \{ \cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2) \} (\cos \theta_3 + i \sin \theta_3)$$

$$= \cos(\theta_1 + \theta_2) \cos \theta_3 + i \sin(\theta_1 + \theta_2) \cos \theta_3 + i \cos(\theta_1 + \theta_2) \sin \theta_3 + i^2 \sin(\theta_1 + \theta_2) \sin \theta_3$$

$$= \cos(\theta_1 + \theta_2) \cos \theta_3 + i \sin(\theta_1 + \theta_2) \cos \theta_3 + i \cos(\theta_1 + \theta_2) \sin \theta_3 - \sin(\theta_1 + \theta_2) \sin \theta_3$$


$$= \cos(\theta_1 + \theta_2) \cos \theta_3 - \sin(\theta_1 + \theta_2) \sin \theta_3 + i \{ \sin(\theta_1 + \theta_2) \cos \theta_3 + \cos(\theta_1 + \theta_2) \sin \theta_3 \}$$

$$= \cos(\theta_1 + \theta_2 + \theta_3) + i \{ \sin(\theta_1 + \theta_2 + \theta_3) \}$$

$$\text{So, } (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2)(\cos \theta_3 + i \sin \theta_3) \text{ } (\cos \theta_n + i \sin \theta_n)$$

$$= \cos(\theta_1 + \theta_2 + \theta_3 + \text{ } + \theta_n) + i \{ \sin(\theta_1 + \theta_2 + \theta_3 + \theta_4 \text{ } + \theta_n) \}$$

$$\text{If } \theta_1 = \theta_2 = \theta_3 = \text{ } = \theta_n = \theta$$



$$\begin{aligned}
 &\text{Then } (\cos \theta_1 + i \sin \theta_1)(\cos \theta_2 + i \sin \theta_2)(\cos \theta_3 + i \sin \theta_3) \dots (\cos \theta_n + i \sin \theta_n) \\
 &= \cos(\theta_1 + \theta_2 + \theta_3 + \dots + \theta_n) + i \{\sin(\theta_1 + \theta_2 + \theta_3 + \dots + \theta_n)\} \\
 &\Rightarrow (\cos \theta + i \sin \theta)(\cos \theta + i \sin \theta)(\cos \theta + i \sin \theta) \dots (\cos \theta + i \sin \theta) \\
 &= \cos(\theta + \theta + \theta + \dots + \theta) + i \{\sin(\theta + \theta + \theta + \dots + \theta)\} \\
 &\Rightarrow (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta
 \end{aligned}$$

Case 2: when n is a negative integer

Let us suppose, $n = -m$, where m is a positive integer.

$$\Rightarrow (\cos \theta + i \sin \theta)^n = (\cos \theta + i \sin \theta)^{-m}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{1}{(\cos \theta + i \sin \theta)^m}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{1}{\cos m\theta + i \sin m\theta}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{(\cos m\theta - i \sin m\theta)}{(\cos m\theta + i \sin m\theta)(\cos m\theta - i \sin m\theta)}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{(\cos m\theta - i \sin m\theta)}{\cos^2 m\theta - i^2 \sin^2 m\theta}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{(\cos m\theta - i \sin m\theta)}{\cos^2 m\theta + \sin^2 m\theta} \quad [i^2 = -1]$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \frac{(\cos m\theta - i \sin m\theta)}{1}$$

$$[\cos^2 \theta + \sin^2 \theta = 1]$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = (\cos m\theta - i \sin m\theta)$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \cos(-m\theta) + i \sin(-m\theta)$$

$$[\cos(-\theta) = \cos \theta; \sin(-\theta) = -\sin \theta]$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta$$

$$[\because n = -m]$$

Case 3: when n is a fraction, positive or negative

Let us suppose, $n = \frac{p}{q}$, where q is a positive integer and p is any integer, positive or negative.

From case 1:

$$\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right)^q = \cos q \frac{\theta}{q} + i \sin q \frac{\theta}{q}$$

$$[(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta]$$

$$\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right)^q = \cos \theta + i \sin \theta$$

Taking the q-th roots on both sides,

$$\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right)^q = \cos \theta + i \sin \theta$$

$$\left\{\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right)^q\right\}^{\frac{1}{q}} = (\cos \theta + i \sin \theta)^{\frac{1}{q}}$$

$$\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right) = (\cos \theta + i \sin \theta)^{\frac{1}{q}}$$

So, $\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}$ is one of the values of $(\cos \theta + i \sin \theta)^{\frac{1}{q}}$

Raising to the p-th power,

$$\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right) = (\cos \theta + i \sin \theta)^{\frac{1}{q}}$$

$$\left\{\left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q}\right)\right\}^p = \left\{(\cos \theta + i \sin \theta)^{\frac{1}{q}}\right\}^p$$

$$\left\{ \left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q} \right) \right\}^p = \left\{ (\cos \theta + i \sin \theta)^{\frac{p}{q}} \right\}$$

$$\left\{ \left(\cos \frac{\theta}{q} + i \sin \frac{\theta}{q} \right) \right\}^p = \left\{ (\cos \theta + i \sin \theta)^{\frac{p}{q}} \right\}$$

$$\Rightarrow \cos p \frac{\theta}{q} + i \sin p \frac{\theta}{q} = \left\{ (\cos \theta + i \sin \theta)^{\frac{p}{q}} \right\}$$

$$\Rightarrow \cos \frac{p}{q} \theta + i \sin \frac{p}{q} \theta = \left\{ (\cos \theta + i \sin \theta)^{\frac{p}{q}} \right\}$$

$$\Rightarrow (\cos \theta + i \sin \theta)^{\frac{p}{q}} = \cos \frac{p}{q} \theta + i \sin \frac{p}{q} \theta$$

$$\Rightarrow (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta \quad \left[n = \frac{p}{q} \right]$$

Q-01: If $x_r = \cos \frac{\pi}{3^r} + i \sin \frac{\pi}{3^r}$, prove that $x_1 x_2 x_3 \dots \dots \dots \text{inf} = i$

Answer: Given,

$$x_r = \cos \frac{\pi}{3^r} + i \sin \frac{\pi}{3^r}$$

Putting $r = 1, 2, 3, 4, \dots \dots \dots$

$$x_1 = \cos \frac{\pi}{3^1} + i \sin \frac{\pi}{3^1}$$

$$x_2 = \cos \frac{\pi}{3^2} + i \sin \frac{\pi}{3^2}$$

$$x_3 = \cos \frac{\pi}{3^3} + i \sin \frac{\pi}{3^3}$$

$$x_4 = \cos \frac{\pi}{3^4} + i \sin \frac{\pi}{3^4}$$

.....

.....

.....

$$\therefore x_1 x_2 x_3 \dots \dots \dots$$

$$(\cos \frac{\pi}{3^1} + i \sin \frac{\pi}{3^1}) (\cos \frac{\pi}{3^2} + i \sin \frac{\pi}{3^2}) (\cos \frac{\pi}{3^3} + i \sin \frac{\pi}{3^3}) (\cos \frac{\pi}{3^4} + i \sin \frac{\pi}{3^4}) \dots$$

$$= \cos(\frac{\pi}{3^1} + \frac{\pi}{3^2} + \frac{\pi}{3^3} + \frac{\pi}{3^4} + \dots) + i \sin(\frac{\pi}{3^1} + \frac{\pi}{3^2} + \frac{\pi}{3^3} + \frac{\pi}{3^4} + \dots)$$

$$= \cos\{\frac{\pi}{3^1}(1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots)\} + i \sin\{\frac{\pi}{3^1}(1 + \frac{1}{3} + \frac{1}{3^2} + \frac{1}{3^3} + \dots)\}$$

$$[\because (1-x)^{-1} = 1 + x + x^2 + x^3 + x^4 + \dots]$$

$$= \cos\{\frac{\pi}{3^1}(1 - \frac{1}{3})^{-1}\} + i \sin\{\frac{\pi}{3^1}(1 - \frac{1}{3})^{-1}\}$$

$$= \cos\{\frac{\pi}{3^1}(\frac{3-1}{3})^{-1}\} + i \sin\{\frac{\pi}{3^1}(\frac{3-1}{3})^{-1}\}$$

$$= \cos\{\frac{\pi}{3^1}(\frac{2}{3})^{-1}\} + i \sin\{\frac{\pi}{3^1}(\frac{2}{3})^{-1}\}$$

$$= \cos\{\frac{\pi}{3^1}(\frac{1}{\frac{3}{2}})\} + i \sin\{\frac{\pi}{3^1}(\frac{1}{\frac{3}{2}})\}$$

$$= \cos\{\frac{\pi}{3^1}(\frac{2}{3})\} + i \sin\{\frac{\pi}{3^1}(\frac{2}{3})\}$$

$$= \cos\{\frac{\pi}{2}\} + i \sin\{\frac{\pi}{2}\}$$

$$= 0 + i.1$$

$$= i$$

Q-02: If $(1 + i \frac{x}{a})(1 + i \frac{x}{b})(1 + i \frac{x}{c}) \dots = A + iB$, Then prove that

$$(i) (1 + \frac{x^2}{a^2})(1 + \frac{x^2}{b^2})(1 + \frac{x^2}{c^2}) \dots = A^2 + B^2$$

$$(ii) \tan^{-1} \frac{x}{a} + \tan^{-1} \frac{x}{b} + \tan^{-1} \frac{x}{c} + \dots = \tan^{-1} \frac{B}{A}$$

Answer:

$$\text{Given, } (1 + i \frac{x}{a})(1 + i \frac{x}{b})(1 + i \frac{x}{c}) \dots = A + iB \text{ -----(i)}$$

Let, $1 = r \cos \alpha, \frac{x}{a} = r \sin \alpha$ So, $\frac{x}{a} = \frac{r \sin \alpha}{r \cos \alpha}$ $\frac{1}{1}$ $\therefore \frac{x}{a} = \tan \alpha$ $\therefore \tan \alpha = \frac{x}{a}$ $\therefore \alpha = \tan^{-1} \frac{x}{a}$	Again, Let, $1 = r \cos \beta, \frac{x}{b} = r \sin \beta$ So, $\frac{x}{b} = \frac{r \sin \beta}{r \cos \beta}$ $\frac{1}{1}$ $\therefore \frac{x}{b} = \tan \beta$ $\therefore \tan \beta = \frac{x}{b}$ $\therefore \beta = \tan^{-1} \frac{x}{b}$	Again, Let, $1 = r \cos \gamma, \frac{x}{c} = r \sin \gamma$ So, $\frac{x}{c} = \frac{r \sin \gamma}{r \cos \gamma}$ $\frac{1}{1}$ $\therefore \frac{x}{c} = \tan \gamma$ etc., $\therefore \tan \gamma = \frac{x}{c}$ $\therefore \gamma = \tan^{-1} \frac{x}{c}$
--	--	---

From (i), we get

$$\text{Given, } (1 + i \frac{x}{a})(1 + i \frac{x}{b})(1 + i \frac{x}{c}) \dots = A + iB$$

$$\Rightarrow (1 + i \tan \alpha)(1 + i \tan \beta)(1 + i \tan \gamma) \dots = A + iB$$

$$\Rightarrow (1 + i \frac{\sin \alpha}{\cos \alpha})(1 + i \frac{\sin \beta}{\cos \beta})(1 + i \frac{\sin \gamma}{\cos \gamma}) \dots = A + iB$$

$$\Rightarrow (\frac{\cos \alpha + i \sin \alpha}{\cos \alpha})(\frac{\cos \beta + i \sin \beta}{\cos \beta})(\frac{\cos \gamma + i \sin \gamma}{\cos \gamma}) \dots = A + iB$$

$$\Rightarrow (\frac{1}{\cos \alpha})(\frac{1}{\cos \beta})(\frac{1}{\cos \gamma}) \dots (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta)(\cos \gamma + i \sin \gamma) \dots = A + iB$$

$$\Rightarrow (\sec \alpha)(\sec \beta)(\sec \gamma) \dots (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta)(\cos \gamma + i \sin \gamma) \dots = A + iB$$

$$\Rightarrow (\sec \alpha)(\sec \beta)(\sec \gamma) \dots [\cos(\alpha + \beta + \gamma + \dots) + i \sin(\alpha + \beta + \gamma + \dots)] = A + iB$$

Equating the real and imaginary part on both sides, we get,

$$(\sec \alpha)(\sec \beta)(\sec \gamma) \dots [\cos(\alpha + \beta + \gamma + \dots)] = A \text{ -----(ii)}$$

$$(\sec \alpha)(\sec \beta)(\sec \gamma) \dots [\sin(\alpha + \beta + \gamma + \dots)] = B \text{ -----(iii)}$$

Squaring (ii) and (iii), we get,

$$(\sec^2 \alpha)(\sec^2 \beta)(\sec^2 \gamma) \dots [\cos^2(\alpha + \beta + \gamma + \dots)] = A^2 \text{ -----(iv)}$$

$$(\sec^2 \alpha)(\sec^2 \beta)(\sec^2 \gamma) \dots [\sin^2(\alpha + \beta + \gamma + \dots)] = B^2 \text{ -----(v)}$$

Adding (iv) & (v)

$$(\sec^2 \alpha \sec^2 \beta \sec^2 \gamma \dots) \cos^2(\alpha + \beta + \gamma + \dots) + (\sec^2 \alpha \sec^2 \beta \sec^2 \gamma \dots) \sin^2(\alpha + \beta + \gamma + \dots) = A^2 + B^2$$

$$\Rightarrow (\sec^2 \alpha \sec^2 \beta \sec^2 \gamma \dots) \{ \cos^2(\alpha + \beta + \gamma + \dots) + \sin^2(\alpha + \beta + \gamma + \dots) \} = A^2 + B^2$$

$$\Rightarrow (\sec^2 \alpha \sec^2 \beta \sec^2 \gamma \dots) 1 = A^2 + B^2$$

$$\Rightarrow \sec^2 \alpha \sec^2 \beta \sec^2 \gamma \dots = A^2 + B^2$$

$$\Rightarrow (1 + \tan^2 \alpha)(1 + \tan^2 \beta)(1 + \tan^2 \gamma) \dots = A^2 + B^2$$

$$\therefore (1 + \frac{x^2}{a^2})(1 + \frac{x^2}{b^2})(1 + \frac{x^2}{c^2}) \dots = A^2 + B^2 \text{ proved (i)}$$

Now, (iii) \div (ii)

$$\frac{(\sec \alpha)(\sec \beta)(\sec \gamma) \dots [\sin(\alpha + \beta + \gamma + \dots)]}{(\sec \alpha)(\sec \beta)(\sec \gamma) \dots [\cos(\alpha + \beta + \gamma + \dots)]} = \frac{B}{A}$$

$$\Rightarrow \frac{[\sin(\alpha + \beta + \gamma + \dots)]}{[\cos(\alpha + \beta + \gamma + \dots)]} = \frac{B}{A}$$

$$\Rightarrow \tan(\alpha + \beta + \gamma + \dots) = \frac{B}{A}$$

$$\Rightarrow \alpha + \beta + \gamma + \dots = \tan^{-1} \frac{B}{A}$$

$$\therefore (\tan^{-1} \frac{x}{a} + \tan^{-1} \frac{x}{b} + \tan^{-1} \frac{x}{c} + \dots) = \tan^{-1} \frac{B}{A} \text{ proved (ii)}$$

Q-03: Using Demoivre's theorem, solve the equation $x^6 + x^5 + x^4 + x^3 + x^2 + x^1 + 1 = 0$

Answer: We have, $x^6 + x^5 + x^4 + x^3 + x^2 + x^1 + 1 = 0$

Multiplying the given equation by (x-1), we get,

$$(x-1)(x^6 + x^5 + x^4 + x^3 + x^2 + x^1 + 1) = 0$$

$$\Rightarrow x^7 - 1 = 0$$


$$\Rightarrow x^7 = 1$$

$$\cos(4\pi + 0)$$

$$= \cos(8.90 + 0)$$

$$= \cos 0$$

$$= 1$$



$$\Rightarrow x^7 = 1$$

$$\Rightarrow x = (1)^{1/7}$$

$$\Rightarrow x = (\cos 0 + i \sin 0)^{1/7}$$

$$\Rightarrow x = \{\cos(2n\pi + 0) + i \sin(2n\pi + 0)\}^{1/7}$$

$$\Rightarrow x = \cos \frac{2n\pi}{7} + i \sin \frac{2n\pi}{7} \quad [(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta]$$

Putting $n = 0, 1, 2, 3, 4, 5$ and 6 , we get roots of equation as

$$\Rightarrow x = \cos 0 + i \sin 0$$

$$\Rightarrow x = \cos \frac{2\pi}{7} + i \sin \frac{2\pi}{7}$$

$$\Rightarrow x = \cos \frac{4\pi}{7} + i \sin \frac{4\pi}{7}$$

$$\Rightarrow x = \cos \frac{6\pi}{7} + i \sin \frac{6\pi}{7}$$

$$\Rightarrow x = \cos \frac{8\pi}{7} + i \sin \frac{8\pi}{7}$$

$$\Rightarrow x = \cos \frac{10\pi}{7} + i \sin \frac{10\pi}{7}$$

$$\Rightarrow x = \cos \frac{12\pi}{7} + i \sin \frac{12\pi}{7}$$

Q-04: Using Demoivre's theorem, find the values of

$$i. (\sqrt{3} + i.1)^{1/5}$$

$$ii. (8i)^{1/3}$$

$$iii. (32)^{1/5}$$

Answer:

$$i. (\sqrt{3} + i.1)^{1/5}$$

$$\text{Let, } \sqrt{3} = r \cos \theta \text{ -----(i)}$$

$$1 = r \sin \theta \text{ -----(ii)}$$

Squaring (i) & (ii),

$$3 = r^2 \cos^2 \theta \text{ -----(iii)}$$

$$1^2 = r^2 \sin^2 \theta \text{ -----(iv)}$$

Adding (iii) & (iv),

$$3 + 1 = r^2 \cos^2 \theta + r^2 \sin^2 \theta$$

$$4 = r^2 (\cos^2 \theta + \sin^2 \theta)$$

$$4 = r^2 . 1$$

$$r = 2 \text{ -----(v)}$$

(ii) \div (i),

$$\frac{1}{\sqrt{3}} = \frac{r \sin \theta}{r \cos \theta}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \tan \theta$$

$$\Rightarrow \tan \frac{\pi}{6} = \tan \theta$$

$$\Rightarrow \tan \theta = \tan \frac{\pi}{6}$$

$$\Rightarrow \theta = \frac{\pi}{6} \text{ -----(vi)}$$

Given

$$(\sqrt{3} + i.1)^{1/5}$$

$$= (r \cos \theta + i.r \sin \theta)^{1/5}$$

$$= \{r(\cos \theta + i \sin \theta)\}^{1/5}$$

$$= \{2(\cos \frac{\pi}{6} + i \sin \frac{\pi}{6})\}^{1/5}$$

$$\begin{aligned}
&= \{2^{1/5}(\cos(2n\pi + \frac{\pi}{6}) + i \sin(2n\pi + \frac{\pi}{6}))\}^{1/5} \\
&= \{2^{1/5}\{\cos(\frac{12n\pi + \pi}{6}) + i \sin(\frac{12n\pi + \pi}{6})\}\}^{1/5} \\
&= \{2^{1/5}\{\cos(\frac{12n\pi + \pi}{30}) + i \sin(\frac{12n\pi + \pi}{30})\}\} [(\cos\theta + i \sin\theta)^n = \cos n\theta + i \sin n\theta]
\end{aligned}$$

Putting $n = 0, 1, 2, 3, 4$

ii) $(8i)^{1/3}$

$$8i = 8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})$$

$$(8i)^{1/3} = \{8(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2})\}^{1/3}$$

$$(8i)^{1/3} = 8^{1/3}\{\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}\}^{1/3}$$

$$(8i)^{1/3} = (2^3)^{1/3}\{\cos \frac{\pi}{2} + i \sin \frac{\pi}{2}\}^{1/3}$$

$$(8i)^{1/3} = 2\{\cos(2n\pi + \frac{\pi}{2}) + i \sin(2n\pi + \frac{\pi}{2})\}^{1/3}$$

$$(8i)^{1/3} = 2\{\cos(\frac{4n\pi + \pi}{2}) + i \sin(\frac{4n\pi + \pi}{2})\}^{1/3} [(\cos\theta + i \sin\theta)^n = \cos n\theta + i \sin n\theta]$$

$$(8i)^{1/3} = 2\{\cos \frac{1}{3}(\frac{4n\pi + \pi}{2}) + i \sin \frac{1}{3}(\frac{4n\pi + \pi}{2})\}$$

$$(8i)^{1/3} = 2\{\cos(\frac{4n\pi + \pi}{6}) + i \sin(\frac{4n\pi + \pi}{6})\}$$

Putting $n = 0, 1, 2$

$$(8i)^{1/3} = 2\{\cos(\frac{4.0\pi + \pi}{6}) + i \sin(\frac{4.0\pi + \pi}{6})\} = 2\{\cos(\frac{\pi}{6}) + i \sin(\frac{\pi}{6})\}$$

$$(8i)^{1/3} = 2\{\cos(\frac{4.1\pi + \pi}{6}) + i \sin(\frac{4.1\pi + \pi}{6})\} = 2\{\cos(\frac{5\pi}{6}) + i \sin(\frac{5\pi}{6})\}$$

$$(8i)^{1/3} = 2\{\cos(\frac{4.2\pi + \pi}{6}) + i \sin(\frac{4.2\pi + \pi}{6})\} = 2\{\cos(\frac{9\pi}{6}) + i \sin(\frac{9\pi}{6})\}$$

iii) $(32)^{1/5}$

$$(32)^{1/5} = \{32(1)\}^{1/5}$$

$$(32)^{1/5} = 32^{1/5} (\cos 0 + i \sin 0)^{1/5}$$

$$(32)^{1/5} = (2^5)^{1/5} (\cos 0 + i \sin 0)^{1/5}$$

$$(32)^{1/5} = 2\{\cos(2n\pi + 0) + i \sin(2n\pi + 0)\}^{1/5} [(\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta]$$

$$(32)^{1/5} = 2\{\cos(\frac{2n\pi}{5}) + i \sin(\frac{2n\pi}{5})\}$$

Putting $n = 0, 1, 2, 3, 4$

$$(32)^{1/5} = 2\{\cos(\frac{2.0.\pi}{5}) + i \sin(\frac{2.0.\pi}{5})\} = 2(\cos 0 + i \sin 0) = 2(1 + 0) = 2$$

$$(32)^{1/5} = 2\{\cos(\frac{2.1.\pi}{5}) + i \sin(\frac{2.1.\pi}{5})\} = 2\{\cos(\frac{2\pi}{5}) + i \sin(\frac{2\pi}{5})\}$$

$$(32)^{1/5} = 2\{\cos(\frac{2.2.\pi}{5}) + i \sin(\frac{2.2.\pi}{5})\} = 2\{\cos(\frac{4\pi}{5}) + i \sin(\frac{4\pi}{5})\}$$

$$(32)^{1/5} = 2\{\cos(\frac{2.3.\pi}{5}) + i \sin(\frac{2.3.\pi}{5})\} = 2\{\cos(\frac{6\pi}{5}) + i \sin(\frac{6\pi}{5})\}$$

$$(32)^{1/5} = 2\{\cos(\frac{2.4.\pi}{5}) + i \sin(\frac{2.4.\pi}{5})\} = 2\{\cos(\frac{8\pi}{5}) + i \sin(\frac{8\pi}{5})\}$$

Q-5: Using Demoivre's theorem find the quadratic equation whose roots are the n th power of the roots of the equation, $x^2 - 2x \cos \theta + 1 = 0$

Answer:

The given equation is $x^2 - 2x \cos \theta + 1 = 0$

$$x = \frac{-(-2 \cos \theta) \pm \sqrt{(-2 \cos \theta)^2 - 4.1.1}}{2.1}$$

$$x = \frac{2 \cos \theta \pm \sqrt{4 \cos^2 \theta - 4}}{2}$$

$$x = \frac{2 \cos \theta \pm \sqrt{-4 + 4 \cos^2 \theta}}{2}$$

$$x = \frac{2 \cos \theta \pm \sqrt{-4(1 - \cos^2 \theta)}}{2}$$

$$\begin{aligned}
 x &= \frac{2 \cos \theta \pm \sqrt{-4 \sin^2 \theta}}{2} \\
 x &= \frac{2 \cos \theta \pm \sqrt{4i^2 \sin^2 \theta}}{2} \\
 x &= \frac{2 \cos \theta \pm 2i\sqrt{\sin^2 \theta}}{2} \\
 x &= \frac{2 \cos \theta \pm 2i \sin \theta}{2} \\
 x &= \frac{2(\cos \theta \pm i \sin \theta)}{2} \\
 x &= \cos \theta \pm i \sin \theta
 \end{aligned}$$

Let α and β are the roots of the equation $x^2 - 2x \cos \theta + 1 = 0$

$$\therefore \alpha = \cos \theta + i \sin \theta \text{ and } \beta = \cos \theta - i \sin \theta$$

We have to form a new equation whose roots are α^n and β^n

We know any equation is $x^2 - (\text{sum of the roots})x + \text{product of the roots} = 0$

$$x^2 - (\alpha^n + \beta^n)x + \alpha^n \beta^n = 0$$

$$x^2 - [(\cos \theta + i \sin \theta)^n + (\cos \theta - i \sin \theta)^n]x + (\cos \theta + i \sin \theta)^n (\cos \theta - i \sin \theta)^n = 0$$

$$[\because (\cos \theta + i \sin \theta)^n = \cos n\theta + i \sin n\theta] [\because (\cos \theta - i \sin \theta)^n = \cos n\theta - i \sin n\theta]$$

$$x^2 - [(\cos n\theta + i \sin n\theta) + (\cos n\theta - i \sin n\theta)]x + (\cos n\theta + i \sin n\theta)(\cos n\theta - i \sin n\theta) = 0$$

$$x^2 - [\cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta]x + [\cos^2 n\theta - (i)^2 \sin^2 n\theta] = 0$$

$$x^2 - [\cos n\theta + i \sin n\theta + \cos n\theta - i \sin n\theta]x + [\cos^2 n\theta + \sin^2 n\theta] = 0$$

$$x^2 - [2 \cos n\theta]x + 1 = 0$$

$$x^2 - 2x \cos n\theta + 1 = 0 \text{ Answer.}$$

