



Human and Ecological Risk Assessment: An International Journal

ISSN: 1080-7039 (Print) 1549-7860 (Online) Journal homepage: <http://www.tandfonline.com/loi/bher20>


Pollution and Health Risk Assessment of Heavy Metals in Urban Soil in China

Lei Liu, Xiuying Zhang & Taiyang Zhong

To cite this article: Lei Liu, Xiuying Zhang & Taiyang Zhong (2015): Pollution and Health Risk Assessment of Heavy Metals in Urban Soil in China, Human and Ecological Risk Assessment: An International Journal, DOI: [10.1080/10807039.2015.1078226](https://doi.org/10.1080/10807039.2015.1078226)

To link to this article: <http://dx.doi.org/10.1080/10807039.2015.1078226>

 View supplementary material 

 Accepted online: 20 Aug 2015.

 Submit your article to this journal 

 Article views: 47

 View related articles 

 View Crossmark data 

Pollution and Health Risk Assessment of Heavy Metals in Urban Soil in China

Lei Liu^a, Xiuying Zhang^{a, b}, Taiyang Zhong^c

^aJiangsu Provincial Key Laboratory of Geographic Information Science and Technology, International Institute for Earth System Science, Nanjing University, Nanjing 210093

^bJiangsu Center for Collaborative Innovation in Geographical Information Resource Development and Application, Nanjing, 210023, China

^cSchool of Geographic and Oceanographic Sciences, Nanjing University, 163 Xianlin Road, Nanjing, Jiangsu Province, 210023, China

*Corresponding author, e-mail: lzhxy77@163.com

Abstract

The pollution and potential health risk due to lifetime exposure to heavy metals in urban soil of China were evaluated, based on the urban soil samples collected from published papers from 2005 to 2014. The contamination levels were in the order of Cd > Hg > Cu > Zn > Pb > As > Ni > Cr, and Hg and Cd fell into the category of “moderately contaminated” to “heavily contaminated”. The non-carcinogenic for different populations varied greatly, among which children faced high risk, and then the adult female and adult male were followed. The hazard index (non-carcinogenic risk) higher than 1.00 occurred in Shanghai, Gansu, Qinghai, Hunan and Anhui, whereas most of those in the northern and western had low risks. For the carcinogenic risk, Anhui and Ningxia provinces had urban soils exceeding the safe reference (1×10^{-6} - 1×10^{-4}). Qinghai and Gansu had high carcinogenic risks since their risk index were much close to the reference, and the others were in low risk.

Key words: Soil heavy metal, pollution, health risk, urban environment, China

Introduction

More than half of people in China are living in cities due to the rapid development of urbanization (Wei and Ye, 2014). Thus it is important to study the influence of urban environment on human health (Cao et al., 2015; Giri and Singh, 2015; Lei et al., 2015; Li et al., 2013). Heavy metals have significant negative effects on human health, ranging from acute reactions to chronic illnesses (Kampa and Castanas, 2008; Pei et al., 2015; Shi et al., 2008; Wang et al., 2015). The mercury (Hg), cadmium (Cd), arsenic (As) and Chromium (Cr), are toxic or carcinogenic even at low concentrations when people are exposed for a long time (Angelone and Udovic, 2014; Khan et al., 2015; Zhang et al., 2015; Zhang et al., 2014). Potentially toxic heavy metal pollution in urban environments has given rise to growing concerns during past decades (Guo et al., 2012; Li et al., 2013; Shi et al., 2008; Wei and Yang, 2010; Xia et al., 2011; Zheng et al., 2015).

Recent studies have focused on monitoring the concentrations, exploring the spatial distribution of heavy metals, identifying the enrichment factors and sources, and assessing the pollution levels in the urban soils. For example, Wei and Yang concluded that about 65% of the investigated cities have high or extremely high contamination levels of heavy metals in urban soils (Wei and Yang, 2010), most of which were in the developed eastern coastal regions and old industrial cities (Yu et al., 2012). The two principle sources of trace metal contamination in urban soil in China were industrial discharges and traffic emissions (Li et al., 2013; Wei and Yang,

2010). The pollution in urban soils can pose directly a significant health risk through soil ingestion, dust inhalation, or dermal contact (Giri and Singh, 2015; Karim and Qureshi, 2014). Thus, it is critical to quantify the potential human health risk due to the exposure to heavy metals in urban soil.

Soil quality criteria based on total metal concentrations might not accurately estimate the real risks (Baize and van Oort, 2014) and are only appropriate for worst-case scenarios (Angelone and Udovic, 2014). Health risk assessment is an effective approach to determine the risk to human health quantitatively posed by several various contaminants through different exposure pathways (Kampa and Castanas, 2008; Luo et al., 2012). The extent of harmful effects depends on the age and structure of the population, and the type of intensity of contact with contaminated soils (Wcisło et al., 2002; Zhang et al., 2015). Few studies concerned the above questions when assessing the health risk assessment of heavy metal contamination of urban soils in cities but Xiamen (Luo et al., 2012), Huludao city (Zheng et al., 2010), Hangzhou, Changxing and Shangyu cities in Zhejiang (Liu et al., 2013) and Gejiu city (Li et al., 2014a). To enact an effective policy to control further harm on human health risk by heavy metals, a comprehensive health risk assessment in urban soil throughout China is needed.

This study collected heavy metal concentrations in urban soil throughout China, from papers published from 2005 to 2014. The aim is to (1) survey heavy metal concentrations, (2) assess the level of heavy metal pollution, and (3) evaluate the health risk in urban soil throughout China.

Materials and methods

Data collection

This study collected 157 published papers on heavy metal concentrations in urban surface soils (0-20 cm) throughout China from 2005 to 2014. The detailed information on the collection process of the published papers and the heavy metal data can be found in the manuscript (Zhang et al., 2014). Eight heavy metals Cu, Pb, Zn, Cd, Ni, As, Cr and Hg were collected in this study. In total, 297 data records on heavy metal concentrations in urban surface soil were collected. Background information on the investigated urban areas is described in Table S1.

Pollution assessment

The geoaccumulation index (I_{geo}) is employed in assessing the contamination levels and the pollution of heavy metals in urban soils in China. This method has been widely applied in European trace metal studies since the late 1960s and it is also employed in pollution assessment of urban soil (Wei and Yang, 2010), which is calculated using the follow equation (Solgi et al., 2012):

$$I_{geo} = \log_2(C_n/1.5B_n) \quad (1)$$

where C_n is the measured concentration of every heavy metal in found in urban soil in China, and B_n is the geochemical background value of the heavy metals found in the soil (mg/kg). In this study, the B_n refers to the heavy metal background concentrations of the province that the n -th data record belongs to (CNEMC, 1990). The background concentrations on a provincial scale

and the 7 classes of I_{geo} are described in Table S2 and S3, respectively (Li et al., 2014b; Wei and Yang, 2010).

Health risk assessment

The risk effects consist of carcinogenic and non-carcinogenic risk assessments for all heavy metals through ingestion and dermal absorption in this study. The average daily intake of heavy metals from soil ingestion (ADI_I) and dermal absorption (ADI_D) are calculated from the equations provided by USEPA (Li et al., 2014b).

A hazard index (HI) approach to assess non-carcinogenic risk is calculated using equation (Yuswir et al., 2014):

$$HI = HQ_i = \sum_{i=1}^n \frac{ADI_i}{RFD_i} \quad (5)$$

where the chronic hazard index (HI) is the sum of hazard quotient (HQ_i) of i -th heavy metal, RFD_i is the chronic reference dose for i -th heavy metal. In this study, RFD for Cu, Pb, Zn, Cd, As, Ni, Cr, Hg were 4×10^{-2} , 1.4×10^{-4} , 3×10^{-1} , 2.5×10^{-5} , 3×10^{-4} , 8×10^{-4} , 1.95×10^{-2} , 1.6×10^{-4} , respectively (Li et al., 2014b).

Carcinogenic risk is estimated (Cao et al., 2015):

$$RISK = ADI \times SF \quad (8)$$

$$SF_{ABS} = \frac{SF_0}{ABS_{GI}} \quad (9)$$

where SF is the carcinogenicity slope factor (per mg/kg-day), SF_{ABS} is the dermally adjusted slope factor (per mg/kg-day), SF_0 is the oral slope factor (per mg/kg-day), and ABS_{GI} is the

gastrointestinal absorption factor(unitless). In this study, only the carcinogenic risk of As is estimated and SF for As is 1.5 (Li et al., 2014b).

Results and discussion

Heavy metal concentrations in urban soil

As shown in Fig. 1, the average concentrations of each heavy metal exceeded the corresponding background values in soils in China (CNEMC, 1990), which were 5.88, 4.15, 2.21, 1.98, 1.83, 1.30, 1.27 and 1.26 times of the background values for Cd, Hg, Cu, Pb, Zn, As, Cr and Ni, respectively. In relative term, the soils in the urban areas were contaminated least by Cr and most by Cd, which was consistent with a previous study (Wei and Yang, 2010). For all metals, the total concentrations showed a large degree of variability, reflecting the non-homogeneous distribution of concentrations of anthropogenically emitted heavy metals (Guo et al., 2012). Firstly, the large urban population generated a large amount of waste, some of which might contained heavy metals (Taghipour et al., 2013). Secondly, vehicle emissions were considered to be the principal source of heavy metal in urban soil (Cheng et al., 2014; Christoforidis and Stamatis, 2009). Thirdly, mining and smelting activities were another source of heavy metal contamination in urban environment, which exposed the minerals to weathering processes during excavation (Yuan et al., 2009).

Pollution assessment of heavy metals in urban soils

The classes of the I_{geo} for the heavy metals are presented in Table 1. The I_{geo} values showed that

all the investigated urban areas for Cr fell below class 3, with 88% falling into class 0, demonstrating that Chinese urban soil was not contaminated by Cr. About 91.72% of Ni soil samples and 76.73% of As fell into class 0, but some of the sampling locations corresponded to heavily contamination by Ni (1.28%) and As (1.99%). More than 85% of the urban soil samples of Cu, Pb and Zn were within class 0 and class 1, indicating moderate level of pollution. However, the I_{geo} values of Cd and Hg were higher than class 3 for 20.76% and 14.07% of urban soil samples respectively, indicating Cd and Hg had been introduced from exterior sources in some regions. Generally, As, Ni and Cr were the least contaminated, with the most of percentage of “uncontaminated” soil samples. Cu, Zn and Pb fell into the category of “uncontaminated” to “moderately contaminated”; Hg and Cd fell into the category of “moderately contaminated” to “heavily contaminated” in China. The contamination levels of these heavy metals were in the order of $Cd > Hg > Cu > Zn > Pb > As > Ni > Cr$, which was consistent with the results obtained by Wei and Yang (2010). In the study of Wei and Yang (2010), Cr and Ni showed the least contamination in the investigated sites, whereas Cd showed the highest I_{geo} values for most cities.

Among the eight social-economic regions (Fig. 2), Northwest region has the least contaminated area with the lowest I_{geo} for all metals. Middle Yellow River region, south coastal region and Middle Yangtze River had high I_{geo} values, indicating urban cities in these regions were moderately to heavily contaminated, particularly for Hg and Cd. In addition, only in Middle

Yangtze region the I_{geo} of Cu was higher than 3, demonstrating the urban soils in this region were significantly contaminated by Cu. Based on the pollution assessment of I_{geo} , Middle Yellow River region, South Coastal region and Middle Yangtze River appeared to be the most polluted and should be selected as the priority control regions and Hg and Cr should be selected as the priority control heavy metals.

Health risk assessment

Non-carcinogenic assessment

The HI for different populations varied strongly, in the order of children > adult female > adult male. About 23.53% of the 297 soil sample records in the South Coastal region with HI > 1 for children were higher than the other social-economic areas. Adults faced higher health risk exposure from ingestion but lower risk from dermal absorption than children did. Similar results were also obtained in the other studies (Li et al., 2014b; Zheng et al., 2010). This is mainly because that heavy metals could be accumulated in bodies over a long time, and especially the non-carcinogenic adverse effects of Cd, Pb and Hg to the tissues of adults are quite serious (Wcislo et al., 2002). Karim and Qureshi (2014) also found the ratio of a child's cumulative HI to an adult for the combined routes is 8.9 times, indicating the children were more susceptible to non-carcinogenic health effects of trace metals compared to adults (Karim and Qureshi, 2014). Particularly, the non-carcinogenic risk of heavy metals to children via oral ingestion needs special attention.

When mapping the non-carcinogenic risk level distributions of the examined urban areas, worst-case scenarios (children) were adopted by applying the non-carcinogenic risk (Fig. 3). The HQ values for Zn, As, Cr and Hg were very low and the highest HQ for these metals was below 0.10 and the highest HQ values for Cd, Ni and Pb were all above 1.00. *HI* values higher than 1 shows that non-carcinogenic risk may occur, while the reverse applies when *HI* lower than 1 (Luo et al., 2012). The *HI* higher than 1.00 occurred in Shanghai, Gansu, Qinghai, Hunan, Inner Mongolia and Anhui, mainly due to mining and smelting activities (He, 2007; Min et al., 1999; Wang et al., 2009), whereas most of the northern and western urban areas' values were lower than 1.00. These results were consistent with that found by Wei and Yang (2010), which reported that environmental pollution and related health effects were concentrated in Middle Yangtze region and East Coastal region as well as Gansu and Qinghai, whereas low pollution levels existed in southwest of China (Wei and Yang, 2010).

A correlation between traffic density and heavy metal deposited in urban soil was found (Li et al., 2013; Wang et al., 2012). Moreover, there was a positive relationship between population and heavy metal contents in urban soil (Charlesworth et al., 2003; Zheng et al., 2010). The main roads of Shanghai city were marked by traffic and heavy commercial vehicles subjected to frequent stop-and-go situations (Karim and Qureshi, 2014). Moreover, the population of Shanghai city was higher than other cities in Gansu, Qinghai, Hunan and Anhui. This suggested that the major contribution to high *HI* values in Shanghai city was from atmospheric deposition,

which traffic density and population made relatively significant contributions. However, this was not the case for Hunan, Gansu, Qinghai, Hunan, and Anhui, where the high HI values in the cities of these provinces were much higher than those of other cities with similar size of population. In smelting district, the heavy metal contamination due to smelting activities pose much higher risk than that from automobiles (Cai et al., 2013; Karim et al., 2014).

Carcinogenic risk assessment

The carcinogenic risks of As for different populations varied greatly for different people groups. The children faced the highest risk, and then adult females and adult males were followed (Fig. 4). The most susceptible subpopulation was young children, through dermal absorption because of their hand-to-mouth activity (Zheng et al., 2010). For adult males and adult females, dermal absorption was the primary pathway of exposure, whereas for children ingestion and dermal absorption both acted as common routes of exposure (Li et al., 2014b). Therefore, the exposure of the heavy metals to children could create more potential health risks.

The acceptable or tolerable risk for regulatory purposes is within the range of 10^{-6} – 10^{-4} (Luo et al., 2012). For children, only 0.62% of the soil samples had carcinogenic risks lower than 1×10^{-6} , indicating these areas were within the absolute safe level. 96.44% of the soil samples in the investigated regions were within the safe range. The remaining 2.47% of the soil samples had the carcinogenic risk values exceeding 1×10^{-4} in the investigated regions. Compared with children, adult health risk due to heavy metals exposure from urban soils is lower. The ingestion resulted

in a carcinogenic risk for As, followed by dermal contact. Moreover, adult health risk due to dermal contact as the route of exposure to urban soils was lower.

To map the carcinogenic risk level of the examined urban areas in China, the worst-case scenario (children) was selected. On a carcinogenic risk map (Fig. 5), Anhui and Ningxia provinces had urban soils exceeding the safe reference, indicating people in the two provinces faced a high carcinogenic risk by As through ingestion and dermal absorption, mainly due to sewage irrigation (Wang et al., 2010), traffic emissions (vehicle exhaust particles, tire wire particles, weathered street surface particles, brake lining wear particles) and urban and industry development (Zhang et al., 1999). Qinghai and Gansu had high carcinogenic risks with the index values close to the reference of 1×10^{-4} , mainly due to irrigation by sewage from smelting or industry plant (Liu, 2008; Nan and Zhao, 2000) and industrial emissions (power plants, coal combustion, metallurgical industry, auto repair shop, chemical plant, etc.) (Nan and Zhao, 2000). But it should be noticed that Hunan, Guangzhou, Jiangxi and Fujian also had the possibility of high carcinogenic risks, while the other provinces were within the low range of carcinogenic risks.

The findings of this study would facilitate the decision-makers to manage and dispose of contaminated soils, and minimize health risks on urban inhabitants. However, there is a need for further study of urban soil exposure parameters that would help reduce the uncertainties associated with the risk calculations. In fact, most of the urban soil samples selected in these

published papers in China were distributed in the developed regions, which covered a disproportionately small fraction of China. In conclusion, there were large uncertainties in the assessment of health risk assessment in China, which resulting from sampling procedures, analysis method and parameters used to calculate the risk index.

Conclusion

In this study, data on heavy metals in urban soils throughout China were collected and heavy metal soil pollution and health risk assessment were evaluated. Northwest region showed the least contaminated, while Middle Yellow River region, south coastal region and Middle Yangtze River were moderately contaminated, especially for Hg and Cd. The urban soils in Middle Yangtze region were significantly contaminated by Cu. Soil pollution by heavy metals continues to pose high carcinogenic and non-carcinogenic risks to the public in some urban areas in China, especially to children and those living in the most heavily polluted regions. Special attention should be paid to the priority control in East Coastal and Middle Yangtze region in order to target the biggest threats to human health. In addition, the protection of vulnerable populations, especially children living in urban areas, should be prioritized. These results would provide valuable information not only for soil quality assessment, but also to enact effective polices to protect human's health.

Acknowledgement

This study is supported by the National Natural Science Foundation of China (No. 41271190), and State Scholarship Fund of China (No. 201208320130).

Reference

- Angelone, M., Udovic, M. 2014. Potentially Harmful Elements in Urban Soils, PHEs, Environment and Human Health. Springer, pp. 221-251.
- Baize, D., van Oort, F. 2014. Potentially harmful elements in forest soils, PHEs, Environment and Human Health. Springer, pp. 151-198.
- Cai, Q.-Y., Mo, C.-H., Li, H.-Q., Lü, H., Zeng, Q.-Y., Li, Y.-W., Wu, X.-L. 2013. Heavy metal contamination of urban soils and dusts in Guangzhou, South China. Environmental monitoring and assessment 185, 1095-1106.
- Cao, S., Duan, X., Zhao, X., Wang, B., Ma, J., Fan, D., Sun, C., He, B., Wei, F., Jiang, G. 2015. Health risk assessment of various metal(loid)s via multiple exposure pathways on children living near a typical lead-acid battery plant, China. Environmental Pollution 200, 16-23.
- Charlesworth, S., Everett, M., McCarthy, R., Ordonez, A., De Miguel, E. 2003. A comparative study of heavy metal concentration and distribution in deposited street dusts in a large and a small urban area: Birmingham and Coventry, West Midlands, UK. Environment International 29, 563-573.
- Cheng, H., Li, M., Zhao, C., Li, K., Peng, M., Qin, A., Cheng, X. 2014. Overview of trace metals in the urban soil of 31 metropolises in China. Journal of Geochemical Exploration 139.
- Christoforidis, A., Stamatis, N. 2009. Heavy metal contamination in street dust and roadside soil along the major national road in Kavala's region, Greece. Geoderma 151, 257-263.

CNEMC. 1990. The Background Values of Elements in Chinese Soils, Environmental Science Press of China, Beijing.

Giri, S., Singh, A.K. 2015. Human health risk assessment via drinking water pathway due to metal contamination in the groundwater of Subarnarekha River Basin, India. *Environmental monitoring and assessment* 187, 1-14.

Guo, G., Wu, F., Xie, F., Zhang, R. 2012. Spatial distribution and pollution assessment of heavy metals in urban soils from southwest China. *Journal of Environmental Sciences* 24, 410-418.

He, M. 2007. Distribution and phytoavailability of antimony at an antimony mining and smelting area, Hunan, China. *Environmental Geochemistry and Health* 29, 209-219.

Kampa, M., Castanas, E. 2008. Human health effects of air pollution. *Environmental pollution* 151, 362-367.

Karim, Z., Qureshi, B.A. 2014. Health risk assessment of heavy metals in urban soil of Karachi, Pakistan. *Human and Ecological Risk Assessment: An International Journal* 20, 658-667.

Karim, Z., Qureshi, B.A., Mumtaz, M., Qureshi, S. 2014. Heavy metal content in urban soils as an indicator of anthropogenic and natural influences on landscape of Karachi—a multivariate spatio-temporal analysis. *Ecological Indicators* 42, 20-31.

Khan, M.U., Malik, R.N., Muhammad, S., Ullah, F., Qadir, A. 2015. Health Risk Assessment of Consumption of Heavy Metals in Market Food Crops from Sialkot and Gujranwala Districts, Pakistan. *Human and Ecological Risk Assessment: An International Journal* 21, 327-337.

Lei, M., Tie, B.-q., Song, Z.-g., Liao, B.-H., Lepo, J.E., Huang, Y.-z. 2015. Heavy metal pollution and potential health risk assessment of white rice around mine areas in Hunan Province, China. *Food Security*, 1-10.

Li, X., Liu, L., Wang, Y., Luo, G., Chen, X., Yang, X., Hall, M.H., Guo, R., Wang, H., Cui, J. 2013. Heavy metal contamination of urban soil in an old industrial city (Shenyang) in Northeast China. *Geoderma* 192, 50-58.

Li, Y., Wang, H., Wang, H., Yin, F., Yang, X., Hu, Y. 2014a. Heavy metal pollution in vegetables grown in the vicinity of a multi-metal mining area in Gejiu, China: total concentrations, speciation analysis, and health risk. *Environmental Science and Pollution Research* 21, 12569-12582.

Li, Z., Ma, Z., van der Kuijp, T.J., Yuan, Z., Huang, L. 2014b. A review of soil heavy metal pollution from mines in China: pollution and health risk assessment. *Science of the Total Environment* 468, 843-853.

Liu, J. 2008. Study on Biological Control of Heavy Metal Pollution of Farmland in Shanxi Province *Journal of Shanxi Agricultural Sciences* 6, 007.

Liu, X., Song, Q., Tang, Y., Li, W., Xu, J., Wu, J., Wang, F., Brookes, P.C. 2013. Human health risk assessment of heavy metals in soil–vegetable system: a multi-medium analysis. *Science of the Total Environment* 463, 530-540.

Luo, X.-S., Ding, J., Xu, B., Wang, Y.-J., Li, H.-B., Yu, S. 2012. Incorporating bioaccessibility

into human health risk assessments of heavy metals in urban park soils. *Science of the Total Environment* 424, 88-96.

Min, D., Wangxiao, M., Lan, L. 1999. ANALYSIS AND RESEARCH ON Pb, Cr, Cd AMOUNT OF 17 VEGETABLES IN SHAANXI [J]. *Journal of Arid Land Resources and Environment* 4, 010.

Nan, Z., Zhao, C. 2000. Heavy metal concentrations in gray calcareous soils of Baiyin region, Gansu province, PR China. *Water, air, and soil pollution* 118, 131-142.

Pei, D., Xie, H., Song, H., Xu, H., Wu, Y. 2015. Bioconcentration Factors and Potential Human Health Risks of Heavy Metals in Cultivated *Lentinus edodes* in Chengdu, People's Republic of China. *Journal of Food Protection*® 78, 390-395.

Shi, G., Chen, Z., Xu, S., Zhang, J., Wang, L., Bi, C., Teng, J. 2008. Potentially toxic metal contamination of urban soils and roadside dust in Shanghai, China. *Environmental Pollution* 156, 251-260.

Solgi, E., Esmaili-Sari, A., Riyahi-Bakhtiari, A., Hadipour, M. 2012. Soil contamination of metals in the three industrial estates, Arak, Iran. *Bulletin of environmental contamination and toxicology* 88, 634-638.

Taghipour, H., Mosaferi, M., Armanfar, F., Gaemmagami, S.J. 2013. Heavy metals pollution in the soils of suburban areas in big cities: a case study. *International Journal of Environmental Science and Technology* 10, 243-250.

Wang, H., Fang, F., Xie, H., Wang, X., Huang, C. 2010. Pollution Evaluation and Source Analysis of Heavy Metal in Urban Soil of Wuhu City. *Urban Environment and Urban Ecology* 23, 36-40.

Wang, L., Chen, L., Zhang, Y., Zhai, C., Zhang, L. 2009. The Evaluation and Sources Analysis of Heavy Metal and Micro-element in Soil for Protected Vegetable in Hebei Province. *Acta Agriculturae Boreali-Sinica*, S2.

Wang, M., Markert, B., Chen, W., Peng, C., Ouyang, Z. 2012. Identification of heavy metal pollutants using multivariate analysis and effects of land uses on their accumulation in urban soils in Beijing, China. *Environmental monitoring and assessment* 184, 5889-5897.

Wang, Q., Liu, J., Cheng, S. 2015. Heavy metals in apple orchard soils and fruits and their health risks in Liaodong Peninsula, Northeast China. *Environmental monitoring and assessment* 187, 1-8.

Wcisło, E., Ioven, D., Kucharski, R., Szdzuj, J. 2002. Human health risk assessment case study: an abandoned metal smelter site in Poland. *Chemosphere* 47, 507-515.

Wei, B., Yang, L. 2010. A review of heavy metal contaminations in urban soils, urban road dusts and agricultural soils from China. *Microchemical Journal* 94, 99-107.

Wei, Y.D., Ye, X. 2014. Urbanization, urban land expansion and environmental change in China. *Stochastic Environmental Research and Risk Assessment* 28, 757-765.

Xia, X., Chen, X., Liu, R., Liu, H. 2011. Heavy metals in urban soils with various types of land

use in Beijing, China. *Journal of Hazardous Materials* 186, 2043-2050.

Yu, S., Zhu, Y.-g., Li, X.-d. 2012. Trace metal contamination in urban soils of China. *Science of the total environment* 421, 17-30.

Yuan, Z., Lau, A.K.H., Shao, M., Louie, P.K., Liu, S.C., Zhu, T. 2009. Source analysis of volatile organic compounds by positive matrix factorization in urban and rural environments in Beijing. *Journal of Geophysical Research: Atmospheres* (1984–2012) 114.

Yuswir, N.S., Praveena, S.M., Aris, A.Z., ISMAIL, S.N.S., Hashim, Z. 2014. Health Risk Assessment on Bioavailability of Heavy Metals in Klang District Urban Surface Soil. *Iranian Journal of Public Health* 43, 167-171.

Zhang, H., Ma, D., Xie, Q., Chen, X. 1999. An approach to studying heavy metal pollution caused by modern city development in Nanjing, China. *Environmental Geology* 38, 223-228.

Zhang, J., Wang, L.-H., Yang, J.-C., Liu, H., Dai, J.-L. 2015. Health risk to residents and stimulation to inherent bacteria of various heavy metals in soil. *Science of The Total Environment* 508, 29-36.

Zhang, X., Chen, D., Zhong, T., Zhang, X., Cheng, M., Li, X. 2014. Assessment of cadmium (Cd) concentration in arable soil in China. *Environmental Science and Pollution Research*, 1-10.

Zheng, L., Tang, Q., Fan, J., Huang, X., Jiang, C., Cheng, H. 2015. Distribution and health risk assessment of mercury in urban street dust from coal energy dominant Huainan City, China. *Environmental Science and Pollution Research*.

Zheng, N., Liu, J., Wang, Q., Liang, Z. 2010. Health risk assessment of heavy metal exposure to street dust in the zinc smelting district, Northeast of China. *Science of the Total Environment* 408, 726-733.

Table 1 Class distribution of geoaccumulation index for heavy metals found in the urban areas in China (%).

	Cu	Pb	Zn	Cd	As	Ni	Cr	Hg
0	50.42	54.58	55.73	27.97	76.73	91.72	87.50	31.11
1	41.18	34.86	33.60	25.42	18.87	5.73	10.08	30.37
2	6.30	7.39	7.51	25.85	2.52	1.27	0.40	24.44
3	0.00	2.11	1.98	11.02	1.26	0.64	2.02	10.37
4	0.84	0.70	0.79	5.93	0.63	0.00	0.00	2.22
5	0.84	0.35	0.40	2.12	0.00	0.64	0.00	0.74
6	0.42	0.00	0.00	1.69	0.00	0.00	0.00	0.74




Fig. 1 Boxplots of the heavy metal concentrations (mg/kg) for the urban areas (BVSC: Background values for soils in China).




Fig. 2. Average I_{geo} values of heavy metals for different types of urban areas.




Fig. 3 Spatial distribution of HI in different types of the examined urban areas. **Notes:** The full names of the provinces are Anhui, Beijing, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Henan, Heilongjiang, Hubei, Hunan, Jilin, Jiangsu, Jiangxi, Liaoning, Neimenggu, Ningxia, Qinghai, Shandong, Shanxi, Shannxi, Shanghai, Sichuan, Tianjin, Xizang, Xinjiang, Yunnan, Zhejiang. Due to lack of data from Taiwan, Hong Kong, Xizang and Macao, these provinces were not considered




Fig. 4 Frequency distribution of Carcinogenic risk index for the three population groups




Fig. 5 Spatial distribution of risks of As for the examined urban areas in China. **Notes:** The full names of the provinces are Anhui, Beijing, Chongqing, Fujian, Gansu, Guangdong, Guangxi, Guizhou, Hainan, Hebei, Henan, Heilongjiang, Hubei, Hunan, Jilin, Jiangsu, Jiangxi, Liaoning, Neimenggu, Ningxia, Qinghai, Shandong, Shanxi, Shannxi, Shanghai, Sichuan, Tianjin, Xizang, Xinjiang, Yunnan, Zhejiang. Due to lack of data from Taiwan, Hong Kong, Xinjiang and Macao, these provinces were not considered.