



RAKSHITA N 2024-IT ▾

R2**Started on** Tuesday, 18 November 2025, 11:04 AM**State** Finished**Completed on** Tuesday, 18 November 2025, 11:04 AM**Time taken** 48 secs**Marks** 1.00/1.00**Grade** 4.00 out of 4.00 (100%)

Question 1 | Correct | Mark 1.00 out of 1.00

Given an array A of sorted integers and another non negative integer k, find if there exists 2 indices i and j such that $A[j] - A[i] = k$, $i \neq j$.

Input Format:

First Line n - Number of elements in an array

Next n Lines - N elements in the array

k - Non - Negative Integer

Output Format:

1 - If pair exists

0 - If no pair exists

Explanation for the given Sample Testcase:

YES as $5 - 1 = 4$

So Return 1.

For example:

Input	Result
3 1 3 5 4	1

Answer: (penalty regime: 0 %)

```

1  #include <stdio.h>
2
3  int main() {
4      int n;
5      scanf("%d", &n);
6
7      int A[n];
8      for (int i = 0; i < n; i++)
9          scanf("%d", &A[i]);
10
11     int k;
12     scanf("%d", &k);
13
14     int i = 0, j = 1;
15     int found = 0;
16
17     while (j < n) {
18         if (i == j) {
19             j++;
20             continue;
21         }
22
23         int diff = A[j] - A[i];
24
25         if (diff == k) {
26             found = 1;
27             break;
28         } else if (diff < k) {
29             j++;
30         } else {
31             i++;
32         }
33     }
34
35     printf("%d\n", found);
36     return 0;
37 }
38

```

	Input	Expected	Got	
✓	3 1 3 5 4	1	1	✓
✓	10 1 4 6 8 12 14 15 20 21 25 1	1	1	✓
✓	10 1 2 3 5 11 14 16 24 28 29 0	0	0	✓
✓	10 0 2 3 7 13 14 15 20 24 25 10	1	1	✓

Passed all tests! ✓

Correct

Marks for this submission: 1.00/1.00.

[Back to Course](#)