Symbolic gate atlas

Exact identities. Explicit phase policies. Native entangler counts.

10 targets × 32 basis panels

Each target has one page with 32 basis panels in alphabetical order. Blue labels denote literal matrix equality. Brown labels denote a projective circuit with its exact scalar factor. Both appear when the projective construction is strictly cheaper. Physical GP operations are forbidden.
E counts native two-qubit primitives, L counts one-qubit operations. Scores are displayed rounded; the archive stores exact expressions. Objective version 2 clamps negative denominator penalties; an exact compatibility audit preserves the earlier retained scores. Only an explicit matching certificate establishes a minimum entangler count. A verified upper bound does not establish minimum local cost. Published-cap notes identify constructions above known universal two-qubit bounds; these remain optimization gaps. Proof conditions, source hashes and attempt histories remain in the archive.
Open cases are shown as open. Proved obstructions preserve the supplied basis and may be conditional for parameterized families. Register constructors use the recorded integer-width or matrix fixtures. These pages do not claim a solution for every register size.
Circuits are drawn by Qiskit's LaTeX drawer. Long circuits continue on successive rows in time order. An explicit $[G]^{\times n}$ box denotes n consecutive identical primitive operations; their full physical count remains in E.
Long exact expressions use numbered η_j labels defined in the expression glossary. No parameter is rounded or dropped.
The separate Clifford volume lists every exact phase in each distinct one-qubit alphabet, with 24 circuits per page. Identical alphabets shared by multiple presets are printed once.

Archive SHA256: 0bb0da1e6a7fe9d555178ac093275c11b0bbcd580b91d99aef08e4ad5e7a0cec

Alphabetical target index

Links lead to the 32-basis page for each target.

C3SX . . . 4
C4X . . . 5
Canonical . . . 6
CZS . . . 7
fSim . . . 8
Peres . . . 9
PhasedFSim . . . 10
SWAPPow . . . 11
SYC . . . 12
TK2 . . . 13

Native basis key

Every panel preserves its supplied alphabet. Native units and macro counts matter.


Preset	Entangler	Local alphabet	Scope / units
aqt_rxx	Rxx	R, Rz	exact continuous
berkeley_b	B	U3	exact continuous
clifford_only	CX	H, S	Clifford analysis/control; not universal.
clifford_t_cx	CX	H, S, T, Tdg	Dense universal; exact only for representable targets, not arbitrary symbolic angles.
clifford_t_cz	CZ	H, S, T, Tdg	approximate discrete
cx_h_phase	CX	H, P	exact continuous
cx_rz_ry	CX	Rz, Ry	exact continuous
cx_u	CX	U3	exact continuous
cz_rz_rx	CZ	Rz, Rx	exact continuous
cz_u	CZ	U3	exact continuous
google_cz	CZ	PhasedXZ	exact continuous
google_fsim	fSim	PhasedXZ	exact continuous
google_sqrt_iswap	SiSWAP	PhasedXZ	exact continuous
google_syc	SYC	PhasedXZ	exact continuous
ibm_cx	CX	Rz, Sx, X	exact continuous
ibm_cz	CZ	Rz, Sx, X	exact continuous
ibm_ecr	ECR	Rz, Sx, X	exact continuous
ionq_aria	IonQMS	GPi, GPi2	Native turns; interaction theta restricted to [0,1/4] on hardware.
ionq_aria_full	IonQMSMax	GPi, GPi2	exact continuous
ionq_forte	IonQZZ	GPi, GPi2	exact continuous
iqm_cz	CZ	R	R(theta,phi) represents PRX in radians; IQM API units require conversion.
iswap_u	iSWAP	U3	exact continuous
quantinuum_zzmax	ZZMax	PhasedX, Rz	exact continuous
quantinuum_zzphase	ZZPhase	PhasedX, Rz	exact continuous
rigetti_cz	CZ	Rz, Sx, X	Sx differs from native Rx(pi/2) by a tracked global phase.
sqrt_iswap_u	SiSWAP	U3	exact continuous
sqrt_swap_u	SSWAP	U3	exact continuous
tket_tk2	TK2	PhasedX, Rz	Compiler macro count; expand TK2 before comparing physical pulse counts.
xx90_u	XX90	U3	exact continuous
xx_rz_ry	Rxx	Rz, Ry	exact continuous
zx_u	Rzx	U3	exact continuous
zz_u	Rzz	U3	exact continuous

C3SX

4 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx

PROJECTIVE E=13 / L=121 / score ≈ 335.188 — E upper bound



$U = -(-1)^{\frac{17}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=40 / L=115 / score ≈ 442.290 — E upper bound



clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: not a clifford

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: determinant and clifford t field

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: determinant and clifford t field


cx_h_phase

EXACT PHASE E=20 / L=35 / score ≈ 166.842 — E upper bound



cx_rz_ry


PROJECTIVE E=20 / L=24 / score ≈ 145.248 — E upper bound



$U = e^{-\frac{31i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cx_u

EXACT PHASE E=20 / L=22 / score ≈ 141.332 — E upper bound



cz_rz_rx


PROJECTIVE E=20 / L=40 / score ≈ 179.680 — E upper bound



$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cz_u

EXACT PHASE E=20 / L=27 / score ≈ 158.962 — E upper bound




google_cz

EXACT PHASE E=20 / L=75 / score ≈ 268.850 — E upper bound




google_fsim

EXACT PHASE E=13 / L=54 / score ≈ 197.376 — E upper bound




google_sqrt_iswap

EXACT PHASE E=40 / L=115 / score ≈ 437.090 — E upper bound




google_syc

EXACT PHASE E=120 / L=115 / score ≈ 749.090 — E upper bound



ibm_cx


PROJECTIVE E=20 / L=26 / score ≈ 149.448 — E upper bound



$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz


PROJECTIVE E=20 / L=84 / score ≈ 266.080 — E upper bound



$U = e^{-\frac{31i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr


PROJECTIVE E=20 / L=88 / score ≈ 274.902 — E upper bound



$U = e^{-\frac{15i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria


PROJECTIVE E=20 / L=170 / score ≈ 479.260 — E upper bound



$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full


PROJECTIVE E=20 / L=170 / score ≈ 471.220 — E upper bound



$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte


PROJECTIVE E=20 / L=190 / score ≈ 527.220 — E upper bound



$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=20 / L=171 / score ≈ 477.884 — E upper bound



$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=26 / L=41 / score ≈ 200.446 — E upper bound




PROJECTIVE E=26 / L=40 / score ≈ 196.440 — E upper bound



$U = \eta_{15}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax


PROJECTIVE E=20 / L=135 / score ≈ 392.778 — E upper bound



$U = \sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase


PROJECTIVE E=13 / L=65 / score ≈ 211.808 — E upper bound



$U = -(-1)^{\frac{17}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz


PROJECTIVE E=20 / L=157 / score ≈ 422.154 — E upper bound



$U = e^{\frac{1i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


sqrt_iswap_u

EXACT PHASE E=40 / L=115 / score ≈ 434.290 — E upper bound




sqrt_swap_u

EXACT PHASE E=40 / L=115 / score ≈ 434.290 — E upper bound



tket_tk2


PROJECTIVE E=13 / L=65 / score ≈ 211.872 — E upper bound



$U = -(-1)^{\frac{17}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


xx90_u

EXACT PHASE E=20 / L=95 / score ≈ 314.170 — E upper bound



xx_rz_ry


PROJECTIVE E=13 / L=67 / score ≈ 214.560 — E upper bound



$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


zx_u

EXACT PHASE E=13 / L=26 / score ≈ 129.782 — E upper bound




zz_u

EXACT PHASE E=13 / L=29 / score ≈ 135.000 — E upper bound



PROJECTIVE E=13 / L=27 / score ≈ 128.988 — E upper bound




$U = e^{\frac{15i\pi}{32}}\, C$. Scalar metadata; no physical GP.

C4X

5 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=27 / L=234 / score ≈ 648.754 — E upper bound



$U = (-1)^{\frac{17}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=72 / L=213 / score ≈ 803.678 — E upper bound



clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: not a clifford

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: determinant and clifford t field

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: determinant and clifford t field


cx_h_phase

EXACT PHASE E=36 / L=69 / score ≈ 307.686 — E upper bound



cx_rz_ry


PROJECTIVE E=36 / L=54 / score ≈ 279.708 — E upper bound



$U = e^{\frac{i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cx_u

EXACT PHASE E=36 / L=41 / score ≈ 253.846 — E upper bound



cz_rz_rx

PROJECTIVE E=36 / L=69 / score ≈ 310.372 — E upper bound




Exact definitions: η_{16}, η_{17} in the expression glossary.

$U = \sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cz_u


EXACT PHASE E=36 / L=46 / score ≈ 275.510 — E upper bound



Exact definitions: η_{18}, η_{19} in the expression glossary.


google_cz

EXACT PHASE E=36 / L=141 / score ≈ 491.646 — E upper bound



google_fsim

EXACT PHASE E=27 / L=114 / score ≈ 398.192 — E upper bound




google_sqrt_iswap

EXACT PHASE E=72 / L=213 / score ≈ 794.478 — E upper bound



google_syc

EXACT PHASE E=216 / L=213 / score ≈ 1356.078 — E upper bound



ibm_cx

PROJECTIVE E=36 / L=59 / score ≈ 288.898 — E upper bound



$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=36 / L=130 / score ≈ 429.746 — E upper bound



$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=36 / L=169 / score ≈ 513.392 — E upper bound

$U = e^{-\frac{7i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria

PROJECTIVE E=36 / L=318 / score ≈ 884.452 — E upper bound

$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full

PROJECTIVE E=36 / L=318 / score ≈ 869.980 — E upper bound

$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=36 / L=354 / score ≈ 970.780 — E upper bound

$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=36 / L=325 / score ≈ 894.500 — E upper bound

$U = -\sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=54 / L=79 / score ≈ 404.674 — E upper bound

Exact definitions: η_{28}, η_{29} in the expression glossary.

PROJECTIVE E=54 / L=78 / score ≈ 400.668 — E upper bound

$U = \eta_{29}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax

PROJECTIVE E=36 / L=241 / score ≈ 698.678 — E upper bound

$U = \sqrt[32]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase

PROJECTIVE E=27 / L=151 / score ≈ 468.080 — E upper bound

$U = (-1)^{\frac{1}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz

PROJECTIVE E=36 / L=130 / score ≈ 429.746 — E upper bound

$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=72 / L=213 / score ≈ 789.278 — E upper bound

sqrt_swap_u

EXACT PHASE E=72 / L=213 / score ≈ 789.278 — E upper bound

tket_tk2

PROJECTIVE E=27 / L=162 / score ≈ 490.246 — E upper bound

$U = (-1)^{\frac{1}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=36 / L=177 / score ≈ 573.062 — E upper bound

xx_rz_ry

PROJECTIVE E=27 / L=147 / score ≈ 453.948 — E upper bound

$U = e^{\frac{17i\pi}{32}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=27 / L=56 / score ≈ 265.190 — E upper bound

PROJECTIVE E=27 / L=54 / score ≈ 259.978 — E upper bound

$U = e^{\frac{7i\pi}{8}}\, C$. Scalar metadata; no physical GP.

zz_u

EXACT PHASE E=27 / L=55 / score ≈ 243.618 — E upper bound

Exact definitions: $\eta_{14}, \eta_{15}, \eta_{16}, \eta_{17}$ in the expression glossary.

PROJECTIVE E=27 / L=53 / score ≈ 237.606 — E upper bound

Exact definitions: $\eta_{14}, \eta_{15}, \eta_{16}, \eta_{17}, \eta_{18}$ in the expression glossary.


$U = (-1)^{\frac{1}{32}}\, C$. Scalar metadata; no physical GP.

Canonical (a, b, c)

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx

EXACT PHASE E=3 / L=41 / score ≈ 102.928 — E upper bound



PROJECTIVE E=3 / L=20 / score ≈ 56.872 — E upper bound



$U = \eta_{36}\, C$. Scalar metadata; no physical GP.


berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 23.242 — conditional E minimum



Exact definitions: $\eta_{27}, \eta_{28}, \eta_{29}, \eta_{30}, \eta_{31}, \eta_{32}, \eta_{33}, \eta_{34}, \eta_{35}, \eta_{36}, \eta_{37}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 21.236 — conditional E minimum



Exact definitions: $\eta_{39}, \eta_{40}, \eta_{41}, \eta_{42}, \eta_{43}, \eta_{44}, \eta_{45}, \eta_{46}, \eta_{47}, \eta_{48}$ in the expression glossary.

$U = \eta_{49}\, C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx


Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz


Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

cx_h_phase

EXACT PHASE E=3 / L=20 / score ≈ 52.620 — conditional E minimum




PROJECTIVE E=3 / L=12 / score ≈ 36.612 — conditional E minimum




$U = e^{i(a-b+c)}\, C$. Scalar metadata; no physical GP.

cx_rz_ry

EXACT PHASE E=4 / L=17 / score ≈ 52.834 — conditional E minimum




PROJECTIVE E=3 / L=11 / score ≈ 36.222 — conditional E minimum




$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

cx_u

EXACT PHASE E=3 / L=7 / score ≈ 28.242 — conditional E minimum



PROJECTIVE E=3 / L=5 / score ≈ 24.230 — conditional E minimum



$U = e^{ic}\, C$. Scalar metadata; no physical GP.

cz_rz_rx

EXACT PHASE E=4 / L=28 / score ≈ 77.256 — conditional E minimum




PROJECTIVE E=3 / L=11 / score ≈ 35.822 — conditional E minimum



$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

cz_u

EXACT PHASE E=3 / L=10 / score ≈ 34.660 — conditional E minimum




PROJECTIVE E=3 / L=8 / score ≈ 30.648 — conditional E minimum



$U = e^{ic}\, C$. Scalar metadata; no physical GP.

google_cz

EXACT PHASE E=3 / L=10 / score ≈ 35.111 — conditional E minimum




PROJECTIVE E=3 / L=8 / score ≈ 30.439 — conditional E minimum



$U = e^{i(a-b+c)}\, C$. Scalar metadata; no physical GP.

google_fsim

EXACT PHASE E=2 / L=6 / score ≈ 21.365 — E upper bound



PROJECTIVE E=2 / L=4 / score ≈ 16.693 — E upper bound



$U = e^{ic}\, C$. Scalar metadata; no physical GP.

google_sqrt_iswap

EXACT PHASE E=6 / L=19 / score ≈ 69.165 — E upper bound



PROJECTIVE E=6 / L=17 / score ≈ 63.893 — E upper bound



$U = \eta_{50}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

google_syc

EXACT PHASE E=18 / L=19 / score ≈ 115.365 — E upper bound



PROJECTIVE E=18 / L=17 / score ≈ 110.693 — E upper bound



$U = \eta_{51}\, C$. Scalar metadata; no physical GP.

ibm_cx

EXACT PHASE E=3 / L=16 / score ≈ 45.216 — conditional E minimum




PROJECTIVE E=3 / L=13 / score ≈ 39.012 — conditional E minimum



$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

ibm_cz

EXACT PHASE E=3 / L=20 / score ≈ 53.620 — conditional E minimum



ibm_ecr

EXACT PHASE E=3 / L=24 / score ≈ 61.018 — conditional E minimum



PROJECTIVE E=3 / L=22 / score ≈ 57.214 — conditional E minimum



$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

ionq_aria

EXACT PHASE E=3 / L=27 / score ≈ 74.463 — E upper bound



PROJECTIVE E=3 / L=27 / score ≈ 74.263 — E upper bound



$U = \frac{\sqrt[3]{-1}\sqrt{2}(-1+i)}{2}\, C$. Scalar metadata; no physical GP.

ionq_aria_full

EXACT PHASE E=3 / L=27 / score ≈ 73.257 — E upper bound




PROJECTIVE E=3 / L=27 / score ≈ 73.057 — E upper bound



$U = \frac{\sqrt[3]{-1}\sqrt{2}(-1+i)}{2}\, C$. Scalar metadata; no physical GP.

ionq_forte

EXACT PHASE E=3 / L=30 / score ≈ 81.457 — E upper bound




iqm_cz

EXACT PHASE E=4 / L=27 / score ≈ 75.708 — conditional E minimum



PROJECTIVE E=3 / L=13 / score ≈ 40.452 — conditional E minimum



$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

iswap_u

EXACT PHASE E=3 / L=13 / score ≈ 41.278 — E upper bound



PROJECTIVE E=3 / L=10 / score ≈ 35.260 — E upper bound



$U = \frac{\sqrt{2}i(1+i)e^{ib}}{2}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax

EXACT PHASE E=4 / L=25 / score ≈ 71.206 — E upper bound




PROJECTIVE E=3 / L=10 / score ≈ 34.293 — conditional E minimum



$U = -iC$. Scalar metadata; no physical GP.


quantinuum_zzphase

EXACT PHASE E=3 / L=18 / score ≈ 53.049 — E upper bound




rigetti_cz

EXACT PHASE E=3 / L=20 / score ≈ 53.620 — conditional E minimum



sqrt_iswap_u

EXACT PHASE E=6 / L=19 / score ≈ 67.914 — E upper bound



PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound



$U = \frac{\sqrt[3]{-1}\sqrt{2}(-1+i)e^{ic}}{2}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

sqrt_swap_u

EXACT PHASE E=6 / L=19 / score ≈ 68.714 — E upper bound



PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound



$U = -\sqrt[4]{-1}e^{ic}\, C$. Scalar metadata; no physical GP.


tket_tk2

EXACT PHASE E=1 / L=0 / score ≈ 4.997 — conditional E minimum



xx90_u

EXACT PHASE E=3 / L=10 / score ≈ 33.860 — conditional E minimum



PROJECTIVE E=3 / L=8 / score ≈ 29.448 — conditional E minimum




$U = ie^{-ib}\, C$. Scalar metadata; no physical GP.

xx_rz_ry

EXACT PHASE E=3 / L=41 / score ≈ 98.888 — E upper bound



PROJECTIVE E=3 / L=31 / score ≈ 80.068 — E upper bound




$U = \eta_{36}\, C$. Scalar metadata; no physical GP.

zx_u

EXACT PHASE E=3 / L=13 / score ≈ 43.284 — E upper bound



PROJECTIVE E=3 / L=11 / score ≈ 39.272 — E upper bound



$U = \frac{\sqrt[3]{-1}\sqrt{2}(1-i)e^{ic}}{2}\, C$. Scalar metadata; no physical GP.

zz_u

EXACT PHASE E=3 / L=12 / score ≈ 38.478 — E upper bound




CZS (θ, ϕ, γ)

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=4 / L=40 / score ≈ 106.532 — E upper bound



$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=2 / L=6 / score ≈ 21.636 — E upper bound



Exact definitions: $\eta_{52}, \eta_{53}, \eta_{54}, \eta_{55}, \eta_{56}$ in the expression glossary.

clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family


cx_h_phase

EXACT PHASE E=4 / L=40 / score ≈ 101.640 — E upper bound



cx_rz_ry


PROJECTIVE E=4 / L=40 / score ≈ 105.680 — E upper bound



$U = e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cx_u

EXACT PHASE E=4 / L=21 / score ≈ 67.726 — E upper bound



cz_rz_rx

PROJECTIVE E=4 / L=42 / score ≈ 110.084 — E upper bound



$U = e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cz_u

EXACT PHASE E=4 / L=29 / score ≈ 85.374 — E upper bound



google_cz

EXACT PHASE E=4 / L=29 / score ≈ 87.268 — E upper bound



google_fsim

EXACT PHASE E=2 / L=9 / score ≈ 32.635 — E upper bound




google_sqrt_iswap

EXACT PHASE E=2 / L=6 / score ≈ 21.757 — E upper bound

Exact definitions: $\eta_{57}, \eta_{58}, \eta_{59}, \eta_{60}, \eta_{61}, \eta_{62}, \eta_{63}, \eta_{64}, \eta_{65}, \eta_{66}, \eta_{67}, \eta_{68}$ in the expression glossary.

google_syc

EXACT PHASE E=24 / L=37 / score ≈ 183.316 — E upper bound



ibm_cx

PROJECTIVE E=4 / L=32 / score ≈ 84.832 — E upper bound

$U = -e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=4 / L=54 / score ≈ 130.448 — E upper bound

$U = -e^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=4 / L=44 / score ≈ 108.832 — E upper bound

$U = -e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria

PROJECTIVE E=4 / L=58 / score ≈ 153.634 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full

PROJECTIVE E=4 / L=58 / score ≈ 152.026 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=4 / L=62 / score ≈ 163.226 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=4 / L=59 / score ≈ 151.836 — E upper bound

$U = e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=4 / L=21 / score ≈ 66.126 — E upper bound

quantinuum_zzmax

PROJECTIVE E=4 / L=38 / score ≈ 102.373 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase

PROJECTIVE E=4 / L=38 / score ≈ 103.181 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz

PROJECTIVE E=4 / L=54 / score ≈ 130.448 — E upper bound

$U = -e^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=2 / L=6 / score ≈ 20.036 — E upper bound

Exact definitions: $\eta_{69}, \eta_{70}, \eta_{71}, \eta_{72}, \eta_{73}, \eta_{74}$ in the expression glossary.

sqrt_swap_u

EXACT PHASE E=8 / L=37 / score ≈ 120.622 — E upper bound

tket_tk2

PROJECTIVE E=2 / L=8 / score ≈ 29.349 — E upper bound

$U = e^{\frac{i(\gamma+\pi)}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=4 / L=33 / score ≈ 96.598 — E upper bound

xx_rz_ry

PROJECTIVE E=4 / L=66 / score ≈ 162.540 — E upper bound

$U = -ie^{\frac{5\pi}{2}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=4 / L=29 / score ≈ 87.782 — E upper bound

zz_u


EXACT PHASE E=4 / L=33 / score ≈ 91.006 — E upper bound

fSim (θ, ϕ)

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=3 / L=22 / score ≈ 61.276 — E upper bound



$U = \eta_{75}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 23.442 — conditional E minimum



Exact definitions: $\eta_{76}, \eta_{77}, \eta_{78}, \eta_{79}, \eta_{80}, \eta_{81}, \eta_{82}, \eta_{83}, \eta_{84}, \eta_{85}, \eta_{86}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 21.436 — conditional E minimum

Exact definitions: $\eta_{78}, \eta_{79}, \eta_{80}, \eta_{81}, \eta_{82}, \eta_{83}, \eta_{84}, \eta_{85}, \eta_{86}, \eta_{87}$ in the expression glossary.

$U = \eta_{88}\, C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family


cx_h_phase

EXACT PHASE E=3 / L=14 / score ≈ 41.216 — E upper bound



cx_rz_ry


PROJECTIVE E=3 / L=13 / score ≈ 40.626 — E upper bound



$U = -e^{-\frac{i(\phi-\pi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cx_u

EXACT PHASE E=3 / L=7 / score ≈ 28.642 — E upper bound



cz_rz_rx


PROJECTIVE E=3 / L=12 / score ≈ 38.024 — E upper bound



$U = -(-1)^{\frac{3}{4}}\, e^{-\frac{i\phi}{2}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cz_u

EXACT PHASE E=3 / L=10 / score ≈ 35.260 — E upper bound



google_cz

EXACT PHASE E=3 / L=10 / score ≈ 35.711 — E upper bound



google_fsim

EXACT PHASE E=1 / L=0 / score ≈ 4.004 — E upper bound



google_sqrt_iswap

EXACT PHASE E=6 / L=21 / score ≈ 74.238 — E upper bound



PROJECTIVE E=6 / L=19 / score ≈ 68.965 — E upper bound

$U = \frac{\sqrt{2}(1-i)e^{-i\theta}}{2}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

google_syc

EXACT PHASE E=18 / L=21 / score ≈ 120.438 — E upper bound



PROJECTIVE E=18 / L=19 / score ≈ 115.765 — E upper bound

$U = -ie^{-i\theta}\, C$. Scalar metadata; no physical GP.

ibm_cx

PROJECTIVE E=3 / L=15 / score ≈ 43.416 — E upper bound

$U = -e^{-\frac{i(\phi-\pi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=3 / L=21 / score ≈ 56.024 — E upper bound



$U = \sqrt[4]{-1}\, e^{-\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=3 / L=23 / score ≈ 59.416 — E upper bound

$U = -ie^{-\frac{i(\phi-\pi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria

PROJECTIVE E=3 / L=31 / score ≈ 84.131 — E upper bound

$U = \eta_{89}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full

PROJECTIVE E=3 / L=31 / score ≈ 82.925 — E upper bound

$U = \eta_{89}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=3 / L=34 / score ≈ 91.325 — E upper bound

$U = \eta_{89}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=3 / L=19 / score ≈ 54.676 — E upper bound

$U = -(-1)^{\frac{3}{4}}\, e^{-\frac{i\phi}{2}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=3 / L=15 / score ≈ 46.690 — E upper bound

PROJECTIVE E=3 / L=12 / score ≈ 39.872 — E upper bound

$U = \eta_{90}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax

PROJECTIVE E=3 / L=12 / score ≈ 38.897 — E upper bound

$U = -ie^{-\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase

PROJECTIVE E=3 / L=20 / score ≈ 57.653 — E upper bound

$U = e^{-\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz

PROJECTIVE E=3 / L=21 / score ≈ 56.024 — E upper bound

$U = \sqrt[4]{-1}\, e^{-\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=6 / L=19 / score ≈ 68.314 — E upper bound

Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

sqrt_swap_u

EXACT PHASE E=6 / L=19 / score ≈ 69.114 — E upper bound

$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

tket_tk2

PROJECTIVE E=1 / L=2 / score ≈ 9.601 — conditional E minimum

$U = e^{-\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=3 / L=12 / score ≈ 39.272 — E upper bound

PROJECTIVE E=3 / L=10 / score ≈ 34.060 — E upper bound

$U = ie^{\frac{i(\phi+2\theta)}{4}}\, C$. Scalar metadata; no physical GP.

xx_rz_ry

PROJECTIVE E=3 / L=32 / score ≈ 82.270 — E upper bound

$U = \eta_{75}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=3 / L=13 / score ≈ 43.684 — E upper bound

zz_u

EXACT PHASE E=3 / L=16 / score ≈ 47.902 — E upper bound

PROJECTIVE E=3 / L=14 / score ≈ 43.090 — E upper bound


$U = e^{\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.

Peres

3 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=5 / L=35 / score ≈ 99.714 — E upper bound



$U = \frac{\sqrt[8]{-1}\sqrt{2}(-1+i)}{2}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=10 / L=29 / score ≈ 109.374 — E upper bound




clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: not a clifford


clifford_t_cx

EXACT PHASE E=5 / L=9 / score ≈ 38.000 — proved E minimum




clifford_t_cz

EXACT PHASE E=5 / L=17 / score ≈ 54.000 — proved E minimum




cx_h_phase

EXACT PHASE E=5 / L=9 / score ≈ 40.814 — proved E minimum



cx_rz_ry


PROJECTIVE E=5 / L=10 / score ≈ 43.220 — proved E minimum



$U = -\sqrt[8]{-1}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cx_u

EXACT PHASE E=5 / L=8 / score ≈ 39.248 — proved E minimum



cz_rz_rx


PROJECTIVE E=5 / L=12 / score ≈ 47.224 — proved E minimum



$U = e^{-\frac{3i\pi}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cz_u


EXACT PHASE E=5 / L=7 / score ≈ 37.242 — proved E minimum



Exact definitions: η_{22} in the expression glossary.


google_cz

EXACT PHASE E=5 / L=19 / score ≈ 65.714 — proved E minimum



google_fsim

EXACT PHASE E=5 / L=19 / score ≈ 65.734 — E upper bound



google_sqrt_iswap


EXACT PHASE E=10 / L=29 / score ≈ 108.374 — E upper bound

PROJECTIVE E=10 / L=29 / score ≈ 107.774 — E upper bound

$U = \frac{\sqrt{2}(1-i)^5}{8}\,C$. Scalar metadata; no physical GP.

google_syc

EXACT PHASE E=30 / L=29 / score ≈ 185.774 — E upper bound



ibm_cx


PROJECTIVE E=5 / L=12 / score ≈ 47.218 — proved E minimum



$U = e^{-\frac{3i\pi}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=5 / L=26 / score ≈ 75.226 — proved E minimum



$U = e^{-\frac{3i\pi}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=5 / L=31 / score ≈ 86.026 — proved E minimum

$U = (-1)^{\frac{3}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria

PROJECTIVE E=5 / L=43 / score ≈ 119.516 — E upper bound

$U = \frac{\sqrt[8]{-1}\sqrt{2}(1+i)^5}{8}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full

PROJECTIVE E=5 / L=43 / score ≈ 117.506 — E upper bound

$U = \frac{\sqrt[8]{-1}\sqrt{2}(1+i)^5}{8}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=5 / L=48 / score ≈ 131.506 — E upper bound

$U = \frac{\sqrt[8]{-1}\sqrt{2}(1+i)}{2}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=5 / L=45 / score ≈ 123.380 — proved E minimum

$U = e^{\frac{i\pi}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=10 / L=18 / score ≈ 84.908 — E upper bound

PROJECTIVE E=10 / L=16 / score ≈ 79.696 — E upper bound

$U = e^{-\frac{i\pi}{8}}\,C$. Scalar metadata; no physical GP.

quantinuum_zzmax

PROJECTIVE E=5 / L=31 / score ≈ 90.486 — proved E minimum

$U = \frac{\sqrt[8]{-1}\sqrt{2}(-1+i)}{2}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase

PROJECTIVE E=5 / L=31 / score ≈ 91.496 — E upper bound

$U = \frac{\sqrt[8]{-1}\sqrt{2}(-1+i)}{2}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz

PROJECTIVE E=5 / L=35 / score ≈ 95.238 — proved E minimum

$U = -(-1)^{\frac{7}{8}}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=10 / L=29 / score ≈ 107.374 — E upper bound

sqrt_swap_u

EXACT PHASE E=10 / L=29 / score ≈ 108.174 — E upper bound

PROJECTIVE E=10 / L=29 / score ≈ 107.374 — E upper bound

$U = e^{-\frac{i\pi}{4}}\,C$. Scalar metadata; no physical GP.

tket_tk2

PROJECTIVE E=5 / L=35 / score ≈ 99.734 — E upper bound

$U = \frac{\sqrt[8]{-1}\sqrt{2}(-1+i)}{2}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=5 / L=24 / score ≈ 77.344 — proved E minimum

xx_rz_ry

PROJECTIVE E=5 / L=35 / score ≈ 94.080 — E upper bound

$U = \eta_{102}\,C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=5 / L=13 / score ≈ 53.288 — E upper bound

Exact definitions: η_{103} in the expression glossary.

PROJECTIVE E=5 / L=11 / score ≈ 48.076 — E upper bound

Exact definitions: η_{101} in the expression glossary.

$U = (-1)^{\frac{3}{8}}\,C$. Scalar metadata; no physical GP.

zz_u

EXACT PHASE E=5 / L=12 / score ≈ 49.082 — E upper bound

Exact definitions: η_{22} in the expression glossary.

PROJECTIVE E=5 / L=10 / score ≈ 44.270 — E upper bound

Exact definitions: η_{22} in the expression glossary.


$U = \frac{\sqrt{2}(1+i)^5}{8}\,C$. Scalar metadata; no physical GP.

PhasedFSim $(\theta, \zeta, \chi, \gamma, \phi)$

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=3 / L=26 / score ≈ 70.084 — E upper bound



$U = \eta_{104}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 23.642 — conditional E minimum



Exact definitions: $\eta_{101}, \eta_{104}, \eta_{76}, \eta_{77}, \eta_{78}, \eta_{80}, \eta_{81}, \eta_{82}, \eta_{84}, \eta_{86}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 21.636 — conditional E minimum



Exact definitions: $\eta_{101}, \eta_{104}, \eta_{76}, \eta_{78}, \eta_{80}, \eta_{81}, \eta_{82}, \eta_{84}, \eta_{86}, \eta_{87}$ in the expression glossary.
$U = \eta_{88}\, C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family


cx_h_phase

EXACT PHASE E=3 / L=15 / score ≈ 43.418 — E upper bound



cx_rz_ry


PROJECTIVE E=3 / L=17 / score ≈ 49.034 — E upper bound




$U = \eta_{107}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cx_u

EXACT PHASE E=3 / L=11 / score ≈ 37.466 — E upper bound



PROJECTIVE E=3 / L=11 / score ≈ 37.266 — E upper bound



$U = e^{\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.

cz_rz_rx


PROJECTIVE E=3 / L=13 / score ≈ 40.226 — E upper bound



Exact definitions: η_{108} in the expression glossary.
$U = \eta_{107}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cz_u

EXACT PHASE E=3 / L=12 / score ≈ 39.672 — E upper bound



google_cz

EXACT PHASE E=3 / L=12 / score ≈ 40.784 — E upper bound



Exact definitions: η_{109}, η_{110} in the expression glossary.

google_fsim

EXACT PHASE E=1 / L=4 / score ≈ 14.149 — E upper bound



google_sqrt_iswap

EXACT PHASE E=6 / L=25 / score ≈ 84.383 — E upper bound

PROJECTIVE E=6 / L=23 / score ≈ 79.111 — E upper bound

$U = \frac{\sqrt{2}(1-i)e^{-i\theta}}{2}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

google_syc

EXACT PHASE E=18 / L=25 / score ≈ 130.583 — E upper bound



PROJECTIVE E=18 / L=23 / score ≈ 125.911 — E upper bound

$U = -ie^{-i\theta}\, C$. Scalar metadata; no physical GP.

ibm_cx

PROJECTIVE E=3 / L=19 / score ≈ 52.224 — E upper bound

$U = -e^{-\frac{i(\phi_3+\phi_0-\pi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz


PROJECTIVE E=3 / L=22 / score ≈ 58.226 — E upper bound



Exact definitions: η_{111} in the expression glossary.
$U = \sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr


PROJECTIVE E=3 / L=27 / score ≈ 68.224 — E upper bound



$U = -ie^{-\frac{i(4+\phi-\pi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria


PROJECTIVE E=3 / L=39 / score ≈ 103.869 — E upper bound



$U = \eta_{112}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full


PROJECTIVE E=3 / L=39 / score ≈ 102.663 — E upper bound



$U = \eta_{112}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte


PROJECTIVE E=3 / L=42 / score ≈ 111.063 — E upper bound



$U = \eta_{112}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz


PROJECTIVE E=3 / L=25 / score ≈ 68.700 — E upper bound



$U = \eta_{107}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=3 / L=17 / score ≈ 51.102 — E upper bound




PROJECTIVE E=3 / L=14 / score ≈ 44.284 — E upper bound



$U = \eta_{90}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax


PROJECTIVE E=3 / L=13 / score ≈ 41.299 — E upper bound



$U = -ie^{-\frac{i(4+\phi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase


PROJECTIVE E=3 / L=22 / score ≈ 62.057 — E upper bound



$U = e^{-\frac{i(4+\phi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz


PROJECTIVE E=3 / L=22 / score ≈ 58.226 — E upper bound



$U = \sqrt[4]{-1}e^{-\frac{i(4+\phi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=6 / L=23 / score ≈ 77.138 — E upper bound



Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

sqrt_swap_u

EXACT PHASE E=6 / L=23 / score ≈ 77.938 — E upper bound




PROJECTIVE E=6 / L=23 / score ≈ 77.138 — E upper bound



$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

tket_tk2


PROJECTIVE E=1 / L=4 / score ≈ 14.005 — conditional E minimum




$U = e^{-\frac{i(4+\phi)}{4}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=3 / L=14 / score ≈ 43.684 — E upper bound




PROJECTIVE E=3 / L=12 / score ≈ 38.472 — E upper bound



$U = ie^{\frac{i(\phi+2\theta)}{4}}\, C$. Scalar metadata; no physical GP.

xx_rz_ry


PROJECTIVE E=3 / L=36 / score ≈ 91.078 — E upper bound



$U = \eta_{104}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=3 / L=17 / score ≈ 52.508 — E upper bound



zz_u

EXACT PHASE E=3 / L=18 / score ≈ 52.314 — E upper bound



PROJECTIVE E=3 / L=16 / score ≈ 47.502 — E upper bound




$U = e^{\frac{i\phi}{4}}\, C$. Scalar metadata; no physical GP.

SWAPPow $(t, global_{shift})$

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=3 / L=20 / score ≈ 56.872 — E upper bound



$U = \eta_{113} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 23.442 — conditional E minimum



Exact definitions: $\eta_{114}, \eta_{115}, \eta_{116}, \eta_{117}, \eta_{118}, \eta_{119}, \eta_{120}, \eta_{121}, \eta_{122}, \eta_{123}, \eta_{124}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 21.436 — conditional E minimum



Exact definitions: $\eta_{116}, \eta_{117}, \eta_{118}, \eta_{119}, \eta_{120}, \eta_{121}, \eta_{122}, \eta_{123}, \eta_{124}, \eta_{125}$ in the expression glossary.

$U = \eta_{126} C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

cx_h_phase

EXACT PHASE E=3 / L=20 / score ≈ 53.220 — E upper bound


PROJECTIVE E=3 / L=12 / score ≈ 37.212 — E upper bound



$U = e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

cx_rz_ry

PROJECTIVE E=3 / L=11 / score ≈ 36.222 — E upper bound




$U = \eta_{127} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cx_u

EXACT PHASE E=3 / L=7 / score ≈ 28.242 — E upper bound



PROJECTIVE E=3 / L=5 / score ≈ 24.230 — E upper bound



$U = e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

cz_rz_rx


PROJECTIVE E=3 / L=11 / score ≈ 36.222 — E upper bound




$U = \eta_{128} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cz_u

EXACT PHASE E=3 / L=10 / score ≈ 35.260 — E upper bound




PROJECTIVE E=3 / L=8 / score ≈ 31.248 — E upper bound



$U = e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

google_cz

EXACT PHASE E=3 / L=10 / score ≈ 34.060 — E upper bound



PROJECTIVE E=3 / L=8 / score ≈ 30.048 — E upper bound



$U = e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

google_fsim

EXACT PHASE E=1 / L=4 / score ≈ 12.628 — conditional E minimum



PROJECTIVE E=1 / L=2 / score ≈ 8.616 — conditional E minimum



$U = e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

google_sqrt_iswap

EXACT PHASE E=6 / L=19 / score ≈ 67.514 — E upper bound




PROJECTIVE E=6 / L=17 / score ≈ 62.902 — E upper bound



$U = \eta_{130} C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

google_syc

EXACT PHASE E=18 / L=19 / score ≈ 113.914 — E upper bound




PROJECTIVE E=18 / L=17 / score ≈ 109.702 — E upper bound



$U = \eta_{132} C$. Scalar metadata; no physical GP.

ibm_cx


PROJECTIVE E=3 / L=13 / score ≈ 39.012 — E upper bound



$U = \eta_{127} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=3 / L=20 / score ≈ 54.020 — E upper bound



$U = -e^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=3 / L=22 / score ≈ 57.214 — E upper bound

$U = \eta_{133} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


ionq_aria

PROJECTIVE E=3 / L=27 / score ≈ 73.272 — E upper bound

$U = \eta_{134} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full


PROJECTIVE E=3 / L=27 / score ≈ 72.066 — E upper bound



$U = \eta_{134} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=3 / L=30 / score ≈ 80.466 — E upper bound



$U = \eta_{134} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=3 / L=13 / score ≈ 41.052 — E upper bound




$U = \eta_{128} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=3 / L=13 / score ≈ 41.878 — E upper bound



PROJECTIVE E=3 / L=10 / score ≈ 35.860 — E upper bound



$U = \eta_{135} C$. Scalar metadata; no physical GP.

quantinuum_zzmax


PROJECTIVE E=3 / L=10 / score ≈ 34.232 — E upper bound



$U = -ie^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase


PROJECTIVE E=3 / L=18 / score ≈ 52.658 — E upper bound



$U = e^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz


PROJECTIVE E=3 / L=20 / score ≈ 54.020 — E upper bound




$U = -e^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=6 / L=19 / score ≈ 67.914 — E upper bound




PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound




$U = \eta_{136} C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

sqrt_swap_u

EXACT PHASE E=6 / L=19 / score ≈ 68.714 — E upper bound



PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound



$U = -\sqrt[4]{-1}e^{i\pi global_{shift}t} C$. Scalar metadata; no physical GP.

tket_tk2

PROJECTIVE E=1 / L=0 / score ≈ 4.606 — conditional E minimum



$U = e^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

xx90_u

EXACT PHASE E=3 / L=10 / score ≈ 34.860 — E upper bound




PROJECTIVE E=3 / L=8 / score ≈ 30.048 — E upper bound



$U = ie^{\frac{i\pi t\left(2global_{shift}+1\right)}{2}} C$. Scalar metadata; no physical GP.

xx_rz_ry


PROJECTIVE E=3 / L=31 / score ≈ 80.068 — E upper bound




$U = \eta_{113} C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=3 / L=13 / score ≈ 43.284 — E upper bound




PROJECTIVE E=3 / L=11 / score ≈ 39.272 — E upper bound




$U = \eta_{137} C$. Scalar metadata; no physical GP.

zz_u

EXACT PHASE E=3 / L=14 / score ≈ 43.890 — E upper bound



PROJECTIVE E=3 / L=12 / score ≈ 39.078 — E upper bound



$U = e^{\frac{i\pi\left(4global_{shift}+1\right)}{4}} C$. Scalar metadata; no physical GP.

SYC

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx


PROJECTIVE E=3 / L=17 / score ≈ 51.809 — E upper bound



$U = \eta_{138}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 24.076 — proved E minimum



Exact definitions: $\eta_{139}, \eta_{140}, \eta_{141}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 22.787 — proved E minimum



$U = e^{\frac{i\pi}{24}}\, C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: not a clifford

clifford_t_cx


Exact: Proved obstruction: determinant group
Projective: Proved obstruction: outside clifford t number field

clifford_t_cz

Exact: Proved obstruction: determinant group
Projective: Proved obstruction: outside clifford t number field


cx_h_phase

EXACT PHASE E=3 / L=3 / score ≈ 20.157 — proved E minimum



cx_rz_ry


PROJECTIVE E=3 / L=3 / score ≈ 20.157 — proved E minimum



$U = e^{-\frac{7i\pi}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

cx_u

EXACT PHASE E=3 / L=3 / score ≈ 20.169 — proved E minimum



cz_rz_rx

PROJECTIVE E=3 / L=12 / score ≈ 39.975 — proved E minimum

$U = (-1)^{\frac{17}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group


cz_u

EXACT PHASE E=3 / L=7 / score ≈ 29.393 — proved E minimum



google_cz

EXACT PHASE E=3 / L=9 / score ≈ 34.605 — proved E minimum



google_fsim

EXACT PHASE E=1 / L=0 / score ≈ 4.721 — proved E minimum

google_sqrt_iswap

EXACT PHASE E=3 / L=9 / score ≈ 39.025 — proved E minimum

Exact definitions: $\eta_{142}, \eta_{143}, \eta_{144}, \eta_{145}, \eta_{146}, \eta_{147}, \eta_{148}, \eta_{149}, \eta_{150}, \eta_{151}$ in the expression glossary.

PROJECTIVE E=3 / L=8 / score ≈ 34.924 — proved E minimum

Exact definitions: $\eta_{144}, \eta_{145}, \eta_{146}, \eta_{147}, \eta_{148}, \eta_{149}, \eta_{150}, \eta_{151}$ in the expression glossary.
$U = \eta_{152}\, C$. Scalar metadata; no physical GP.

google_syc

EXACT PHASE E=1 / L=0 / score ≈ 4.000 — proved E minimum

ibm_cx

PROJECTIVE E=3 / L=3 / score ≈ 20.157 — proved E minimum

$U = e^{-\frac{7i\pi}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_cz

PROJECTIVE E=3 / L=14 / score ≈ 42.763 — proved E minimum

$U = e^{-\frac{19i\pi}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ibm_ecr

PROJECTIVE E=3 / L=15 / score ≈ 44.763 — proved E minimum

$U = (-1)^{\frac{17}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria

PROJECTIVE E=3 / L=25 / score ≈ 70.219 — E upper bound

$U = \eta_{138}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_aria_full

PROJECTIVE E=3 / L=21 / score ≈ 60.405 — E upper bound

$U = \eta_{153}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

ionq_forte

PROJECTIVE E=3 / L=28 / score ≈ 77.413 — E upper bound

$U = \eta_{138}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iqm_cz

PROJECTIVE E=3 / L=19 / score ≈ 56.627 — proved E minimum

$U = (-1)^{\frac{17}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

iswap_u

EXACT PHASE E=3 / L=9 / score ≈ 37.156 — proved E minimum

PROJECTIVE E=3 / L=7 / score ≈ 31.310 — proved E minimum

$U = e^{\frac{5i}{24}}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax

PROJECTIVE E=3 / L=12 / score ≈ 40.187 — proved E minimum

$U = -(-1)^{\frac{11}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

quantinuum_zzphase

PROJECTIVE E=3 / L=16 / score ≈ 49.801 — E upper bound

$U = \eta_{138}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

rigetti_cz

PROJECTIVE E=3 / L=14 / score ≈ 42.763 — proved E minimum

$U = e^{-\frac{19i\pi}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

sqrt_iswap_u

EXACT PHASE E=3 / L=9 / score ≈ 36.322 — proved E minimum

Exact definitions: $\eta_{154}, \eta_{155}, \eta_{156}, \eta_{157}, \eta_{158}, \eta_{159}, \eta_{160}, \eta_{161}$ in the expression glossary.

PROJECTIVE E=3 / L=8 / score ≈ 32.682 — proved E minimum

Exact definitions: $\eta_{156}, \eta_{157}, \eta_{158}, \eta_{159}, \eta_{160}, \eta_{161}$ in the expression glossary.
$U = \eta_{162}\, C$. Scalar metadata; no physical GP.

sqrt_swap_u

EXACT PHASE E=6 / L=15 / score ≈ 60.641 — E upper bound

PROJECTIVE E=6 / L=15 / score ≈ 59.841 — E upper bound

$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

xx90_u

EXACT PHASE E=3 / L=12 / score ≈ 40.023 — proved E minimum

Exact definitions: η_{163}, η_{164} in the expression glossary.

PROJECTIVE E=3 / L=10 / score ≈ 36.011 — proved E minimum

$U = -(-1)^{\frac{19}{24}}\, C$. Scalar metadata; no physical GP.

xx_rz_ry

PROJECTIVE E=3 / L=18 / score ≈ 52.993 — E upper bound

$U = e^{\frac{11i\pi}{24}}\, C$. Scalar metadata; no physical GP.
Exact: Proved obstruction: determinant group

zx_u

EXACT PHASE E=3 / L=8 / score ≈ 33.005 — E upper bound

zz_u

EXACT PHASE E=3 / L=10 / score ≈ 38.017 — E upper bound

PROJECTIVE E=3 / L=8 / score ≈ 33.205 — E upper bound


$U = \frac{\sqrt{2}(1+i)^3}{4}\, C$. Scalar metadata; no physical GP.

TK2 (a, b, c)

2 qubits / 32 preserved native bases / alphabetical panel order

aqt_rxx

EXACT PHASE E=3 / L=41 / score ≈ 102.928 — E upper bound



PROJECTIVE E=3 / L=20 / score ≈ 56.872 — E upper bound



$U = \eta_{36}\, C$. Scalar metadata; no physical GP.

berkeley_b

EXACT PHASE E=2 / L=7 / score ≈ 23.242 — conditional E minimum



Exact definitions: $\eta_{165}, \eta_{166}, \eta_{167}, \eta_{168}, \eta_{169}, \eta_{170}, \eta_{171}, \eta_{172}, \eta_{173}, \eta_{174}, \eta_{175}$ in the expression glossary.

PROJECTIVE E=2 / L=6 / score ≈ 21.236 — conditional E minimum



Exact definitions: $\eta_{167}, \eta_{168}, \eta_{169}, \eta_{170}, \eta_{171}, \eta_{172}, \eta_{173}, \eta_{174}, \eta_{175}, \eta_{176}$ in the expression glossary.

$U = \eta_{177}\, C$. Scalar metadata; no physical GP.

clifford_only

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cx

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

clifford_t_cz

Exact: Proved obstruction: discrete alphabet nonconstant family
Projective: Proved obstruction: discrete alphabet nonconstant family

cx_h_phase

EXACT PHASE E=3 / L=20 / score ≈ 53.020 — E upper bound




PROJECTIVE E=3 / L=12 / score ≈ 36.612 — E upper bound


$U = e^{-\frac{i\pi(a-b+c)}{2}}\, C$. Scalar metadata; no physical GP.

cx_rz_ry

EXACT PHASE E=4 / L=17 / score ≈ 52.834 — E upper bound




PROJECTIVE E=3 / L=11 / score ≈ 36.222 — E upper bound




$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

cx_u

EXACT PHASE E=3 / L=7 / score ≈ 28.242 — E upper bound



PROJECTIVE E=3 / L=5 / score ≈ 24.230 — E upper bound




$U = e^{-\frac{i\pi c}{2}}\, C$. Scalar metadata; no physical GP.

cz_rz_rx

EXACT PHASE E=4 / L=28 / score ≈ 77.256 — E upper bound




PROJECTIVE E=3 / L=11 / score ≈ 35.822 — E upper bound




$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

cz_u

EXACT PHASE E=3 / L=10 / score ≈ 35.060 — E upper bound




PROJECTIVE E=3 / L=8 / score ≈ 30.648 — E upper bound




$U = e^{-\frac{i\pi c}{2}}\, C$. Scalar metadata; no physical GP.

google_cz

EXACT PHASE E=3 / L=10 / score ≈ 33.660 — E upper bound



PROJECTIVE E=3 / L=8 / score ≈ 29.448 — E upper bound




$U = e^{\frac{i\pi(a-b+c)}{2}}\, C$. Scalar metadata; no physical GP.

google_fsim

EXACT PHASE E=2 / L=6 / score ≈ 20.644 — E upper bound



PROJECTIVE E=2 / L=4 / score ≈ 16.432 — E upper bound



$U = e^{-\frac{i\pi c}{2}}\, C$. Scalar metadata; no physical GP.

google_sqrt_iswap

EXACT PHASE E=6 / L=19 / score ≈ 67.514 — E upper bound



PROJECTIVE E=6 / L=17 / score ≈ 62.902 — E upper bound

$U = \eta_{178}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

google_syc

EXACT PHASE E=18 / L=19 / score ≈ 113.914 — E upper bound



PROJECTIVE E=18 / L=17 / score ≈ 109.702 — E upper bound

$U = e^{\frac{i\pi(-a-b-c-1)}{2}}\, C$. Scalar metadata; no physical GP.

ibm_cx

EXACT PHASE E=3 / L=16 / score ≈ 45.216 — E upper bound



PROJECTIVE E=3 / L=13 / score ≈ 39.012 — E upper bound

$U = -\sqrt[4]{-1}\, C$. Scalar metadata; no physical GP.

ibm_cz

EXACT PHASE E=3 / L=20 / score ≈ 53.620 — E upper bound

ibm_ecr

EXACT PHASE E=3 / L=24 / score ≈ 61.018 — E upper bound

PROJECTIVE E=3 / L=22 / score ≈ 57.214 — E upper bound

$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

ionq_aria

EXACT PHASE E=3 / L=27 / score ≈ 73.472 — E upper bound

PROJECTIVE E=3 / L=27 / score ≈ 73.272 — E upper bound

$U = \frac{\sqrt[4]{-1}\sqrt{2}(-1+i)}{2}\, C$. Scalar metadata; no physical GP.

ionq_aria_full

EXACT PHASE E=3 / L=27 / score ≈ 72.266 — E upper bound

PROJECTIVE E=3 / L=27 / score ≈ 72.066 — E upper bound

$U = \frac{\sqrt[4]{-1}\sqrt{2}(-1+i)}{2}\, C$. Scalar metadata; no physical GP.

ionq_forte

EXACT PHASE E=3 / L=30 / score ≈ 80.466 — E upper bound

iqm_cz

EXACT PHASE E=4 / L=27 / score ≈ 75.708 — E upper bound

PROJECTIVE E=3 / L=13 / score ≈ 40.452 — E upper bound

$U = -(-1)^{\frac{3}{4}}\, C$. Scalar metadata; no physical GP.

iswap_u

EXACT PHASE E=3 / L=12 / score ≈ 40.072 — E upper bound

PROJECTIVE E=3 / L=10 / score ≈ 35.260 — E upper bound

$U = \eta_{179}\, C$. Scalar metadata; no physical GP.

quantinuum_zzmax

EXACT PHASE E=4 / L=25 / score ≈ 70.876 — E upper bound

PROJECTIVE E=3 / L=10 / score ≈ 33.632 — E upper bound

$U = -i\, C$. Scalar metadata; no physical GP.

quantinuum_zzphase

EXACT PHASE E=3 / L=18 / score ≈ 52.058 — E upper bound

rigetti_cz

EXACT PHASE E=3 / L=20 / score ≈ 53.620 — E upper bound

sqrt_iswap_u

EXACT PHASE E=6 / L=19 / score ≈ 68.314 — E upper bound

PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound

$U = \eta_{180}\, C$. Scalar metadata; no physical GP.
Published uniform cap: $E \leq 3$. Current construction exceeds it; optimization remains.

sqrt_swap_u

EXACT PHASE E=6 / L=19 / score ≈ 69.114 — E upper bound

PROJECTIVE E=6 / L=17 / score ≈ 63.902 — E upper bound

$U = -\sqrt[4]{-1}e^{-\frac{i\pi c}{2}}\, C$. Scalar metadata; no physical GP.

tket_tk2

EXACT PHASE E=1 / L=0 / score ≈ 4.006 — E upper bound

xx90_u

EXACT PHASE E=3 / L=10 / score ≈ 34.260 — E upper bound

PROJECTIVE E=3 / L=8 / score ≈ 29.448 — E upper bound

$U = ie^{-\frac{i\pi b}{2}}\, C$. Scalar metadata; no physical GP.

xx_rz_ry

EXACT PHASE E=3 / L=41 / score ≈ 98.888 — E upper bound

PROJECTIVE E=3 / L=31 / score ≈ 80.068 — E upper bound

$U = \eta_{36}\, C$. Scalar metadata; no physical GP.

zx_u

EXACT PHASE E=3 / L=13 / score ≈ 43.684 — E upper bound

PROJECTIVE E=3 / L=11 / score ≈ 39.272 — E upper bound

$U = \eta_{181}\, C$. Scalar metadata; no physical GP.

zz_u

EXACT PHASE E=3 / L=12 / score ≈ 38.478 — E upper bound

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{1} = -2\pi + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)}\right)$$

$$\eta_{2} = -\frac{\pi}{32} + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right)$$

$$\eta_{3} = \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right) + \frac{31\pi}{32}$$

$$\eta_{4} = -2i\log\left(\frac{\frac{\sqrt{2}}{2} + \frac{\sqrt{2}i\sqrt{\frac{(-1)^{\frac{1}{16}}\left(\sin\left(\frac{\pi}{16}\right) - i\cos\left(\frac{\pi}{16}\right)\right)}{2}}}{2}}{\sqrt{\frac{1}{2} + \frac{(-1)^{\frac{1}{16}}\left(\sin\left(\frac{\pi}{16}\right) - i\cos\left(\frac{\pi}{16}\right)\right)}{2}}}\right)$$

$$\eta_{5} = \operatorname{atan}\left(\frac{\sin\left(\frac{7\pi}{16}\right)}{\cos\left(\frac{7\pi}{16}\right)}\right)$$

$$\eta_{6} = -2i\log\left(\frac{\frac{\sqrt{\frac{-(-1)^{\frac{1}{16}}\left(1+e^{\frac{i\pi}{16}}\right)^{2}}{2}} + \frac{i\sqrt{\frac{(-1)^{\frac{1}{16}}\left(1-1^{\frac{1}{8}}(-1)\right)\left(1-e^{\frac{i\pi}{16}}\right)}{2}}}{2}}{\sqrt{\frac{(-1)^{\frac{1}{16}}\left(1-1^{\frac{1}{8}}(-1)\right)\left(1-e^{\frac{i\pi}{16}}\right)}{4} - \frac{(-1)^{\frac{1}{16}}\left(1+e^{\frac{i\pi}{16}}\right)^{2}}{4}}}\right)$$

$$\eta_{7} = -\operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{1-\cos\left(\frac{\pi}{16}\right)}\right) - \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right) + \pi$$

$$\eta_{8} = -2i\log\left(\frac{\frac{\sqrt{2}}{2} + \frac{\sqrt{2}i\sqrt{\frac{1^{\frac{1}{8}}(-1+e^{-\frac{i\pi}{16}})}{2}}}{2}}{\sqrt{\frac{1}{2} + \frac{1^{\frac{1}{8}}(-1+e^{-\frac{i\pi}{16}})}{2}}}\right)$$

$$\eta_{9} = \operatorname{atan}\left(\frac{-\sqrt{2-\sqrt{2}}}{-2+\sqrt{\sqrt{2}+2}}\right) + \pi$$

$$\eta_{10} = -2i\log\left(\frac{\frac{\sqrt{\left(1-(-1)^{\frac{1}{16}}\right)\left(1+e^{\frac{i\pi}{16}}\right)}}{2} + \frac{i\sqrt{\left(1+(-1)^{\frac{1}{16}}\right)\left(1-e^{\frac{i\pi}{16}}\right)}}{2}}{\sqrt{\frac{\left(1+(-1)^{\frac{1}{16}}\right)\left(1-e^{\frac{i\pi}{16}}\right)}{4} + \frac{\left(1-(-1)^{\frac{1}{16}}\right)\left(1+e^{\frac{i\pi}{16}}\right)}{4}}}\right)$$

$$\eta_{11} = -\pi + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right) + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{1-\cos\left(\frac{\pi}{16}\right)}\right)$$

$$\eta_{12} = -2i\log\left(\frac{\frac{\sqrt{\frac{-(-1)^{\frac{1}{16}}\left(1+e^{\frac{i\pi}{16}}\right)^{2}}{2}} + \frac{i\sqrt{\frac{\left(1-1^{\frac{1}{8}}(-1)\right)\left(1-e^{-\frac{i\pi}{16}}\right)}{2}}}{2}}{\sqrt{\frac{\left(1-1^{\frac{1}{8}}(-1)\right)\left(1-e^{-\frac{i\pi}{16}}\right)}{4} - \frac{(-1)^{\frac{1}{16}}\left(1+e^{\frac{i\pi}{16}}\right)^{2}}{4}}}\right)$$

$$\eta_{13} = -\operatorname{atan}\left(\frac{\sin\left(\frac{7\pi}{16}\right)}{1-\cos\left(\frac{7\pi}{16}\right)}\right) - \operatorname{atan}\left(\frac{\sin\left(\frac{7\pi}{16}\right)}{\cos\left(\frac{7\pi}{16}\right)+1}\right)$$

$$\eta_{14} = -\pi + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)}\right)$$

$$\eta_{15} = e^{i\left(-\frac{\pi}{32} + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right)\right)}$$

$$\eta_{16} = -\pi - \operatorname{acos}\left(\frac{\sqrt{3}}{3}\right)$$

$$\eta_{17} = -\pi + \operatorname{acos}\left(\frac{\sqrt{3}}{3}\right)$$

$$\eta_{18} = -\frac{\pi}{2} + \operatorname{acos}\left(\frac{\sqrt{3}}{3}\right)$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{19} = -\frac{\pi}{2} - \operatorname{acos}\left(\frac{\sqrt{3}}{3}\right)$$

$$\eta_{20} = -\frac{37\pi}{32} + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right)$$

$$\eta_{21} = -\frac{5\pi}{32} + \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right)$$

$$\eta_{22} = 2\operatorname{atan}\left(\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right)$$

$$\eta_{23} = -2i\log\left(\frac{\frac{\sqrt{2}}{2}+\frac{\sqrt[8]{2}i\sqrt{\sqrt[4]{-1}(1-i)}}{2}}{\sqrt{\frac{1}{2}+\frac{\sqrt[8]{-1}\sqrt{2}(1-i)}{4}}}\right)$$

$$\eta_{24} = -2i\log\left(\frac{\frac{\sqrt{2}\sqrt{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}}{4}+\frac{\sqrt{2}i\sqrt{\left(-i-(-1)^{\frac{3}{4}}\right)\left(\sqrt{2}+\sqrt{2}i+2i\right)}}{4}}{\sqrt{\frac{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}{8}-\frac{\left(-1\right)^{\frac{3}{4}}+i\right)\left(\sqrt{2}+\sqrt{2}i+2i\right)}{8}}}\right)$$

$$\eta_{25} = -2i\log\left(\frac{\frac{\sqrt{2}\sqrt{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}}{4}+\frac{\sqrt{2}i\sqrt{\left(1-(-1)^{\frac{3}{4}}\right)\left(\sqrt{2}+2+\sqrt{2}i\right)}}{4}}{\sqrt{\frac{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}{8}+\frac{\left(1-(-1)^{\frac{3}{4}}\right)\left(\sqrt{2}+2+\sqrt{2}i\right)}{8}}}\right)$$

$$\eta_{26} = -2i\log\left(\frac{\frac{\sqrt{-(-1)^{\frac{3}{4}}\left(1+e^{\frac{i\pi}{4}}\right)^2}}{2}+\frac{i\sqrt{(-1)^{\frac{3}{4}}\left(1-\sqrt[4]{-1}\right)\left(1-e^{\frac{i\pi}{4}}\right)}}{2}}{\sqrt{\frac{(-1)^{\frac{3}{4}}\left(1-\sqrt[4]{-1}\right)\left(1-e^{\frac{i\pi}{4}}\right)}{4}-\frac{(-1)^{\frac{3}{4}}\left(1+e^{\frac{i\pi}{4}}\right)^2}{4}}}\right)$$

$$\eta_{27} = -2i\log\left(\frac{\frac{\sqrt{2}\sqrt{\left(1-(-1)^{\frac{3}{4}}\right)\left(\sqrt{2}+2+\sqrt{2}i\right)}}{4}+\frac{\sqrt{2}i\sqrt{\left(-1-(-1)^{\frac{3}{4}}\right)\left(-2+\sqrt{2}+\sqrt{2}i\right)}}{4}}{\sqrt{\frac{\left(-1-(-1)^{\frac{3}{4}}\right)\left(-2+\sqrt{2}+\sqrt{2}i\right)}{8}+\frac{\left(1-(-1)^{\frac{3}{4}}\right)\left(\sqrt{2}+2+\sqrt{2}i\right)}{8}}}\right)$$

$$\eta_{28} = -2i\log\left(\frac{\frac{\sqrt{2}\sqrt{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}}{4}+\frac{\sqrt{2}i\sqrt{\left(1+\sqrt[4]{-1}\right)\left(\sqrt{2}+2-\sqrt{2}i\right)}}{4}}{\sqrt{\frac{\left(1-\sqrt[4]{-1}\right)\left(-\sqrt{2}+2+\sqrt{2}i\right)}{8}+\frac{\left(1+\sqrt[4]{-1}\right)\left(\sqrt{2}+2-\sqrt{2}i\right)}{8}}}\right)$$

$$\eta_{29} = e^{i\left(-\frac{37\pi}{32}+\operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)+1}\right)\right)}$$

$$\eta_{30} = -\operatorname{atan}\left(\frac{\cos\left(\frac{\pi}{16}\right)}{\sin\left(\frac{\pi}{16}\right)}\right)$$

$$\eta_{31} = \operatorname{atan}\left(\frac{-2+\sqrt{\sqrt{2}+2}}{\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{32} = \operatorname{atan}\left(\frac{\sin\left(\frac{\pi}{16}\right)}{\cos\left(\frac{\pi}{16}\right)}\right)$$

$$\eta_{33} = \operatorname{atan}\left(\frac{2-\sqrt{\sqrt{2}+2}}{\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{34} = -\operatorname{atan}\left(\frac{2-\sqrt{\sqrt{2}+2}}{\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{35} = -\pi + \operatorname{atan}\left(\frac{2-\sqrt{\sqrt{2}+2}}{\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{36} = \frac{(-1)^{\frac{3}{4}}\sqrt{2}(1+i)^3}{4}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{37} = \frac{\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor+\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)+2\sin^2\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)}=0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sqrt{2}\sin\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}$$

$$\eta_{38} = \frac{\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor+\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)+2\sin^2\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)}=0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sqrt{2}\sin\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} + \frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \left(\frac{3\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right)+3\pi$$

$$\eta_{39} = -\frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \pi$$

$$\eta_{40} = -2\operatorname{atan_2}\left(\sqrt{2}\sin\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right),\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)+2\sin^2\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)}\right)$$

$$\eta_{41} = -\begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)+2\sin^2\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)}=0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sqrt{2}\sin\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}$$

$$\eta_{42} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{43} = \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{44} = -\frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) - 2\pi$$

$$\eta_{45} = \left((1-\left(2\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)\left(\frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \pi\right) + \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{46} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{47} = \left(\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \frac{\pi}{2}$$

$$\eta_{48} = -\frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \pi$$

$$\eta_{49} = e^{i\left(\frac{\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor+\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)+2\sin^2\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)}=0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sqrt{2}\sin\left(b-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(c-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} + \frac{\begin{cases} 0 & \text{for } \cos\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(b+c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}-\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)=0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}\cos(2b)\cos(2c)},\sin\left(b-c-\frac{\pi\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor}{2}+\frac{\pi\left\lfloor \frac{2c}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \left(\frac{3\pi\left(\left\lfloor \frac{2b}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right)+2\pi\right)}$$

$$\eta_{50} = \frac{\sqrt{2}(-1+i)e^{i(a+b+c)}}{2}$$

$$\eta_{51} = \sin(a+b+c) - i\cos(a+b+c)$$

$$\eta_{52} = \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)+\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}=0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi),-\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2}+\frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2+\sin^2\left(\frac{\gamma}{2}\right)}=0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2}+\frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2},-\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{5\pi}{4}$$

$$\eta_{53} = -\frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)+\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}=0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi),-\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2}+\frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2+\sin^2\left(\frac{\gamma}{2}\right)}=0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2}+\frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2},-\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\pi}{4}$$

$$\eta_{54} = -\operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)+\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)},\sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2}+\frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2+\sin^2\left(\frac{\gamma}{2}\right)}\right)$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{55} = \frac{\gamma}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\pi}{4}$$

$$\eta_{56} = \frac{\gamma}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\pi}{4}$$

$$\eta_{57} = \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} + \frac{3\pi}{4}$$

$$\eta_{58} = \frac{-\frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} - \frac{3\pi}{4} + \frac{1}{2}$$

$$\eta_{59} = \frac{-\frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} - \frac{\gamma}{4}$$

$$\eta_{60} = \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} + \frac{\gamma}{4} + \frac{1}{2}$$

$$\eta_{61} = \frac{\pi - \operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right)}{\pi}$$

$$\eta_{62} = \frac{\operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right) - \pi}{\pi} + \frac{1}{2}$$

$$\eta_{63} = \frac{\operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right)}{\pi}$$

$$\eta_{64} = -\frac{\operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right)}{\pi} + \frac{1}{2}$$

$$\eta_{65} = \frac{\frac{\gamma}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{3\pi}{4}$$

$$\eta_{66} = \frac{-\frac{\gamma}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} - \frac{3\pi}{4} + \frac{1}{2}$$

$$\eta_{67} = \frac{\frac{\gamma}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{3\pi}{4}$$

$$\eta_{68} = \frac{-\frac{\gamma}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2}}{\pi} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{3\pi}{4} + \frac{1}{2}$$

$$\eta_{69} = \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{3\pi}{4}$$

$$\eta_{70} = -\frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\pi}{4}$$

$$\eta_{71} = \pi - \operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right)$$

$$\eta_{72} = \operatorname{atan_2}\left(\sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)}, \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)}\right)$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{73} = \frac{\gamma}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{3\pi}{4}$$

$$\eta_{74} = \frac{\gamma}{2} + \frac{\begin{cases} 0 & \text{for } \sqrt{\sin^2(\phi)\sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right) + \sin^2(\theta)\cos^2\left(\frac{\gamma}{2}\right)\cos^2(\phi)} = 0 \\ \operatorname{atan_2}\left(-\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\cos(\phi), -\sin(\phi)\sin(\theta)\cos\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \sqrt{\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}\right)^2 + \sin^2\left(\frac{\gamma}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\frac{\cos\left(\frac{\gamma}{2}-\theta\right)}{2} + \frac{\cos\left(\frac{\gamma}{2}+\theta\right)}{2}, -\sin\left(\frac{\gamma}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{3\pi}{4}$$

$$\eta_{75} = \frac{\sqrt{2}i(1+i)^3 e^{\frac{i(\pi-\phi)}{4}}}{4}$$

$$\eta_{76} = \frac{\phi}{4} + \frac{\pi\left(\left\lfloor -\frac{\phi}{2\pi} + \frac{1}{2}\right\rfloor + \left\lfloor -\frac{\theta}{\pi} + \frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta) + 2\sin^2\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sqrt{2}\sin\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} - \left(\frac{3\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + \pi$$

$$\eta_{77} = \frac{\phi}{4} + \frac{\pi\left(\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor + \left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta) + 2\sin^2\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sqrt{2}\sin\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} + \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \left(\frac{3\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + 3\pi$$

$$\eta_{78} = -\frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \pi$$

$$\eta_{79} = -2\operatorname{atan_2}\left(-\sqrt{2}\sin\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right), \sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta) + 2\sin^2\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)}\right)$$

$$\eta_{80} = -\begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta) + 2\sin^2\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sqrt{2}\sin\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}$$

$$\eta_{81} = \left(-\frac{\pi\left(1 - \left(2\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{82} = -\frac{\phi}{2} + \left(\pi\left(\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{83} = -\frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) - 2\pi$$

$$\eta_{84} = \left(\left(1 - \left(2\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)\left(\frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \pi\right)\right) + \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{85} = -\frac{\phi}{2} - \left(\frac{\pi\left(1 - \left(2\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{86} = \left(\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right) + \frac{\pi}{2}$$

$$\eta_{87} = -\frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \pi$$

$$\eta_{88} = \epsilon\left(\frac{\phi}{2} + \frac{\pi\left(\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor + \left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor\right)}{2} + \begin{cases} 0 & \text{for } \sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta) + 2\sin^2\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sqrt{2}\sin\left(\frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\phi}{4} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} + \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, -\sin\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} + \frac{\begin{cases} 0 & \text{for } \cos\left(\frac{\phi}{4} + \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} + \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\sqrt{(-1)^{\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}(-1)^{\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}\cos\left(\frac{\phi}{2}\right)\cos(\theta)}, \sin\left(\frac{\phi}{4} - \frac{\theta}{2} + \frac{\pi\left\lfloor -\frac{\phi}{2\pi}+\frac{1}{2}\right\rfloor}{2} - \frac{\pi\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}}{2} - \left(\frac{2\pi\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + 2\pi\right)$$

$$\eta_{89} = \frac{\sqrt{2}(1+i)^3 e^{\frac{i(\pi-\phi)}{4}}}{4}$$

$$\eta_{90} = \frac{\sqrt{2}i(1+i)e^{\frac{i(\phi-2\theta)}{4}}}{2}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{91} = \operatorname{atan}\left(\frac{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}\right) + \operatorname{atan}\left(\frac{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right)$$

$$\eta_{92} = -2i\log\left(\frac{\frac{\sqrt{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)}}{2}+\frac{i\sqrt{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}}{2}}{\sqrt{\frac{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}{4}}+\frac{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)}{4}}}\right)$$

$$\eta_{93} = \operatorname{atan}\left(\frac{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right) - \operatorname{atan}\left(\frac{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{94} = -\pi - \operatorname{atan}\left(\frac{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right) - \operatorname{atan}\left(\frac{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{95} = -2i\log\left(\frac{\frac{\sqrt{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}}{2}+\frac{i\sqrt{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)}}{2}}{\sqrt{\frac{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}{4}}+\frac{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)}{4}}}\right)$$

$$\eta_{96} = \operatorname{atan}\left(\frac{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}\right) - \operatorname{atan}\left(\frac{-\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right) + \pi$$

$$\eta_{97} = -2i\log\left(\frac{\frac{\sqrt{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}}{2}+\frac{i\sqrt{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-\sqrt{-2-\sqrt{2}}\right)}}{2}}{\sqrt{\frac{\sqrt[8]{-1}\left(\sqrt{\sqrt{2}+2}-i\sqrt{2-\sqrt{2}}\right)}{4}}+\frac{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-\sqrt{-2-\sqrt{2}}\right)}{4}}}\right)$$

$$\eta_{98} = -\pi + 2\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}}{\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{99} = -\pi - \operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}+(1+\sqrt{2})\sqrt{2-\sqrt{2}}}{-\sqrt{2-\sqrt{2}}+(1+\sqrt{2})\sqrt{\sqrt{2}+2}}\right) - \operatorname{atan}\left(\frac{-\sqrt{2-\sqrt{2}}+(1-\sqrt{2})\sqrt{\sqrt{2}+2}}{(1-\sqrt{2})\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right)$$

$$\eta_{100} = -2i\log\left(\frac{\frac{\sqrt{2}\sqrt{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-\sqrt{-2-\sqrt{2}}\right)}\left((-1)^{\frac{3}{4}}(1+i)+1-\right)\left(-\sqrt{2}i(1-i)(1+i)+2+2i\right)}}{8}+\frac{\sqrt{2}i\sqrt{-(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)(2+\sqrt{2}(1-i)(1+i)+2i)\left((-1)^{\frac{3}{4}}(1+i)-1+\right)}}{8}}{\sqrt{\frac{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-\sqrt{-2-\sqrt{2}}\right)\left((-1)^{\frac{3}{4}}(1+i)+1-i\right)\left(-\sqrt{2}i(1-i)(1+i)+2+2i\right)}{32}}-\frac{(-1)^{\frac{3}{8}}\left(\sqrt{2-\sqrt{2}}-i\sqrt{\sqrt{2}+2}\right)(2+\sqrt{2}i(1-i)(1+i)+2i)\left((-1)^{\frac{3}{4}}(1+i)-1+\right)}{32}}}\right)$$

$$\eta_{101} = -\pi - \operatorname{atan}\left(\frac{-\sqrt{2-\sqrt{2}}+(1-\sqrt{2})\sqrt{\sqrt{2}+2}}{(1-\sqrt{2})\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right) + \operatorname{atan}\left(\frac{-\sqrt{2-\sqrt{2}}+(1+\sqrt{2})\sqrt{\sqrt{2}+2}}{\sqrt{\sqrt{2}+2}+(1+\sqrt{2})\sqrt{2-\sqrt{2}}}\right)$$

$$\eta_{102} = e^{i\left(-\frac{\operatorname{atan}\left(\frac{\sqrt{\sqrt{2}+2}+(1+\sqrt{2})\sqrt{2-\sqrt{2}}}{-\sqrt{2-\sqrt{2}}+(1+\sqrt{2})\sqrt{\sqrt{2}+2}}\right)}{2}+\frac{\operatorname{atan}\left(\frac{-\sqrt{2-\sqrt{2}}+(1+\sqrt{2})\sqrt{\sqrt{2}+2}}{\sqrt{\sqrt{2}+2}+(1+\sqrt{2})\sqrt{2-\sqrt{2}}}\right)}{2}+\frac{7\pi}{8}\right)}$$

$$\eta_{103} = 2\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}\right)$$

$$\eta_{104} = \frac{\sqrt{2}i(1+i)^{\frac{3}{4}}e^{\frac{i(-4\gamma-\phi+\pi)}{4}}}{4}$$

$$\eta_{105} = -\tfrac{\lambda}{2} - \gamma - \tfrac{\phi}{2} + \tfrac{\zeta}{2} + \left(\pi\left(\left\lfloor -\tfrac{\phi}{2\pi}+\tfrac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor -\tfrac{\theta}{\pi}+\tfrac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{106} = \tfrac{\lambda}{2} - \gamma - \tfrac{\phi}{2} - \tfrac{\zeta}{2} - \left(\tfrac{\pi\left(1-\left(2\left(\left\lfloor -\frac{\theta}{\pi}+\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor -\tfrac{\phi}{2\pi}+\tfrac{1}{2}\right\rfloor \bmod 2\right)\right) + \left(\pi\left(\left\lfloor -\tfrac{\theta}{\pi}+\tfrac{1}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{107} = -(-1)^{\frac{3}{4}}e^{-\frac{i(4\gamma+\phi)}{4}}$$

$$\eta_{108} = \tfrac{\lambda}{2} - \gamma - \tfrac{\phi}{2} - \tfrac{\zeta}{2} - \tfrac{\pi i}{2}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{109} = \frac{\frac{\chi}{2} - \gamma - \frac{\phi}{2} - \frac{\zeta}{2}}{\pi}$$

$$\eta_{110} = \frac{-\frac{\chi}{2} - \gamma - \frac{\phi}{2} + \frac{\zeta}{2}}{\pi}$$

$$\eta_{111} = \frac{\chi}{2} - \gamma - \frac{\phi}{2} - \frac{\zeta}{2} - \frac{3\pi}{2}$$

$$\eta_{112} = \frac{\sqrt{2}(1+i)^2 e^{\frac{i(-\zeta_{\mathrm{shift}}-\phi+\pi)}{4}}}{4}$$

$$\eta_{113} = \frac{\sqrt{2}i(1+i)^2 e^{\frac{i\pi\left(t\left(4\mathrm{global}_{\mathrm{shift}}+1\right)+1\right)}{4}}}{4}$$

$$\eta_{114} = \frac{\pi t\left(4\mathrm{global}_{\mathrm{shift}}+1\right)}{4} + \begin{cases} 0 & \text{for } \sqrt{2\sin^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)+\cos^2\left(\frac{\pi t}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\left|\cos\left(\frac{\pi t}{2}\right)\right|, -\sqrt{2}\sin\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} - \left(\frac{3\pi\left(\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + \pi\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor + \pi$$

$$\eta_{115} = \frac{\pi t\left(4\mathrm{global}_{\mathrm{shift}}+1\right)}{4} + \frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} + \begin{cases} 0 & \text{for } \sqrt{2\sin^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)+\cos^2\left(\frac{\pi t}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\left|\cos\left(\frac{\pi t}{2}\right)\right|, -\sqrt{2}\sin\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} - \left(\frac{3\pi\left(\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} + \pi\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor + 3\pi$$

$$\eta_{116} = \frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} + \pi$$

$$\eta_{117} = -2\operatorname{atan_2}\left(-\sqrt{2}\sin\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right), \sqrt{2\sin^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)+\cos^2\left(\frac{\pi t}{2}\right)}\right)$$

$$\eta_{118} = -\begin{cases} 0 & \text{for } \sqrt{2\sin^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)+\cos^2\left(\frac{\pi t}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\left|\cos\left(\frac{\pi t}{2}\right)\right|, -\sqrt{2}\sin\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases}$$

$$\eta_{119} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{120} = 2\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)$$

$$\eta_{121} = -\frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} + \left(\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right) - \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} - 2\pi$$

$$\eta_{122} = \left(\left(1-\left(2\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right)\right)\left(-\frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} + \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} + \pi\right)\right) + \left(\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{123} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor \bmod 2\right)\right)\right)}{2}\right) + \left(2\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right)$$

$$\eta_{124} = \left(\pi\left(\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor \bmod 2\right)\right) + \frac{\pi}{2}$$

$$\eta_{125} = -\frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} - \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} - \pi$$

$$\eta_{126} = e^{i\left(\frac{\pi t\left(4\mathrm{global}_{\mathrm{shift}}+1\right)}{4} + \frac{\begin{cases} 0 & \text{for } (-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\cos\left(\frac{\pi t}{2}\right) = 0 \\ \operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, 0\right) & \text{otherwise} \end{cases}}{2} + \begin{cases} 0 & \text{for } \sqrt{2\sin^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos^2\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)+\cos^2\left(\frac{\pi t}{2}\right)} = 0 \\ \operatorname{atan_2}\left(\left|\cos\left(\frac{\pi t}{2}\right)\right|, -\sqrt{2}\sin\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\cos\left(\frac{\pi t}{4}+\frac{\pi\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}{2}\right)\right) & \text{otherwise} \end{cases} - \left(\frac{3\pi\left(\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor \bmod 2\right)}{2}\right) + \frac{\operatorname{atan_2}\left(-\left|\cos\left(\frac{\pi t}{2}\right)\right|, -(-1)^{\left\lfloor\frac{\chi}{2\pi}-\frac{1}{2}\right\rfloor}\sin\left(\frac{\pi t}{2}\right)\right)}{2} + \pi\left\lfloor\frac{1}{2}-\frac{\phi}{2}\right\rfloor + 2\pi\right)}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{127} = -e^{\frac{i\pi\left(t\left(4g\mathrm{lobal}_{shift}+1\right)+1\right)}{4}}$$

$$\eta_{128} = -\left(-1\right)^{\frac{3}{4}} e^{\frac{i\pi t\left(4g\mathrm{lobal}_{shift}+1\right)}{4}}$$

$$\eta_{129} = -\frac{t(2g\mathrm{lobal}_{shift}-1)}{2} - \frac{3}{4}$$

$$\eta_{130} = \frac{\sqrt{2}(-1+i)e^{\frac{i\pi t\left(g\mathrm{lobal}_{shift}-\frac{1}{2}\right)}{2}}}{2}$$

$$\eta_{131} = -\frac{\frac{\pi t\left(2g\mathrm{lobal}_{shift}-1\right)}{2}}{\pi} - \pi$$

$$\eta_{132} = e^{\frac{i\pi\left(t\left(4g\mathrm{lobal}_{shift}+1\right)-3t-2\right)}{4}}$$

$$\eta_{133} = -ie^{\frac{i\pi\left(t\left(4g\mathrm{lobal}_{shift}+1\right)+1\right)}{4}}$$

$$\eta_{134} = \frac{\sqrt{2}(1+i)^2 e^{\frac{i\pi\left(t\left(4g\mathrm{lobal}_{shift}+1\right)+1\right)}{4}}}{4}$$

$$\eta_{135} = \frac{\sqrt{2}i(1+i)e^{ig\mathrm{lobal}_{shift}t}}{2}$$

$$\eta_{136} = \frac{\sqrt[4]{-1}\sqrt{2}(-1+i)e^{ig\mathrm{lobal}_{shift}t}}{2}$$

$$\eta_{137} = \frac{\sqrt[4]{-1}\sqrt{2}(1-i)e^{ig\mathrm{lobal}_{shift}t}}{2}$$

$$\eta_{138} = \frac{(-1)^{\frac{3}{4}}\sqrt{2}(1-i)}{2}$$

$$\eta_{139} = -\pi + \operatorname{atan}\left(\frac{-\sqrt{3\sqrt{2}+6}+\sqrt{2-\sqrt{2}}+\sqrt{6-3\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)$$

$$\eta_{140} = \operatorname{atan}\left(\frac{-\sqrt{3\sqrt{2}+6}+\sqrt{2-\sqrt{2}}+\sqrt{6-3\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right) + \pi$$

$$\eta_{141} = \pi - 2\operatorname{atan}\left(\frac{-\sqrt{3}\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)$$

$$\eta_{142} = \frac{\operatorname{atan}\left(\frac{2\sqrt{\sqrt{2}+6}}{-\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{2-\sqrt{2}}}\right)+\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+6}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)+\frac{7\pi}{12}-2\operatorname{atan}\left(\frac{2\sqrt[4]{3}\sqrt{1+\sqrt{3}}}{-\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{2-\sqrt{2}}}\right)}{\pi}$$

$$\eta_{143} = \frac{-\frac{13\pi}{24}+\operatorname{atan}\left(\frac{2\sqrt[4]{3}\sqrt{1+\sqrt{3}}}{-\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{2-\sqrt{2}}}\right)-\frac{\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+\sqrt{6}}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)}{2}-\frac{\operatorname{atan}\left(\frac{2\sqrt{\sqrt{3}+\sqrt{6}}}{-\sqrt{\sqrt{3}+2}+\sqrt{3}\sqrt{2-\sqrt{3}}}\right)}{2}}{\pi}+\frac{1}{2}$$

$$\eta_{144} = \frac{-\frac{\pi}{3}-\frac{\operatorname{atan}\left(\frac{\sqrt{\sqrt{3}+\frac{\sqrt{6}}{2}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{3}}{4}}}{2}+\sqrt{\frac{\sqrt{3}}{2}+\frac{1}{2}}}\right)}{\pi}-\frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{3}}{2}+\frac{\sqrt{6}}{2}}}{\sqrt{\frac{1}{2}-\frac{\sqrt{3}}{4}}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{3}}{2}+\frac{1}{2}}}{2}}\right)}{\pi}}{\pi}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{145} = \frac{1}{2} - \frac{-\frac{\pi}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} - \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi}$$

$$\eta_{146} = \frac{-\frac{\pi}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi}$$

$$\eta_{147} = \frac{-\frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} + \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} + \frac{\pi}{2}}{\pi} + \frac{1}{2}$$

$$\eta_{148} = \frac{-\operatorname{atan}\left(\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right) - \operatorname{atan}\left(\frac{\sqrt{3}\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right)}{\pi}$$

$$\eta_{149} = \frac{-\operatorname{atan}\left(\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right) + \operatorname{atan}\left(\frac{\sqrt{3}\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right)}{\pi}$$

$$\eta_{150} = \frac{\frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi}}{\pi}$$

$$\eta_{151} = \frac{-\frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{\pi}}{\pi}$$

$$\eta_{152} = e^{i\left(\frac{\operatorname{atan}\left(\frac{2\sqrt{\sqrt{2}+\sqrt{6}}}{-\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{2-\sqrt{2}}}\right)}{2} + \frac{\pi}{24} + \frac{\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+\sqrt{6}}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)}{2} - \operatorname{atan}\left(\frac{2\sqrt[4]{3}\sqrt{1+\sqrt{3}}}{-\sqrt{\sqrt{2}+2}+\sqrt{3}\sqrt{2-\sqrt{2}}}\right)\right)}$$

$$\eta_{153} = \frac{(-1)^{\frac{3}{24}}\sqrt{2}(1+i)}{2}$$

$$\eta_{154} = \frac{\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+\sqrt{6}}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)}{2} + \frac{13\pi}{24}$$

$$\eta_{155} = \frac{\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+\sqrt{6}}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)}{2} + \frac{25\pi}{24}$$

$$\eta_{156} = -\frac{\pi}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2}$$

$$\eta_{157} = -\frac{\pi}{2} - \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2}$$

$$\eta_{158} = -\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right) - \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right)$$

$$\eta_{159} = -\operatorname{atan}\left(\frac{\sqrt{2}\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right) + \operatorname{atan}\left(\frac{\sqrt{2}\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}\right)$$

$$\eta_{160} = \frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2}$$

$$\eta_{161} = -\frac{\operatorname{atan}\left(\frac{\sqrt{-\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}+\frac{\sqrt{3}\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2} + \frac{\operatorname{atan}\left(\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{\sqrt{6}}{4}}}{-\frac{\sqrt{3}\sqrt{\frac{1}{2}-\frac{\sqrt{2}}{4}}}{2}+\frac{\sqrt{\frac{\sqrt{2}}{4}+\frac{1}{2}}}{2}}\right)}{2}$$

$$\eta_{162} = e^{i\left(\frac{\pi}{24} + \frac{\operatorname{atan}\left(\frac{2\sqrt{-\sqrt{2}+\sqrt{6}}}{-\sqrt{2-\sqrt{2}}+\sqrt{3}\sqrt{\sqrt{2}+2}}\right)}{2}\right)}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{163} = \operatorname{atan}\left(\frac{-\sqrt{3}\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right)$$

$$\eta_{164} = \operatorname{atan}\left(\frac{-\sqrt{3}\sqrt{\sqrt{2}+2}+\sqrt{2-\sqrt{2}}}{\sqrt{3}\sqrt{2-\sqrt{2}}+\sqrt{\sqrt{2}+2}}\right)+\pi$$

$$\eta_{165} = \frac{\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor+\left\lfloor\frac{1}{2}-c\right\rfloor\right)}{2}+\begin{cases}0 & \text{for } \sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)+2\sin^2\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos^2\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)}=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sqrt{2}\sin\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}-\left(\frac{3\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)}{2}\right)+\pi$$

$$\eta_{166} = \frac{\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor+\left\lfloor\frac{1}{2}-c\right\rfloor\right)}{2}+\begin{cases}0 & \text{for } \sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)+2\sin^2\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos^2\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)}=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sqrt{2}\sin\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}+\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\left(\frac{3\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)}{2}\right)+3\pi$$

$$\eta_{167} = -\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\pi$$

$$\eta_{168} = -2\operatorname{atan}_2\left(-\sqrt{2}\sin\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\cos\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right),\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)+2\sin^2\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos^2\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)}\right)$$

$$\eta_{169} = -\begin{cases}0 & \text{for } \sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)+2\sin^2\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos^2\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)}=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sqrt{2}\sin\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}$$

$$\eta_{170} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)\right)}{2}\right)+\left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)$$

$$\eta_{171} = \left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)+\left(\pi\left(\left\lfloor\frac{1}{2}-c\right\rfloor \bmod 2\right)\right)$$

$$\eta_{172} = -\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)-2\pi$$

$$\eta_{173} = \left(\left(1-\left(2\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)\right)\left(\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\pi\right)\right)+\left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)$$

$$\eta_{174} = \left(-\frac{\pi\left(1-\left(2\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)\right)}{2}\right)+\left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)+\left(\pi\left(\left\lfloor\frac{1}{2}-c\right\rfloor \bmod 2\right)\right)$$

$$\eta_{175} = \left(\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)\right)+\frac{\pi}{2}$$

$$\eta_{176} = -\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\pi$$

$$\eta_{177} = e^{i\left(\frac{\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor+\left\lfloor\frac{1}{2}-c\right\rfloor\right)}{2}+\begin{cases}0 & \text{for } \sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)+2\sin^2\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos^2\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)}=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sqrt{2}\sin\left(\frac{\pi b}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}\right)\cos\left(\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}+\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}+\frac{\begin{cases}0 & \text{for } \cos\left(\frac{\pi b}{2}+\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}+\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)=0\\ \operatorname{atan}_2\left(-\sqrt{(-1)^{\left\lfloor\frac{1}{2}-b\right\rfloor}(-1)^{\left\lfloor\frac{1}{2}-c\right\rfloor}\cos(\pi b)\cos(\pi c)},-\sin\left(\frac{\pi b}{2}-\frac{\pi c}{2}+\frac{\pi\left\lfloor\frac{1}{2}-b\right\rfloor}{2}-\frac{\pi\left\lfloor\frac{1}{2}-c\right\rfloor}{2}\right)\right) & \text{otherwise}\end{cases}}{2}-\left(\frac{3\pi\left(\left\lfloor\frac{1}{2}-b\right\rfloor \bmod 2\right)}{2}\right)+2\pi\right)}$$

$$\eta_{178} = \frac{\sqrt{2}(1-i)^3 e^{-\frac{i\pi(a+b+c+1)}{2}}}{4}$$

$$\eta_{179} = \frac{\sqrt{2}i(1+i)e^{-\frac{i\pi b}{2}}}{2}$$

$$\eta_{180} = \frac{\sqrt[4]{-1}\sqrt{2}(-1+i)e^{-\frac{i\pi c}{2}}}{2}$$

Exact expression glossary

Definitions shared by the numbered circuit parameters and scalar factors.

$$\eta_{181} = \frac{\sqrt[4]{-1}\sqrt{2}(1-i)e^{-\frac{i\pi}{2}}}{2}$$