# Paternal Age and De Novo Mutation Rate

*De novo single-nucleotide mutations in offspring increase linearly with paternal age at conception; this establishes a biological germline mutation rate, not an individual psychiatric or cognitive forecast for a child.*

**HIGH-MUTATION-RATE; LOW-FORECAST | STATUS: STABLE | DOMAIN: PRECONCEPTION, PREGNANCY, GENETICS | REPLICATION: V**

## WHAT WE KNOW (Narrowest Supported Conclusion)

Male spermatogenesis involves continuous germline cell divisions throughout post-pubertal life (approximately 23 divisions per year), in contrast to female oogenesis where mitotic divisions cease prior to birth. In whole-genome sequencing of parent-offspring trios, the number of de novo single-nucleotide mutations transmitted to offspring increases linearly with the father's age at conception, adding roughly 1.5 to 2.0 new mutations per year of paternal age. The overwhelming majority of these mutations occur in non-coding or functionally silent genomic regions and produce no clinical phenotype.

## EVIDENCE AND CAUSAL ASSESSMENT

Kong et al. (2012), Nature, sequenced 78 Icelandic parent-offspring trios and demonstrated that paternal age at conception accounted for the dominant share of variation in offspring de novo mutation counts, estimating an increase of approximately 2.0 mutations per year of paternal age, while maternal age contributed no significant linear increase.

Jónsson et al. (2017), Nature, independently replicated and substantially scaled these findings across 1,548 Icelandic trios (identifying 108,778 de novo mutations). The multi-generational analysis established a precise genome-wide paternal increase of 1.51 mutations per year of father's age (95% CI [1.45, 1.57]), alongside a smaller maternal effect of 0.37 mutations per year (95% CI [0.32, 0.43]).

The causal mechanism is biological and well-established: continuous DNA replication errors and progressive replication-independent DNA damage in spermatogonia drive the mutational clock.

## EFFECT SIZE AND PERSISTENCE

The quantitative effect is approximately 1.51 de novo single-nucleotide mutations per year of paternal age. Across a 15-year difference in father's age (e.g., age 20 vs. age 35), an offspring inherits roughly 23 additional de novo point mutations out of 3.2 billion base pairs. Because over 98% of the human genome is non-coding and only a tiny fraction of coding mutations are deleterious, absolute clinical risk for specific neurodevelopmental conditions remains very low.

## BIAS, HETEROGENEITY, AND CONFOUNDING

GI attenuation: n/a for the germline biological mutation rate (measured directly via whole-genome sequence trios).

In contrast, observational epidemiological associations between advanced paternal age and child psychiatric outcomes (autism, schizophrenia) are largely confounded (>50%) by delayed reproductive selection, parental subclinical psychiatric traits, assortative mating, and socioeconomic indicators. Sibling-comparison studies demonstrate that population-level psychiatric risk shrinks substantially when comparing siblings born to the same father at different ages.

## WHAT THIS DOES NOT MEAN (Strict Boundary Conditions)

Does not mean that men in their late 30s or 40s should avoid having children out of fear of severe genetic disease. Does not mean older fathers produce children with impaired intelligence. A biological fact about germline replication does not license clinical alarmism or eugenic reproductive ranking.

## WHAT ACTION IS REASONABLY JUSTIFIED

Clinical genetic counselors should provide accurate absolute risk perspectives rather than inflated relative risk statistics when prospective parents inquire about paternal age. Routine whole-exome or whole-genome screening of asymptomatic children based solely on paternal age is not clinically indicated.

## VERIFIED SOURCES & DOI IDENTIFIERS

- Kong et al. (2012), Nature (DOI: 10.1038/nature11396 (https://doi.org/10.1038/nature11396))
- Jónsson et al. (2017), Nature (DOI: 10.1038/nature24018 (https://doi.org/10.1038/nature24018))