

TD – Semantics and Verification
X– Duality and Ultrafilter Extension
Friday 1st April 2022

TA: Ralph Sarkis
ralph.sarkis@ens-lyon.fr

Duality

For a fixed set Act , recall that $\mathfrak{L}(\text{HML})$ is the set of HML-formulas, it is a boolean algebra with the order $\varphi \leq \psi \Leftrightarrow \varphi \rightarrow \psi \equiv \top$ and the syntactically evident operations. Moreover, we let

- $\alpha \in \text{Act}$, the functions $[\alpha] : \mathfrak{L}(\text{HML}) \rightarrow \mathfrak{L}(\text{HML})$ and $\langle \alpha \rangle : \mathfrak{L}(\text{HML}) \rightarrow \mathfrak{L}(\text{HML})$ are defined by
 - $[\alpha](\phi) = [\alpha]\phi$
 - $\langle \alpha \rangle(\phi) = \langle \alpha \rangle\phi$
- for $TS = (S, \text{Act}, \rightarrow, I, \text{AP}, L)$, the functions $\llbracket [\alpha] \rrbracket$ and $\llbracket \langle \alpha \rangle \rrbracket$ are defined by
 - $\llbracket [\alpha] \rrbracket(A) = \{s \in S \mid \forall s' \in \text{Succ}^\alpha(s), s' \in A\}$
 - $\llbracket \langle \alpha \rangle \rrbracket(A) = \{s \in S \mid \exists s' \in \text{Succ}^\alpha(s), s' \in A\}$

For two Boolean algebras B and B' and a function $f : B \rightarrow B'$, the **dual** of f is the function $f^\partial : B \rightarrow B'$ defined by $f^\partial(b) = \neg' f(\neg b)$.

Exercise 1.

1. Consider a transition system $TS = (S, \text{Act}, \rightarrow, I, \text{AP}, L)$ and $\alpha \in \text{Act}$. Show that:
 - $\llbracket [\alpha] \rrbracket = \llbracket \langle \alpha \rangle \rrbracket^\partial$
 - $\llbracket \langle \alpha \rangle \rrbracket = \llbracket [\alpha] \rrbracket^\partial$
2. Let $\alpha \in \text{Act}$. Show that:
 - $[\alpha] = \langle \alpha \rangle^\partial$
 - $\langle \alpha \rangle = [\alpha]^\partial$

Exercise 2.

Let B and B' be two Boolean algebras and $f : B \rightarrow B'$. Show that:

1. $f^{\partial\partial} = f$
2. If f is map of join (resp. meet) semilattices, then f^∂ is map of meet (resp. join) semilattices.
3. If f is a map of lattices, then $f^\partial = f$.

Ultrafilter extension

Recall that:

- for a BAO $B^+ = (B, (f_\alpha)_{\alpha \in \text{Act}})$, the ultrafilter frame $\mathcal{U}f(B^+)$ is defined as
 - the states are the ultrafilter over B , and
 - given \mathcal{F}, \mathcal{H} two ultrafilters, $\mathcal{F} \xrightarrow{\alpha} \mathcal{H}$ iff $\forall b \in B, b \in \mathcal{H} \Rightarrow f_\alpha(b) \in \mathcal{F}$.
- for a set X , we define the function $\pi : X \rightarrow \mathcal{U}f(X)$ by $\pi(x) = \{A \in \mathcal{P}(X) \mid x \in A\}$. When X is finite, all ultrafilters are principal, so π is a bijection between X and $\mathcal{U}f(X)$.
- for a $TS = (S, \text{Act}, \rightarrow, I, \text{AP}, L)$, the ultrafilter extension $\mathcal{U}f(TS)$ is the transition system where
 - the states are the ultrafilters $\mathcal{U}f(S)$ on S
 - $\mathcal{F} \xrightarrow{\alpha} \mathcal{H}$ iff $\llbracket \langle \alpha \rangle \rrbracket(A) \in \mathcal{F}$ whenever $A \in \mathcal{H}$
 - $a \in L_{\mathcal{U}f}(\mathcal{F})$ iff $\{s \in S \mid a \in L(s)\} \in \mathcal{F}$
 - the initial states are $\{\pi(s) \mid s \in I\}$

Exercise 3.

Consider a BAO $B^+ = (B, (f_\alpha)_{\alpha \in \text{Act}})$. Show that in the ultrafilter frame $\mathcal{U}f(B^+)$, we have

$$\mathcal{F} \xrightarrow{\alpha} \mathcal{H} \text{ iff } \forall b \in B, f_\alpha(b) \in \mathcal{F} \Rightarrow b \in \mathcal{H}$$

Exercise 4.

Consider a $TS = (S, \text{Act}, \rightarrow, I, \text{AP}, L)$. Show that:

1. Given $s \in S$ and $a \in \text{AP}$, $a \in L(s)$ iff $a \in L_{\mathcal{U}f}(\pi(s))$.
2. Show that

$$\mathcal{F} \xrightarrow{\alpha} \mathcal{H} \text{ iff } \llbracket [\alpha] \rrbracket(A) \in \mathcal{F} \Rightarrow A \in \mathcal{H}$$

3. Given $s, s' \in S$ and $\alpha \in \text{Act}$, $s \xrightarrow{\alpha} s'$ in TS iff $\pi(s) \xrightarrow{\alpha} \pi(s')$ in $\mathcal{U}f(TS)$.