

Modelling Nondeterminism, Probability and Termination

The story of Ralph's internship with his amazing supervisors Matteo and Valeria.

Ralph Sarkis

ENS de Lyon

March 25th, 2021

Overview of the presentation:

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling
 - ▶ Nondeterministic choices

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices

Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination


Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability


Overview of the presentation:

- ▶ **Set** monads and equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ **1Met** monads and quantitative equational theories modelling
 - ▶ Nondeterministic choices
 - ▶ Probabilistic choices
 - ▶ Termination
 - ▶ Nondeterminism and probability
- ▶ Combining nondeterminism, probability and termination **NEW!**

Motivation




Motivation



Questions

- Are the two systems equivalent?


Motivation



Questions

- ▶ Are the two systems equivalent?
- ▶ Are they close to each other?


Motivation



Questions

- ▶ Are the two systems equivalent?
- ▶ Are they close to each other?
- ▶ What if the transition $x \rightarrow y$ is always chosen?

Motivation



Questions

- ▶ Are the two systems equivalent?
- ▶ Are they close to each other?
- ▶ What if the transition $x \rightarrow y$ is always chosen?
- ▶ What if it is never chosen?

Categorically

Categorically

- ▶ We use **monads**: functors with additional structure that is closely related to computation

Equationally

Categorically

- ▶ We use **monads**: functors with additional structure that is closely related to computation

Equationally

- ▶ We use **equational theories**: sets of operation symbols with axioms they satisfy

Categorically

- ▶ We use **monads**: functors with additional structure that is closely related to computation

Equationally

- ▶ We use **equational theories**: sets of operation symbols with axioms they satisfy

A link between these two pictures: **algebraic presentations of monads**.

Nondeterministic Choices (in **Set**)

Categorically

Nondeterministic Choices (in **Set**)

Categorically

- Powerset monad \mathcal{P} :

$$\mathcal{P}(X) = \{\text{non-empty finite subsets of } X\}$$

$$\eta_X = x \mapsto \{x\}$$

$$\mu_X = \mathcal{F} \mapsto \bigcup_{S \in \mathcal{F}} S$$

Nondeterministic Choices (in **Set**)

Categorically

Equationally

- Powerset monad \mathcal{P} :

$$\mathcal{P}(X) = \{\text{non-empty finite subsets of } X\}$$

$$\eta_X = x \mapsto \{x\}$$

$$\mu_X = \mathcal{F} \mapsto \bigcup_{S \in \mathcal{F}} S$$

- Transition system $t : X \rightarrow \mathcal{P}(X)$

$$t(x) = \{y \in X \mid x \xrightarrow{t} y\}$$

Nondeterministic Choices (in **Set**)

Categorically

- Powerset monad \mathcal{P} :

$$\mathcal{P}(X) = \{\text{non-empty finite subsets of } X\}$$

$$\eta_X = x \mapsto \{x\}$$

$$\mu_X = \mathcal{F} \mapsto \bigcup_{S \in \mathcal{F}} S$$

- Transition system $t : X \rightarrow \mathcal{P}(X)$

$$t(x) = \{y \in X \mid x \xrightarrow{t} y\}$$

Equationally

- Theory of (sup-)semilattices:
A binary operation \oplus satisfying

$$x \oplus x = x \quad I$$

$$x \oplus y = y \oplus x \quad C$$

$$(x \oplus y) \oplus z = x \oplus (y \oplus z). \quad A$$

Nondeterministic Choices (in **Set**)

Categorically

- Powerset monad \mathcal{P} :

$$\mathcal{P}(X) = \{\text{non-empty finite subsets of } X\}$$

$$\eta_X = x \mapsto \{x\}$$

$$\mu_X = \mathcal{F} \mapsto \bigcup_{S \in \mathcal{F}} S$$

- Transition system $t : X \rightarrow \mathcal{P}(X)$

$$t(x) = \{y \in X \mid x \xrightarrow{t} y\}$$

Equationally

- Theory of (sup-)semilattices:
A binary operation \oplus satisfying

$$x \oplus x = x \quad I$$

$$x \oplus y = y \oplus x \quad C$$

$$(x \oplus y) \oplus z = x \oplus (y \oplus z). \quad A$$

- Operational semantics

$$\frac{q \xrightarrow{a} r \quad q \xrightarrow{a} s}{q \xrightarrow{a} r \oplus s} \quad \frac{q \xrightarrow{a} r \quad q' \xrightarrow{a} r'}{q \oplus q' \xrightarrow{a} r \oplus r'} \\ \frac{q \downarrow \text{accept}}{q \oplus r \downarrow \text{accept}}$$

Probabilistic Choices (in **Set**)

Categorically

Probabilistic Choices (in **Set**)

Categorically

- Distribution monad \mathcal{D} :

$$\mathcal{D}(X) = \{\text{finitely supported} \\ \text{probability distributions on } X\}$$

$$\eta_X = x \mapsto \delta_x \text{ (Dirac)}$$

$$\mu_X = \Phi \mapsto \sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi) \cdot \varphi$$

Categorically

Equationally

- Distribution monad \mathcal{D} :

$$\mathcal{D}(X) = \{\text{finitely supported probability distributions on } X\}$$

$$\eta_X = x \mapsto \delta_x \text{ (Dirac)}$$

$$\mu_X = \Phi \mapsto \sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi) \cdot \varphi$$

- Transition system $t : X \rightarrow \mathcal{D}(X)$

$$t(x)(y) = \mathbb{P}[x \xrightarrow{t} y]$$

Probabilistic Choices (in **Set**)

Categorically

- Distribution monad \mathcal{D} :

$\mathcal{D}(X) = \{\text{finitely supported probability distributions on } X\}$

$$\eta_X = x \mapsto \delta_x \text{ (Dirac)}$$

$$\mu_X = \Phi \mapsto \sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi) \cdot \varphi$$

- Transition system $t : X \rightarrow \mathcal{D}(X)$

$$t(x)(y) = \mathbb{P}[x \xrightarrow{t} y]$$

Equationally

- Theory of convex algebras:
For each $p \in (0, 1)$, a binary operation $+_p$ satisfying

$$x +_p x = x \quad I_p$$

$$x +_p y = y +_{1-p} x \quad C_p$$

$$(x +_q y) +_p z = x +_{pq} (y +_{\frac{pq}{pq}} z) \quad A_p$$

Probabilistic Choices (in **Set**)

Categorically

- Distribution monad \mathcal{D} :

$\mathcal{D}(X) = \{\text{finitely supported probability distributions on } X\}$

$$\eta_X = x \mapsto \delta_x \text{ (Dirac)}$$

$$\mu_X = \Phi \mapsto \sum_{\varphi \in \text{supp}(\Phi)} \Phi(\varphi) \cdot \varphi$$

- Transition system $t : X \rightarrow \mathcal{D}(X)$

$$t(x)(y) = \mathbb{P}[x \xrightarrow{t} y]$$

Equationally

- Theory of convex algebras:
For each $p \in (0, 1)$, a binary operation $+_p$ satisfying

$$x +_p x = x \quad I_p$$

$$x +_p y = y +_{1-p} x \quad C_p$$

$$(x +_q y) +_p z = x +_{pq} (y +_{\frac{pq}{p-q}} z) \quad A_p$$

- Operational semantics

$$\frac{\frac{q \xrightarrow{a,p} r \quad q \xrightarrow{a,1-p} s}{q \xrightarrow{a} r +_p s} \quad \frac{q \xrightarrow{a} r \quad q' \xrightarrow{a} r'}{q +_p q' \xrightarrow{a} r +_p r'}}{\frac{q \downarrow o(q) \quad r \downarrow o(r)}{q +_p r \downarrow po(q) + (1-p)o(r)}}$$

Termination (in **Set**)

Categorically

Termination (in **Set**)

Categorically

- Maybe monad $\cdot + \mathbf{1}$:

$$X + \mathbf{1} = X \sqcup \{\star\}$$

$$\eta_X = \text{inl}^{X+\mathbf{1}}$$

$$\mu_X = [\text{inl}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}]$$

Termination (in **Set**)

Categorically

- Maybe monad $\cdot + \mathbf{1}$:

$$X + \mathbf{1} = X \sqcup \{\star\}$$

$$\eta_X = \text{inl}^{X+\mathbf{1}}$$

$$\mu_X = [\text{inl}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}]$$

- Transition system $t : X \rightarrow X + \mathbf{1}$

$$t(x) = \begin{cases} y & x \xrightarrow{t} y \\ \star & x \nrightarrow \end{cases}$$

Equationally

Termination (in **Set**)

Categorically

- Maybe monad $\cdot + \mathbf{1}$:

$$X + \mathbf{1} = X \sqcup \{\star\}$$

$$\eta_X = \text{inl}^{X+\mathbf{1}}$$

$$\mu_X = [\text{inl}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}]$$

- Transition system $t : X \rightarrow X + \mathbf{1}$

$$t(x) = \begin{cases} y & x \xrightarrow{t} y \\ \star & x \nrightarrow \end{cases}$$

Equationally

- Theory of pointed sets: A constant (0-ary) \star with no axioms.

Termination (in **Set**)

Categorically

- ▶ Maybe monad $\cdot + \mathbf{1}$:

$$X + \mathbf{1} = X \sqcup \{\star\}$$

$$\eta_X = \text{inl}^{X+\mathbf{1}}$$

$$\mu_X = [\text{inl}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}, \text{inr}^{X+\mathbf{1}}]$$

- ▶ Transition system $t : X \rightarrow X + \mathbf{1}$

$$t(x) = \begin{cases} y & x \xrightarrow{t} y \\ \star & x \nrightarrow \end{cases}$$

Equationally

- ▶ Theory of pointed sets: A constant (0-ary) \star with no axioms.
- ▶ Operational semantics

$$\frac{q \nrightarrow}{q \rightarrow \star}$$

Nondeterminism and Probability (in **Set**)

Categorically

Nondeterminism and Probability (in Set)

Categorically

► Monad \mathcal{C} :

$\mathcal{C}(X) = \{\text{non-empty finitely generated}$
 $\text{convex sets of distributions on } X\}$

$$\eta_X = x \mapsto \{\delta_x\}$$

$$\mu_X = F \mapsto \bigcup_{\Phi \in F} \left\{ \sum_{S \in \text{supp}(\Phi)} \Phi(S) \cdot \varphi_S \mid \varphi_S \in S \right\}$$

Nondeterminism and Probability (in **Set**)

Categorically

Equationally

► Monad \mathcal{C} :

$\mathcal{C}(X) = \{\text{non-empty finitely generated convex sets of distributions on } X\}$

$$\eta_X = x \mapsto \{\delta_x\}$$

$$\mu_X = F \mapsto \bigcup_{\Phi \in F} \left\{ \sum_{S \in \text{supp}(\Phi)} \Phi(S) \cdot \varphi_S \mid \varphi_S \in S \right\}$$

► Transition system $t : X \rightarrow \mathcal{C}(X)$

$$t(x) = \text{conv}\{\varphi \in \mathcal{D}(X) \mid x \xrightarrow{t} \varphi\},$$

where conv is the convex closure.

Nondeterminism and Probability (in Set)

Categorically

- Monad \mathcal{C} :

$\mathcal{C}(X) = \{\text{non-empty finitely generated convex sets of distributions on } X\}$

$$\eta_X = x \mapsto \{\delta_x\}$$

$$\mu_X = F \mapsto \bigcup_{\Phi \in F} \left\{ \sum_{S \in \text{supp}(\Phi)} \Phi(S) \cdot \varphi_S \mid \varphi_S \in S \right\}$$

- Transition system $t : X \rightarrow \mathcal{C}(X)$

$$t(x) = \text{conv}\{\varphi \in \mathcal{D}(X) \mid x \xrightarrow{t} \varphi\},$$

where conv is the convex closure.

Equationally

- Theory of convex semilattices: A semilattice operation and convex algebra operations that distribute

$$(x \oplus y) +_p z = (x +_p z) \oplus (y +_p z) \quad (D_p)$$

Nondeterminism and Probability (in Set)

Categorically

- Monad \mathcal{C} :

$\mathcal{C}(X) = \{\text{non-empty finitely generated convex sets of distributions on } X\}$

$$\eta_X = x \mapsto \{\delta_x\}$$

$$\mu_X = F \mapsto \bigcup_{\Phi \in F} \left\{ \sum_{S \in \text{supp}(\Phi)} \Phi(S) \cdot \varphi_S \mid \varphi_S \in S \right\}$$

- Transition system $t : X \rightarrow \mathcal{C}(X)$

$$t(x) = \text{conv}\{\varphi \in \mathcal{D}(X) \mid x \xrightarrow{t} \varphi\},$$

where conv is the convex closure.

Equationally

- Theory of convex semilattices: A semilattice operation and convex algebra operations that distribute

$$(x \oplus y) +_p z = (x +_p z) \oplus (y +_p z) \quad (D_p)$$

- Operational semantics: Combine all previous rules.

[BSV19]

Categorically

Categorically

- ▶ We switch from **Set** to **1Met**.

Categorically

- ▶ We switch from **Set** to **1Met**.
- ▶ Sets become metric spaces (X, d) which are **1-bounded**, i.e.:
 $\forall x, y \in X, d(x, y) \leq 1.$

Categorically

- ▶ We switch from **Set** to **1Met**.
- ▶ Sets become metric spaces (X, d) which are **1-bounded**, i.e.:
 $\forall x, y \in X, d(x, y) \leq 1.$
- ▶ Functions become **non-expansive** maps, i.e.: $f : (X, d_X) \rightarrow (Y, d_Y)$ satisfies

$$\forall x, x' \in X, d_Y(f(x), f(x')) \leq d_X(x, x').$$

Equationally

Categorically

- ▶ We switch from **Set** to **1Met**.
- ▶ Sets become metric spaces (X, d) which are **1-bounded**, i.e.:
 $\forall x, y \in X, d(x, y) \leq 1$.
- ▶ Functions become **non-expansive** maps, i.e.: $f : (X, d_X) \rightarrow (Y, d_Y)$ satisfies

$$\forall x, x' \in X, d_Y(f(x), f(x')) \leq d_X(x, x').$$

Equationally

- ▶ Equations become quantitative

$$x =_\varepsilon y \quad (\varepsilon \in [0, 1])$$

Categorically

- ▶ We switch from **Set** to **1Met**.
- ▶ Sets become metric spaces (X, d) which are **1-bounded**, i.e.:
 $\forall x, y \in X, d(x, y) \leq 1$.
- ▶ Functions become **non-expansive** maps, i.e.: $f : (X, d_X) \rightarrow (Y, d_Y)$ satisfies

$$\forall x, x' \in X, d_Y(f(x), f(x')) \leq d_X(x, x').$$

Equationally

- ▶ Equations become quantitative

$$x =_\varepsilon y \quad (\varepsilon \in [0, 1])$$

- ▶ Quantitative equational logic

Quantitative Equational Logic

$\vdash t =_0 t$	Refl
$\vdash t =_1 s$	1-bounded
$t =_\varepsilon s \vdash s =_\varepsilon t$	Symm
$t =_\varepsilon s, s =_\delta u \vdash t =_{\varepsilon+\delta} u$	Triang
$t =_\varepsilon s \vdash t =_{\varepsilon+\delta} s$	Max (for $\delta > 0$)
$\forall \varepsilon > \delta, t =_\varepsilon s \vdash t =_\delta s$	Arch (for $\delta \geq 0$)
$\forall 1 \leq i \leq n, t_i =_\varepsilon s_i \vdash f(t_1, \dots, t_n) =_\varepsilon f(s_1, \dots, s_n)$	Nexp (for f an n -ary)
$\Gamma \vdash t =_\varepsilon s \implies \sigma(\Gamma) \vdash \sigma(t) =_\varepsilon \sigma(s)$	Subst (for σ a substitution)
$\phi \in \Gamma \implies \Gamma \vdash \phi$	Assumption

Introduced by Mardare, Panangaden and Plotkin [MPP16].

Nondeterminism Choices (in **1Met**)

Categorically

Nondeterminism Choices (in **1Met**)

Categorically

- Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ into $H(d) : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow [0, 1]$:

$$H(d)(S, T) = \max \left\{ \sup_{s \in S} \inf_{t \in T} d(s, t), \sup_{t \in T} \inf_{s \in S} d(s, t) \right\}$$

Nondeterminism Choices (in **1Met**)

Categorically

- ▶ Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ into $H(d) : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow [0, 1]$:

$$H(d)(S, T) = \max \left\{ \sup_{s \in S} \inf_{t \in T} d(s, t), \sup_{t \in T} \inf_{s \in S} d(s, t) \right\}$$

- ▶ Hausdorff lifting of \mathcal{P} : $\widehat{\mathcal{P}}(X, d) = (\mathcal{P}(X), H(d))$.

Nondeterminism Choices (in **1Met**)

Categorically

- ▶ Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ into $H(d) : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow [0, 1]$:

$$H(d)(S, T) = \max \left\{ \sup_{s \in S} \inf_{t \in T} d(s, t), \sup_{t \in T} \inf_{s \in S} d(s, t) \right\}$$

- ▶ Hausdorff lifting of \mathcal{P} : $\widehat{\mathcal{P}}(X, d) = (\mathcal{P}(X), H(d))$.

Equationally

Nondeterminism Choices (in 1Met)

Categorically

- Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ into $H(d) : \mathcal{P}(X) \times \mathcal{P}(X) \rightarrow [0, 1]$:

$$H(d)(S, T) = \max \left\{ \sup_{s \in S} \inf_{t \in T} d(s, t), \sup_{t \in T} \inf_{s \in S} d(s, t) \right\}$$

- Hausdorff lifting of \mathcal{P} : $\widehat{\mathcal{P}}(X, d) = (\mathcal{P}(X), H(d))$.

Equationally

- Theory of quantitative (sup-)semilattices: binary operation \oplus satisfying

$$\vdash x \oplus x =_0 x \quad I$$

$$\vdash x \oplus y =_0 y \oplus x \quad C$$

$$\vdash (x \oplus y) \oplus z =_0 x \oplus (y \oplus z) \quad A$$

$$x_1 =_{\varepsilon_1} y_1, x_2 =_{\varepsilon_2} y_2 \vdash x_1 \oplus x_2 =_{\max\{\varepsilon_1, \varepsilon_2\}} y_1 \oplus y_2 \quad H$$

Probabilistic choices (in **1Met**)

Categorically

Probabilistic choices (in **1Met**)

Categorically

- Kantorovitch lifting of $d : X \times X \rightarrow [0, 1]$ into $K(d) : \mathcal{D}(X) \times \mathcal{D}(X) \rightarrow [0, 1]$:

$$K(d)(\varphi, \psi) = \inf_{\omega} \sum_{x_1, x_2 \in X} \omega(x_1, x_2) \cdot d(x_1, x_2),$$

where ω ranges over couplings of φ and ψ .

Probabilistic choices (in **1Met**)

Categorically

- ▶ Kantorovitch lifting of $d : X \times X \rightarrow [0, 1]$ into $K(d) : \mathcal{D}(X) \times \mathcal{D}(X) \rightarrow [0, 1]$:

$$K(d)(\varphi, \psi) = \inf_{\omega} \sum_{x_1, x_2 \in X} \omega(x_1, x_2) \cdot d(x_1, x_2),$$

where ω ranges over couplings of φ and ψ .

- ▶ Kantorovitch lifting of \mathcal{D} : $\widehat{\mathcal{D}}(X, d) = (\mathcal{D}(X), K(d))$.

Probabilistic choices (in **1Met**)

Categorically

- ▶ Kantorovitch lifting of $d : X \times X \rightarrow [0, 1]$ into $K(d) : \mathcal{D}(X) \times \mathcal{D}(X) \rightarrow [0, 1]$:

$$K(d)(\varphi, \psi) = \inf_{\omega} \sum_{x_1, x_2 \in X} \omega(x_1, x_2) \cdot d(x_1, x_2),$$

where ω ranges over couplings of φ and ψ .

- ▶ Kantorovitch lifting of \mathcal{D} : $\widehat{\mathcal{D}}(X, d) = (\mathcal{D}(X), K(d))$.

Equationally

Probabilistic choices (in **1Met**)

Categorically

- Kantorovitch lifting of $d : X \times X \rightarrow [0, 1]$ into $K(d) : \mathcal{D}(X) \times \mathcal{D}(X) \rightarrow [0, 1]$:

$$K(d)(\varphi, \psi) = \inf_{\omega} \sum_{x_1, x_2 \in X} \omega(x_1, x_2) \cdot d(x_1, x_2),$$

where ω ranges over couplings of φ and ψ .

- Kantorovitch lifting of \mathcal{D} : $\widehat{\mathcal{D}}(X, d) = (\mathcal{D}(X), K(d))$.

Equationally

- Theory of quantitative convex algebras

$$\vdash x +_p x =_0 x \quad I_p$$

$$\vdash x +_p y =_0 y +_p x \quad C_p$$

$$\vdash (x +_q y) +_p z =_0 x +_{pq} (y +_{\frac{p\bar{q}}{pq}} z) \quad A_p$$

$$x_1 =_{\varepsilon_1} y_1, x_2 =_{\varepsilon_2} y_2 \vdash x_1 +_p x_2 =_{p\varepsilon_1 + (1-p)\varepsilon_2} y_1 +_p y_2 \quad K_p$$

Termination (in **1Met**)

Categorically

Termination (in **1Met**)

Categorically

- ▶ The maybe monad $\cdot + \hat{\mathbf{1}}$ puts \star at distance 1 from every other point:

$$X + \hat{\mathbf{1}} = X \sqcup \{\star\}$$

$$(d + d_{\hat{\mathbf{1}}})(x, y) = \begin{cases} 1 & x =_0 \star \vee y = \star \\ d(x, y) & \text{otherwise} \end{cases}$$

Equationally

Termination (in **1Met**)

Categorically

- ▶ The maybe monad $\cdot + \hat{\mathbf{1}}$ puts \star at distance 1 from every other point:

$$X + \hat{\mathbf{1}} = X \sqcup \{\star\}$$

$$(d + d_{\hat{\mathbf{1}}})(x, y) = \begin{cases} 1 & x =_0 \star \vee y = \star \\ d(x, y) & \text{otherwise} \end{cases}$$

Equationally

- ▶ Theory of pointed spaces:

$$\vdash x =_1 \star$$

Nondeterminism and Probability (in **1Met**)

Categorically

Categorically

- ▶ Combine Kantorovitch and Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ to obtain $\widehat{\mathcal{C}}(X, d) = (\mathcal{C}(X), HK(d))$.

Nondeterminism and Probability (in **1Met**)

Categorically

- Combine Kantorovitch and Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ to obtain $\widehat{\mathcal{C}}(X, d) = (\mathcal{C}(X), HK(d))$.

Equationally

Nondeterminism and Probability (in **1Met**)

Categorically

- Combine Kantorovitch and Hausdorff lifting of $d : X \times X \rightarrow [0, 1]$ to obtain $\widehat{\mathcal{C}}(X, d) = (\mathcal{C}(X), HK(d))$.

Equationally

- Quantitative theory of convex semilattices: combine all previous axioms plus

$$\vdash (x \oplus y) +_p z =_0 (x +_p z) \oplus (y +_p z) \qquad D_p$$

[MV20]

Adding Termination (easy)

Categorically

Adding Termination (easy)

Categorically

- For any monad M , $M(\cdot + \mathbf{1})$ is a monad.

Adding Termination (easy)

Categorically

- ▶ For any monad M , $M(\cdot + \mathbf{1})$ is a monad.
- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is the monad of non-empty finitely generated convex sets of *subdistributions* on X .

Adding Termination (easy)

Categorically

- ▶ For any monad M , $M(\cdot + \mathbf{1})$ is a monad.
- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is the monad of non-empty finitely generated convex sets of *subdistributions* on X .
- ▶ Similarly on **1Met**: the monad is $\widehat{\mathcal{C}}(\cdot + \widehat{\mathbf{1}})$.

Equationally

Adding Termination (easy)

Categorically

- ▶ For any monad M , $M(\cdot + \mathbf{1})$ is a monad.
- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is the monad of non-empty finitely generated convex sets of *subdistributions* on X .
- ▶ Similarly on **1Met**: the monad is $\hat{\mathcal{C}}(\cdot + \hat{\mathbf{1}})$.

Equationally

- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is presented by the theory of pointed convex semilattices.

Adding Termination (easy)

Categorically

- ▶ For any monad M , $M(\cdot + \mathbf{1})$ is a monad.
- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is the monad of non-empty finitely generated convex sets of *subdistributions* on X .
- ▶ Similarly on **1Met**: the monad is $\widehat{\mathcal{C}}(\cdot + \widehat{\mathbf{1}})$.

Equationally

- ▶ On **Set**: $\mathcal{C}(\cdot + \mathbf{1})$ is presented by the theory of pointed convex semilattices.
- ▶ On **1Met**: $\widehat{\mathcal{C}}(\cdot + \widehat{\mathbf{1}})$ is presented by the theory of quantitative pointed convex semilattices.

Adding Termination (medium)

On Set

Adding Termination (medium)

On Set

- ▶ We define the monad $\mathcal{C} + \mathbf{1}$ for *possibly empty* finitely generated convex sets of distributions on X .

Adding Termination (medium)

On Set

- ▶ We define the monad $\mathcal{C} + \mathbf{1}$ for *possibly empty* finitely generated convex sets of distributions on X .
- ▶ \star behaves like the empty set, i.e.: $\mathcal{C} + \mathbf{1}$ is presented by convex semilattices with *bottom* and *black-hole*:

$$x \oplus \star = x \quad (\perp)$$

$$x +_p \star = \star \quad (BH_p)$$

Adding Termination (medium)

On Set

- ▶ We define the monad $\mathcal{C} + \mathbf{1}$ for *possibly empty* finitely generated convex sets of distributions on X .
- ▶ \star behaves like the empty set, i.e.: $\mathcal{C} + \mathbf{1}$ is presented by convex semilattices with *bottom* and *black-hole*:

$$x \oplus \star = x$$

$$(\perp)$$

$$x +_p \star = \star$$

$$(BH_p)$$

On 1Met

Adding Termination (medium)

On Set

- ▶ We define the monad $\mathcal{C} + \mathbf{1}$ for *possibly empty* finitely generated convex sets of distributions on X .
- ▶ \star behaves like the empty set, i.e.: $\mathcal{C} + \mathbf{1}$ is presented by convex semilattices with *bottom* and *black-hole*:

$$x \oplus \star = x \quad (\perp)$$

$$x +_p \star = \star \quad (BH_p)$$

On 1Met

- ▶ The black-hole equation trivializes the theory.

Adding Termination (medium)

On Set

- ▶ We define the monad $\mathcal{C} + \mathbf{1}$ for *possibly empty* finitely generated convex sets of distributions on X .
- ▶ \star behaves like the empty set, i.e.: $\mathcal{C} + \mathbf{1}$ is presented by convex semilattices with *bottom* and *black-hole*:

$$x \oplus \star = x$$

$$(\perp)$$

$$x +_p \star = \star$$

$$(BH_p)$$

On **1Met**

- ▶ The black-hole equation trivializes the theory.
- ▶ There is (probably) no monad $\mathcal{C} + \mathbf{1}$ on **1Met**.

Adding Termination (hard)

On Set

Adding Termination (hard)

On Set

- ▶ Remove the black-hole equation.
- ▶ This theory presents the monad \mathcal{C}^\downarrow for \perp -closed finitely generated convex sets of subdistributions on X .

Adding Termination (hard)

On Set

- ▶ Remove the black-hole equation.
- ▶ This theory presents the monad \mathcal{C}^\downarrow for \perp -closed finitely generated convex sets of subdistributions on X .
- ▶ $S \in \mathcal{C}(X + \mathbf{1})$ is \perp -**closed** iff $\forall \varphi \in S$,
 $\{\psi \in \mathcal{C}(X + \mathbf{1}) \mid \forall x \in X, \psi(x) \leq \varphi(x)\} \subseteq S$

Adding Termination (hard)

On Set

- ▶ Remove the black-hole equation.
- ▶ This theory presents the monad \mathcal{C}^\downarrow for \perp -closed finitely generated convex sets of subdistributions on X .
- ▶ $S \in \mathcal{C}(X + \mathbf{1})$ is \perp -**closed** iff $\forall \varphi \in S$,
 $\{\psi \in \mathcal{C}(X + \mathbf{1}) \mid \forall x \in X, \psi(x) \leq \varphi(x)\} \subseteq S$




Figure 1: Examples of \perp -closed sets.

Adding Termination (hard)

On Set

- ▶ Remove the black-hole equation.
- ▶ This theory presents the monad \mathcal{C}^\downarrow for \perp -closed finitely generated convex sets of subdistributions on X .
- ▶ $S \in \mathcal{C}(X + \mathbf{1})$ is \perp -**closed** iff $\forall \varphi \in S$,
 $\{\psi \in \mathcal{C}(X + \mathbf{1}) \mid \forall x \in X, \psi(x) \leq \varphi(x)\} \subseteq S$

On 1Met




Figure 1: Examples of \perp -closed sets.

Adding Termination (hard)

On Set

- ▶ Remove the black-hole equation.
- ▶ This theory presents the monad \mathcal{C}^\downarrow for \perp -closed finitely generated convex sets of subdistributions on X .
- ▶ $S \in \mathcal{C}(X + \mathbf{1})$ is \perp -**closed** iff $\forall \varphi \in S, \{\psi \in \mathcal{C}(X + \mathbf{1}) \mid \forall x \in X, \psi(x) \leq \varphi(x)\} \subseteq S$

On 1Met

- ▶ The theory of quantitative convex semilattices with bottom corresponds to \mathcal{C}^\downarrow with the Kantorovitch and Hausdorff liftings of the metrics.




Figure 1: Examples of \perp -closed sets.

Summary

On Set		On 1Met	
\mathcal{P}	Semilattices	$\hat{\mathcal{P}}$	Quantitative semilattices
\mathcal{D}	Convex algebras	$\hat{\mathcal{D}}$	Quantitative convex algebras
$\cdot + \mathbf{1}$	Pointed sets	$\cdot + \hat{\mathbf{1}}$	Pointed spaces
\mathcal{C}	Convex semilattices (CS)	$\hat{\mathcal{C}}$	Quantitative CS
$\mathcal{C}(\cdot + \mathbf{1})$	Pointed CS	$\hat{\mathcal{C}}(\cdot + \hat{\mathbf{1}})$	Quantitative pointed CS
$\mathcal{C} + \mathbf{1}$	CS with \perp and black-hole	Trivial	Quantitative CS with \perp and black-hole
\mathcal{C}^\downarrow	CS with \perp	$\hat{\mathcal{C}}^\downarrow$	Quantitative CS with \perp

See Section VI of preprint.

- ▶ Can our results be obtained *compositionally*?

- ▶ Can our results be obtained *compositionally*? Close to answering yes with Daniela Petrişan.

- ▶ Can our results be obtained *compositionally*? Close to answering yes with Daniela Petrişan.
- ▶ Where can quantitative algebraic reasoning be applied?

References

- [BSV19] Filippo Bonchi, Ana Sokolova, and Valeria Vignudelli. “The Theory of Traces for Systems with Nondeterminism and Probability”. In: *34th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2019, Vancouver, BC, Canada, June 24-27, 2019*. IEEE, 2019, pp. 1–14. DOI: [10.1109/LICS.2019.8785673](https://doi.org/10.1109/LICS.2019.8785673).
- [MPP16] Radu Mardare, Prakash Panangaden, and Gordon D. Plotkin. “Quantitative Algebraic Reasoning”. In: *Proceedings of the 31st Annual ACM/IEEE Symposium on Logic in Computer Science, LICS '16, New York, NY, USA, July 5-8, 2016*. Ed. by Martin Grohe, Eric Koskinen, and Natarajan Shankar. ACM, 2016, pp. 700–709. DOI: [10.1145/2933575.2934518](https://doi.org/10.1145/2933575.2934518). URL: <https://doi.org/10.1145/2933575.2934518>.
- [MV20] Matteo Mio and Valeria Vignudelli. “Monads and Quantitative Equational Theories for Nondeterminism and Probability”. In: *CoRR abs/2005.07509* (2020). arXiv: 2005.07509. URL: <https://arxiv.org/abs/2005.07509>.

Merci !