

Math 207: Statistics

Chapter 21: The Accuracy of Percentages

Population (parameters)

Sample (statistics)

Dr. Ralph Wojtowicz

Mathematics Department



SHENANDOAH
UNIVERSITY

1 EV, SE and the Central Limit Theorem

2 Examples



- Example I
- Example I

3 Confidence Intervals

- Inferential Statistics
- Confidence Intervals
- Example

Expected Value, Standard Error, Central Limit Theorem

- Many statistics problems are modeled as samples from a box of numbered tickets.
- Solution procedure:
 - Formulate a box model.
 - Compute the average and SD of the contents of the box.
 - Determine if you are computing a sum or average (% in a 0/1 box is an average).
 - Use the appropriate formulas to compute the EV and SE for the sample.
 - Use the normal curve to compute chances of the sample being in a specified range.



Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

$$\boxed{253,785 \text{ } \boxed{0} \text{ s } 433,211 \text{ } \boxed{1} \text{ s}}$$

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$ and $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$ and $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$ and $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$.

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

$$\boxed{253,785 \text{ } \boxed{0} \text{ s } 433,211 \text{ } \boxed{1} \text{ s}}$$

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$ and $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$ and $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$.
- Compute the z scores for 61% and 65%:
 $z_1 = (61 - 63.1)/4.83 = -0.435$
 $z_2 = (65 - 63.1)/4.83 = 0.393$

Example: Box Contents Known

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size $n = 100$ from the box below.

$$\boxed{253,785 \text{ } \boxed{0} \text{ s } 433,211 \text{ } \boxed{1} \text{ s}}$$

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$ and $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$ and $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$.
- Compute the z scores for 61% and 65%:
 $z_1 = (61 - 63.1)/4.83 = -0.435$
 $z_2 = (65 - 63.1)/4.83 = 0.393$
- Area under the normal curve: $\text{pnorm}(0.393) - \text{pnorm}(-0.435) = 0.321 = 32.1\%$

Example: Box Contents Unknown

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.

???	0	s	???	1	s
-----	---	---	-----	---	---

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.

??? 0s ??? 1s

- Suppose our sample has 63 1s.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.

??? 0s ??? 1s

- Suppose our sample has 63 1s.
- Estimate the fraction of 1s in the box and put a plus or minus on the estimate.
- $AV_{\text{box}} = ???$ Use 63% as our estimate.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.

???	0	s	???	1	s
-----	---	---	-----	---	---

- Suppose our sample has 63 1s.
- Estimate the fraction of 1s in the box and put a plus or minus on the estimate.
- $AV_{\text{box}} = ???$ Use 63% as our estimate.
- $SD_{\text{box}} = ???$. Use $(1 - 0)\sqrt{0.63(1 - 0.63)} \approx 0.483$ as our estimate.
- $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.

$$\boxed{??? \ 0 \text{ s} \quad ??? \ 1 \text{ s}}$$


- Suppose our sample has 63 $\boxed{1}$ s.
- Estimate the fraction of $\boxed{1}$ s in the box and put a plus or minus on the estimate.
- $AV_{\text{box}} = ???$ Use 63% as our estimate.
- $SD_{\text{box}} = ???$. Use $(1 - 0)\sqrt{0.63(1 - 0.63)} \approx 0.483$ as our estimate.
- $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$.
- If we assume that the samples follow a normal curve, then

$$63\% \pm 2 \cdot (4.83\%) = 63\% \pm 9.7\%$$

is **95% confidence interval** on the fraction of $\boxed{1}$ s.

Inferential Statistics

- In real situations, we typically don't know the contents of the box!
- That is where **inferential statistics** is used.



Confidence Intervals

- In real situations, we typically don't know the contents of the box!

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only 0s and 1s.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only 0s and 1s.
- Take a random sample of size n .

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only $\boxed{0}$ s and $\boxed{1}$ s.
- Take a random sample of size n .
- The fraction p of $\boxed{1}$ s in the sample estimates the fraction of $\boxed{1}$ s in the box.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only $\boxed{0}$ s and $\boxed{1}$ s.
- Take a random sample of size n .
- The fraction p of $\boxed{1}$ s in the sample estimates the fraction of $\boxed{1}$ s in the box.
- Use the shortcut method to estimate the SD: $SD_{box} \approx (1 - 0)\sqrt{p(1 - p)}$.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only $\boxed{0}$ s and $\boxed{1}$ s.
- Take a random sample of size n .
- The fraction p of $\boxed{1}$ s in the sample estimates the fraction of $\boxed{1}$ s in the box.
- Use the shortcut method to estimate the SD: $SD_{box} \approx (1 - 0)\sqrt{p(1 - p)}$.
- Then compute the SE.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only $\boxed{0}$ s and $\boxed{1}$ s.
- Take a random sample of size n .
- The fraction p of $\boxed{1}$ s in the sample estimates the fraction of $\boxed{1}$ s in the box.
- Use the shortcut method to estimate the SD: $SD_{box} \approx (1 - 0)\sqrt{p(1 - p)}$.
- Then compute the SE.
- The 95% confidence interval is:

$$(\text{observed fraction}) \pm 2 \cdot (\text{SE estimate})$$

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{box} \quad SE_{av} = \frac{SD_{box}}{\sqrt{n}}$$

- Assume the box contains only $\boxed{0}$ s and $\boxed{1}$ s.
- Take a random sample of size n .
- The fraction p of $\boxed{1}$ s in the sample estimates the fraction of $\boxed{1}$ s in the box.
- Use the shortcut method to estimate the SD: $SD_{box} \approx (1 - 0)\sqrt{p(1 - p)}$.
- Then compute the SE.
- The 95% confidence interval is:
(observed fraction) $\pm 2 \cdot$ (SE estimate)
- Prior to taking the sample, we are 95% confident that this procedure will give an interval that contains the true average.

Example 1 Page 378

- A university has 25,000 registered students.

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - a for every student who lives at home
 - a for others

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - a 1 for every student who lives at home
 - a 0 for others
- Observed fraction of 1s: $317/400$.

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - a for every student who lives at home
 - a for others
- Observed fraction of s: $317/400$.
- Estimated fraction of s in the box: $317/400 = 79.3\%$.

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - a 1 for every student who lives at home
 - a 0 for others
- Observed fraction of 1s: $317/400$.
- Estimated fraction of 1s in the box: $317/400 = 79.3\%$.
- Now compute a \pm on this estimate:
 - $SD_{\text{box}} \approx \sqrt{0.793 \cdot (1 - 0.793)} = 0.41$.
 - $SE_{\text{av}} = 0.41/\sqrt{400} \approx 0.020 = 2.0\%$

Example 1 Page 378

- A university has 25,000 registered students.
- Estimate the fraction who live at home by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - a 1 for every student who lives at home
 - a 0 for others
- Observed fraction of 1s: $317/400$.
- Estimated fraction of 1s in the box: $317/400 = 79.3\%$.
- Now compute a \pm on this estimate:
 - $SD_{\text{box}} \approx \sqrt{0.793 \cdot (1 - 0.793)} = 0.41$.
 - $SE_{\text{av}} = 0.41/\sqrt{400} \approx 0.020 = 2.0\%$
- 95% confidence interval:

$$79.3\% \pm 2 \cdot (2.0\%) = 79.3\% \pm 4\%$$