

Math 207: Statistics

Chapter 23: The Accuracy of Averages

Population (parameters)

Sample (statistics)

Dr. Ralph Wojtowicz

Mathematics Department



SHENANDOAH
UNIVERSITY

1 EV, SE and the Central Limit Theorem

2 Examples



- Example I
- Example I

3 Confidence Intervals

- Inferential Statistics
- Confidence Intervals
- Example

Expected Value, Standard Error, Central Limit Theorem

- Many statistics problems are modeled as samples from a box of numbered tickets.
- Solution procedure:
 - Formulate a box model.
 - Compute the average and SD of the contents of the box.
 - Determine if you are computing a sum or average (% in a 0/1 box is an average).
 - Use the appropriate formulas to compute the EV and SE for the sample.
 - Use the normal curve to compute chances of the sample being in a specified range.



Example: Box Contents Known

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2.$

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2.$
- $EV_{\text{av}} = AV_{\text{box}} = 4$ and $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{2}{\sqrt{100}} = 0.2.$

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2.$
- $EV_{\text{av}} = AV_{\text{box}} = 4$ and $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{2}{\sqrt{100}} = 0.4.$
- We expect the average of the draws to be around 4 give or take 0.4 or so.

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2.$
- $EV_{\text{av}} = AV_{\text{box}} = 4$ and $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{2}{\sqrt{100}} = 0.4.$
- We expect the average of the draws to be around 4 give or take 0.4 or so.
- What is the chance that the average will be between 3.5 and 4.5?

Example: Box Contents Known


- Take a random sample of size $n = 100$ from the box below.

1	2	3	4	5	6	7
---	---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2.$
- $EV_{\text{av}} = AV_{\text{box}} = 4$ and $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{2}{\sqrt{100}} = 0.4.$
- We expect the average of the draws to be around 4 give or take 0.4 or so.
- What is the chance that the average will be between 3.5 and 4.5?
- Compute the z scores for 61% and 65%:
 $z_1 = (3.5 - 4)/0.4 = -1.25$
 $z_2 = (4.5 - 4)/0.4 = 1.25$

Example: Box Contents Known

- Take a random sample of size $n = 100$ from the box below.




- $AV_{\text{box}} = \frac{1+2+\dots+7}{7} = 4$ and $SD_{\text{box}} = \sqrt{\frac{(1-4)^2+(2-4)^2+\dots+(7-4)^2}{7}} = \sqrt{\frac{28}{7}} = \sqrt{4} = 2$.
- $EV_{\text{av}} = AV_{\text{box}} = 4$ and $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{2}{\sqrt{100}} = 0.4$.
- We expect the average of the draws to be around 4 give or take 0.4 or so.
- What is the chance that the average will be between 3.5 and 4.5?
- Compute the z scores for 61% and 65%:
 $z_1 = (3.5 - 4)/0.4 = -1.25$
 $z_2 = (4.5 - 4)/0.4 = 1.25$
- Area under the normal curve: $\text{pnorm}(1.25) - \text{pnorm}(-1.25) = 0.789 = 78.9\%$

Example: Box Contents Unknown

Example: Box Contents Unknown


- Take a random sample of size $n = 100$ from the box below.



We don't know what numbers are in the box!

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.




We don't know what numbers are in the box!

- Suppose our sample of size 100 has an average of 10.2 and an SD of 0.8.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.




We don't know what numbers are in the box!

- Suppose our sample of size 100 has an average of 10.2 and an SD of 0.8.
- Estimate the true average of the tickets to be 10.2 but also give a plus or minus on the estimate.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.




We don't know what numbers are in the box!

- Suppose our sample of size 100 has an average of 10.2 and an SD of 0.8.
- Estimate the true average of the tickets to be 10.2 but also give a plus or minus on the estimate.
- Estimate $SD_{\text{box}} = 0.8$.
- $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.8}{\sqrt{100}} = 0.08$.

Example: Box Contents Unknown

- Take a random sample of size $n = 100$ from the box below.



We don't know what numbers are in the box!


- Suppose our sample of size 100 has an average of 10.2 and an SD of 0.8.
- Estimate the true average of the tickets to be 10.2 but also give a plus or minus on the estimate.
- Estimate $SD_{\text{box}} = 0.8$.
- $SE_{\text{av}} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.8}{\sqrt{100}} = 0.08$.
- If we assume that the samples follow a normal curve, then

$$10.2 \pm 2 \cdot (0.08) = 10.2 \pm 0.16$$

is 95% confidence interval on the average of the box.

Inferential Statistics

- In real situations, we typically don't know the contents of the box!
- That is where **inferential statistics** is used.



Confidence Intervals

- In real situations, we typically don't know the contents of the box!

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .
- Compute the average and SD of the sample.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .
- Compute the average and SD of the sample.
- Use the SD of the sample to approximate the SD of the box.

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .
- Compute the average and SD of the sample.
- Use the SD of the sample to approximate the SD of the box.
- Then compute the SE_{av}

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .
- Compute the average and SD of the sample.
- Use the SD of the sample to approximate the SD of the box.
- Then compute the SE_{av}
- The 95% confidence interval is:

$$(\text{observed average}) \pm 2 \cdot (\text{SE estimate})$$

Confidence Intervals

- In real situations, we typically don't know the contents of the box!
- We can take a sample and compute a **confidence interval** on the average of the numbers in the box.
- This gives an estimate of the average and puts a plus-or-minus on the estimate.

$$EV_{av} = AV_{\text{box}} \quad SE_{av} = \frac{SD_{\text{box}}}{\sqrt{n}}$$

- Take a random sample of size n .
- Compute the average and SD of the sample.
- Use the SD of the sample to approximate the SD of the box.
- Then compute the SE_{av}
- The 95% confidence interval is:

$$(\text{observed average}) \pm 2 \cdot (\text{SE estimate})$$

- Prior to taking the sample, we are 95% confident that this procedure will give an interval that contains the true average.

Example Page 415

- A city has 25,000 residents.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - Each ticket has the income of one resident.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - Each ticket has the income of one resident.
- Observed average: $\$62,396,714/1000 = \$62,400$.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - Each ticket has the income of one resident.
- Observed average: $\$62,396,714/1000 = \$62,400$.
- SD of the sample: \$53,000 (given).
- Estimated average income: \$62,400.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - Each ticket has the income of one resident.
- Observed average: $\$62,396,714/1000 = \$62,400$.
- SD of the sample: \$53,000 (given).
- Estimated average income: \$62,400.
- Now compute a \pm on this estimate:
 - $SD_{\text{box}} \approx \$53,000$.
 - $SE_{\text{av}} = \$53,000/\sqrt{1000} \approx \$1,700$.

Example Page 415

- A city has 25,000 residents.
- Estimate the average income of the residents by taking a random sample.
- Box model:
 - 25,000 numbered tickets
 - Each ticket has the income of one resident.
- Observed average: $\$62,396,714/1000 = \$62,400$.
- SD of the sample: $\$53,000$ (given).
- Estimated average income: $\$62,400$.
- Now compute a \pm on this estimate:
 - $SD_{\text{box}} \approx \$53,000$.
 - $SE_{\text{av}} = \$53,000/\sqrt{1000} \approx \$1,700$.
- 95% confidence interval:

$$\$53,000 \pm 2 \cdot (1,700) = \$53,000 \pm \$3,400$$