

# Math 207: Statistics

## Chapter 20: Chance Error in Sampling

Population (parameters)

Sample (statistics)

Dr. Ralph Wojtowicz

Mathematics Department



**SHENANDOAH**  
UNIVERSITY



## 1 EV, SE and the Central Limit Theorem

## 2 Examples

- Example I
- Example II
- Example III

# Expected Value, Standard Error, Central Limit Theorem

- Many statistics problems are modeled as samples from a box of numbered tickets.
- Solution procedure:
  - Formulate a box model.
  - Compute the average and SD of the contents of the box.
  - Determine if you are computing a sum or average (% in a 0/1 box is an average).
  - Use the appropriate formulas to compute the EV and SE for the sample.
  - Use the normal curve to compute chances of the sample being in a specified range.



## Example (I)

- Running a political poll is like sampling from a box of number tickets.

## Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

## Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?

# Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

$$\boxed{253,785 \text{ } \boxed{0} \text{ s } 433,211 \text{ } \boxed{1} \text{ s}}$$

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$  and  $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$

# Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$  and  $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$ .



# Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$  and  $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$ .
- Compute the  $z$  scores for 61% and 65%:  
 $z_1 = (61 - 63.1)/4.83 = -0.435$   
 $z_2 = (65 - 63.1)/4.83 = 0.393$

# Example (I)

- Running a political poll is like sampling from a box of number tickets.
- Take a random sample of size  $n = 100$  from the box below.

253,785	0	s	433,211	1	s
---------	---	---	---------	---	---

- What is the chance that the percent of 1s is between 61% and 65%?
- $AV_{\text{box}} = \frac{433,211}{433,211+253,785} \approx 0.631$  and  $SD_{\text{box}} = (1 - 0)\sqrt{0.631(1 - 0.631)} \approx 0.483$
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{100}} = 0.0483 = 4.83\%$ .
- Compute the  $z$  scores for 61% and 65%:  
 $z_1 = (61 - 63.1)/4.83 = -0.435$   
 $z_2 = (65 - 63.1)/4.83 = 0.393$
- Area under the normal curve:  $\text{pnorm}(0.393) - \text{pnorm}(-0.435) = 0.321 = 32.1\%$

## Example (II)

- What if we increase the sample size to 1000?

## Example (II)

- What if we increase the sample size to 1000?
- $AV_{\text{box}} \approx 0.631$  and  $SD_{\text{box}} \approx 0.483$  don't change.

## Example (II)

- What if we increase the sample size to 1000?
- $AV_{\text{box}} \approx 0.631$  and  $SD_{\text{box}} \approx 0.483$  don't change.
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{1000}} = 0.0153 = 1.53\%$ .

## Example (II)



- What if we increase the sample size to 1000?
- $AV_{\text{box}} \approx 0.631$  and  $SD_{\text{box}} \approx 0.483$  don't change.
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{1000}} = 0.0153 = 1.53\%$ .
- Compute the z scores for 61% and 65%:  
 $z_1 = (61 - 63.1)/1.53 = -1.37$   
 $z_2 = (65 - 63.1)/1.53 = 1.24$

## Example (II)

- What if we increase the sample size to 1000?
- $AV_{\text{box}} \approx 0.631$  and  $SD_{\text{box}} \approx 0.483$  don't change.
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{1000}} = 0.0153 = 1.53\%$ .
- Compute the z scores for 61% and 65%:  
 $z_1 = (61 - 63.1)/1.53 = -1.37$   
 $z_2 = (65 - 63.1)/1.53 = 1.24$
- Area under the normal curve:  $\text{pnorm}(1.24) - \text{pnorm}(-1.37) = 0.808 = 80.8\%$

## Example (II)

- What if we increase the sample size to 1000?
- $AV_{\text{box}} \approx 0.631$  and  $SD_{\text{box}} \approx 0.483$  don't change.
- $EV_{\%} = AV_{\text{box}} = 0.631 = 63.1\%$  and  $SE_{\%} = \frac{SD_{\text{box}}}{\sqrt{n}} \approx \frac{0.483}{\sqrt{1000}} = 0.0153 = 1.53\%$ .
- Compute the z scores for 61% and 65%:  
 $z_1 = (61 - 63.1)/1.53 = -1.37$   
 $z_2 = (65 - 63.1)/1.53 = 1.24$
- Area under the normal curve:  $\text{pnorm}(1.24) - \text{pnorm}(-1.37) = 0.808 = 80.8\%$





## Example (III)

- Suppose we roll 100 dice.

## Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?

## Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?
- The sample is like the sum of 100 draws from:

1	2	3	4	5	6
---	---	---	---	---	---

## Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?
- The sample is like the sum of 100 draws from:

1
2
3
4
5
6

- $AV_{\text{box}} = \frac{1+2+\dots+6}{6} = 3.5$  and  $SD_{\text{box}} = \sqrt{\frac{(1-3.5)^2 + (2-3.5)^2 + \dots + (6-3.5)^2}{6}} \approx 1.71$

## Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?
- The sample is like the sum of 100 draws from:

1 2 3 4 5 6

- $AV_{\text{box}} = \frac{1+2+\dots+6}{6} = 3.5$  and  $SD_{\text{box}} = \sqrt{\frac{(1-3.5)^2+(2-3.5)^2+\dots+(6-3.5)^2}{6}} \approx 1.71$
- $EV_{\text{sum}} = n \cdot AV_{\text{box}} = 350$  and  $SE_{\text{sum}} = \sqrt{n} \cdot SD_{\text{box}} \approx 17.1$ .

# Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?
- The sample is like the sum of 100 draws from:

1	2	3	4	5	6
---	---	---	---	---	---

- $AV_{\text{box}} = \frac{1+2+\dots+6}{6} = 3.5$  and  $SD_{\text{box}} = \sqrt{\frac{(1-3.5)^2+(2-3.5)^2+\dots+(6-3.5)^2}{6}} \approx 1.71$
- $EV_{\text{sum}} = n \cdot AV_{\text{box}} = 350$  and  $SE_{\text{sum}} = \sqrt{n} \cdot SD_{\text{box}} \approx 17.1$ .
- Compute the z scores for 375:  
 $z = (375 - 350)/17.1 = 1.46$

## Example (III)

- Suppose we roll 100 dice.
- What is the chance that the total will be greater than 375?
- The sample is like the sum of 100 draws from:

$$\boxed{1} \boxed{2} \boxed{3} \boxed{4} \boxed{5} \boxed{6}$$

- $AV_{\text{box}} = \frac{1+2+\dots+6}{6} = 3.5$  and  $SD_{\text{box}} = \sqrt{\frac{(1-3.5)^2+(2-3.5)^2+\dots+(6-3.5)^2}{6}} \approx 1.71$
- $EV_{\text{sum}} = n \cdot AV_{\text{box}} = 350$  and  $SE_{\text{sum}} = \sqrt{n} \cdot SD_{\text{box}} \approx 17.1$ .
- Compute the z scores for 375:  
 $z = (375 - 350)/17.1 = 1.46$
- Area under the normal curve:  $1 - \text{pnorm}(1.46) \approx 0.0721 = 7.21\%$