

An updated systematic overview of triple combination therapy in antiretroviral-naïve HIV-infected adults

John A. Bartlett^a, Michael J. Fath^b, Ralph DeMasi^c, Ashwaq Hermes^b, Joseph Quinn^d, Elsa Mondou^d and Franck Rousseau^d

Objective: To compare the effectiveness of three drug combination antiretroviral therapy (ART) in treatment-naïve HIV-infected persons, and identify the predictors of responses.

Design and methods: Overview of trials identified by searching public domain publications and conference presentations. The three-drug combination therapy was defined as two nucleoside reverse transcriptase inhibitors (NRTI) or nucleotide and NRTI, and either: (1) a protease inhibitor (PI); (2) a non-nucleoside RTI (NNRTI); (3) a third NRTI; or (4) a ritonavir-boosted PI (BPI). Week 24 and 48 results for the proportions of patients with plasma HIV RNA levels < 400 and < 50 copies/ml, and change in CD4+ cell counts were recorded.

Results: Fifty-three trials met the entry criteria, and enrolled 14 264 patients into 90 treatment arms. Overall 55% of patients had plasma HIV RNA levels < 50 copies/ml at week 48 and this percentage increased with later publication dates. In unadjusted pairwise comparisons at week 48, significantly greater percentages of patients receiving NNRTI (64%) and BPI (64%) had RNA < 50 copies/ml than NRTI (54%) or PI (43%), and CD4+ cell count increases were significantly greater in the BPI group (+200 cells/ μ l) than the PI (+179), NNRTI (+173), or NRTI (+161) groups. Pill count and percentage of patients with week 48 plasma HIV RNA levels < 50 copies/ml were correlated in the univariate analysis ($P=0.0053$; $r=-0.323$), but pill count was not a significant predictor in the multivariate analyses. Drug class and baseline CD4+ cell counts were significant predictors, but explained only a modest amount of the treatment effect, ($R^2=0.355$).

Conclusions: NNRTI and BPI-containing regimens offer superior virologic suppression over 48 weeks, supporting existing guidelines for the choice of initial ART. Pill count was not a consistent predictor of virologic suppression.

© 2006 Lippincott Williams & Wilkins

AIDS 2006, **20**:2051–2064

Keywords: systematic overview, antiretroviral therapy, treatment-naïve

Introduction

Antiretroviral therapy with three agents provides potent suppression of plasma HIV RNA levels below detectable limits in many treated persons, with resulting immunologic improvements, decreases in HIV-related complications, and prolonged survival. Twenty individual antiretroviral drugs, four fixed dose two-drug combinations,

and two fixed dose three-drug combinations are currently available, resulting in 364 potential three-drug combinations for treatment. The ideal drug combinations will maximize antiretroviral potency, immunologic reconstitution, and quality of life while minimizing medication-related toxicities and the potential for drug resistance. Published guidelines utilize the currently available evidence base to make specific

From the ^aDuke University Medical Center, Durham, North Carolina, ^bAbbott Laboratories, Abbott Park, Illinois, ^cTrimeris Inc, Morrisville, North Carolina, and ^dGilead Sciences, Durham, North Carolina, USA.

Correspondence to John A. Bartlett, Box 3152, Durham, North Carolina, 27710, USA.

E-mail: jab5@duke.edu

Received: 3 March 2006; accepted: 26 June 2006.

recommendations regarding treatment [1,2], but obviously not all three-drug combinations can be compared directly in clinical trials.

Although prospective, randomized trials provide the most robust evidence for regimen comparisons, systematic overviews may assist in assessing the relative attributes of alternative initial treatment regimens. A previous systematic overview identified the need for improved potency in initial treatment regimens, a lack of differences in virologic suppression and CD4+ cell count increases between treatment regimens using unboosted protease inhibitors (PI), non-nucleoside reverse transcriptase inhibitors (NNRTI), and a third nucleoside reverse transcriptase inhibitor (NRTI) with two NRTI in three-drug regimens, and the predictive value of pill count for virologic responses [3]. However, since the publication of this systematic overview additional clinical trials have been completed evaluating novel classes of antiretroviral agents [boosted protease inhibitors (BPI), nucleotide reverse transcriptase inhibitors and fusion inhibitors], and simpler treatment regimens with lower pill counts due to improved formulations and fixed dose combinations.

This systematic overview was performed to re-examine the activity of initial three-drug combinations in antiretroviral-naïve persons with contemporary agents, the predictors of activity, and the temporal changes in activity over time.

Methods

Study selection

A search of public domain and recent conference presentations was conducted for the period from 1994 to July 2004. A MEDLINE search was undertaken with the following keywords: clinical trial, plasma HIV-1 RNA, HAART, antiretroviral therapy, and naïve. In addition, the package inserts for US Food and Drug Administration (FDA)-approved antiretroviral drugs were examined and listed trials were reviewed. Proceedings from the following conferences were reviewed: Conference on Retroviruses and Opportunistic Infections (CROI), the World AIDS Conference, the International AIDS Society (IAS) Conference on HIV Pathogenesis and Treatment, the European Conference on Clinical Aspects and Treatment of HIV Infection, the Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC), the International Conference on Drug Therapy in HIV Infection (ICDT), and the Annual Meeting of the Infectious Diseases Society of America (IDSA). Triple combination therapy was defined as either two nucleoside or a nucleoside and a nucleotide reverse transcriptase inhibitor and either: (1) a protease inhibitor (PI); (2) a non-nucleoside reverse transcriptase inhibitor (NNRTI); (3) a third nucleoside reverse transcriptase

inhibitor (NRTI); or (4) a ritonavir-boosted protease inhibitor (BPI). Regimens were assigned to these four categories based upon the class of the third agent in the regimen, (PI, NNRTI, NRTI and BPI; the inclusion of BPI-containing regimens represents an addition since the previous overview). Trials were included if they were at least 24 weeks in duration, had at least 30 patients per treatment arm, included chronically HIV-infected patients who were treatment-naïve or had very limited prior exposure to antiretroviral therapy (< 2 weeks prior NRTI exposure), and reported the percentages of patients with undetectable plasma HIV RNA levels (< 400 and < 50 copies/ml) using an intent-to-treat missing equals failure analysis, (ITT:M = F). Trials were excluded if they studied induction/maintenance strategies, acutely or recently infected patient populations (< 6 months since seroconversion), or experimental treatments which did not successfully receive US FDA approval.

Data collection

Following inclusion of a trial in the overview, all published or presented source documentation was reviewed for each trial, (journal citations, conference abstract books, posters, slides, package inserts, etc.). Information on each trial was abstracted on the following trial characteristics and results: (1) trial design: treatment regimens, open-label or blinded, randomization if used, control groups, and average daily pill count (defined as the number of pills or tablets/day including placebo) in the treatment regimen; (2) baseline characteristics: number of individuals enrolled, percentage male, race, age, log₁₀ plasma HIV RNA, and CD4+ cell counts; and (3) response rates: percentages of patients with plasma HIV RNA < 400 and < 50 copies/ml at 24 and 48 weeks in the ITT:M = F populations, and change in CD4+ cell counts at 24 and 48 weeks in the as-treated populations. The date of a study's first presentation or publication was defined as the 'publication date'. Publication dates were divided into the following intervals; prior to 1998, 1999–2000, 2001–2002 and 2003–2004. Abstractions were performed by one reviewer and were confirmed by a second; any discrepancies were reconciled by conference with the study team. If median values were unavailable then the mean was used. All trials reported plasma HIV RNA results with the Roche Amplicor assay (< 400 copies/ml) and the Roche Ultrasensitive assay (< 50 copies/ml) (Roche Diagnostics, Indianapolis, Indiana, USA) with the exception of the START I and II trials and the VIRGO I/II trials, where the results were reported as the percentage < 500 copies/ml with the Chiron bDNA assay (Chiron Corporation, Emeryville, California, USA).

Statistical methods

For plasma HIV RNA responses, the primary analysis utilized reported results for ITT:M = F population, and for changes in CD4+ cell counts, the primary analysis used reported results for the as-treated population. When

ITT:M=F results were unavailable, the ITT:M=F results were calculated using the number of patients with plasma HIV RNA undetectable at weeks 24 or 48 divided by the number of enrolled patients. Response rates for each of the four drug classes (PI, NNRTI, NRTI and BPI) were estimated with the number of treatment groups and patients contributing to the estimate. These resulting estimates were then compared by constructing confidence intervals (CI) for the differences in response rates between drug classes. Correlation analyses and weighted least squares multivariate linear regression analyses (MLR) using backwards stepwise selection procedures were used to assess the variability in treatment group response rates as a function of the following factors; baseline \log_{10} plasma HIV RNA level and CD4+ cell count, drug class, and average daily pill count. For the MLR analysis of categorical virologic endpoints, the natural log of the response rate was used. Statistical significance was determined using $\alpha=0.05$. Bubble plots were constructed, and the results for each treatment arm were weighted in direct proportion to the numbers of patients enrolled. Sensitivity analyses were conducted to evaluate different factors including dichotomous pill counts ($\leq 10/\text{day}$ versus $> 10/\text{day}$), analyses of continuous pill counts over these ranges (3–10/day and 11–22/day), and pill counts as predictors of response within each regimen category.

Results

Descriptive characteristics of included trials

Fifty-three trials met the inclusion criteria and are listed in Table 1, (representing the addition of 30 trials since the previously published overview). These 53 trials enrolled 14 264 patients into 90 independent treatment arms. The median baseline plasma HIV RNA level was $4.75 \log_{10}$ copies/ml and the median baseline CD4+ cell count was $315/\mu\text{l}$. Seventy-six percent of patients were male.

Design characteristics and regimens studied

Thirty-four trials were randomized and 19 included only one treatment arm. Thirty-two trials were open-label, 19 were double-blinded, and two were partially blinded. A total of 90 treatment arms were included in these 53 trials with 40 unique treatment regimens. NNRTI-containing regimens were most commonly studied ($n=38$), followed by PI-containing ($n=32$), NRTI-containing ($n=12$) and BPI-containing regimens ($n=8$). Efavirenz-containing regimens were most commonly studied; they were included in 20 different trials and 29 treatment regimens in 5798 individuals. In addition, efavirenz-containing regimens were combined with more NRTI combinations than any other drug, (nine stavudine and lamivudine, eight zidovudine and lamivudine, four abacavir and lamivudine, three stavudine and didanosine, two didanosine and lamivudine, two didanosine and emtricitabine, one

tenofovir and lamivudine). In comparison with the previous overview, there was a slight decrease in the representation of PI-containing regimens [15/31 regimens in 2001 (48%) versus 32/90 in 2005 (36%)], no difference in the representation of NNRTI-containing regimens [13/31 regimens in 2001 (41%) versus 38/90 in 2005 (42%)], a substantial increase in the number of efavirenz-containing regimens (five in 2001 versus 29 in 2005), and the addition of BPI-containing regimens. The median pill count was 10/day, with a range of 3–22/day.

Temporal trends in baseline characteristics and responses by publication date

In the previous overview, the median baseline plasma HIV RNA levels and CD4+ cell counts were $4.69 \log_{10}$ copies/ml and $375/\mu\text{l}$. In the current update, the values were $4.75 \log_{10}$ copies/ml ($P=0.10$) and $315/\mu\text{l}$ ($P=0.001$). Over the period of observation, the weighted mean responses for the percentage of patients with plasma HIV RNA levels < 50 copies/ml at week 48 steadily improved, (prior to 1998, 41%; 1999–2000, 50%; 2001–2002, 56%; 2003–2004, 64%), (Fig. 1). During the same periods of time the weighted mean responses for increases in CD4+ cell counts at week 48 were similar, (prior to 1998, +173 cells/ μl ; 1999–2000, +174 cells/ μl ; 2001–2002, +180 cells/ μl ; 2003–2004, +169 cells/ μl) (Fig. 1).

Virologic and immunologic responses

The virologic and immunologic responses at weeks 24 and 48 are shown in Table 2 with the number of treatment arms and patients, the weighted means, and the 95% CI. These responses are unadjusted for baseline characteristics, and are presented as the percentages with plasma HIV RNA levels below 400 and 50 copies/ml and the change from baseline in CD4+ cell counts by drug class. Overall, 55% of patients had plasma HIV RNA levels below 50 copies/ml at week 48 with a CD4+ cell count change from baseline of +176 cells/ μl . In unadjusted pairwise comparisons of the proportions of patients with plasma HIV RNA < 50 copies/ml at week 48, the BPI and NNRTI groups were each superior to the PI and NRTI groups (BPI versus PI, $P<0.001$; BPI versus NRTI, $P=0.017$; NNRTI versus PI, $P<0.001$; NNRTI versus NRTI, $P=0.001$), but were not significantly different from each other, (BPI versus NNRTI, $P=0.945$). The NRTI group was also superior to the PI group in pairwise comparisons ($P=0.001$). In pairwise comparisons of the unadjusted changes from baseline in CD4+ cell counts, the BPI group was superior to NNRTI, PI and NRTI groups (BPI versus NNRTI, $P<0.001$; BPI versus PI, $P=0.004$; BPI versus NRTI, $P<0.001$). The PI and NNRTI groups did not have significantly greater increases in CD4+ cell counts than the NRTI group, and were not significantly different from each other.

Figure 2 displays a bubble plot of the relationship between pill count and plasma HIV RNA levels < 50 copies/ml at

Table 1. Studies included in the analysis.

Study name/number	Class	Regimens	No patients	BL RNA	BL CD4
Study 096 Brett-Smith, 2002 [4]; Pollard, 2002 [6]; Brett-Smith, 2003 [5]	NNRTI	EFV + 3TC + d4T(ir)	76	4.65	285
Study 099 Brett-Smith, 2002 [4]; Pollard, 2002 [6]; Brett-Smith, 2003 [5]	NNRTI	EFV + 3TC + d4T(xr)	74	4.65	285
	NNRTI	EFV + 3TC + d4T(ir)	391	4.8	277
2NN van Leth, 2003 [8]; Van Leeuwen, [7] 2002; van Leth, [9] 2004	NNRTI	EFV + 3TC + d4T(xr)	392	4.8	277
	NNRTI	NVP qd + d4T + 3TC	220	4.7	200
ACTG 384 Shafer, 2003 [12]; Robbins, 2002 [10]; Robbins, 2002 [11]	NNRTI	NVP bid + d4T + 3TC	387	4.7	170
	NNRTI	EFV qd + d4T + 3TC	400	4.7	190
	NNRTI	EFV + ZDV + 3TC	155	4.9	278
ACTG 5095 Gulick, 2004 [14]; Gulick, 2003 [13]; NIAID, 2003 [15]; NIAID, 2003 [15]	NRTI	ZDV + 3TC + ABC	382	4.9	234
AI424-007 Sanne, 2003 [17]; Squires, 2001 [18]; Gatell, 2001 [16]	PI	ZDV (400) + ddI + d4T	78	4.65	357
AI424-008 Sanne, 2001 [24]; Cahn, 2001 [20]; Murphy, 2003 [21]; Pantaleo, 2001 [23]; Wood, 2004 [25]; Bristol-Myers Squibb Company, 2004 [19]; Murphy, 2003 [22]	PI	NFV + ddI + d4T	82	4.79	341
	PI	ZDV (400) + d4T + 3TC	181	4.74	294
	PI	NFV + d4T + 3TC	91	4.73	283
AI424-034 Squires, 2002 [28]; Delfraissy, 2002 [27]; Bristol-Myers Squibb Company, 2004 [26]	NNRTI	EFV qd + COM	405	4.8	321
AI454-148 Gathe, 1999 [29]; Bristol Myers Squibb, 2004 [26]	PI	ZDV qd + COM	405	4.8	321
	PI	NFV + ZDV + 3TC	253	4.69	340
AI454-152 Badaro, 2001 [30]; Gathe, 2001 [31]; Bristol Myers Squibb, 2004 [26]	PI	NFV + ddI + d4T	503	4.69	340
	PI	NFV + COM	253	4.71	411
ANRS 091 (MONTANA) Molina, 2000 [32]; Molina, 2001 [33]; Molina, 2000 [34]; Molina, 2003 [36]; Molina, 2000 [35]	PI	NFV + ddI-EC + d4T	258	4.71	411
	NNRTI	EFV + ddI + FTC	40	4.8	396
ANRS 12-04 Landman, 2002 [37]; Landman, 2003 [38]	NNRTI	EFV + ddI + 3TC	40	5.4	164
Atlantic Murphy, 1999 [39]; van Leeuwen, 2003 [40]	NNRTI	NVP + ddI + d4T arm 3	89	4.33	394
AVANTI 2 Avanti Study Group, 2000 [41]; Hill, 1998 [43]; Goebel, 1997 [42]	NRTI	3TC + ddI + d4T arm 2	109	4.2	396
	PI	IDV + ddI + d4T arm 1	100	4.3	417
	PI	IDV + ZDV + 3TC	52	4.7	281
AVANTI 3 Gartland, 2001 [45]; Clumeck, 1998 [44]	PI	NFV + ZDV + 3TC	53	5	295
CLASS Bartlett, 2002 [46]; Bartlett, 2004 [132]	BPI	APV/RTV + ABC + 3TC	96	4.8	296
CNA 30021(ZODIAC) Gazzard, 2003 [47]; Glaxo-Smith-Kline, 2005 [48]; Piliero, 2003 [49]	NNRTI	EFV + ABC + 3TC	97	4.9	307
	NRTI	ABC + d4T + 3TC	98	4.9	299
	NNRTI	EFV + ABC qd + 3TC	384	4.89	264
CNA 3003 (CNAA3003/CNAB3003) Fischl, 1999 [50]; Fischl, 1998 [51]	NNRTI	EFV + ABC + 3TC	386	4.89	259
	NRTI	ABC + ZDV + 3TC	87	4.5	473
CNA30024 Dejesus, 2003 [52]; Glaxo-Smith-Kline, 2005 [48]	NNRTI	EFV + ABC + 3TC	324	4.79	267
	NNRTI	EFV + ZDV + 3TC	325	4.79	258

Table 1 (continued)

Study name/number	Class	Regimens	No patients	BL RNA	BL CD4
CNA3005 (CNAB3005) Staszewski, 1999 [53]; Staszewski, 1999 [54]; Staszewski, 1999 [55]; Staszewski, 2001 [56]	NRTI	ABC + ZDV + 3TC	282	4.85	359
	PI	IDV + ZDV + 3TC	280	4.82	360
CNA3014 Vibhagool, 2001 [58]; Jordan J, [57] 2002; Vibhagool, [59] 2004	NRTI	ABC + COM	164	4.78	331
	PI	IDV + COM	165	4.82	299
CNAF3007 Matheron, [60] 2003	NRTI	ABC + COM	95	4.2	387
	PI	NFV + COM	91	4.1	449
COMBINE Podzamczar, 2002 [129]; Podzamczar, 2000 [62]; Podzamczar, 2001 [61]	NNRTI	NVP + COM	72	4.77	361
	PI	NFV + COM	70	4.81	351
Dart I Dejesus, 2003 [63]; Ward, 2004 [64]	NNRTI	EFV + ddI-EC + 3TC	65	4.8	311
DART II Brett-Smith, 2003 [5]; Ward, 2004 [64]; Jayaweera, 2004 [66]; Felizarta, 2004 [65]	NNRTI	d4T-XR + 3TC + EFV	70	4.54	351
DMP 266-005 Manion, 1999 [69]; Manion, 1999 [70]; Hicks, 1998 [68]; Haas, 1998 [67]	NNRTI	EFV + ZDV + 3TC	34	4.64	386
DMP266-006 Staszewski, 1999 [69]	NNRTI	EFV + ZDV + 3TC	154	4.77	345
	PI	IDV + ZDV + 3TC	148	4.79	341
DMP266-043 Luskin-Hawk, 1999 [72]; Cohen, 2000 [71]	NNRTI	EFV + d4T + 3TC	68	4.85	375
DMP266-044 Luskin-Hawk, 1999 [72]; Cohen, 2000 [71]	NNRTI	EFV + ddI + d4T	65	4.89	289
Domula Domula, 2002 [73]	NNRTI	EFV + COM	35	4.9	275
	NNRTI	EFV + d4T + 3TC	35	4.9	163
	NNRTI	EFV + ddI + d4T	38	4.9	165
	PI	IDV + d4T + 3TC	32	4.38	708
Earth-2 Garcia, 1999 [74]; Garcia, 2000 [75]	NNRTI	3TC qd + ZDV + EFV	278	4.64	340
EPV 20001 Dejesus, 2004 [76]	NNRTI	3TC bd + ZDV + EFV	276	4.69	386
	NRTI	ZDV + 3TC 300 qd + ABC 300 bid	50	4.8	380
	NRTI	ZDV + 3TC 150 bid + ABC 600 qd	51	4.8	380
	NRTI	ZDV + 3TC 150 bid + ABC 300 bid	50	4.8	380
FTC 301A Saag, 2002 [80]; Raffi, 2003 [79]; Cahn, 2003 [78]; Saag, 2002 [80]	NNRTI	EVF + ddI + FTC qd	286	4.9	282
	NNRTI	EFV + ddI + d4T	285	4.9	300
Gerstoft, Danish study Katzenstein, 2000 [81]; Gerstoft, 2003 [82]; Hill, 2004 [83]	NRTI	ABC + d4T + ddI	60	5.00	190
	PI	3TC + ZDV + SQV/r (400/400 bid)	60	5.00	152
GPO Bangkok Anekthananon, 2004 [84]	NNRTI	d4T + 3TC + NVP	101	5.40	59
GS 903 Gallant, 2002 [85]; Staszewski, 2003 [86]	NNRTI	EFV + 3TC + TDF + PLA	299	4.9	276
	NNRTI	EFV + d4T + 3TC + PLA	301	4.9	283
ICC-002 Skowron, 1997 [87]	NNRTI	NVP + ZDV + ddI	53		
	NRTI	ZDV + ddI + 3TC	53		
IMEA-01 Saimot, 1998 [88]	PI	RTV + ddI + d4T	36	4.86	236
INCAS Myers, 1996 [91]; Montaner, 1998 [89]; Montaner, 1998 [90]	NNRTI	NVP + ZDV + ddI	51	4.25	395
M/3331/0013C Wood, 1999 [92]; Wood, 2004 [25]	NNRTI	DLV + ZDV + 3TC	74	5.33	185
M02-418 Gathe, 2004 [93]; Podzamczar, 2003 [94]	BPI	LPV/RTV + FTC + TDF (qd)	115	4.8	214
	BPI	LPV/RTV + FTC + TDF (bid)	75	4.6	232
M97-720 (ABT 378/R) Murphy, 2001 [100]; Murphy, 2002 [99]; Eron, 1999 [95]; Hicks, 1999 [97]; White, 2001 [101]; Landay, 2002 [98]; Gulick, 2000 [96]	BPI	LPV/RTV + d4T + 3TC	68	4.9	301

Table 1 (continued)

Study name/number	Class	Regimens	No patients	BL RNA	BL CD4
M98-863 Walmsley, 2002 [134]; Luo, 2002 [104]; Cernohous, 2002 [102]; King, 2002 [103]	BPI	LPV/RTV + d4T + 3TC	326	5.01	232
Merck Study Schranz, 2001 [105]	PI	NFV + d4T + 3TC	327	4.98	232
NEAT (APV 30001) Nadler, 2003 [106]; Rodriguez-French, 2002 [107]; Rodriguez-French, 2004 [108]; Vertex Pharmaceuticals, 2004 [109]	BPI	IDV-RTV + d4T + 3TC	89	5.03	238
	PI	fosAPV + 3TC + ABC	166	4.82	214
Ozcombo I Carr, 1999 [110]	PI	NFV + ABC + 3TC	83	4.85	212
	PI	IDV + ZDV + 3TC	35	5.01	267
	PI	IDV + d4T + 3TC	34	5.21	313
	PI	IDV + ddl + d4T	37	5	277
	PI	APV + ZDV + 3TC	116	4.64	442
PROAB 3001, 141W94 Goodgame, [111] 1999; Goodgame, [112] 2000	BPI	fosAPV/RTV + 3TC + ABC	322	4.78	166
SOLO (APV 30002, GW433908) Schurmann, 2002 [113]; Gathe, 2004 [93]	PI	NFV + 3TC + ABC	327	4.83	177
START I Squires, 1999 [114]; Squires, 2000 [115]	PI	IDV + d4T + 3TC	101	4.59	408
START II Eron, 1998 [116]; Murphy, 2003 [21]; Eron, 2000 [117]	PI	IDV + ZDV + 3TC	103	4.47	391
	PI	IDV + ZDV + 3TC	103	4.59	409
SUN Sension, 2001 [118]	PI	IDV + d4T + ddl	102	4.47	433
VIRGO Raffi, 1999 [119]; Reliquet, 2001 [120]	PI	SQV + ZDV + 3TC	42	4.8	419
	NNRTI	NVP + ddl + d4T	100	4.7	413

NRTI, nucleoside reverse transcriptase inhibitor; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI, protease inhibitor; BPI, boosted PI; ir, immediate release; xr, extended release; qd, once daily; bid, twice daily; 3TC, lamivudine; d4T, stavudine; d4T-XR, extended release stavudine; ddl, didanosine; ddi-EC, enteric coated didanosine; ABC, abacavir; APV, amprenavir; COM, fixed dose combination zidovudine and lamivudine; EFV, efavirenz; fosAPV, fos-amprenavir; FTC, emtricitabine; LPV, lopinavir; NFV, nelfinavir; NVP, nevirapine; PLA, placebo; RTV, ritonavir; TDF, tenofovir; ZDV, zidovudine.

week 48. In a univariate analysis this relationship was significant ($P=0.0053$, $r=-0.323$). However, when evaluated using multivariable linear regression analysis, this relationship was no longer significant. Additional analyses were performed to adjust for baseline differences and identify predictors of virologic and immunologic responses (Table 3). Lower baseline CD4+ cell counts were associated with increased percentages of plasma HIV RNA < 50 copies/ml at weeks 24 and 48, and greater CD4+ cell count increases at week 48. Baseline plasma HIV RNA levels were not consistently associated with plasma HIV RNA responses or CD4+ cell count increases. Drug class was significantly associated with plasma HIV RNA levels < 400 copies/ml at week 24,

plasma HIV RNA levels < 50 copies/ml at weeks 24 and 48, and with increases in CD4+ cell counts at week 48. Lower pill count was associated with higher percentages of plasma HIV RNA levels < 50 copies/ml at week 24 but not with plasma HIV RNA < 50 copies/ml at week 48. Higher pill counts were associated with a greater percentage of patients having plasma HIV RNA < 400 copies/ml at week 48. Publication interval was only predictive of plasma HIV RNA levels < 400 copies/ml at week 24. The modeling of all factors had limited predictive value with a maximum R^2 value of 0.436 for plasma HIV RNA levels < 400 copies/ml at week 48, and an R^2 value of 0.355 for plasma HIV RNA levels < 50 copies/ml at week 48.




Fig. 1. Week 48 virologic and immunologic responses by publication date. Weighted mean responses of (a) percentages of patients with plasma HIV RNA levels < 50 copies/ml and (b) increases in CD4+ cell counts/μl from baseline at week 48, analyzed by publication date, (defined as the date when a study was first presented or published).

Table 2. Virologic and immunologic results at 24 and 48 weeks.

Drug class	Variable	Week 24 results			Week 48 results		
		Percentage of subjects with RNA < 400	Percentage of subjects with RNA < 50	Change in CD4 cell count	Percentage of subjects with RNA < 400	Percentage of subjects with RNA < 50	Change in CD4 cell count
PI	No. treatment arms	20	18	17	27	32	32
	No. subjects	3564	2572	2674	4474	4735	4602
	Weighted mean	66	49	136	57	43	179
	(95% CI)	62, 69	45, 52	121, 150	54, 60	40, 46	167, 191
BPI	No. treatment arms	4	5	4	4	6	6
	No. subjects	8.5	559	347	812	1002	1002
	Weighted mean	78	64	103	73	64	200
	(95% CI)	71, 85	56, 71	99, 162	66, 80	57, 70	189, 211
NNRTI	No. treatment arms	18	23	20	28	33	36
	No. subjects	2712	4286	3361	5476	6544	6705
	Weighted mean	77	62	130	73	64	173
	(95% CI)	74, 81	59, 65	119, 140	70, 75	61, 66	164, 183
NRTI	No. treatment arms	6	4	6	7	8	8
	No. subjects	629	573	629	1175	1277	1166
	Weighted mean	66	56	106	64	54	161
	(95% CI)	59, 74	48, 64	91, 121	58, 69	48, 59	141, 181
Overall	No. treatment arms	48	50	47	66	79	82
	No. subjects	7710	8100	7011	11937	13558	13475
	Weighted mean	71	57	130	66	55	176
	(95% CI)	69, 73	55, 60	123, 137	64, 68	54, 57	170, 183

NRTI, nucleoside reverse transcriptase inhibitor; NNRTI, non-nucleoside reverse transcriptase inhibitor; PI, protease inhibitor; BPI, boosted PI; CI, confidence interval.

Further analyses were undertaken to explore the relationship between pill count and the weighted mean response of week 48 plasma HIV RNA levels < 50 copies/ml. When pill count was analyzed as a continuous variable in all patients with week 48 results (79 treatment arms and 13 558 participants), no significant overall relationship was identified. Pill counts were next analyzed by drug class dichotomously

(≤ 10 pills/day and > 10 pills/day), continuously within these two categories (≤ 10 or > 10 pills/day), when one study with disproportionately low response rates (AI424-034) was excluded in a sensitivity analysis, and as a continuous variable within each drug class. In six of nine secondary analyses no significant association was identified; in the analyses examining dichotomous categories (≤ 10 versus > 10) and within the NRTI




Fig. 2. Bubble plot displaying the relationship between the percentages of subjects with plasma HIV RNA levels < 50 copies/ml at week 48 and pill counts. The size of the bubble reflects the numbers of subjects, and the bubbles are color-coded by regimen type. Univariate analysis demonstrated a significant relationship between lower pill counts and virologic responses, ($P = 0.0053$; $r = -0.323$). However, after adjustment in the multivariate linear regression, the correlation between pill count and response was no longer significant. ●, nucleoside reverse transcriptase inhibitor; ●, non-nucleoside reverse transcriptase inhibitor; ●, protease inhibitor; ●, boosted protease inhibitor.

Table 3. Multivariate linear regression analysis: results at 24 and 48 weeks.

Factor	Week 24 results			Week 48 results		
	Percentage of subjects with RNA < 400	Percentage of subjects with RNA < 50	Change in CD4 cell count	Percentage of subjects with RNA < 400	Percentage of subjects with RNA < 50	Change in CD4
Baseline CD4+ Cell count	NS	−0.617	NS	NS	−0.335	−25.785
Baseline HIV RNA level	NS	−0.584	NS	+0.382	NS	NS
Pill count	NS	+0.294	NS	−0.182	NS	NS
Drug class	−0.031	−0.158	NS	NS	−0.115	5.879
Publication interval	0.126	NS	NS	NS	NS	NS
Model R ²	0.445	0.334	0.000	0.436	0.355	0.108

NS, not statistically significant at $\alpha = 0.05$; for significant factors, numbers reported are parameter estimates from multiple regression analyses. Note that a (+) indicates a positive correlation between the variable and improved response, while (−) indicates a negative correlation between the variable and improved response.

class results favored lower pill counts, and in the PI class the results favored higher pill counts.

Discussion

The results of this expanded overview offer important new observations on the success of contemporary antiretroviral treatment regimens in trials studying ART-naïve populations. The percentages of patients achieving plasma HIV RNA levels < 50 copies/ml at 48 weeks is increasing over time, and in reports from most recent trials, more than 80% of patients may reach this threshold [121–123]. The improving virologic responses were seen despite the enrollment of patient populations with lower CD4+ cell counts at entry, and surprisingly, lower entry CD4+ cell counts were associated with improved responses in some analyses. Drug class was the most significant predictor of regimen potency, with the best responses achieved by NNRTI and BPI-containing regimens. Pill count was not consistently identified as a predictor of virologic response, in contrast to our previously published overview.

The improving response rates over time are encouraging evidence for the identification of new treatment regimens and their strategic use in ART-naïve patients. Similar observations have been reported in numerous observational cohorts [124–127] and a portion of the declining risk of virologic failure over time could be attributed to the choice of starting regimen [127]. The multivariable linear regression in our study offered the opportunity to assess the predictive value of publication date for plasma HIV RNA levels < 50 copies/ml in comparison with other factors such as drug class. The results demonstrate that improving virologic success is due to the use of more potent drug classes, and not simply a function of publication date. The superiority of NNRTI and BPI-containing regimens over PI and NRTI-containing regimens in suppressing plasma HIV RNA levels below 50 copies/ml observed in this overview is supported by the results of smaller randomized clinical trials [128–131] and provides an additional evidence base consistent with

current therapeutic guidelines. Greater CD4+cell count increases were seen in patients receiving BPI-containing than NNRTI, PI and NRTI-containing regimens after 48 weeks. Importantly, the superior CD4+ cell count increases were seen in both the unadjusted analyses and were associated with drug class in the multivariable linear regression analyses. The potential mechanisms for the greater CD4+cell count increases on BPI-containing regimens are uncertain, and must be confirmed in additional randomized studies, especially those directly comparing BPI and NNRTI-containing regimens given the apparent lack of differences between them in observed antiretroviral potency.

Additional factors such as drug-related toxicities and the resistance consequences of first regimen failure may also influence the choice of first line treatment regimens. This analysis considered drug-related toxicities only to the extent that toxicities resulted in treatment discontinuation, and therefore affected the ITT:M=F results. Certainly, many of the toxicities of NNRTI and BPI-containing regimens [132–134] may be non-overlapping, thus allowing for individualization of treatment choices. Nonetheless, the most robust recommendations on drug-related toxicities should come from directly comparative, randomized clinical trials. Drug resistance appears to be less commonly reported from randomized clinical trials among patients receiving BPI-containing regimens, interestingly in both the protease and reverse transcriptase genes. A recent systematic overview identified a lower resistance cost associated with the use of BPI versus NNRTI-containing regimens [135]. However, once again these analyses cannot substitute for resistance results generated from randomized clinical trials comparing the two approaches.

In contrast to our previously published overview, pill count was not identified as a significant predictor of plasma HIV RNA levels below 50 copies/ml at week 48 in the multivariable analysis. However, pill count should not be forgotten as a potential factor influencing the outcome of treatment interventions; pill count was predictive in the current univariable and dichotomous

(≤ 10 versus > 10) analyses. Why did the current analysis not duplicate the results from the previous overview? A number of factors must be considered in assessing the contrasting results. The current overview is much larger with more participants, more trials, more unique treatment regimens, and a broader range of pill counts, and as a result, findings from the current overview are more robust. The current overview included a greatly expanded number of patients receiving NNRTI-containing regimens and BPI-containing regimens, identified as the most potent drug classes and therefore diminishing the impact of pill counts.

There are a number of important limitations to the current overview. Overviews cannot replace randomized clinical trials, which provide the most robust evidence to guide clinical practice. The data used for this overview were generated from population-based results and not individual patients, and therefore the overview may lack the ability to discern some predictors of treatment response, especially baseline characteristics. Analyses focused on the contribution of a third agent to regimen success, and it is possible that the nucleoside or nucleotide reverse transcriptase inhibitor component may affect these results. A recently completed trial highlighted the potential importance of these agents [123]. In the multivariable logistic regression analysis, the R^2 value for the model predicting plasma HIV RNA levels < 50 copies/ml at week 48 was 0.355, reflecting the importance of other unmeasured factors in contributing to treatment responses. In addition, some trials which prematurely closed such as ESS 30009 are not analyzed in this overview [136]. Finally, the time span of included trials now exceeds 10 years, and during this period substantial improvements in care have been implemented beyond the choice of treatment regimen such as improved care access and delivery, monitoring, adherence education, and support for comorbidities.

Efavirenz was studied in the greatest number of trials, in the largest number of patients, and in more diverse regimens than any other single agent. Efavirenz-containing regimens have provided impressive results in recent trials with suppression of plasma HIV RNA levels below 50 copies/ml in approximately 80% of patients in ITT:M=F analysis, [121–123]. However, NNRTI-containing regimens have been associated with higher resistance costs when virologic failure occurs [135], which may compromise response to subsequent treatment regimens. Important results will be generated from randomized trials which are currently underway comparing directly NNRTI-containing versus BPI-containing regimens (ACTG 5142, 5202).

In summary, this overview has identified improving temporal trends in virologic responses, superior virologic responses for NNRTI and BPI-containing regimens, and

an inconsistent and less pivotal role for lowered pill counts.

Acknowledgements

Donna Walczak and Leslie Groves contributed through data collection, and Donna Walczak and Vladimir Mats performed many of the statistical analyses.

Sponsorship: This research was supported by Abbott Laboratories. J.A.B. is supported by NIH Awards K-24 AI01744, AIDS Clinical Trials Group grant A1136156, CFAR grant P30 AI64518, ISAAC U01 AI062563, and the Fogarty International Center D43 TW06732.

References

1. Deng SC, Chen MY, Hsieh SM, Sheng WH, Hsiao CF, Hung CC, Chang SC. **Response to efavirenz plus two nucleoside reverse-transcriptase inhibitors in patients with advanced stage human immunodeficiency virus-1 infection in Taiwan.** *J Microbiol Immunol Infect* 2003; **36**:10–14.
2. Yeni PG, Hammer SM, Hirsch MS, Saag MS, Schechter M, Carpenter CC, et al. **Treatment for adult HIV infection: 2004 recommendations of the International AIDS Society–USA Panel.** *JAMA* 2004; **292**:251–265.
3. Bartlett JA, DeMasi R, Quinn J, Moxham C, Rousseau F. **Overview of the effectiveness of triple combination therapy in antiretroviral-naïve HIV-1 infected adults.** *AIDS* 2001; **15**:1369–1377.
4. Brett-Smith H, Reynolds L. **48 week efficacy results: stavudine extended-release (XR) formulation as compared to stavudine immediate-release (IR) formulation.** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, California, 2002.
5. Brett-Smith H, Reynolds L, Bessen L, Rutkiewicz V. **Two-year analysis of stavudine extended-release/prolonged-release capsules (XR/PRC) as compared to stavudine immediate-release (IR): efficacy and safety.** In: *43rd Interscience Conference on Antimicrobial Agents and Chemotherapy*, Chicago, 2003.
6. Pollard R, Ive P, Farthing C, Whelden M, Thompson S, Brett-Smith H. **Stavudine XR vs stavudine IR as part of potent antiretroviral combination therapy: 24-week safety and antiviral efficacy.** In: *Ninth Conference on Retroviruses and Opportunistic Infections*. Seattle, WA; 2002.
7. Van Leeuwen R. **Design of the 2NN study: a large-scale, randomized comparison of nevirapine and efavirenz.** In: *Sixth International Congress on Drug Therapy in HIV Infection*, Glasgow, 2002.
8. van Leth F, Hassink E, Phanuphak P, Miller S, Gazzard B, Cahn P, et al. **Results of the 2NN Study: a randomized comparative trial of first-line antiretroviral therapy with regimens containing either nevirapine alone, efavirenz alone or both drugs combined, together with stavudine and lamivudine.** In: *Tenth Conference on Retroviruses and Opportunistic Infections*, Boston, 2003.
9. van Leth F, Phanuphak P, Ruxrungtham K, Baraldi E, Miller S, Gazzard B, et al. **Comparison of first-line antiretroviral therapy with regimens including nevirapine, efavirenz, or both drugs, plus stavudine and lamivudine: a randomised open-label trial, the 2NN Study.** *Lancet* 2004; **363**:1253–1263.
10. Robbins G, Shafer R, Smeaton L, De Gruttola V, Pettinelli C, Snyder S, et al. **Antiretroviral strategies in naïve HIV+ subjects: comparison of 4-drug versus sequential 3-drug regimens (ACTG 384).** In: *XIV International AIDS Society Conference*, Barcelona, Spain, International AIDS Society, 2002 [abstract LbOr20B].
11. Robbins G, Shafer R, Smeaton L, De Gruttola V, Pettinelli C, Snyder S, et al. **Antiretroviral strategies in naïve HIV+ subjects: comparison of sequential 3-drug regimens (ACTG 384).** In: *XIV International AIDS Society Conference*, Barcelona, Spain, 2002 [abstract LbOr20A].

12. Shafer RW, Smeaton LM, Robbins GK, De Gruttola V, Snyder SW, D'Aquila RT, et al. **Comparison of four-drug regimens and pairs of sequential three-drug regimens as initial therapy for HIV-1 infection.** *N Engl J Med* 2003; **349**:2304–2315.
13. Gulick R, Ribaud HJ, Shikuma CM, Lustgarten S, Squires KE, Meyer WA, et al. **Triple-nucleoside regimens versus efavirenz-containing regimens for the initial treatment of HIV-1 infection.** *N Engl J Med* 2004; **350**:1850–1861.
14. Gulick RM, Meibohm A, Havlir D, Eron JJ, Mosley A, Chodakewitz JA, et al. **Six-year follow-up of HIV-1-infected adults in a clinical trial of antiretroviral therapy with indinavir, zidovudine, and lamivudine 1.** *AIDS* 2003; **17**:2345–2349.
15. Gulick RM, Ribaud HJ, Shikuma CM, Lalama C, Schackman BR, Meyer WA, et al. **Three-vs four-drug antiretroviral regimens for the initial treatment of HIV-1 infection: A randomized controlled trial.** *JAMA* 2006; **296**:769–781.
16. Gatell J, Squires K, Piliero P, Sanne I, Wood R, Shelton S, et al. **Atazanavir (BMS-232632): absence of serum lipid changes after 48 weeks of treatment-naïve HIV-positive subjects (Trial A1424-007).** In: *Eighth European Conference on Clinical Aspects and Treatment of HIV Infection*, Athens, Greece, 2001.
17. Sanne I, Piliero P, Squires K, Thiry A, Schnittman S. **Results of a phase 2 clinical trial at 48 weeks (A1424-007): a dose-ranging, safety, and efficacy comparative trial of atazanavir at three doses in combination with didanosine and stavudine in antiretroviral-naïve subjects 20.** *J Acquir Immune Defic Syndr* 2003; **32**:18–29.
18. Squires K, Gatell J, Piliero P, Sanne I, Wood R, Schnittman SM. **48-week safety and efficacy results from a phase II study of a once-daily HIV-1 protease inhibitor (PI), BMS-232632.** In: *Eighth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL; 2001.
19. Bristol-Myers Squibb Company. *Reyataz (atazanavir sulfate) capsules prescribing information.* New York: Bristol-Myers Squibb Company; 2004.
20. Cahn P, Percival L, Phanuphak P, Sanne I, Kelleher T, Giordano M. **Phase II 24-week data from study A1424-008: Comparative results of BMS-232632, stavudine, Lamivudine as HAART for treatment-naïve HIV-infected patients.** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, 2001.
21. Murphy R, Pokrovsky V, Rozenbaum W, Wood R, Percival L, Odesloo L, Giordano M. **Long-term efficacy and safety of atazanavir with stavudine and lamivudine in patients previously treated with nelfinavir or ATV: 108-week results of BMS study 008/044.** In: *Tenth Conference on Retroviruses and Opportunistic Infections*, Boston, MA, 2003.
22. Murphy RL, Sanne I, Cahn P, Phanuphak P, Percival L, Kelleher T, Giordano M. **Dose-ranging, randomized, clinical trial of atazanavir with lamivudine and stavudine in antiretroviral-naïve subjects: 48-week results.** *AIDS* 2003; **17**:2603–2614.
23. Pantaleo G, Sanne I, Cahn P, Percival L, Phanuphak P, Kelleher T, et al. **Atazanavir (BMS-232632): 48-week safety and efficacy vs nelfinavir, each in combination with stavudine and lamivudine, in treatment-naïve, HIV-positive subjects (A1424-008).** In: *Eighth European Conference on Clinical Aspects and Treatment of HIV – Infection*, Athens, Greece, 2001.
24. Sanne I, Cahn P, Percival L, Phanuphak P, Kelleher T, Giordano M, Pantaleo G. **Comparative results (Phase II 48-Week): BMS-232632, stavudine, lamivudine as HAART for treatment-naïve HIV(+) patients (A1424-008).** In: *41st Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC 2001)*, Chicago, IL, 2001.
25. Wood R, Phanuphak P, Cahn P, Pokrovskiy V, Rozenbaum W, Pantaleo G, et al. **Long-term efficacy and safety of atazanavir with stavudine and lamivudine in patients previously treated with nelfinavir or atazanavir.** *J Acquir Immune Defic Syndr* 2004; **36**:684–692.
26. Bristol Myers Squibb Company. *Videx (didanosine) prescribing information.* New York: Bristol Myers Squibb Company; 2004.
27. Delfraissy JF, Rivero A, Yakovlev A, Squires K, Thiry A, Giordano M. **Comparison of antiretroviral efficacy and safety of atazanavir qd and efavirenz qd with fixed-dose ZDV + 3TC through week 24 (A1424-034).** In: *Sixth International Congress on Drug Therapy in HIV Infection*, Glasgow, UK, 2002.
28. Squires K, Thiry A, Giordano M. **Atazanavir (ATV) QD and efavirenz (EFV) QD with fixed-dose ZDV+3TC: comparison of antiviral efficacy and safety through wk 24 (A1424-034).** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, CA, 2002.
29. Gathe J, Benetucci J, Lupo S, Connaughton E, Kelleher T, McLaren C. **Comparison of a triple regimen containing once-daily didanosine vs a regimen of ZDV/3TC/NLF.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
30. Badaro R, Gathe J, Grimwood J, McLaren C, Kleszczewski K. **Evaluation of a triple combination regime containing enteric coated didanosine administered once-daily.** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, Argentina, 2001.
31. Gathe J, Badaro R, Grimwood J, Abrams D, Kleszczewski K, McLaren C. **Comparison of a triple combination regimen containing an enteric coated formulation of didanosine administered once-daily versus a regimen of combivir plus nelfinavir.** In: *Eighth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 2001.
32. Molina J. **Once-daily therapy with emtricitabine, didanosine and efavirenz in treatment naïve HIV-infected adults: 48-week follow-up of The ANRS 091 Trial.** In: *38th Annual Meeting of the IDSA (Infectious Diseases Society of America)*, New Orleans, Louisiana, 2000.
33. Molina JF, Journot V, Maillard A, Noe E, Raffi F, Rozenbaum W, et al. and the Montana Study Group. **Once-daily combination therapy with emtricitabine, didanosine and efavirenz in treatment naïve HIV-infected adults: 96-week follow-up of the ANRS 091 trial.** In: *Eighth Conference on Retroviruses and Opportunistic Infections*, Athens, 2001.
34. Molina JM, Ferchal F, Rancinan C, Raffi F, Rozenbaum W, Sereni D, et al. **Once-daily combination therapy with emtricitabine, didanosine, and efavirenz in human immunodeficiency virus-infected patients 2.** *J Infect Dis* 2000; **182**:599–602.
35. Molina JM, Ferchal F, Rancinan C, Raffi F, Rozenbaum W, Sereni D, et al. **A pilot study of FTC + ddI + efavirenz in treatment-naïve HIV-infected adults: a potent and convenient once-a-day HAART (ANRS 091 trial).** In: *Seventh Conference on Retroviruses and Opportunistic Infections*, 2000, [abstract 518].
36. Molina JN, Raffi F, Rozenbaum W, Morlat P, Rancinan C, Journot V, et al. and the Montana Study Group. **Once-daily combination therapy with emtricitabine (FTC), didanosine (ddI) and efavirenz in treatment naïve HIV-infected adults: 3-year follow-up of the MONTANA (ANRS 091) trial.** In: *Second IAS Conference on HIV Pathogenesis and Treatment*, 2003.
37. Landman R, Schiemann R, Thiam S, Vray M, Canestri A, Mboup S, et al. **Once-a-day highly active antiretroviral therapy in treatment-naïve HIV-1-infected adults in Senegal.** *AIDS* 2003; **17**:1017–1022.
38. Landman R, Thiam S, Canestri A, Delaporte E, Mboup S, Vray M, et al. **Long-term evaluation (15 Months) of ddI, 3TC and efavirenz once-daily regimen in naïve patients in Senegal: ANRS 12-04/IMEA 011 study.** In: *Ninth Conference on Retroviruses and Opportunistic Infections*, Seattle, WA, 2002.
39. Murphy RL, Katlama C, Johnson V, Squires K, Horban A, Gatell JM, et al. **The Atlantic Study: A randomized, open-label trial comparing two protease inhibitor (PI)-sparing antiretroviral strategies versus a standard PI-containing regimen.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
40. van Leeuwen R, Katlama C, Murphy RL, Squires K, Gatell J, Horban A, et al. **A randomized trial to study first-line combination therapy with or without a protease inhibitor in HIV-1-infected patients 9.** *AIDS* 2003; **17**:987–999.
41. Avanti Study Group. **AVANTI 2. Randomized, double-blind trial to evaluate the efficacy and safety of zidovudine plus lamivudine versus zidovudine plus lamivudine plus indinavir in HIV-infected antiretroviral-naïve patients 7.** *AIDS* 2000; **14**:367–374.
42. Goebel FD. **AVANTI 2. A randomized, double blind, comparative trial to evaluate combination antiretroviral regimens for the treatment of HIV-1 infection: AZT/3TC/indinavir in anti-retroviral naïve patients.** In: *Sixth European Conference on Clinical Aspects and Treatment of HIV-Infection*, Hamburg, Germany, 1997 [abstract 211].

43. Hill A, Demasi R, Kuhn M. **Different analyses give highly variable estimates of HIV-1 RNA undetectability and log reduction in clinical trials.** In: *Twelfth World AIDS Conference*, Geneva, 1998.
44. Clumeck N. **Avanti 3: A randomized, double blind, comparative trial to evaluate the efficacy, safety and tolerance of AZT/3TC vs AZT/3TC/nelfinavir in antiretroviral naive patients.** In: *Fifth Conference on Retrovirus and Opportunistic Infections*, Chicago, IL, 1998.
45. Gartland M. **AVANTI 3: a randomized, double-blind trial to compare the efficacy and safety of lamivudine plus zidovudine versus lamivudine plus zidovudine plus nelfinavir in HIV-1-infected antiretroviral-naïve patients.** *Antivir Ther* 2001; 6:127-134.
46. Bartlett JA, Johnson J, Shaefer MS, Herrera G, Sosa N, Rodriguez AE. **Abacavir/lamivudine (ABC/3TC) in combination with efavirenz (NNRTI), amprenavir/ritonavir (PI) or stavudine (NRTI): ESS40001 (CLASS) preliminary 48 week results.** In: *14th International AIDS Conference*, Barcelona, Spain, 2002.
47. Gazzard B, Dejesus E, Cahn P, Castillo S, Zhao H, Gordon D, et al. **Abacavir (ABC) once daily (OAD) plus lamivudine (3TC) OAD in combination with efavirenz (EFV) OAD is well-tolerated and effective in the treatment of antiretroviral therapy (ART) naïve adults with HIV-1 infection (ZODIAC study: CNA30021).** In: *43rd Annual Interscience Conference on Antimicrobial Agents and Chemotherapy*, Chicago, Illinois, 2003.
48. Glaxo-Smith-Kline. **Ziagen (abacavir sulfate) package insert.** Research Triangle Park, NC: Glaxo-Smith-Kline; 2005.
49. Piliero P. **Is Abacavir a once daily drug.** In: *43rd Interscience Conference on Antimicrobial Agents and Chemotherapy*, Chicago, IL, 2003.
50. Fischl M, Greenberg S, Clumeck N, Peters B, Rubio R, Gould J, et al. **Ziagen (abacavir, ABC, 1592) combined with 3TC & ZDV is highly effective and durable through 48 weeks in HIV-1 infected antiretroviral-therapy-naïve subjects (CNA3003).** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
51. Fischl M, Greenberg S, Clumeck N, Peters B, Rubio R, Pobiner B, Verity L. **Safety and activity of abacavir (ABC, 1592) with 3TC/ZDV in antiretroviral naïve subjects.** In: *Twelfth International AIDS Conference*, Geneva, 1998.
52. Dejesus E, Herrera G, Teofilo E, Castillo S, Bonny T, Thorpe D, et al. **Efficacy and safety of abacavir (ABC) versus zidovudine (ZDV) in antiretroviral therapy naïve adults with HIV-1 infection (Study CNA30024).** In: *43rd Interscience Conference on Antimicrobial Agents and Chemotherapy*, Chicago, IL, 2003.
53. Staszewski S, Keiser P, Gathe J, Haas D, Montaner J, Hammer S, et al. **Ziagen/combivir is equivalent to indinavir/combivir in antiretroviral therapy (ART) naïve adults at 24 weeks (CNA3005).** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
54. Staszewski S, Keiser P, Gathe J, Haas D, Montaner J, Johnson M, et al. **Comparison of antiviral response with abacavir/combivir to indinavir/combivir in therapy-naïve adults at 48 weeks (CNA3005).** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC)*, San Francisco, 1999.
55. Staszewski S, Keiser P, Gathe J, Haas D, Montaner J, Johnson M, et al. **Comparison of antiviral response with abacavir/3TC/ZDV to indinavir/3TC/ZDV in therapy naïve adults at 48 weeks (CNA3005).** In: *Seventh European Conference on Clinical Aspects and Treatment of HIV-Infection*, Lisbon, Portugal, 1999.
56. Staszewski S, Keiser P, Montaner J, Raffi F, Gathe J, Brotas V, et al. **Abacavir-lamivudine-zidovudine vs indinavir-lamivudine-zidovudine in antiretroviral-naïve HIV-infected adults: A randomized equivalence trial 85.** *JAMA* 2001; 285:1155-1163.
57. Jordan J CP, Vibhagool A. **Predictors of adherence and efficacy in HIV-1-infected patients treated with abacavir/combivir (ABC/COM) or indinavir/combivir (IDV/COM): final 48-week dData from CNA3014.** 543-T. In: *Ninth Conference on Retroviruses and Opportunistic Infections*, Seattle, 2002.
58. Vibhagool A, Cahn P, Schechter M, Smail F, Soto-Ramirez L, Carosi G, et al. **Abacavir/combivir is comparable to indinavir/Combivir(tm) (IDV/COM) in HIV-1-infected antiretroviral therapy naïve adults: preliminary results of a 48-week open-label study (CNA3014).** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, 2001.
59. Vibhagool A, Cahn P, Schechter M, Smail F, Soto-Ramirez L, Carosi G, et al. **Triple nucleoside treatment with abacavir plus the lamivudine/zidovudine combination tablet (COM) compared to indinavir/COM in antiretroviral therapy-naïve adults: results of a 48-week open-label, equivalence trial (CNA3014).** *Curr Med Res Opin* 2004; 20:1103-1114.
60. Matheron S, Descamps D, Boue F, Livrozet JM, Lafeuillade A, Aquilina C, et al. **Triple nucleoside combination zidovudine/lamivudine/abacavir versus zidovudine/lamivudine/nelfinavir as first-line therapy in HIV-1-infected adults: a randomized trial.** *Antivir Ther* 2003; 8:163-171.
61. Podzamczar D, Ferrer E, Consiglio E, Gatell J, Perez P, Perez J, et al. **Final 12-month results from the COMBINE study: A randomized, open, multicenter trial comparing combivir plus nelfinavir or nevirapine in naïve patients.** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, Argentina, 2001 [abstract 7].
62. Podzamczar D, Lupo S. **Preliminary results of a randomized multicenter study comparing combivir (ZDV/3TC) plus nelfinavir or nevirapine in HIV-infected naïve patients (COMBINE Study).** In: *Seventh Conference on Retroviruses and Opportunistic Infections*; 2000 [abstract 510].
63. Dejesus E, Cohen C, Sax P, Nahass R, Morris A, Gothelf S, et al. **Efficacy and safety of didanosine-EC (DDI-EC), lamivudine (3TC), and efavirenz (EFV) as a once daily regimen for treatment-naïve HIV patients: 24 week interim results from the DART I trial.** In: *41st Annual Meeting of IDSA*, San Diego, 2003.
64. Ward D, Cohen C, Keiser P, Morris A, Sax P, Dezii C, et al. **Efficacy and safety of efavirenz (EFV), lamivudine (3TC), didanosine-EC (DDI-EC), as a once daily regimen for treatment-naïve HIV patients: 48 week results from the DART I trial.** In: *15th International AIDS Conference*, Bangkok, Thailand, 2004.
65. Felizarta F, Becker S, Bellos N, Jayaweera D, Sands M, Slater L, et al. **Adherence and efficacy with a once-daily efavirenz-based regimen: 48 week results from the daily antiretroviral therapy II (DART II) study.** In: *15th International AIDS Conference*, Bangkok, Thailand, 2004.
66. Jayaweera D, Becker S, Bellos N, Felizarta F, Sands M, Slater L, et al. **A once daily efavirenz-based regimen for treatment-naïve HIV patients: 48 week efficacy and safety results from the DART II trial.** University of Miami School of Medicine, Miami, FL, United States. In: *15th International AIDS Conference*, Bangkok, Thailand, 2004.
67. Haas D, Seekins D, Cooper D, Gallant J, Tashima K, Hicks C, Manion DJ. **A phase II, double-blind, placebo-controlled, dose ranging study to assess the antiretroviral activity and safety of efavirenz (EFV, SUSTIVA™, DMP 266) in combination with open-label zidovudine (ZDV) with lamivudine (3TC) at 36 weeks [DMP 266-005].** In: *12th World AIDS Conference*, Geneva, 1998.
68. Hicks C, Haas D, Seekins D, Cooper R, Gallant J, Carpenter C, et al. **A phase II, double-blind, placebo-controlled, dose ranging study to assess the antiretroviral activity and safety of DMP 266 (Efavirenz, SUSTIVA™) in combination with open-label zidovudine (ZDV) with lamivudine (3TC) [DMP 266-005].** In: *Fifth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1998.
69. Manion DJ, Faulkner E, Saxton TD, Labriola DF, Ruiz NM. **Durability of response of efavirenz (SUSTIVA™, EFV)-containing regimens: report of the post-control period results of studies with EFV.** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
70. Manion DJ, Labriola DF, Ruiz NM. **Efficacy of efavirenz (SUSTIVA™) containing regimens in patients with baseline plasma HIV-RNA viral loads exceeding 100,000 copies/ml.** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
71. Cohen C, Elion R, Green S, Sathasivam K, Waterhouse B, Villano S, Rui NM. **Sustiva (efavirenz) is highly effective and well tolerated in combination with the nucleosides stavudine and lamivudine (Study DMP 266-043).** In: *The 40th Annual Meeting of the Interscience Conference on Antimicrobial Agents and Chemotherapy*, Toronto, ONT, 2000.
72. Luskin-Hawk R, Cohen C, Lang J, Pegram S, Elion R, Green S, et al. **Efavirenz is well-tolerated and highly efficacious in combination with the nucleoside stavudine (d4T) + lamivudine (3TC).** In: *Infectious Disease Society of America (IDSA)*; 1999.

73. Domula MR, Wasmuth JC, Jutte A, Fatkenheuer G, Oette M, Theisen A, et al. **A comparison of AZT/3TC vs 3TC/d4T and ddI/d4T in combination with efavirenz as first line therapy: efficacy and safety results after 48 weeks.** In: *Ninth Conference on Retroviruses and Opportunistic Infections*, Seattle, WA, 2002.
74. Garcia F, Alonso M, Romeu J, Knobel H, Arrizabalaga J, Podzamczar D, et al. **An open randomized trial of indinavir, lamivudine, and stavudine versus stavudine/zidovudine and lamivudine in HIV-1 infected patients with CD4+ T cell count above 500/mm³.** Spanish Earth-2 Study. In: *Seventh European Conference on Clinical Aspects and Treatment of HIV-Infection*. Lisbon, Portugal; 1999.
75. Garcia F, Alonso MM, Romeu J, Knobel H, Arrizabalaga J, Ferrer E, et al. **Comparison of immunologic restoration and virologic response in plasma, tonsillar tissue, and cerebrospinal fluid in HIV-1-infected patients treated with double versus triple antiretroviral therapy in very early stages: The Spanish EARTH-2 Study. Early Anti-Retroviral Therapy Study.** *J Acquir Immune Defic Syndr* 2000; 25:26–35.
76. DeJesus E, McCarty D, Farthing CF, Shortino DD, Grinsztejn B, Thomas DA, et al. **Once-daily versus twice-daily lamivudine, in combination with zidovudine and efavirenz, for the treatment of antiretroviral-naïve adults with HIV infection: a randomized equivalence trial.** *Clin Infect Dis* 2004; 39:411–418.
77. Bowonwatanuwong C, Mootsikapun P, Supparatpinoy K, Tansuphaswadikul S, Athisegran R, Pasook P, Jones A. **A randomised, open label study to investigate abacavir (ABC) and lamivudine (3TC) as once daily (QD) components of a triple combination regimen (EPV40001).** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, 2001.
78. Cahn P, Raffi F, Saag M, Wolff M, Pearce D, Molina JM, et al. **Virologic efficacy and patterns of resistance mutations in ART-naïve patients receiving combination therapy with once-daily emtricitabine compared to twice-daily stavudine in a randomized, double-blind, multi-center clinical trial.** In: *10th Conference on Retroviruses and Opportunistic Infections*, Boston, MA, 2003.
79. Raffi F, Saag M, Cahn P, Wolff M, Pearce D, Molina J, et al. **Stavudine plus didanosine and nevirapine in antiretroviral-naïve HIV-infected adults: preliminary safety and efficacy results.** VIRGO Study Team. In: *Second IAS Conference on HIV Pathogenesis and Treatment*; 2003.
80. Saag M, Cahn P, Raffi F, Wolff M, Pearce D, Molina JM, et al. **A randomized, double-blind, multicenter comparison of emtricitabine QD to stavudine BID.** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, 2002.
81. Katzenstein TL, Kirk O, Pedersen C, Lundgren JD, Nielsen H, Obel N, et al. **The Danish protease inhibitor study: a randomized study comparing the virological efficacy of 3 protease inhibitor-containing regimens for the treatment of human immunodeficiency virus type 1 infection.** *J Infect Dis* 2000; 182:744–750.
82. Gerstoft J, Kirk O, Obel N, Pedersen C, Mathiesen L, Nielsen H, et al. **Low efficacy and high frequency of adverse events in a randomized trial of the triple nucleoside regimen abacavir, stavudine and didanosine 1.** *AIDS* 2003; 17: 2045–2052.
83. Hill A, Moyle G. **A meta-analysis of the virologic efficacy of regimens containing four versus three active antiretroviral agents as initial therapy for HIV-1 infection.** In: *XIII International HIV Drug Resistance Workshop*, Tenerife, Canary Islands, 2004.
84. Anekthananon T, Ratanasuwana A, Techasathit W, Sonjai A, Suwanagool S. **Safety and efficacy of a simplified fixed-dose combination of stavudine, lamivudine, and nevirapine (GPO-VIR) for the treatment of HIV-infected patients.** In: *XV International AIDS Conference*, Bangkok, Thailand, 2004.
85. Gallant J, Staszewski S, Pozniak A, Lu B, Miller MD, Coakley DF, Cheng A. **Favorable lipid and mitochondrial (mt) DNA profile for tenofovir disoproxil fumarate (TDF) compared to stavudine (d4T) in combination with lamivudine (3TC) and efavirenz (EFV) in antiretroviral therapy (ART) naïve patients: a 48 week interim analysis.** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, CA, 2002.
86. Staszewski S, Gallant J, Pozniak A, Suleiman JMAH, DeJesus E, Lu B, et al. **Efficacy and safety of tenofovir DF (TDF) versus stavudine (d4T) when used in combination with lamivudine and efavirenz in antiretroviral naïve patients: 96-week preliminary interim results.** In: *10th Conference on Retroviruses and Opportunistic Infections*, Boston, MA, 2003.
87. Skowron G, Myers M, Pedneault L, Smiley L, Warburg M, Odorisio M. **A protocol to evaluate the safety and efficacy of multidrug combination antiretroviral therapy with zidovudine (ZDV) and didanosine (ddI) with or without lamivudine (3TC) or nevirapine (NVP) for the treatment of HIV infection.** In: *37th Interscience Conference on Antimicrobial Agents and Chemotherapy*, Toronto, ONT, 1997.
88. Saimot AG, Landman R, Damond F, Mathes D, Gaudebout C, Liebowitch J, et al. **Stavudine, didanosine, and ritonavir as a triple therapy in antiretroviral-naïve patients: results at 72 weeks.** In: *12th World AIDS Conference*, Geneva, 1998.
89. Montaner J, Reiss P, Cooper D, Vella S, Dohnanyi C, Harris M, et al. **Long-term follow-up of patients treated with nevirapine (NVP) based combination therapy within the INCAS Trial.** In: *Fifth Conference on Retrovirus and Opportunistic Infections*, Chicago, IL, 1998.
90. Montaner JS, Reiss P, Cooper D, Vella S, Harris M, Conway B, et al. **A randomized, double-blind trial comparing combinations of nevirapine, didanosine, and zidovudine for HIV-infected patients: the INCAS Trial.** Italy, The Netherlands, Canada and Australia Study. *JAMA* 1998; 279:930–937.
91. Myers M, Montaner J. **A Randomized, double-blinded comparative trial of the effects of zidovudine, didanosine and nevirapine combinations in antiretroviral naïve, AIDS-free, HIV-infected patients with CD4 counts 200–600/mm³.** In: *XI International Conference on AIDS*; 1996.
92. Wood R, Hawkins DA, Moyle G, De Cian W, Ingrosso A, Greenwald C. **Second placebo-controlled study in naïve individuals confirms the role of delavirdine in highly active antiretroviral, protease-sparing treatment.** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
93. Gathe J, Podzamczar D, Johnson M, Schwartz R, Yeh V, Travers N, et al. **Once-daily vs twice-daily lopinavir/ritonavir in antiretroviral-naïve patients: 48-week results.** In: *11th Conference on Retroviruses and Opportunistic Infections*, San Francisco, CA, 2004.
94. Podzamczar D, Gathe J, Johnson M, Schwartz R, Villacian J, Marsh T, et al. **F13 – efficacy and safety of once-daily lopinavir/ritonavir vs. twice-daily lopinavir/ritonavir in antiretroviral-naïve patients: 24-week results.** In: *Ninth European Aids Conference (EACS)*, Warsaw, Poland, 2003.
95. Eron J, King M, Xu Y, Brun S, Real K, Murphy R, et al. **ABT-378/ritonavir (ABT-387/r) Suppresses HIV RNA to >400 copies/ml in 95% of treatment-naïve patients and in 78% of PI-experienced patients at 36 weeks.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
96. Gulick R, King M, Brun S, Real K, Murphy R, Hicks C, et al. **ABT-378/ritonavir (ABT-378/r) in antiretroviral naïve HIV+ patients: 72 weeks.** In: *Seventh Conference on Retroviruses and Opportunistic Infections*, San Francisco, CA, 2000.
97. Hicks C, King M, Brun S, Real K, Murphy R, Eron J, et al. **ABT-378/ritonavir (ABT-378/R) in antiretroviral naïve HIV+ patients: 48 weeks.** In: *Seventh European Conference on Clinical Aspects and Treatment of HIV-Infection*, Lisbon, Portugal, 1999.
98. Landay A, Brun S, King M, Marsh T, Murphy R, Hicks C, et al. **Three-year immunologic responses in antiretroviral-naïve HIV+ patients treated with lopinavir/ritonavir (Kaletra) based therapy.** In: *XIV International AIDS Conference*, Barcelona, 2002.
99. Murphy R, Brun S, King M, Manning L, Hicks C, Eron J, et al. **Lopinavir/ritonavir (Kaletra) in antiretroviral naïve HIV+ patients: 4 year follow-up.** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, CA, 2002.
100. Murphy RL, Brun S, Hicks C, Eron JJ, Gulick R, King M, et al. **ABT-378/ritonavir plus stavudine and lamivudine for the treatment of antiretroviral-naïve adults with HIV-1 infection: 48-week results.** *AIDS* 2001; 15:F1–F9.

101. White C, Brun S, King M, Murphy R, Hicks C, Eron J, Sun E. **Lopinavir/ritonavir (Kaletra) in antiretroviral naive HIV+ patients: 144 week follow up.** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, Argentina, 2001.
102. Cernohous P, Bernstein B, Mosley J, King M, Bauer E, Sun E. **Safety and efficacy of lopinavir/ritonavir in women in a phase III of antiretroviral-naïve subjects.** In: *XIV International AIDS Conference*, Barcelona, Spain, 2002.
103. King M, Bernstein B, Cernohous P, Moseley J, Bauer E, Sun E. **Impact of baseline CD4 cell count and viral load on durability of virologic response through 96 weeks for lopinavir/ritonavir (LPV/r) and nelfinavir (NFV) in a phase III clinical trial.** In: *Ninth Conference on Retroviruses and Opportunistic Infections*, Seattle, Washington, 2002.
104. Luo M, Boggs R, Bernstein B, Sun E, Ashraf T. **Assessing treatment pattern and cost consequence of lopinavir/ritonavir vs nelfinavir therapy in HIV patients.** In: *Sixth International Congress Drug Therapy in HIV Infection*, Glasgow, UK, 2002.
105. Schranz J. **A multicenter, open-label, 24-week pilot study with a 24-week extension to evaluate the safety, tolerability and efficacy of indinavir (IDV)-ritonavir (RTV) 800/100 BID combination with D4t plus 3TC in HIV-infected individuals.** In: *First IAS Conference on HIV Pathogenesis and Treatment*, Buenos Aires, Argentina, 2001.
106. Nadler J, Rodriguez-French A, Millard J, Wannamaker P. **The NEAT Study: GW433908 efficacy and safety in ART-naïve subjects, final 48-week analysis.** In: *10th Conference on Retroviruses and Opportunistic Infections*, Boston, MA, 2003.
107. Rodriguez-French A, Boghossian J, Gray GE, Nadler JP, Quinones AR, Sepulveda GE, et al. **The NEAT study: a 48-week open-label study to compare the antiviral efficacy and safety of GW433908 versus nelfinavir in antiretroviral therapy-naïve HIV-1-infected patients.** *J Acquir Immune Defic Syndr* 2004; **35**:22–32.
108. Rodriguez-French A, Nadler J. **The NEAT Study: GW433908 efficacy and safety in anti-retroviral therapy (ART) naïve subjects, preliminary 24-week results.** In: *42nd Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, CA, 2002.
109. Vertex Pharmaceuticals Inc. *Lexiva (fosamprenavir calcium) prescribing information*. Cambridge, MA, USA: Vertex Pharmaceuticals Inc.; 2004.
110. Carr A, Hudson J, Hoy J, Chuah J, French M, Ozcombo I. **Control of HIV replication in treatment naïve patients in a randomized trial of HAART.** In: *Sixth Conference on Retroviruses and Opportunistic Infections*, Chicago, IL, 1999.
111. Goodgame J, Hanson C, Vafidis I, Stein A, Jablonowski H. **Amprenavir (141W94, APV)/3TC/ZDV exerts durable antiviral activity in HIV-1-infected antiretroviral therapy-naïve subjects through 48 weeks of therapy.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
112. Goodgame JC, Pottage JC Jr, Jablonowski H, Hardy WD, Stein A, Fischl M, et al. **Amprenavir in combination with lamivudine and zidovudine versus lamivudine and zidovudine alone in HIV-1-infected antiretroviral-naïve adults. Amprenavir PROAB3001 International Study Team.** *Antivir Ther* 2000; **5**:215–225.
113. Schurmann D, Gathe J, Sanne I, Wood I. **Efficacy and safety of GW433908/ritonavir once daily in therapy naïve subjects, 48 week results: the SOLO Study.** In: *Sixth International Congress on Drug Therapy in HIV Infection*, Glasgow, UK, 2002.
114. Squires K. **An open-label, randomized, comparative study of stavudine (d4T) + lamivudine (3TC) + indinavir (IDV) versus zidovudine (ZDV) + 3TC + IDV in treatment of HIV-infected patients: START I.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
115. Squires KE, Gulick R, Tebas P, Santana J, Mulanovich V, Clark R, et al. **A comparison of stavudine plus lamivudine versus zidovudine plus lamivudine in combination with indinavir in antiretroviral naïve individuals with HIV infection: selection of thymidine analog regimen therapy (START I).** *AIDS* 2000; **14**:1591–1600.
116. Eron J, Peterson D, Murphy R, Pottage J, Mauney J, Schoellkopf N, Stevens M. **An open-label, randomized, comparative study of d4t + ddI + IDV versus ZDV + 3TC + IDV in treatment naïve HIV-infected patients (START II).** In: *12th World AIDS Conference*, Geneva, 1998.
117. Eron JJ Jr, Murphy RL, Peterson D, Pottage J, Parenti DM, Jemsek J, et al. **A comparison of stavudine, didanosine and indinavir with zidovudine, lamivudine and indinavir for the initial treatment of HIV-1 infected individuals: selection of thymidine analog regimen therapy (START II) 7.** *AIDS* 2000; **14**:1601–1610.
118. Sension MG, Farthing C, Shaffer AG, Graham E, Siemon-Hryczyk P, Pilson RS. **Challenges of antiretroviral treatment in transient and drug-using populations: the SUN study 86.** *AIDS Patient Care STDs* 2001; **15**:129–136.
119. Raffi F, Reliquet V, Ferre V, Besnier JM, Zucman D, Raguin G, et al. **d4T + qd ddI + nevirapine (bid or qd) in antiretroviral-naïve HIV-1-infected patients: 1-year results of the VIRGO Study.** In: *39th Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Francisco, CA, 1999.
120. Reliquet V, Ferre V, Audrain M, Raffi F. **Long-term follow-up of antiretroviral-naïve HIV-infected patients treated with D4T + QD DDI + nevirapine (BID OR QD) in the VIRGO Study.** In: *Eighth European Conference on Clinic Aspects and Treatment of HIV – Infection*, Athens, Greece, 2001.
121. Gallant JE, Staszewski S, Pozniak AL, DeJesus E, Suleiman JM, Miller MD, et al. **Efficacy and safety of tenofovir DF vs stavudine in combination therapy in antiretroviral-naïve patients: a 3-year randomized trial.** *JAMA* 2004; **292**:191–201.
122. Gulick RM, Ribaudo HJ, Shikuma CM, Lustgarten S, Squires KE, Meyer WA 3rd et al. **Triple-nucleoside regimens versus efavirenz-containing regimens for the initial treatment of HIV-1 infection.** *N Engl J Med* 2004; **350**:1850–1861.
123. Pozniak A, Gallant JE, DeJesus E, Campo R, Arribas J, Gazzard BG, et al. **Superior outcome for tenofovir DF, emtricitabine and efavirenz compared to fixed dose zidovudine/lamivudine and EFV in antiretroviral naïve patients.** In: *Third IAS Conference on HIV Pathogenesis and Treatment*, Rio de Janeiro, Brazil, 2005 [abstract We.OA 02.02].
124. Bartlett JG, Gallant JE. *Medical management of HIV infection*. Baltimore: Johns Hopkins University; 2005.
125. DHHS. *Guidelines for the use of antiretroviral agents in HIV-1-infected adults and adolescents*. Department of Health and Human Services <http://aidsinfo.nih.gov/guidelines/guideline-detail.asp> Accessed: May 4, 2006.
126. Egger M, May M, Chene G, Phillips AN, Ledergerber B, Dabis F, et al. **Prognosis of HIV-1-infected patients starting highly active antiretroviral therapy: a collaborative analysis of prospective studies.** *Lancet* 2002; **360**:119–129.
127. Phillips A. **Short-term risk of AIDS according to current CD4 cell count and viral load in antiretroviral drug-naïve individuals and those treated in the monotherapy era.** *AIDS* 2004; **18**:51–58.
128. Cooper D, Yeni PG. **Virologic and immunological outcomes at 3 years following initiation of ART with regimens containing a NNRTI or PI or both: the INITIO trial.** In: *Twelfth Conference on Retrovirus and Opportunistic Infections*, Boston, MA, 2005 [abstract 165 LB].
129. Podzamczar D, Ferrer E, Consiglio E, Gatell JM, Perez P, Perez JL, et al. **A randomized clinical trial comparing nelfinavir or nevirapine associated to zidovudine/lamivudine in HIV-infected naïve patients (the Combine Study).** *Antivir Ther* 2002; **7**:81–90.
130. Robbins GK, De Gruttola V, Shafer RW, Smeaton LM, Snyder SW, Pettinelli C, et al. **Comparison of sequential three-drug regimens as initial therapy for HIV-1 infection.** *N Engl J Med* 2003; **349**:2293–2303.
131. Staszewski S, Morales-Ramirez J, Tashima KT, Rachlis A, Skiest D, Stanford J, et al. **Efavirenz plus zidovudine and lamivudine, efavirenz plus indinavir, and indinavir plus zidovudine and lamivudine in the treatment of HIV-1 infection in adults. Study 006 Team.** *N Engl J Med* 1999; **341**:1865–1873.
132. Bartlett JA, Johnson J, Herrera G, Sosa N, Rodriguez A, Liao Q, et al. **Long-term results of initial therapy with abacavir and lamivudine combined with efavirenz, amprenavir/ritonavir, or stavudine.** *J Acquir Immune Defic Syndr* 2006; in press.

133. Gathe JC Jr, Ive P, Wood R, Schurmann D, Bellos NC, Dejesus E, *et al.* **SOLO: 48-week efficacy and safety comparison of once-daily fosamprenavir /ritonavir versus twice-daily nelfinavir in naive HIV-1-infected patients.** *AIDS* 2004; **18**:1529–1537.
134. Walmsley S, Bernstein B, King M, Arribas J, Beall G, Ruane P, *et al.* **Lopinavir–ritonavir versus nelfinavir for the initial treatment of HIV infection.** *N Engl J Med* 2002; **346**:2039–2046.
135. Bartlett JA, Buda J, von Scheele B, Mauskopf J, Davis E, Elston R, *et al.* **Minimizing resistance consequences after virologic failure on initial combination therapy: A systematic overview.** *J Acquir Immune Defic Syndr* 2006; **41**:323–331.
136. Gallant JE, Rodriguez AE, Weinberg WG, Young B, Berger DS, Lim ML, *et al.* **Early virologic nonresponse to tenofovir, abacavir, and lamivudine in HIV-infected antiretroviral-naïve subjects.** *J Infect Dis* 2005; **192**:1921–1930.