

Using HIV-1 sequence variability to explore virus biology

Kathryn M. McGrath ^{a,b}, Noah G. Hoffman ^{a,d}, Wolfgang Resch ^{a,c},
Julie A.E. Nelson ^a, Ronald Swanstrom ^{a,c,*}

^a *UNC Center for AIDS Research, University of North Carolina at Chapel Hill, 22-062 Lineberger Cancer Center, CB # 7295, Chapel Hill, NC 27599-7295, USA*

^b *Curriculum in Genetics and Molecular Biology, University of North Carolina at Chapel Hill, 22-062 Lineberger Cancer Center, CB # 7295, Chapel Hill, NC 27599-7295, USA*

^c *Department of Biochemistry and Biophysics, University of North Carolina at Chapel Hill, 22-062 Lineberger Cancer Center, CB # 7295, Chapel Hill, NC 27599-7295, USA*

^d *Department of Microbiology and Immunology, University of North Carolina at Chapel Hill, 22-062 Lineberger Cancer Center, CB # 7295, Chapel Hill, NC 27599-7295, USA*

Received in revised form 29 March 2001; accepted 29 March 2001

Abstract

Human immunodeficiency virus type 1 (HIV-1) only recently established an epidemic world-wide infection in the human population. The virus persists in the human host through active replication and is able to avoid clearance by the immune system. Active replication is an important component of the rapid evolutionary potential of HIV-1, a potential which manifests itself in the evolution of immune escape variants, drug resistant variants, and variants with the ability to use different cell surface coreceptors in conjunction with CD4. Multiple zoonotic introductions, compartmentalization of virus replication in the body, and genetic bottlenecks associated with sampling during transmission, antiretroviral therapy, and geographic and/or host population isolation further contribute to the range of sequences present in extant viruses. The sum of the history of all of these phenomena is reflected in HIV-1 sequence variability, and most of these phenomena are ongoing today. Here we review the use of HIV-1 sequence variability to explore its underlying biology. © 2001 Elsevier Science B.V. All rights reserved.

Keywords: Human immunodeficiency virus (HIV); Sequence variability; Evolution

1. Introduction

HIV-1 is a member of the Lentivirus genus of the *Retroviridae* family. In the early 1980s, recog-

nition of a new immunodeficiency disease that appeared to be communicable (CDC, 1981; Gottlieb et al., 1981; Masur et al., 1981) led to the original isolation of HIV-1 (Barre-Sinoussi et al., 1983) and the demonstration that serum samples from AIDS patients could detect the virus (Sarnadharan et al., 1984; Schupbach et al., 1984). The earliest known isolate of HIV-1 is from a

* Corresponding author. Tel.: +1-919-9665710; fax: +1-919-9668212.

E-mail address: risunc@med.unc.edu (R. Swanstrom).

1959 plasma sample, which suggests that the current epidemic started in this century (Zhu et al., 1998). As of the end of 2000, over 36 million people are currently infected with HIV-1 worldwide (AIDS Epidemic Update: December 2000 [<http://www.us.unaids.org>]).

As independent HIV-1 isolates were initially identified and characterized, it became evident that genomic heterogeneity existed among different isolates (Benn et al., 1985; Hahn et al., 1985; Wong-Staal et al., 1985). Sequence analysis confirmed these results and showed that diversity was not evenly distributed throughout the genome, but was greatest in the envelope (*env*) gene (Hahn et al., 1985; Starcich et al., 1986). Analysis of serial isolates from patients demonstrated that inpatient isolates were more related than outpatient isolates and that inpatient isolates exhibited evolution over time (Hahn et al., 1986). Broader groupings of HIV-1 isolates have been recognized; analysis of HIV-1 isolates from around the world has revealed eight distinct subtypes of HIV-1, which cluster together phylogenetically to form group M (for main), which is distinct from groups O (for outlier) and N (for non-M/non-O) (Robertson et al., 1999).

Like HIV-1, HIV-2 isolates exhibit genetic variability similar to what has been observed among HIV-1 isolates (Gao et al., 1994; Chen et al., 1997). HIV-2 was initially isolated from patients in West Africa who exhibited AIDS-like symptoms, but were seronegative for HIV-1 (Clavel et al., 1986). Isolation of the virus and sequencing of the genome revealed that while the genetic organization of HIV-2 was highly similar to that of HIV-1, the two viruses exhibited distant sequence similarity at the nucleotide level (Guyader et al., 1987). One difference in the genetic organization of these two viruses is that the accessory gene *vpu* is unique to HIV-1 isolates (Cohen et al., 1988), whereas *vpx* is only found in HIV-2 isolates (Henderson et al., 1988; Kappes et al., 1988).

The mechanisms that result in retroviral sequence diversity have been studied for many years (Temin, 1992), though much attention has recently been focused on HIV-1. The sequence diversity generated by these mechanisms represents the source of variants that can evade the host

immune responses, that can give rise to drug-resistant strains, that can acquire altered cell tropism, and that give rise to many other biological phenomena. Collectively, these allow the persistent viral replication that ultimately weakens the host immune system and results in immunodeficiency. The goal of this review is to provide a broad overview of the use of sequence variability to study HIV-1 evolution, and to discuss the viral mechanisms and the host responses that shape this evolution.

2. Evolution of HIV in the human population

2.1. Introduction of primate lentiviruses into humans

Genetic distance analysis of the known primate lentiviruses indicates that they cluster into five distinct lineages, with HIV-1 and HIV-2 showing the highest degree of similarity with chimpanzee and sooty mangabey lentiviruses, respectively (Fig. 1). A consideration of the genetic distances within and among the five lineages has led to the view that these two human viruses originated from distinct non-human primate lentiviruses that had been evolving in their natural hosts for a long period of time before independently entering the human population (reviewed in Beer et al., 1999).

The primate reservoir that was the source of HIV-2 has been well documented as the sooty mangabey (*Cercocebus atys*) (Gao et al., 1992; Hirsch et al., 1989). Sooty mangabeys are the natural host of SIVsm and are indigenous to west Africa, which is also the epicenter of HIV-2 infection in humans. Furthermore, SIVsm is found in approximately 30% of sooty mangabeys living in the wild (Sharp et al., 1995). Phylogenetic analysis suggests that HIV-2 and SIVsm are closely related, as is SIVmac, which likely arose from the inadvertent infection of rhesus macaques with SIVsm in captivity (Fig. 1) (Hirsch et al., 1989). In a large sampling of HIV-2 and SIVsm isolates, the two lineages were found to be phylogenetically interspersed. A geographic link has been established between some phylogenetically related SIVsm and HIV-2 strains (Chen et al., 1996; Chen

et al., 1997). This suggests that there have been multiple introductions of SIVsm into the human population, representing the range of sequence diversity of the SIVsm extant population (Gao et al., 1994; Chen et al., 1997).

Recent work has indicated that the source of HIV-1 is from a subspecies of chimpanzees, *Pan troglodytes troglodytes*. Initial work with chimpanzee lentiviruses (SIVcpz) did not demonstrate a conclusive link to HIV-1. Wild-captured chim-

panzees have a low seroprevalence of SIVcpz, approximately 3% (Peeters et al., 1989, 1992). Also, some SIVcpz sequences, such as SIVcpz-ANT, show less sequence similarity with HIV-1 than expected for closely related virus strains (Fig. 1) (Vanden Haesevelde et al., 1996). However, more recent work has suggested that SIVcpz strains more phylogenetically similar to HIV-1, such as SIVcpzGAB1, SIVcpzGAB2, and SIVcpzUS, were restricted to the chimpanzee sub-




Fig. 1. Phylogenetic analysis of *pol* region from primate lentiviruses. Nucleotide sequences were aligned by using the pileup function in the GCG (Genetics Computer Group) package (University of Wisconsin, Madison). A phylogenetic tree was constructed using the neighbor-joining method [with DNAdist (Kimura 2-parameter) and Neighbor from the PHYLIP package (version 3.5c) (Felsenstein, 1993)]. The five primate lentivirus groups are numbered 1 through 5 and the three HIV-1 groups are identified. The letters A through D indicate viral clades for HIV-1 and HIV-2. The sequences were obtained from the HIV Sequence Database website (<http://hiv-web.lanl.gov>) under the following accession numbers: HIV-1A (U455) (M62320); HIV-1B (HXB2) (K03455, M38432), HIV-1C (IN21068) (AF067155), HIV-1D (NDK) (M27323), HIV-1ant70 (L20587), HIV-1mvp51 (L20571), SIVcpzANT (U42720), HIV-1YBF30 (AJ006022), SIVcpzUS (AF103818), SIVcpzGAB1 (X52154), SIVmac239 (M33262), SIVsmmh4 (X14307), HIV-2A (ST) (M31113), HIV-2B (UC1) (L07625), SIVmndgb1 (M27470, X15781), SIVl'hoest (AF075269), SIVagm677a (M58410), SIVsyk173 (L06042).

species *P.t. troglodytes*, whereas SIVcpz strains that are less similar were isolated from other subspecies (Gao et al., 1999; Corbet et al., 2000). This work has provided convincing evidence that HIV-1 was introduced into humans by zoonotic transmissions from a single sub-species of chimpanzees.

A current hypothesis holds that the three most divergent groupings of HIV-1, called M, N, and O, are the result of three independent introductions of SIVcpz into the human population. Group M viruses are responsible for the majority of the human epidemic, and viral isolates in this group are divided into several subtypes or clades (Sharp et al., 1995; McCutchan et al., 1996). Group O is represented by a small number of viral isolates from Cameroon (Gürtler et al., 1996), and group N has just one known isolate, YBF30, from Cameroon (Simon et al., 1998). The genetic distances between the three groups are large, with group M and O envelope (Env) identities as low as 53% (Gao et al., 1996).

Determining when HIV-1 entered the human population has been difficult because of the many problems associated with calculating the rate of HIV-1 diversification. A key piece of evidence for the time of origin comes from the phylogenetic analysis of an early HIV-1 group M isolate from an African plasma sample stored since 1959 (Zhu et al., 1998). With its position on the group M phylogenetic tree and current evolutionary models, group M HIV-1 was probably introduced into the human population between 1915 and 1941, likely around 1931, and then continued to diversify in the human population (Korber et al., 2000b). However, there is a claim suggesting that HIV-1 was transmitted to the human population via the oral polio vaccine (OPV) in the late 1950s (Hooper, 1999). It has been suggested that some OPV stocks were grown in kidney cells prepared from SIVcpz-infected chimpanzees, which resulted in SIVcpz transmission during immunization. While it is difficult to determine exactly what happened at that time, the available evidence refutes this claim (Hillis, 2000; Korber et al., 2000a,b; Plotkin and Koprowski, 2000).

2.2. Spread and diversification in the human population

Following the introduction of group M virus into the human population, it is thought that rapid diversification and early genetic isolation of subepidemics led to the establishment of recognizable subtypes. According to criteria defined recently, eight circulating subtypes (or clades) of group M HIV-1 have been identified: A, B, C, D, F, G, H, J, and K (Robertson et al., 1999). Phylogenetic analysis of a large number of HIV-1 sequences revealed distinct sequence clusters which is the basis of the designations of distinct subtypes (Fig. 2). Within a subtype, there can exist sub-subtypes, which are distinct lineages that are very closely related to one subtype. For example, subtype F is comprised of two sub-subtypes, F1 and F2, and subtypes B and D could have been classified as sub-subtypes (Robertson et al., 1999). It appears likely that the epicenter of the group M pandemic is in Central Africa, where the earliest known group M ancestral isolate was identified and where all currently identified group M subtypes are cocirculating with high levels of genetic diversity (Zhu et al., 1998; Vidal et al., 2000).

Recombination has also contributed to the diversification of HIV-1. Retroviruses contain two copies of genomic RNA per virion, and a cell dually infected with different virus strains can produce virions containing heterodimeric RNAs. Reverse transcription of such heterodimeric RNA in a newly infected cell provides the opportunity for recombination since reverse transcriptase (RT) is able to jump between the two RNA strands during minus strand synthesis (Hu and Temin, 1990; Zhang et al., 2000). Recombinant viruses have been found involving every group and nearly every subtype of HIV-1 and HIV-2 (Kuiken et al., 1999). However, there have been no reports of recombination between HIV-1 and HIV-2, despite a large number of dual infections (Pieniazek et al., 1991; George et al., 1992). Most recombinant viruses first appeared in Africa, where all HIV-1 subtypes circulate (Robertson et al., 1998). It has been estimated that as many as 10% of all HIV-1 isolates may be intersubtype viral mosaics (Robertson et al., 1995).




Fig. 2. Phylogenetic analysis of *env* and *gag* using representatives of the eight subtypes of HIV-1 group M. Trees were constructed as in Fig. 1. The sequences are in the HIV Sequence Database under the following accession numbers: SE7253 (AF069670), Q2317 (AF004885), JRFL (U63632), RF (M17451, M12508), ETH2220 (U46016), 92BR025.8 (U52953), ELI (K03454, X04414), 94UG114.1 (U88824), FIN9363 (AF075703), VI850 (AF077336), 93BR020.1 (AF005494), DR-CBL (AF084936), SE6165 (AF061642), HH8793.1.1 (AF061640), VI991 (VI991), VI997 (VI997), 90CF056.1 (AF005496), SE9280.9 (AF082394), SE9173.3 (AF082395).

Although there are many HIV-1 recombinants, not all of them play a major role in the HIV-1 pandemic. Mosaic strains that are involved in expanding HIV-1 infections have been referred to as 'circulating recombinant forms' (CRF) (Carr et al., 1998). At present, there are eleven recognized CRFs throughout the world (HIV Sequence Database [<http://hiv-web.lanl.gov>]). For example, an AE recombinant is found extensively in south-east Asia, an AG recombinant in west and central Africa, and an AB recombinant in Russia (Fig. 3)

(Robertson et al., 1999). It is possible that AE is not a recombinant, as some statistical tests suggest that AE is monophyletic (Anderson et al., 2000). Continuous monitoring will be necessary to establish whether these recombinants come to represent a more significant part in the HIV-1 pandemic and what role they might play in limiting vaccine efficacy.

3. HIV-1 evolution within an individual

HIV-1 evolution during chronic infection is potentiated by the large number of rounds of virus replication and the inherent error-prone nature of reverse transcription. At any given time, the virus population of an HIV-1 infected individual is heterogeneous and dynamic, with the cross-sectional diversity of the *env* population initially increasing at the earlier stages of infection, but declining toward the later stages of infection (Delwart et al., 1997; Shankarappa et al., 1999). This section focuses on the viral mechanisms that fuel HIV-1 evolution and the host characteristics that help to shape the HIV-1 population within an individual.

3.1. Viral mechanisms of HIV-1 diversity

There are two viral mechanisms that contribute to the generation of variability of HIV-1 within an individual. One is the introduction of mutations into the viral genome by the error-prone reverse transcriptase (RT), and the other is recombination between pre-existing viral populations. The rate of HIV-1 evolution, which is dependent on these two mechanisms, will also be discussed in this section.

3.1.1. Mutation rate of HIV-1

Retrovirus replication involves the use of three separate polymerase systems, each with its own distinct potential for introducing errors. In the integrated (proviral) state of viral DNA, the cellular DNA replication machinery carries out replication of the viral genetic information, presumably with high fidelity. Synthesis of viral genomic RNA by the host RNA polymerase II is

a potentially error-prone step, although at present the contribution of this step in introducing mutations is poorly understood. More often, error-prone DNA synthesis by the viral DNA polymerase, RT, is envisioned as the ultimate source of viral sequence diversity. Numerous studies have documented that retroviral polymerases are error-prone *in vitro* (reviewed in Preston and Dougherty, 1996). Most studies of purified RT have over-estimated the potential error rate *in vivo*, which is significantly lower than that observed *in vitro* (Varela-Echavarria et al., 1992; Mansky and Temin, 1995), but still significantly higher than the fidelity associated with host DNA synthesis.

Several different retroviruses have been examined to establish the range of error rates per round of replication. From these studies, the rate

of all types of mutations combined is in the range of 0.1–1 mutation per genome per round of replication (reviewed in Preston and Dougherty, 1996). A major contribution to the propensity of RT to incorporate point mutations is the absence of an exonuclease activity that would permit proof-reading of misincorporated bases, although potential rudimentary proofreading mechanisms involving pyrophosphorolysis (Arion et al., 1998) or dinucleotide polyphosphate synthesis (Meyer et al., 1998) have been uncovered.

In studies performed using HIV-1, the *in vivo* rate for point mutations was found to be 3.4×10^{-5} (Mansky and Temin, 1995; Mansky, 1996a). Thus a majority of HIV-1 daughter genomes are faithful copies of the parental genome, but on average between three and four out of ten will carry a new point mutation. Several factors have




Fig. 3. (A) Differential clustering of a HIV-1 recombinant subtype in the *env* (subtype B) and *gag* (subtype A) regions. The noted sequences in the trees are A/B recombinants. Trees were constructed as in Fig. 1. The sequences came from the HIV Sequence Database: Kal153 (AF193276) and RU98001 (AF193277). (B) Schematic representation of HIV-1 A/B recombinant genome. Hatched areas have subtype A-like sequence and black areas have subtype B-like sequence. Adapted from Robertson et al. (1999).

been identified that can alter the mutation rate of HIV-1. Mutations in *vpr* increase the error rate approximately threefold but do not change the spectrum of mutations (Mansky, 1996b). The Vpr protein has been reported to bind to several cellular proteins, in particular uracil *N*-glycosylase (UNG) and the human homologue of yeast RAD23 (HHR23A), which are both involved in DNA repair (Zhao et al., 1994; Refaeli et al., 1995). The ability of Vpr to bind to UNG correlates with its ability to affect the mutation rate (Mansky et al., 2000). Hypermutation, which is the high density accumulation of a specific type of nucleotide substitution, can also contribute to the mutation spectrum (Vartanian et al., 1991; Kim et al., 1996). Several different types of hypermutation have been noted (although G to A is the most common), and each may arise under conditions of deoxynucleotide pool imbalance (Martinez et al., 1994). The presence of either RT inhibitor AZT or 3TC has been shown to increase mutation rates in vivo (Julias et al., 1997; Mansky and Bernard, 2000). Furthermore, AZT-resistant RT has an increased mutation rate (Mansky and Bernard, 2000).

The types of mutations that are introduced by RT are far from random, although an insufficient number of mutations have been characterized to understand all of the determinants of mutation bias. The majority of mutations are substitutions, with G to A transitions by far the most common. Frameshift mutations occur several fold less frequently than substitution mutations and most often represent the addition or deletion of a nucleotide in a homopolymeric run (Pathak and Temin, 1990a,b). Approximately one in 10 genomes suffers a significant internal deletion during a single round of replication (Olsen et al., 1990; Parthasarathi et al., 1995). Deletions are strongly potentiated by the presence of direct repeats (Delviks and Pathak, 1999, and others within).

3.1.2. Recombination

Recombination between related viral variants within an infected individual provides another source of variability. Interclade recombination, as described above, results in large evolutionary steps, whereas intraclade recombination results in smaller evolutionary changes among related vari-

ants. There have been reports of indirect evidence of intraclade recombination, in which different regions of the genome appear to be evolving independently (Simmonds et al., 1991; Morris et al., 1999). Demonstration of recombination in an individual infected with subtype B HIV-1 was found in a comparison of infectious molecular clones from proviral DNA (Groenink et al., 1992). Recombination can be an efficient mechanism for producing new combinations of sequences; for example, a rhesus monkey had predominantly wild type SIVmac239 only 2 weeks after being infected with two SIVmac239 strains that were attenuated in different genes and were injected into different legs (Wooley et al., 1997). We have found evidence of recombination between related variants in an individual infected with subtype B HIV-1, where recombination events in the V1/V2 region of *env* generated new sequence variants that rapidly achieved fixation, possibly the result of immune selection (McGrath and Swanstrom, unpublished results).

3.1.3. Rate of HIV-1 evolution

The observed rate of HIV-1 evolution in vivo is determined by the rate of mutation per generation (resulting from RT errors and recombination), the number of generations, and the fixation rate of mutations (through selection or genetic bottlenecks). Because the rate of fixation and number of generations are difficult to observe directly, the rate of HIV-1 evolution during human infection is usually estimated from the slope of a line that best fits a plot of nucleotide sequence divergence from an ancestral sequence versus time. Using branch lengths of a phylogenetic tree generated by maximum-likelihood methods as a measure of divergence, Korber and colleagues recently estimated the rate of evolution to be 0.24% (0.18–0.28%) substitutions per base pair per year for *env* coding sequences and 0.19% (0.09–0.27%) for *gag* (Korber et al., 2000b). However, previously published rates of divergence approaching 1% per year cannot be ruled out because of the variability of rate estimates (discussed below) (Lukashov et al., 1995; Zhang et al., 1997a; Leitner and Albert, 1999; Shankarappa et al., 1999).

One complication to determining the rate of HIV-1 divergence is that models of evolution and phylogenetic techniques used to calculate divergence typically do not take recombination into account (Korber et al., 1998). Rates of evolution have also been shown or are hypothesized to differ: (1) from one genomic region to another (Leitner and Albert, 1999); (2) among different sublineages of virus within a host (Korber et al., 1998); (3) during different stages of infection (Shankarappa et al., 1999); and (4) between hosts (Lukashov et al., 1995; Zhang et al., 1997a). In addition, estimates of the evolutionary rate depend on the alignment methods and the evolutionary model used (Korber et al., 2000b). Thus, the assignment of a rate of HIV-1 evolution is context-dependent, and generalizations must be made with care.

3.2. The HIV-1 population in an infected individual

All mechanisms described thus far result in the replication of HIV-1 as a group of closely related but non-identical sequences, a type of population that has been termed a quasispecies (Eigen, 1971; reviewed in Domingo and Holland, 1997). The behavior of such a quasispecies depends on the fraction of individual viruses that actively partake in successful replication. The following section presents different arguments pertaining to the size of the actively replicating HIV-1 population within an individual. The effect of compartmentalization of viral replication on diversity and the link between disease progression and viral diversity also will be examined.

3.2.1. Effective population size

Only a fraction of the total viral population within an individual contributes to the gene pool of the next generation of virus. The size of this fraction is referred to as the effective population size, as opposed to the total population size. The effective population size is an important determinant of the influence of stochastic versus deterministic effects in viral evolution. In a population of small effective size, the fate of viral variants can be influenced by random events, independent

of their fitness compared to other variants present, i.e. stochastic behavior. In large populations undergoing many rounds of replication, the abundance of variants is not subject to random events, but is governed only by their fitness, i.e. deterministic behavior. The magnitude of a variant's selective advantage is also important in determining its fate in the total viral population. Variants with small selective advantages or neutral mutations tend to be governed by stochastic effects, whereas variants with large selective advantages behave more deterministically (Rouzine and Coffin, 1999).

The effective size of the HIV-1 population has yet to be determined conclusively. In one view there is a large HIV-1 effective population size within an individual, which may be greater than 3×10^5 , the inverse of the HIV-1 mutation rate (Coffin, 1995). If this is true, every possible point mutation is generated every day, and relative fitness determines the level at which a mutant sequence persists in the viral population. The high number of productively infected cells during chronic infection, between 10^7 and 10^8 (Haase et al., 1996), and the generation of an estimated 10^{10} virions per day (Perelson et al., 1996) support this model. Further evidence for a large effective population size was recently obtained from the analysis of *pro* sequences, in which the population size was estimated to be at least 10^5 (Rouzine and Coffin, 1999).

Other observations seem more consistent with a smaller effective population size. Viral population size has been estimated from the analysis of *env* sequences to be between 1×10^3 and 2×10^3 , approximately 4–5 orders of magnitude lower than the number of infected cells (Leigh Brown, 1997; Leigh Brown and Richman, 1997). The small effective population size compared to the large number of infected cells could be due to a subdivision of the total population into independent populations, frequent bottleneck events, or the production of virus particles by a large fraction of cells that do not contribute to the next round of infection. The complex antiretroviral resistance patterns that have been observed in infected individuals may lend support to the stochastic model of evolution (Leigh Brown and Richman, 1997).

The large and small effective population size models have been generally treated as mutually exclusive. However, it has been suggested that neither model alone can describe the evolution of HIV-1 under all conditions (Rouzine and Coffin, 1999). HIV-1 may behave mostly deterministically during the steady state of infection, but events such as bottlenecks during drug treatment or transmission may increase the importance of stochastic effects. At this time the effective population size remains both a critically important parameter for understanding HIV evolution and a source of ongoing discussion.

3.2.2. Compartmentalization

The HIV-1 population within an infected individual is heterogeneous, although this heterogeneity may not be evenly distributed in the body. Disparity has been found between viral sub-populations that develop in different compartments of the body, particularly in the brain, blood, genital tract, oral cavity, and lymphoid tissue. Several ideas have been presented to explain the compartmentalization phenomenon, such as physical isolation of a tissue compartment, selective migration of infected cells, distinct types of target cells, and localized selective pressures. Understanding the differences in viral evolution in distinct compartments is important for studying issues such as sexual transmission, bioavailability of antiviral drugs, and virus–host interactions.

Since the predominant mode of HIV-1 transmission is sexual, it is important to understand the biology of HIV-1 in genital secretions. There are several lines of evidence to suggest that the blood and the male genital tract are at least partially distinct compartments. First, there is at most a weak correlation between the levels of either CD4⁺ T cells or viral RNA load in blood and viral RNA load in semen (Liuzzi et al., 1996; Kiessling et al., 1998; Vernazza et al., 1999). Second, responses to antivirals can differ between the blood and in the male genital tract, indicating that drug bioavailability may vary between these sites (Coombs et al., 1998; Eron et al., 1998; Kashuba et al., 1999). Third, the distribution of *env* gene species in blood and male genital tract is sometimes discordant (Vernazza et al., 1994;

Coombs et al., 1998; Ping et al., 2000). Last, the majority variant in blood is not always the majority variant found in semen, whether comparing DNA or RNA (Delwart et al., 1998; Eron et al., 1998; Kiessling et al., 1998).

There are similar lines of evidence that the blood and female genital tract are not a single, well mixed compartment, such as discordant viral loads (Rasheed, 1998), differential responses to nucleoside RT inhibitors (Rasheed et al., 1996), and diverse populations in both compartments (Poss et al., 1995; Overbaugh et al., 1996; Poss et al., 1998). Increased HIV-1 shedding in the genital tracts of men and women has been associated with localized inflammation, indicating that variants may be established through the migration of infected lymphocytes (Clemetson et al., 1993; Kreiss et al., 1994; Anderson et al., 1998). Understanding the selection of certain viral variants in the genital tract is a necessary part of understanding the genetic bottlenecks of infection and of HIV-1 transmission.

The brain can be considered a separate compartment for HIV-1 because of the blood–brain barrier. When brain-derived sequences have been compared to sequences found in other parts of the body they usually form a separate cluster (Steuler et al., 1992; Korber et al., 1994; Wong et al., 1997). Previous work has focused on comparing *env* sequences in the brain to *env* sequences from other compartments of the same individual, because the viral surface protein is often the determinant of virulence or tropism in different cell types (Power et al., 1994). Viruses replicating in the brain are macrophage-tropic, which is consistent with the type of infected cells found in the brain (Epstein et al., 1991; Keys et al., 1993; Korber et al., 1994; Power et al., 1994).

Virus in the oral cavity appears to be largely from the same compartment as the blood. Several studies have compared viral RNA loads in blood versus saliva, and most have found that they are positively correlated, despite technical differences between studies (Phillips et al., 1994; Liuzzi et al., 1996; Shugars et al., 2000). Analysis of V3 *env* sequences in blood and saliva at the viral RNA and proviral DNA levels have shown that the same variants usually exist in both compartments,

although there are a few instances of differences (Kakizawa et al., 1996; Freel et al., 2001). Recent evidence suggests that inflammation in the oral cavity is associated with an increase in viral loads, similar to what has been seen in the genital tract (Uccini et al., 1999; Shugars et al., 2000).

Virus isolated from the lymph nodes also appears to represent the same compartment as the blood. The majority of HIV-1 virions are produced in the lymphoid organs, and virions not trapped by the follicular dendritic cells can enter the bloodstream (reviewed in Fauci and Desrosiers, 1997). Several studies have demonstrated that variants in the blood are similar to those in lymph nodes (Ball et al., 1994; Donaldson et al., 1994). Other studies have shown similarities in variants from lymph nodes and in non-lymphoid organs such as the brain and lungs, suggesting that lymphocytes that were infected in the lymph nodes have infiltrated these organs and are the source of virus (Reddy et al., 1996; Hughes et al., 1997; Morris et al., 1999). One study looked at the distribution of proviral DNA variants across the spleen and found a non-homogenous distribution (Delassus et al., 1992). One difference has been found in the blood plasma/lymphoid cell pools of virus, specifically between the viral RNA from virions and the proviral DNA from the infected lymphocytes from the same blood sample. Typically, variants in the viral RNA at one timepoint are found in the proviral DNA at later timepoints (Simmonds et al., 1991; Ball et al., 1994), which suggests that a significant fraction of the infected lymphocytes represent a potential reservoir for archival viral sequences. Similarly, drug resistance markers show a delayed appearance in viral DNA compared to viral RNA, indicating that actively replicating sequences represent only a fraction of the total viral DNA in PBMCs (Smith et al., 1993; Lineberger et al., 1995; Wei et al., 1995).

3.2.3. *Viral diversity versus progression rate*

The rate of progression to AIDS within an individual can be predicted by the steady state viral load, or set point (Gupta et al., 1993; Mellors et al., 1995, 1996). Steady state is achieved when immune responses are able to partially con-

trol viral replication following primary infection (reviewed in Coffin, 1996). Higher viral loads have been shown to be associated with faster disease progression (Mellors et al., 1996), and this prediction can be made as soon as 6–12 months following seroconversion (Mellors et al., 1995; Lefrere et al., 1998).

The extent to which host factors versus viral factors determine the rate of disease progression is unknown. Several studies have focused on differences between slow and rapid progressors. Typically, infected individuals progress to AIDS within 10 years, but some individuals progress much more rapidly and some individuals remain AIDS-free for more than 10 years. Slow progressors usually have higher neutralizing antibody titers and stronger cytotoxic T-lymphocyte (CTL) responses to HIV-1 than patients with typical progression (Pantaleo et al., 1995; Montefiori et al., 1996; Betts et al., 1999). In contrast, rapid progressors typically have weak or no neutralizing antibodies toward HIV-1 (Cao et al., 1995; Liu et al., 1997).

Viral divergence and diversity also appear to correlate with the rate of disease progression. Patients with slower disease progression have higher ratios of nonsynonymous to synonymous site mutations as well as higher diversity of viral quasiespecies (Lukashov et al., 1995; Wolinsky et al., 1996; Delwart et al., 1997; Liu et al., 1997). The antigenic diversity threshold model, which is based on mathematical models, attempts to explain the difference in evolution in slow and rapid progressors by suggesting that the immune system will be exhausted when the accumulation of viral variants reaches a certain threshold (Wodarz and Nowak, 1999). The threshold for each individual is different, which could explain the differences seen between rapid and slow progressors. The continuous virus adaptation model (Lukashov and Goudsmit, 1998), on the other hand, suggests that a competent immune system causes the virus to constantly evolve and adapt to the changing environment. This ability to adapt allows the virus to undergo continuous replication which leads to the depletion of CD4 + T cells and, ultimately, the development of AIDS. The former model would predict similar rates of evolution




Fig. 4. Summary of sequence variability in V3. Subtype B V3 sequences were classified as SI-like based on the presence of at least one non-consensus basic substitution listed in line (2), plus a change away from an acidic amino-acid at position 25; other sequences were considered NSI-like. A statistical test was designed to assess the distribution of substitutions in the resulting two sequence sets. Frequencies of substitutions indicated in lines (3) and (4) deviated significantly from expected values, and these changes were thus classified as either SI-associated or NSI-associated, respectively. Substitutions indicated in line (1) were not significantly associated with either group, although underlined amino acids were overrepresented among SI-like V3 sequences. Adapted from Milich et al. (1997).

until the threshold is reached. This prediction is inconsistent with data comparing slow and rapid progressors (Delwart et al., 1997).

4. Selective pressures within the host

HIV-1 evolution within an individual is directed by selective pressures placed on the virus by the host that act on different parts of the virus at different times during infection. This section focuses on the interplay between pressures imposed on viral replication within the host environment and compensatory changes in viral populations.

4.1. Cell-type specificity

Generally, HIV-1 can only infect cells that express CD4 and one of two chemokine receptors, CCR5 or CXCR4 (reviewed in Moore et al., 1997; Berger et al., 1999). The preferred phenotypic classification for HIV-1 isolates is defined by this CD4-dependent chemokine receptor usage: R5 for isolates using CCR5 for entry, X4 for those using CXCR4, and R5X4 for those capable of using both coreceptors (Berger et al., 1998). Other common phenotypic classifications that generally correlate with the R5 and X4 designations, respectively, are growth rate in vitro (slow/low vs.

rapid/high), cell tropism (macrophage-tropic vs. T cell line-tropic), or the ability to induce multinucleated giant cell formation in cultured T-cell lines (non-syncytium inducing/NSI vs. syncytium inducing/SI) (Asjo et al., 1986; Tersmette et al., 1988; Schuitemaker et al., 1992). Coreceptor usage does not fully explain tropism, however, since most X4 isolates cannot infect macrophages despite the expression of CXCR4 by those cells (Cheng-Mayer et al., 1997). There are also at least two mechanisms for dual-tropism: usage of either CXCR4 or CCR5 to enter either macrophages or T-cells (dual-tropic R5X4), and exclusive usage of CXCR4 to enter either of these cell types (dual-tropic X4) (Yi et al., 1999). These observations raise questions about the possible roles of post-entry factors in determining tropism.

To whatever extent growth kinetics, syncytium induction, and tropism are tied to coreceptor usage, all are determined to a large extent by sequences in the third variable loop (V3) of the HIV-1 gp120 envelope glycoprotein (Hwang et al., 1991; Cann et al., 1992; de Jong et al., 1992; Cocchi et al., 1996; Trkola et al., 1996). CXCR4 usage is most often associated with an accumulation of positively-charged amino acid substitutions in V3, particularly at positions 11 and 25 (Fig. 4) (Chesebro et al., 1992; Fouchier et al., 1992; Milich et al., 1993; Chesebro et al., 1996). In

many contexts, a change in V3 is both necessary and sufficient to produce a change in coreceptor usage (Xiao et al., 1998). However, there is some suggestion that sites outside of V3, particularly in the C4 region, vary with V3 and influence coreceptor usage by modifying either the internal stability of gp120, or coreceptor binding (Carrillo and Ratner, 1996; Milich et al., 1997).

In most patients, only R5/NSI viruses can be isolated during acute HIV infection and immediately after seroconversion (Roos et al., 1992; Zhu et al., 1993). Although occasionally X4/SI isolates can be detected early, they usually appear later in infection and are associated with progression to AIDS (Tersmette et al., 1989). It has been suggested that the presence of X4 variants may be transient, which could hinder their detection during disease progression (Shankarappa et al., 1999). In a large prospective study, Koot et al. (1993) reported that HIV-1-infected patients were over six-fold more likely to progress to AIDS within 30 months when SI isolates were detected. X4/SI detection is also predictive of rapid CD4 cell decline (Koot et al., 1993; Connor et al., 1997). The low frequency of transmission of X4 variants is unexplained and likely results from one or more of the following explanations: (i) poor representation of X4 variants in genital secretions; (ii) few target cells at mucosal surfaces; (iii) selective clearance of X4 variants during primary infection; and (iv) a predominance of CCR5-expressing activated T cells at lymphoid tissue sites that sustain the infection (Harouse et al., 1999).

The use of a V3-specific heteroduplex tracking assay (V3-HTA) (Nelson et al., 1997) has allowed the quantitative assessment of the dynamic interaction between R5 and X4 variants. In this analysis, these variants have much more complex interactions, including the persistence of R5 variants as the major species even in the presence of X4 variants, and the persistence of low-abundance variants with highly evolved V3 sequences (Nelson et al., 2000). The association of X4 variants with disease progression, and the maintenance of R5 variants throughout infection suggest that X4 viruses are either a cause of, or evolve in response to, progressive immunosuppression, while R5

variants make up the reservoir responsible for persistent infection (Schuitemaker et al., 1992).

4.2. Transmission

Transmission of HIV-1 is typically accompanied by a loss of both genotypic and phenotypic viral heterogeneity (Wolfs et al., 1992). Patients with longstanding HIV infections contain complex viral populations that display a range of phenotypes, whereas recent seroconverters contain a more genotypically homogenous virus that is uniformly R5/NSI (Zhu et al., 1993; Shankarappa et al., 1998). X4/SI variants are rarely sampled in recipients even when detected in the transmitter (Roos et al., 1992). Studies of epidemiologically related patients have shown that the predominant species detected in recently infected recipients are usually sampled as minor variants present in the blood of the transmitter (Wolinsky et al., 1992; Zhu et al., 1993). Viral heterogeneity early in infection may be influenced by the sex of the recipient; in one study of recently infected Africans who had acquired HIV-1 infections via heterosexual contact, women were found to harbor multiple viral variants, while the virus in men was more homogeneous (Long et al., 2000).

The purifying effect of transmission is most pronounced in *env*, particularly in the V3 region (Pang et al., 1992; Zhu et al., 1993). Soon after infection, V3 is less variable than other regions of the genome, such as V4–V5 of gp120 (Wolinsky et al., 1992) and p17/MA (Zhang et al., 1993), while in the chronically infected individual, gp120 and the V3 loop are of above-average diversity compared to the rest of the genome. In addition, high ratios of the frequency of synonymous to non-synonymous mutations (K_s/K_a) in V3 relative to V4–V5 soon after infection indicate the suppression of variants bearing coding changes in V3 during initial infection (Wolinsky et al., 1992). During chronic infection, there are low K_s/K_a ratios for V3, indicating selection for diversification in that region (Seibert et al., 1995; Wolinsky et al., 1996; Zhang et al., 1997a).

After transmission, both the timing and nature of the selective pressures responsible for the re-

duction in sequence variability remain poorly defined. Selection likely occurs very soon after transmission, but before seroconversion (Zhang et al., 1993). Strong early selection on V3 accompanied by recombination among members of an initially diverse population is consistent with the observed homogeneity of V3 and heterogeneity of other genomic regions (Zhang et al., 1993). The nature of this selective pressure remains controversial. It has been suggested that V3 loops of R5 viruses are more constrained in sequence than X4 viruses by the functional requirements of the interaction with the coreceptor (Sato et al., 1999). Selective replication of R5 viruses might also be a result of tropism requirements, since R5 variants have been shown to preferentially replicate in sub-optimally activated T-cells (Vicenzi et al., 1999). This *in vitro* observation was not borne out in HIV-infected patients, however, where naïve T cells contained culturable virus and proviral DNA representing both X4 and R5 variants at all stages of disease (Ostrowski et al., 1999).

4.3. Immune responses

Anti-HIV-1 immune responses drive viral evolution by selecting variants with reduced sensitivity to neutralizing antibodies and CTLs. The emergence of CD8⁺ CTL directed against HIV-1 epitopes coincides with a significant drop in initial viremia during primary infection, with most individuals having a measurable anti-HIV-1 CTL response at the time that they present with symptoms of acute HIV-1 infection (Borrow et al., 1994; Koup et al., 1994). The initial CTL response is important for containing primary infection because a lack of a sustained CTL response against several HIV-1 epitopes results in rapid progression to AIDS (Hay et al., 1999). In one study, a broader CTL response against HIV-1 was associated with a slower rate of CD4⁺ T cell loss (Pantaleo et al., 1997). Anti-HIV-1 memory CTLs are typically maintained at high levels during asymptomatic infection, and the loss of this CTL response is correlated with progression to AIDS (Rinaldo et al., 1995). Similarly, ablation of CD8⁺ cells in the macaque/SIV model leads to dramatic increases in viral load (Jin et al., 1999; Schmitz et al., 1999).

The best evidence of the potency of the CTL response, as well as a measurement of the selective pressure exerted on HIV-1 by the CTL response, is the emergence of CTL escape mutants. Escape mutants were first described in chronically infected individuals who were followed over several years (Phillips et al., 1991). In two separate studies of individuals with primary infection, CTL escape mutants were found just 22–30 days after the onset of symptoms of acute infection (Borrow et al., 1997; Price et al., 1997). In both studies, the escape mutants rapidly became the dominant variants *in vivo*, and these individuals had rapid CD4⁺ T cell losses indicative of rapid disease progression. In one SIV-infected macaque study, it was demonstrated that CTL responses specific for Tat, a viral protein expressed early in the life cycle, selected for viral escape variants (Allen et al., 2000). In long-term asymptomatic HIV-1-infected individuals and in SIV-infected rhesus macaques, the appearance of CTL escape mutants is associated with disease progression (Goulder et al., 1997; Evans et al., 1999). Correlations with viral load for both the initial anti-HIV-1 CTL response and the emergence of CTL escape mutants suggest that the effectiveness of CTLs against HIV-1 may determine the length of the asymptomatic period of infection.

The neutralizing antibody response takes longer to develop than HIV-1-specific CTL responses. Neutralizing antibody responses against autologous HIV-1 strains have been detected as soon as 5 weeks after the onset of primary infection symptoms, although they are typically seen any time up to 1 year after seroconversion (Arendrup et al., 1992; Pilgrim et al., 1997). Long-term non-progressors tend to have broader neutralizing antibody responses against HIV-1, as measured against panels of primary isolates, than do short-term non-progressors or progressors (Pilgrim et al., 1997). It is unknown, however, whether the broader neutralizing antibodies are the cause or an effect of long-term non-progression.

Neutralization escape mutants have been described in several studies, including a longitudinal study of SIVcpzANT in a chimpanzee. During chronic asymptomatic infection, escape mutants emerged an average of 15 months after a neutral-

izing antibody response was detected, and new neutralizing antibodies to the escape mutants were detected an average of 8 months later (Nyambi et al., 1997). There is a similar lag time between neutralization and escape in humans; sera are able to neutralize viral isolates from earlier timepoints, but not the contemporaneous isolate (Arendrup et al., 1992; Lathey et al., 1997). Another study found HIV-1 infected hu-PBL-SCID mice passively administered with high levels of potent neutralizing antibodies developed neutralizing escape mutants within a few days, which correlated with an increase in viral load (Poignard et al., 1999). Neutralization escape mutants do arise in long-term non-progressors (Bradney et al., 1999), indicating that, unlike CTL escape, escape from antibodies is not correlated with progression. The existence of escape mutants indicates that neutralizing antibodies exert selective pressure on HIV-1, although the low titers of antibodies and long time periods for escape mutants to emerge suggest that the selective pressure is likely weak.

4.4. Drug therapy

Combinations of antiretroviral drugs can be very potent in controlling HIV replication. Current antiretroviral therapy most commonly targets two viral enzymes, RT and protease (PR). In response to suboptimal drug therapy, variants are selected that have mutations in the targeted genes which lead to a reduction of drug sensitivity. Individuals in whom these variants arise then fail the drug therapy. Approximately 30–50% of HIV-1 infected individuals outside of clinical trials fail highly active antiretroviral therapy (Fatkenheuer et al., 1997; Deeks et al., 1999). This high rate of failure raises the issue of the spread of drug-resistant viral strains through the human population, which would undermine currently available treatment options.

Transmission of HIV-1 strains resistant to RT inhibitors (Erice et al., 1993; Quigg et al., 1997) or to both RT and PR inhibitors (Hecht et al., 1998) has been documented. Multiple studies have investigated the frequency of resistance-related mutations in HIV-1 from recently infected individuals (Birk and Sonnerborg, 1998; Fontaine

et al., 1998; Gomez-Cano et al., 1998; Pellegrin et al., 1999). Drug susceptibility testing has shown that up to 26% of therapy-naïve individuals harbored populations with a modest (3-to 10-fold) reduction in susceptibility to at least one drug (Boden et al., 1999; Little et al., 1999; Yerly et al., 1999), although the clinical significance of this observation is unclear. Because known primary resistance mutations are frequently absent in population-based sequencing of these infected individuals, isolates with such modest resistance likely represent background variability of drug sensitivity (Little et al., 1999). However, between 2 and 16% of the individuals studied did have viral populations with greater than 10-fold reduced susceptibility to one or more drugs, along with primary resistance mutations. These latter data suggest that drug sensitivity testing prior to treatment of drug-naïve individuals may become warranted.

4.5. Compensatory mutations

Initial adaptation of an HIV-1 population to strong selective pressures can result in a compromised replication rate. In subsequent rounds of replication under continued selection, the occurrence of second-site mutations that compensate for part or all of the lost fitness can be observed. This phenomenon has been most clearly demonstrated for the evolution of resistance to protease inhibitors. Some mutations implicated in resistance to protease inhibitors reduce viral fitness (Croteau et al., 1997; Mammano et al., 1998; Martinez-Picado et al., 1999). A possible explanation for the lower fitness associated with lower susceptibility to protease inhibitors is the reduced binding affinity of the protease mutants for the substrate as well as the inhibitor (Schock et al., 1996). Viral genomes with mutant protease genes can regain fitness by two mechanisms. First, mutations outside the protease active site can increase the catalytic efficiency of protease by increasing the turnover number without changing the reduced inhibitor binding (Pazhanisamy et al., 1996; Schock et al., 1996). Second, mutations at one or more protease cleavage sites can occur to enhance the rate of cleavage of the substrate by

the attenuated enzyme, as has been shown for the Gag cleavage sites in nucleocapsid-p1-p6 (Doyon et al., 1996; Zhang et al., 1997b; Mammano et al., 1998). Compensation for reduced activity due to inhibitor resistance has also been demonstrated for RT (Boyer et al., 1998).

It should also be pointed out that the *in vitro* generation of second-site revertants of mutations introduced into the HIV genome is a powerful genetic approach to map physical or functional interactions between different elements of HIV. For example, this method has been used to map RNA secondary structures (Garcia et al., 1989; Emiliani et al., 1996; Berkhout et al., 1997), and to examine interactions between p17/MA and gp41 (Mammano et al., 1995) and between HIV genomic RNA and Gag (Liang et al., 1999; Pailart and Gottlinger, 1999).

5. Summary

The study of HIV-1 sequence variability will continue to represent an important strategy for understanding the selective pressures on the virus and the mechanisms of escape from those selective pressures. Virus evolution over time and the presence of co-existing viral populations within a person point to dynamic interactions between the host and virus. These interactions include host immune responses, evolution of coreceptor use, and compartmentalization. Initial studies have shown a clear correlation between viral diversity and disease progression, and an understanding of this relationship will become an integral part of our perception of the disease of AIDS.

Escape from selective pressure will always be represented by specific sequence changes. Thus, genetic analysis will continue to be a primary tool for defining the nature of drug resistance and will become increasingly linked to understanding the host immune response. An important underlying question is the extent to which a small effective population size may be fixing mutations through stochastic processes as opposed to a large effective size driven only by selection for fitness.

Finally, the study of sequence variability will assume an important role in vaccine trials. An

important component of understanding vaccine efficacy will be correlating sequence divergence within the transmitting population with vaccine failures to define the role of virus heterogeneity in vaccine success/failure. Thus, the study of HIV sequence diversity within and between people will continue to provide us with important insights as we struggle against this pathogen.

Acknowledgements

The authors were supported by grants from the NIH.

References

- Allen, T.M., O'Connor, D.H., Jing, P., Dzuris, J.L., Mothe, B.R., Vogel, T.U., et al., 2000. Tat-specific cytotoxic T lymphocytes select for SIV escape variants during resolution of primary viraemia. *Nature* 407 (6802), 386–390.
- Anderson, D.J., Politch, J.A., Tucker, L.D., Fichorova, R., Haimovici, F., Tuomala, R.E., et al., 1998. Quantitation of mediators of inflammation and immunity in genital tract secretions and their relevance to HIV type 1 transmission. *AIDS Res. Hum. Retroviruses* 14 (Suppl. 1), S43–S49.
- Anderson, J.P., Rodrigo, A.G., Learn, G.H., Madan, A., Delahunty, C., Coon, M., et al., 2000. Testing the hypothesis of a recombinant origin of human immunodeficiency virus type 1 subtype E. *J. Virol.* 74 (22), 10752–10765.
- Arendrup, M., Nielsen, C., Hansen, J.E., Pedersen, C., Mathiesen, L., Nielsen, J.O., 1992. Autologous HIV-1 neutralizing antibodies: emergence of neutralization-resistant escape virus and subsequent development of escape virus neutralizing antibodies. *J. Acquired Immune Defic. Syndr.* 5 (3), 303–307.
- Arion, D., Kaushik, N., McCormick, S., Borkow, G., Parniak, M.A., 1998. Phenotypic mechanism of HIV-1 resistance to 3'-azido-3'-deoxythymidine (AZT): increased polymerization processivity and enhanced sensitivity to pyrophosphate of the mutant viral reverse transcriptase. *Biochemistry* 37 (45), 15908–15917.
- Asjo, B., Morfeldt-Manson, L., Albert, J., Biberfeld, G., Karlsson, A., Lidman, K., et al., 1986. Replicative capacity of human immunodeficiency virus from patients with varying severity of HIV infection. *Lancet* ii (8508), 660–662.
- Ball, J.K., Holmes, E.C., Whitwell, H., Desselberger, U., 1994. Genomic variation of human immunodeficiency virus type 1 (HIV-1): molecular analyses of HIV-1 in sequential blood samples and various organs obtained at autopsy. *J. Gen. Virol.* 75 (Pt 4), 67–79.
- Barre-Sinoussi, F., Chermann, J.C., Rey, F., Nugeyre, M.T., Chamaret, S., Gruest, J., et al., 1983. Isolation of a T-

- lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS). *Science* 220 (4599), 868–871.
- Beer, B.E., Bailes, E., Sharp, P.M., Hirsch, V.M., 1999. Diversity and evolution of primate lentiviruses. In: Kuiken, C.L., Foley, B., Hahn, B., Korber, B., McCutchan, F., Marx, P.A., Mellors, J.W., Mullins, J.I., Sodroski, J., Wolinsky, S. (Eds.), *Human Retroviruses and AIDS 1999: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM.
- Benn, S., Rutledge, R., Folks, T., Gold, J., Baker, L., McCormick, J., et al., 1985. Genomic heterogeneity of AIDS retroviral isolates from North America and Zaire. *Science* 230 (4728), 949–951.
- Berger, E.A., Doms, R.W., Fenyo, E.M., Korber, B.T., Littman, D.R., Moore, J.P., et al., 1998. A new classification for HIV-1. *Nature* 391 (6664), 240.
- Berger, E.A., Murphy, P.M., Farber, J.M., 1999. Chemokine receptors as HIV-1 coreceptors: roles in viral entry, tropism, and disease. *Annu. Rev. Immunol.* 17, 657–700.
- Berkhout, B., Klaver, B., Das, A.T., 1997. Forced evolution of a regulatory RNA helix in the HIV-1 genome. *Nucleic Acids Res.* 25 (5), 940–947.
- Betts, M.R., Krowka, J.F., Kepler, T.B., Davidian, M., Christopherson, C., Kwok, S., et al., 1999. Human immunodeficiency virus type 1-specific cytotoxic T lymphocyte activity is inversely correlated with HIV type 1 viral load in HIV type 1-infected long-term survivors. *AIDS Res. Hum. Retroviruses* 15 (13), 1219–1228.
- Birk, M., Sonnerborg, A., 1998. Variations in HIV-1 *pol* gene associated with reduced sensitivity to antiretroviral drugs in treatment-naïve patients. *AIDS* 12 (18), 2369–2375.
- Boden, D., Hurley, A., Zhang, L., Cao, Y., Guo, Y., Jones, E., et al., 1999. HIV-1 drug resistance in newly infected individuals. *J. Am. Med. Assoc.* 282 (12), 1135–1141.
- Borrow, P., Lewicki, H., Hahn, B.H., Shaw, G.M., Oldstone, M.B., 1994. Virus-specific CD8⁺ cytotoxic T-lymphocyte activity associated with control of viremia in primary human immunodeficiency virus type 1 infection. *J. Virol.* 68 (9), 6103–6110.
- Borrow, P., Lewicki, H., Wei, X., Horwitz, M.S., Pfeffer, N., Meyers, H., et al., 1997. Antiviral pressure exerted by HIV-1-specific cytotoxic T lymphocytes (CTLs) during primary infection demonstrated by rapid selection of CTL escape virus. *Nat. Med.* 3 (2), 205–211.
- Boyer, P.L., Gao, H.Q., Hughes, S.H., 1998. A mutation at position 190 of human immunodeficiency virus type 1 reverse transcriptase interacts with mutations at positions 74 and 75 via the template primer. *Antimicrob. Agents Chemother.* 42 (2), 447–452.
- Bradney, A.P., Scheer, S., Crawford, J.M., Buchbinder, S.P., Montefiori, D.C., 1999. Neutralization escape in human immunodeficiency virus type 1-infected long-term nonprogressors. *J. Infect. Dis.* 179 (5), 1264–1267.
- Cann, A.J., Churcher, M.J., Boyd, M., O'Brien, W., Zhao, J.Q., Zack, J., et al., 1992. The region of the envelope gene of human immunodeficiency virus type 1 responsible for determination of cell tropism. *J. Virol.* 66 (1), 305–309.
- Cao, Y., Qin, L., Zhang, L., Safritz, J., Ho, D.D., 1995. Virologic and immunologic characterization of long-term survivors of human immunodeficiency virus type 1 infection. *N. Engl. J. Med.* 332 (4), 201–208.
- Carr, J.K., Foley, B.T., Leitner, T., Salminen, M., Korber, B., McCutchan, F., 1998. Reference sequences representing the principal genetic diversity of HIV-1 in the pandemic. In: Korber, B., Kuiken, C.L., Foley, B., Hahn, B., McCutchan, F., Mellors, J.W., Sodroski, J. (Eds.), *Human Retroviruses and AIDS 1998: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM.
- Carrillo, A., Ratner, L., 1996. Human immunodeficiency virus type 1 tropism for T-lymphoid cell lines: role of the V3 loop and C4 envelope determinants. *J. Virol.* 70 (2), 1301–1309.
- CDC, 1981. Kaposi's sarcoma and Pneumocystis pneumonia among homosexual men — New York City and California. *MMWR Morb. Mortal. Wkly. Rep.* 30 (25), 305–308.
- Chen, Z., Telfer, P., Gettie, A., Reed, P., Zhang, L., Ho, D.D., Marx, P.A., 1996. Genetic characterization of new west African simian immunodeficiency virus SIVsm: geographic clustering of household-derived SIV strains with human immunodeficiency virus type 2 subtypes and genetically diverse viruses from a single feral sooty mangabey troop. *J. Virol.* 70 (6), 3617–3627.
- Chen, Z., Luckay, A., Sodora, D.L., Telfer, P., Reed, P., Gettie, A., et al., 1997. Human immunodeficiency virus type 2 (HIV-2) seroprevalence and characterization of a distinct HIV-2 genetic subtype from the natural range of simian immunodeficiency virus-infected sooty mangabeys. *J. Virol.* 71 (5), 3953–3960.
- Cheng-Mayer, C., Liu, R., Landau, N.R., Stamatatos, L., 1997. Macrophage tropism of human immunodeficiency virus type 1 and utilization of the CC-CKR5 coreceptor. *J. Virol.* 71 (2), 1657–1661.
- Chesebro, B., Wehrly, K., Nishio, J., Perryman, S., 1992. Macrophage-tropic human immunodeficiency virus isolates from different patients exhibit unusual V3 envelope sequence homogeneity in comparison with T-cell-tropic isolates: definition of critical amino acids involved in cell tropism. *J. Virol.* 66 (11), 6547–6554.
- Chesebro, B., Wehrly, K., Nishio, J., Perryman, S., 1996. Mapping of independent V3 envelope determinants of human immunodeficiency virus type 1 macrophage tropism and syncytium formation in lymphocytes. *J. Virol.* 70 (12), 9055–9059.
- Clavel, F., Guetard, D., Brun-Vezinet, F., Chamaret, S., Rey, M.A., Santos-Ferreira, M.O., et al., 1986. Isolation of a new human retrovirus from West African patients with AIDS. *Science* 233 (4761), 343–346.

- Clemetson, D.B., Moss, G.B., Willerford, D.M., Hensel, M., Emonyi, W., Holmes, K.K., et al., 1993. Detection of HIV DNA in cervical and vaginal secretions. Prevalence and correlates among women in Nairobi, Kenya. *J. Am. Med. Assoc.* 269 (22), 2860–2864.
- Cocchi, F., DeVico, A.L., Garzino-Demo, A., Cara, A., Gallo, R.C., Lusso, P., 1996. The V3 domain of the HIV-1 gp120 envelope glycoprotein is critical for chemokine-mediated blockade of infection. *Nat. Med.* 2 (11), 1244–1247.
- Coffin, J.M., 1995. HIV population dynamics in vivo: implications for genetic variation, pathogenesis, and therapy. *Science* 267 (5197), 483–489.
- Coffin, J.M., 1996. HIV viral dynamics. *AIDS* 10 (Suppl. 3), S75–S84.
- Cohen, E.A., Terwilliger, E.F., Sodroski, J.G., Haseltine, W.A., 1988. Identification of a protein encoded by the *vpu* gene of HIV-1. *Nature* 334 (6182), 532–534.
- Connor, R.I., Sheridan, K.E., Ceradini, D., Choe, S., Landau, N.R., 1997. Change in coreceptor use correlates with disease progression in HIV-1-infected individuals. *J. Exp. Med.* 185 (4), 621–628.
- Coombs, R.W., Speck, C.E., Hughes, J.P., Lee, W., Sampoleo, R., Ross, S.O., et al., 1998. Association between culturable human immunodeficiency virus type 1 (HIV-1) in semen and HIV-1 RNA levels in semen and blood: evidence for compartmentalization of HIV-1 between semen and blood. *J. Infect. Dis.* 177 (2), 320–330.
- Corbet, S., Muller-Trutwin, M.C., Versmisse, P., Delarue, S., Ayoub, A., Lewis, J., et al., 2000. *env* sequences of simian immunodeficiency viruses from chimpanzees in Cameroon are strongly related to those of human immunodeficiency virus group N from the same geographic area. *J. Virol.* 74 (1), 529–534.
- Croteau, G., Doyon, L., Thibeault, D., McKercher, G., Pilote, L., Lamarre, D., 1997. Impaired fitness of human immunodeficiency virus type 1 variants with high-level resistance to protease inhibitors. *J. Virol.* 71 (2), 1089–1096.
- de Jong, J.J., Goudsmit, J., Keulen, W., Klaver, B., Krone, W., Tersmette, M., et al., 1992. Human immunodeficiency virus type 1 clones chimeric for the envelope V3 domain differ in syncytium formation and replication capacity. *J. Virol.* 66 (2), 757–765.
- Deeks, S.G., Hecht, F.M., Swanson, M., Elbeik, T., Loftus, R., Cohen, P.T., et al., 1999. HIV RNA and CD4 cell count response to protease inhibitor therapy in an urban AIDS clinic: response to both initial and salvage therapy. *AIDS* 13 (6), F35–F43.
- Delassus, S., Cheynier, R., Wain-Hobson, S., 1992. Nonhomogeneous distribution of human immunodeficiency virus type 1 proviruses in the spleen. *J. Virol.* 66 (9), 5642–5645.
- Delviks, K.A., Pathak, V.K., 1999. Effect of distance between homologous sequences and 3' homology on the frequency of retroviral reverse transcriptase template switching. *J. Virol.* 73 (10), 7923–7932.
- Delwart, E.L., Mullins, J.I., Gupta, P., Learn, G.H. Jr, Holodniy, M., Katzenstein, D., et al., 1998. Human immunodeficiency virus type 1 populations in blood and semen. *J. Virol.* 72 (1), 617–623.
- Delwart, E.L., Pan, H., Sheppard, H.W., Wolpert, D., Neumann, A.U., Korber, B., et al., 1997. Slower evolution of human immunodeficiency virus type 1 quasispecies during progression to AIDS. *J. Virol.* 71 (10), 7498–7508.
- Domingo, E., Holland, J.J., 1997. RNA virus mutations and fitness for survival. *Annu. Rev. Microbiol.* 51, 151–178.
- Donaldson, Y.K., Bell, J.E., Holmes, E.C., Hughes, E.S., Brown, H.K., Simmonds, P., 1994. In vivo distribution and cytopathology of variants of human immunodeficiency virus type 1 showing restricted sequence variability in the V3 loop. *J. Virol.* 68 (9), 5991–6005.
- Doyon, L., Croteau, G., Thibeault, D., Poulin, F., Pilote, L., Lamarre, D., 1996. Second locus involved in human immunodeficiency virus type 1 resistance to protease inhibitors. *J. Virol.* 70 (6), 3763–3769.
- Eigen, M., 1971. Selforganization of matter and the evolution of biological macromolecules. *Naturwissenschaften* 58 (10), 465–523.
- Emiliani, S., Van Lint, C., Fischle, W., Paras, P. Jr, Ott, M., Brady, J., Verdin, E., 1996. A point mutation in the HIV-1 Tat responsive element is associated with postintegration latency. *Proc. Natl. Acad. Sci. USA* 93 (13), 6377–6381.
- Epstein, L.G., Kuiken, C., Blumberg, B.M., Hartman, S., Sharer, L.R., Clement, M., et al., 1991. HIV-1 V3 domain variation in brain and spleen of children with AIDS: tissue-specific evolution within host-determined quasispecies. *Virology* 180 (2), 583–590.
- Erice, A., Mayers, D.L., Strike, D.G., Sannerud, K.J., McCutchan, F.E., Henry, K., et al., 1993. Brief report: primary infection with zidovudine-resistant human immunodeficiency virus type 1. *N. Engl. J. Med.* 328 (16), 1163–1165.
- Eron, J.J., Vernazza, P.L., Johnston, D.M., Seillier-Moisewitsch, F., Alcorn, T.M., Fiscus, S.A., et al., 1998. Resistance of HIV-1 to antiretroviral agents in blood and seminal plasma: implications for transmission. *AIDS* 12 (15), F181–F189.
- Evans, D.T., O'Conner, D.H., Jing, P., Dzuris, J.L., Sidney, J., da Silva, J., et al., 1999. Virus-specific cytotoxic T-lymphocyte responses select for amino-acid variation in simian immunodeficiency virus Env and Nef. *Nat. Med.* 5 (11), 1270–1276.
- Fatkenheuer, G., Theisen, A., Rockstroh, J., Grabow, T., Wicke, C., Becker, K., et al., 1997. Virological treatment failure of protease inhibitor therapy in an unselected cohort of HIV-infected patients. *AIDS* 11 (14), F113–F116.
- Fauci, A.S., Desrosiers, R.C., 1997. Pathogenesis of HIV and SIV. In: Coffin, J.M., Hughes, S.H., Varmus, H.E. (Eds.), *Retroviruses*. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 587–636.
- Felsenstein, J., 1993. PHYLIP — phylogeny inference package (Version 3.2). *Cladistics* 5, 164–166.
- Fontaine, E., Lambert, C., Servais, J., Ninove, D., Plessier, J.M., Staub, T., et al., 1998. Fast genotypic detection of drug resistance mutations in the HIV-1 reverse transcriptase gene of treatment-naïve patients. *J. Hum. Virol.* 1 (7), 451–456.

- Fouchier, R.A.M., Groenink, M., Kootstra, N.A., Tersmette, M., Huisman, H.G., Miedema, F., et al., 1992. Phenotype-associated sequence variation in the third variable domain of the human immunodeficiency virus type 1 gp120 molecule. *J. Virol.* 66 (5), 3183–3187.
- Freel, S.A., Williams, J.M., Nelson, J.A.E., Patton, L.L., Fiscus, S.A., Swanstrom, R. et al. (2001) Characterization of human immunodeficiency virus type 1 in saliva and blood plasma by V3-specific heteroduplex tracking assay and genotype analyses *J. Virol.* 75 (10), 4936–4940.
- Gao, F., Bailes, E., Robertson, D.L., Chen, Y., Rodenburg, C.M., Michael, S.F., et al., 1999. Origin of HIV-1 in the chimpanzee *Pan troglodytes troglodytes*. *Nature* 397 (6718), 436–441.
- Gao, F., Morrison, S.G., Robertson, D.L., Thornton, C.L., Craig, S., Karlsson, G., et al., 1996. Molecular cloning and analysis of functional envelope genes from human immunodeficiency virus type 1 sequence subtypes A through G. *J. Virol.* 70 (3), 1651–1667.
- Gao, F., Yue, L., Robertson, D.L., Hill, S.C., Hui, H., Biggar, R.J., et al., 1994. Genetic diversity of human immunodeficiency virus type 2: evidence for distinct sequence subtypes with differences in virus biology. *J. Virol.* 68 (11), 7433–7447.
- Gao, F., Yue, L., White, A.T., Pappas, P.G., Barchue, J., Hanson, A.P., et al., 1992. Human infection by genetically diverse SIVsm-related HIV-2 in West Africa. *Nature* 358 (6386), 495–499.
- Garcia, J.A., Harrich, D., Sultankis, E., Wu, F., Mitsuyasu, R., Gaynor, R.B., 1989. Human immunodeficiency virus type 1 LTR TATA and TAR region sequences required for transcriptional regulation. *EMBO J.* 8 (3), 765–778.
- George, J.R., Ou, C.Y., Parekh, B., Brattegaard, K., Brown, V., Boateng, E., et al., 1992. Prevalence of HIV-1 and HIV-2 mixed infections in Cote d'Ivoire. *Lancet* 340 (8815), 337–339.
- Gomez-Cano, M., Rubio, A., Puig, T., Perez-Olmeda, M., Ruiz, L., Soriano, V., et al., 1998. Prevalence of genotypic resistance to nucleoside analogues in antiretroviral-naïve and antiretroviral-experienced HIV-infected patients in Spain. *AIDS* 12 (9), 1015–1020.
- Gottlieb, M.S., Schroff, R., Schanker, H.M., Weisman, J.D., Fan, P.T., Wolf, R.A., et al., 1981. *Pneumocystis carinii* pneumonia and mucosal candidiasis in previously healthy homosexual men: evidence of a new acquired cellular immunodeficiency. *N. Engl. J. Med.* 305 (24), 1425–1431.
- Goulder, P.J.R., Phillips, R.E., Colbert, R.A., McAdam, S., Ogg, G., Nowak, M.A., et al., 1997. Late escape from an immunodominant cytotoxic T-lymphocyte response associated with progression to AIDS. *Nat. Med.* 3 (2), 212–217.
- Groenink, M., Andeweg, A.C., Fouchier, R.A.M., Broersen, S., van der Jagt, R.C., Schuitemaker, H., et al., 1992. Phenotype-associated *env* gene variation among eight related human immunodeficiency virus type 1 clones: evidence for in vivo recombination and determinants of cytotropism outside the V3 domain. *J. Virol.* 66 (10), 6175–6180.
- Gupta, P., Kingsley, L., Armstrong, J., Ding, M., Cottrill, M., Rinaldo, C., 1993. Enhanced expression of human immunodeficiency virus type 1 correlates with development of AIDS. *Virology* 196 (2), 586–595.
- Gürtler, L.G., Zekeng, L., Tsague, J.M., van Brunn, A., Afane Ze, E., Eberle, J., et al., 1996. HIV-1 subtype O: epidemiology, pathogenesis, diagnosis, and perspectives of the evolution of HIV. *Arch. Virol. Suppl.* 11, 195–202.
- Guyader, M., Emerman, M., Sonigo, P., Clavel, F., Montagnier, L., Alizon, M., 1987. Genome organization and transactivation of the human immunodeficiency virus type 2. *Nature* 326 (6114), 662–669.
- Haase, A.T., Henry, K., Zupancic, M., Sedgewick, G., Faust, R.A., Melroe, H., et al., 1996. Quantitative image analysis of HIV-1 infection in lymphoid tissue. *Science* 274 (5289), 985–989.
- Hahn, B.H., Gonda, M.A., Shaw, G.M., Popovic, M., Hoxie, J.A., Gallo, R.C., et al., 1985. Genomic diversity of the acquired immune deficiency syndrome virus HTLV-III: different viruses exhibit greatest divergence in their envelope genes. *Proc. Natl. Acad. Sci. USA* 82 (14), 4813–4817.
- Hahn, B.H., Shaw, G.M., Taylor, M.E., Redfield, R.R., Markham, P.D., Salahuddin, S.Z., et al., 1986. Genetic variation in HTLV-III/LAV over time in patients with AIDS or at risk for AIDS. *Science* 232 (4757), 1548–1553.
- Harouse, J.M., Gettie, A., Tan, R.C., Blanchard, J., Cheng-Mayer, C., 1999. Distinct pathogenic sequela in rhesus macaques infected with CCR5 or CXCR4 utilizing SHIVs. *Science* 284 (5415), 816–819.
- Hay, C.M., Ruhl, D.J., Basgoz, N.O., Wilson, C.C., Billingsley, J.M., DePasquale, M.P., et al., 1999. Lack of viral escape and defective in vivo activation of human immunodeficiency virus type 1-specific cytotoxic T lymphocytes in rapidly progressive infection. *J. Virol.* 73 (7), 5509–5519.
- Hecht, F.M., Grant, R.M., Petropoulos, C.J., Dillon, B., Chesney, M.A., Tian, H., et al., 1998. Sexual transmission of an HIV-1 variant resistant to multiple reverse transcriptase and protease inhibitors. *N. Engl. J. Med.* 339 (5), 307–311.
- Henderson, L.E., Sowder, R.C., Copeland, T.D., Benveniste, R.E., Oroszlan, S., 1988. Isolation and characterization of a novel protein (X-ORF product) from SIV and HIV-2. *Science* 241 (4862), 199–201.
- Hillis, D.M., 2000. How to resolve the debate on the origin of AIDS. *Science* 289 (5486), 1877–1878.
- Hirsch, V.M., Olmsted, R.A., Murphey-Corb, M., Purcell, R.H., Johnson, P.R., 1989. An African primate lentivirus (SIVsm) closely related to HIV-2. *Nature* 339 (6223), 389–392.
- Hooper, E., 1999. The River: A Journey to the Source of HIV and AIDS. Little Brown and Company 1104 pp.
- Hu, W.S., Temin, H.M., 1990. Genetic consequences of packaging two RNA genomes in one retroviral particle: pseudodiploidy and high rate of genetic recombination. *Proc. Natl. Acad. Sci. USA* 87 (4), 1556–1560.

- Hughes, E.S., Bell, J.E., Simmonds, P., 1997. Investigation of population diversity of human immunodeficiency virus type 1 in vivo by nucleotide sequencing and length polymorphism analysis of the V1/V2 hypervariable region of *env*. *J. Gen. Virol.* 78 (Pt 11), 2871–2882.
- Hwang, S.S., Boyle, T.J., Lyster, H.K., Cullen, B.R., 1991. Identification of the envelope V3 loop as the primary determinant of cell tropism in HIV-1. *Science* 253 (5015), 71–74.
- Jin, X., Bauer, D.E., Tuttleton, S.E., Lewin, S., Gettie, A., Blanchard, J., et al., 1999. Dramatic rise in plasma viremia after CD8(+) T cell depletion in simian immunodeficiency virus-infected macaques. *J. Exp. Med.* 189 (6), 991–998.
- Julias, J.G., Kim, T., Arnold, G., Pathak, V.K., 1997. The antiretrovirus drug 3'-azido-3'-deoxythymidine increases the retrovirus mutation rate. *J. Virol.* 71 (6), 4254–4263.
- Kakizawa, J., Ushijima, H., Morishita, Y., Oka, S., Ikeda, Y., Muller, W.E., 1996. Diversity of HIV type 1 envelope V3 loop region in saliva. *AIDS Res. Hum. Retroviruses* 12 (6), 561–563.
- Kappes, J.C., Morrow, C.D., Lee, S.W., Jameson, B.A., Kent, S.B., Hood, L.E., et al., 1988. Identification of a novel retroviral gene unique to human immunodeficiency virus type 2 and simian immunodeficiency virus SIV_{MAC}. *J. Virol.* 62 (9), 3501–3505.
- Kashuba, A.D., Dyer, J.R., Kramer, L.M., Raasch, R.H., Eron, J.J., Cohen, M.S., 1999. Antiretroviral-drug concentrations in semen: implications for sexual transmission of human immunodeficiency virus type 1. *Antimicrob. Agents Chemother.* 43 (8), 1817–1826.
- Keys, B., Karis, J., Fadeel, B., Valentin, A., Norkrans, G., Hagberg, L., et al., 1993. V3 sequences of paired HIV-1 isolates from blood and cerebrospinal fluid cluster according to host and show variation related to the clinical stage of disease. *Virology* 196 (2), 475–483.
- Kiessling, A.A., Fitzgerald, L.M., Zhang, D., Chhay, H., Brettler, D., Eyre, R.C., et al., 1998. Human immunodeficiency virus in semen arises from a genetically distinct virus reservoir. *AIDS Res. Hum. Retroviruses* 14 (Suppl. 1), S33–S41.
- Kim, T., Mudry, R.A. Jr, Rexrode, C.A. 2nd, Pathak, V.K., 1996. Retroviral mutation rates and A-to-G hypermutations during different stages of retroviral replication. *J. Virol.* 70 (11), 7594–7602.
- Koot, M., Keet, I.P., Vos, A.H., de Goede, R.E.Y., Roos, M.T., Coutinho, R.A., et al., 1993. Prognostic value of HIV-1 syncytium-inducing phenotype for rate of CD4+ cell depletion and progression to AIDS. *Ann. Intern. Med.* 118 (9), 681–688.
- Korber, B., Bhattacharya, T., Theiler, J., Gupta, J., Lapedes, A., Hahn, B., et al., 2000a. Search for the origin of HIV and AIDS. *Science* 289 (5482), 1140.
- Korber, B., Muldoon, M., Theiler, J., Gao, F., Gupta, R., Lapedes, A., et al., 2000b. Timing the ancestor of the HIV-1 pandemic strains. *Science* 288 (5472), 1789–1796.
- Korber, B., Theiler, J., Wolinsky, S., 1998. Limitations of a molecular clock applied to considerations of the origin of HIV-1. *Science* 280 (5371), 1868–1871.
- Korber, B.T., Kunstman, K.J., Patterson, B.K., Furtado, M., McEvilly, M.M., Levy, R., et al., 1994. Genetic differences between blood- and brain-derived viral sequences from human immunodeficiency virus type 1-infected patients: evidence of conserved elements in the V3 region of the envelope protein of brain-derived sequences. *J. Virol.* 68 (11), 7467–7481.
- Koup, R.A., Safrit, J.T., Cao, Y., Andrews, C.A., McLeod, G., Borkowsky, W., et al., 1994. Temporal association of cellular immune responses with the initial control of viremia in primary human immunodeficiency virus type 1 syndrome. *J. Virol.* 68 (7), 4650–4655.
- Kreiss, J., Willerford, D.M., Hensel, M., Emonyi, W., Plummer, F., Ndinya-Achola, J., et al., 1994. Association between cervical inflammation and cervical shedding of human immunodeficiency virus DNA. *J. Infect. Dis.* 170 (6), 1597–1601.
- Kuiken, C.L., Foley, B., Hahn, B., Korber, B., McCutchan, F., Marx, P.A., et al. (Eds.), 1999. *Human Retroviruses and AIDS 1999: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM.
- Lathey, J.L., Pratt, R.D., Spector, S.A., 1997. Appearance of autologous neutralizing antibody correlates with reduction in virus load and phenotype switch during primary infection with human immunodeficiency virus type 1. *J. Infect. Dis.* 175, 231–232.
- Lefrere, J.J., Roudot-Thoraval, F., Mariotti, M., Thauvin, M., Lerable, J., Salpêtrier, J., et al., 1998. The risk of disease progression is determined during the first year of human immunodeficiency virus type 1 infection. *J. Infect. Dis.* 177 (6), 1541–1548.
- Leigh Brown, A.J., 1997. Analysis of HIV-1 *env* gene sequences reveals evidence for a low effective number in the viral population. *Proc. Natl. Acad. Sci. USA* 94 (5), 1862–1865.
- Leigh Brown, A.J., Richman, D.D., 1997. HIV-1: gambling on the evolution of drug resistance? *Nat. Med.* 3 (3), 268–271.
- Leitner, T., Albert, J., 1999. The molecular clock of HIV-1 unveiled through analysis of a known transmission history. *Proc. Natl. Acad. Sci. USA* 96 (19), 10752–10757.
- Liang, C., Rong, L., Quan, Y., Laughrea, M., Kleiman, L., Wainberg, M.A., 1999. Mutations within four distinct Gag proteins are required to restore replication of human immunodeficiency virus type 1 after deletion mutagenesis within the dimerization initiation site. *J. Virol.* 73 (8), 7014–7020.
- Lineberger, D.W., Kessler, J.A., Waterbury, J.A., Byrnes, V.W., Massari, F., Staszewski, S., et al., 1995. Turnover of circulating virion RNA and of cell-associated viral DNA reflects active viral replication in human immunodeficiency virus type 1-infected individuals. *J. Virol.* 69 (4), 2637–2639.
- Little, S.J., Daar, E.S., D'Aquila, R.T., Keiser, P.H., Connick, E., Whitcomb, J.M., et al., 1999. Reduced antiretroviral drug susceptibility among patients with primary HIV infection. *J. Am. Med. Assoc.* 282 (12), 1142–1149.

- Liu, S.L., Schacker, T., Musey, L., Shriner, D., McElrath, M.J., Corey, L., et al., 1997. Divergent patterns of progression to AIDS after infection from the same source: human immunodeficiency virus type 1 evolution and antiviral responses. *J. Virol.* 71 (6), 4284–4295.
- Liu, G., Chirrianni, A., Clementi, M., Bagnarelli, P., Valenza, A., Cataldo, P.T., et al., 1996. Analysis of HIV-1 load in blood, semen and saliva: evidence for different viral compartments in a cross-sectional and longitudinal study. *AIDS* 10 (14), F51–F56.
- Long, E.M., Martin, H.L. Jr, Kreiss, J.K., Rainwater, S.M., Lavreys, L., Jackson, D.J., et al., 2000. Gender differences in HIV-1 diversity at time of infection. *Nat. Med.* 6 (1), 71–75.
- Lukashov, V.V., Goudsmit, J., 1998. HIV heterogeneity and disease progression in AIDS: a model of continuous virus adaptation. *AIDS* 12 (Suppl. A), S43–S52.
- Lukashov, V.V., Kuiken, C.L., Goudsmit, J., 1995. Intrahost human immunodeficiency virus type 1 evolution is related to length of the immunocompetent period. *J. Virol.* 69 (11), 6911–6916.
- Mammano, F., Kondo, E., Sodroski, J., Bukovsky, A., Gottlinger, H.G., 1995. Rescue of human immunodeficiency virus type 1 matrix protein mutants by envelope glycoproteins with short cytoplasmic domains. *J. Virol.* 69 (6), 3824–3830.
- Mammano, F., Petit, C., Clavel, F., 1998. Resistance-associated loss of viral fitness in human immunodeficiency virus type 1: phenotypic analysis of protease and *gag* coevolution in protease inhibitor-treated patients. *J. Virol.* 72 (9), 7632–7637.
- Mansky, L.M., 1996a. Forward mutation rate of human immunodeficiency virus type 1 in a T lymphoid cell line. *AIDS Res. Hum. Retroviruses* 12 (4), 307–314.
- Mansky, L.M., 1996b. The mutation rate of human immunodeficiency virus type 1 is influenced by the *vpr* gene. *Virology* 222 (2), 391–400.
- Mansky, L.M., Bernard, L.C., 2000. 3'-Azido-3'-deoxythymidine (AZT) and AZT-resistant reverse transcriptase can increase the in vivo mutation rate of human immunodeficiency virus type 1. *J. Virol.* 74 (20), 9532–9539.
- Mansky, L.M., Preveral, S., Selig, L., Benarous, R., Benichou, S., 2000. The interaction of VPR with uracil DNA glycosylase modulates the human immunodeficiency virus type 1 in vivo mutation rate. *J. Virol.* 74 (15), 7039–7047.
- Mansky, L.M., Temin, H.M., 1995. Lower in vivo mutation rate of human immunodeficiency virus type 1 than that predicted from the fidelity of purified reverse transcriptase. *J. Virol.* 69 (8), 5087–5094.
- Martinez, M.A., Vartanian, J.P., Wain-Hobson, S., 1994. Hypermutagenesis of RNA using human immunodeficiency virus type 1 reverse transcriptase and biased dNTP concentrations. *Proc. Natl. Acad. Sci. USA* 91 (25), 11787–11791.
- Martinez-Picado, J., Savara, A.V., Sutton, L., D'Aquila, R.T., 1999. Replicative fitness of protease inhibitor-resistant mutants of human immunodeficiency virus type 1. *J. Virol.* 73 (5), 3744–3752.
- Masur, H., Michelis, M.A., Greene, J.B., Onorato, I., Stouwe, R.A., Holzman, R.S., et al., 1981. An outbreak of community-acquired *Pneumocystis carinii* pneumonia: initial manifestation of cellular immune dysfunction. *N. Engl. J. Med.* 305 (24), 1431–1438.
- McCutchan, F.E., Salminen, M.O., Carr, J.K., Burke, D.S., 1996. HIV-1 genetic diversity. *AIDS* 10 (Suppl.3), S13–S20.
- Mellors, J.W., Kingsley, L.A., Rinaldo, C.R. Jr, Todd, J.A., Hoo, B.S., Kokka, R.P., et al., 1995. Quantitation of HIV-1 RNA in plasma predicts outcome after seroconversion. *Ann. Intern. Med.* 122 (8), 573–579.
- Mellors, J.W., Rinaldo, C.R. Jr, Gupta, P., White, R.M., Todd, J.A., Kingsley, L.A., 1996. Prognosis in HIV-1 infection predicted by the quantity of virus in plasma. *Science* 272 (5265), 1167–1170.
- Meyer, P.R., Matsuura, S.E., So, A.G., Scott, W.A., 1998. Unblocking of chain-terminated primer by HIV-1 reverse transcriptase through a nucleotide-dependent mechanism. *Proc. Natl. Acad. Sci. USA* 95 (23), 13471–13476.
- Milich, L., Margolin, B., Swanstrom, R., 1993. V3 loop of the human immunodeficiency virus type 1 Env protein: interpreting sequence variability. *J. Virol.* 67 (9), 5623–5634.
- Milich, L., Margolin, B.H., Swanstrom, R., 1997. Patterns of amino acid variability in NSI-like and SI-like V3 sequences and a linked change in the CD4-binding domain of the HIV-1 Env protein. *Virology* 239 (1), 108–118.
- Montefiori, D.C., Pantaleo, G., Fink, L.M., Zhou, J.T., Zhou, J.Y., Bilska, M., et al., 1996. Neutralizing and infection-enhancing antibody responses to human immunodeficiency virus type 1 in long-term nonprogressors. *J. Infect. Dis.* 173 (1), 60–67.
- Moore, J.P., Trkola, A., Dragic, T., 1997. Co-receptors for HIV-1 entry. *Curr. Opin. Immunol.* 9, 551–562.
- Morris, A., Marsden, M., Halcrow, K., Hughes, E.S., Brett, R.P., Bell, J.E., et al., 1999. Mosaic structure of the human immunodeficiency virus type 1 genome infecting lymphoid cells and the brain: evidence for frequent in vivo recombination events in the evolution of regional populations. *J. Virol.* 73 (10), 8720–8731.
- Nelson, J.A.E., Baribaud, F., Edwards, T., Swanstrom, R., 2000. Rapid changes in HIV-1 populations in the absence of external selective pressure. *J. Virol.* 74 (18), 8494–8501.
- Nelson, J.A.E., Fiscus, S.A., Swanstrom, R., 1997. Evolutionary variants of the human immunodeficiency virus type 1 V3 region characterized by using a heteroduplex tracking assay. *J. Virol.* 71 (11), 8750–8758.
- Nyambi, P.N., Lewi, P., Peeters, M., Janssens, W., Heyndrickx, L., Fransen, K., et al., 1997. Study of the dynamics of neutralization escape mutants in a chimpanzee naturally infected with the simian immunodeficiency virus SIVcpz-ant. *J. Virol.* 71 (3), 2320–2330.
- Olsen, J.C., Bova-Hill, C., Grandgenett, D.P., Quinn, T.P., Manfredi, J.P., Swanstrom, R., 1990. Rearrangements in unintegrated retroviral DNA are complex and are the result of multiple genetic determinants. *J. Virol.* 64 (11), 5475–5484.

- Ostrowski, M.A., Chun, T.W., Justement, S.J., Motola, I., Spinelli, M.A., Adelsberger, J., et al., 1999. Both memory and CD45RA⁺/CD62L⁺ naive CD4⁺ T cells are infected in human immunodeficiency virus type 1-infected individuals. *J. Virol.* 73 (8), 6430–6435.
- Overbaugh, J., Anderson, R.J., Ndinya-Achola, J.O., Kreiss, J.K., 1996. Distinct but related human immunodeficiency virus type 1 variant populations in genital secretions and blood. *AIDS Res. Hum. Retroviruses* 12 (2), 107–115.
- Paillart, J.C., Gottlinger, H.G., 1999. Opposing effects of human immunodeficiency virus type 1 matrix mutations support a myristyl switch model of Gag membrane targeting. *J. Virol.* 73 (4), 2604–2612.
- Pang, S., Shlesinger, Y., Daar, E.S., Moudgil, T., Ho, D.D., Chen, I.S.Y., 1992. Rapid generation of sequence variation during primary HIV-1 infection. *AIDS* 6 (5), 453–460.
- Pantaleo, G., Menzo, S., Vaccarezza, M., Graziosi, C., Cohen, O.J., Demarest, J.F., et al., 1995. Studies in subjects with long-term nonprogressive human immunodeficiency virus infection. *N. Engl. J. Med.* 332 (4), 209–216.
- Pantaleo, G., Soudeyns, H., Demarest, J.F., Vaccarezza, M., Graziosi, C., Paolucci, S., et al., 1997. Accumulation of human immunodeficiency virus-specific cytotoxic T lymphocytes away from the predominant site of virus replication during primary infection. *Eur. J. Immunol.* 27 (12), 3166–3173.
- Parthasarathi, S., Varela-Echavarria, A., Ron, Y., Preston, B.D., Dougherty, J.P., 1995. Genetic rearrangements occurring during a single cycle of murine leukemia virus vector replication: characterization and implications. *J. Virol.* 69 (12), 7991–8000.
- Pathak, V.K., Temin, H.M., 1990a. Broad spectrum of in vivo forward mutations, hypermutations, and mutational hotspots in a retroviral shuttle vector after a single replication cycle: deletions and deletions with insertions. *Proc. Natl. Acad. Sci. USA* 87 (16), 6024–6028.
- Pathak, V.K., Temin, H.M., 1990b. Broad spectrum of in vivo forward mutations, hypermutations, and mutational hotspots in a retroviral shuttle vector after a single replication cycle: substitutions, frameshifts, and hypermutations. *Proc. Natl. Acad. Sci. USA* 87 (16), 6019–6023.
- Pazhanisamy, S., Stuver, C.M., Cullinan, A.B., Margolin, N., Rao, B.G., Livingston, D.J., 1996. Kinetic characterization of human immunodeficiency virus type-1 protease-resistant variants. *J. Biol. Chem.* 271 (30), 17979–17985.
- Peeters, M., Fransen, K., Delaporte, E., Vanden Haesevelde, M., Gershy-Damet, G.M., Kestens, L., et al., 1992. Isolation and characterization of a new chimpanzee lentivirus (simian immunodeficiency virus isolate cpz-ant) from a wild-captured chimpanzee. *AIDS* 6 (5), 447–451.
- Peeters, M., Honore, C., Huet, T., Bedjabaga, L., Ossari, S., Bussi, P., et al., 1989. Isolation and partial characterization of an HIV-related virus occurring naturally in chimpanzees in Gabon. *AIDS* 3 (10), 625–630.
- Pellegrin, I., Izopet, J., Reynes, J., Denayrolles, M., Montes, B., Pellegrin, J.L., et al., 1999. Emergence of zidovudine and multidrug-resistance mutations in the HIV-1 reverse transcriptase gene in therapy-naïve patients receiving stavudine plus didanosine combination therapy. *AIDS* 13 (13), 1705–1709.
- Perelson, A.S., Neumann, A.U., Markowitz, M., Leonard, J.M., Ho, D.D., 1996. HIV-1 dynamics in vivo: virion clearance rate, infected cell life-span, and viral generation time. *Science* 271 (5255), 1582–1586.
- Phillips, J., Qureshi, N., Barr, C., Henrard, D.R., 1994. Low level of cell-free virus detected at high frequency in saliva from HIV-1-infected individuals. *AIDS* 8 (7), 1011–1012.
- Phillips, R.E., Rowland-Jones, S., Nixon, D.F., Gotch, F.M., Edwards, J.P., Ogunlesi, A.O., et al., 1991. Human immunodeficiency virus genetic variation that can escape cytotoxic T cell recognition. *Nature* 354 (6353), 453–459.
- Pieniazek, D., Peralta, J.M., Ferreira, J.A., Krebs, J.W., Owen, S.M., Sion, F.S., et al., 1991. Identification of mixed HIV-1/HIV-2 infections in Brazil by polymerase chain reaction. *AIDS* 5 (11), 1293–1299.
- Pilgrim, A.K., Pantaleo, G., Cohen, O.J., Fink, L.M., Zhou, J.Y., Zhou, J.T., et al., 1997. Neutralizing antibody responses to human immunodeficiency virus type 1 in primary infection and long-term-nonprogressive infection. *J. Infect. Dis.* 176 (4), 924–932.
- Ping, L., Cohen, M.S., Hoffman, I., Vernazza, P., Seillier-Moisewitsch, F., Chakraborty, H., et al., 2000. Effects of genital tract inflammation on HIV-1 V3 populations in blood and semen. *J. Virol.* 74 (19), 8946–8952.
- Plotkin, S.A., Koprowski, H., 2000. Search for the origin of HIV and AIDS. *Science* 289 (5482), 1141.
- Poignard, P., Sabbe, R., Picchio, G.R., Wang, M., Gulizia, R.J., Katinger, H., et al., 1999. Neutralizing antibodies have limited effects on the control of established HIV-1 infection in vivo. *Immunity* 10 (4), 431–438.
- Poss, M., Martin, H.L., Kreiss, J.K., Granville, L., Chohan, B., Nyange, P., et al., 1995. Diversity in virus populations from genital secretions and peripheral blood from women recently infected with human immunodeficiency virus type 1. *J. Virol.* 69 (12), 8118–8122.
- Poss, M., Rodrigo, A.G., Gosink, J.J., Learn, G.H., De Vange Panteleeff, D., et al., 1998. Evolution of envelope sequences from the genital tract and peripheral blood of women infected with clade A human immunodeficiency virus type 1. *J. Virol.* 72 (10), 8240–8251.
- Power, C., McArthur, J.C., Johnson, R.T., Griffin, D.E., Glass, J.D., Perryman, S., et al., 1994. Demented and nondemented patients with AIDS differ in brain-derived human immunodeficiency virus type 1 envelope sequences. *J. Virol.* 68 (7), 4643–4649.
- Preston, B.D., Dougherty, J.P., 1996. Mechanisms of retroviral mutation. *Trends Microbiol.* 4 (1), 16–21.
- Price, D.A., Goulder, P.J.R., Klenerman, P., Sewell, A.K., Easterbrook, P.J., Troop, M., et al., 1997. Positive selection of HIV-1 cytotoxic T lymphocyte escape variants during primary infection. *Proc. Natl. Acad. Sci. USA* 94 (5), 1890–1895.
- Quigg, M., Rebus, S., France, A.J., McMenamin, J., Darby, G., Leigh Brown, A.J., 1997. Mutations associated with

- zidovudine resistance in HIV-1 among recent seroconverters. *AIDS* 11 (6), 835–836.
- Rasheed, S., 1998. Infectivity and dynamics of HIV type 1 replication in the blood and reproductive tract of HIV type 1-infected women. *AIDS Res. Hum. Retroviruses* 14 (Suppl. 1), S105–S118.
- Rasheed, S., Li, Z., Xu, D., Kovacs, A., 1996. Presence of cell-free human immunodeficiency virus in cervicovaginal secretions is independent of viral load in the blood of human immunodeficiency virus-infected women. *Am. J. Obstet. Gynecol.* 175 (1), 122–129.
- Reddy, R.T., Achim, C.L., Sirko, D.A., Tehranchi, S., Kraus, F.G., Wong-Staal, F., et al., 1996. Sequence analysis of the V3 loop in brain and spleen of patients with HIV encephalitis. *AIDS Res. Hum. Retroviruses* 12 (6), 477–482.
- Rafaeli, Y., Levy, D.N., Weiner, D.B., 1995. The glucocorticoid receptor type II complex is a target of the HIV-1 vpr gene product. *Proc. Natl. Acad. Sci. USA* 92 (8), 3621–3625.
- Rinaldo, C.R., Beltz, L.A., Huang, X.L., Gupta, P., Fan, Z., Torpey, D.J. III, 1995. Anti-HIV type 1 cytotoxic T lymphocyte effector activity and disease progression in the first 8 years of HIV type 1 infection of homosexual men. *AIDS Res. Hum. Retroviruses* 11 (4), 481–489.
- Robertson, D.L., Anderson, J.P., Bradac, J.A., Carr, J.K., Foley, B., Funkhouser, R.K., et al., 1999. HIV-1 nomenclature proposal: A reference guide to HIV-1 classification. In: Kuiken, C.L., Foley, B., Hahn, B., Korber, B., McCutchan, F., Marx, P.A., Mellors, J.W., Mullins, J.I., Sodroski, J., Welinksky, S. (Eds.), *Human Retroviruses and AIDS 1999: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM.
- Robertson, D.L., Gao, F., Hahn, B.H., Sharp, P.M., 1998. Intersubtype recombinant HIV-1 sequences. In: Korber, B., Kuiken, C.L., Foley, B., Hahn, B., McCutchan, F., Mellors, J.W., Sodroski, J. (Eds.), *Human Retroviruses and AIDS 1998: A Compilation and Analysis of Nucleic Acid and Amino Acid Sequences*. Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM.
- Robertson, D.L., Sharp, P.M., McCutchan, F.E., Hahn, B.H., 1995. Recombination in HIV-1. *Nature* 374 (6518), 124–126.
- Roos, M.T., Lange, J.M., de Goede, R.E.Y., Coutinho, R.A., Schellekens, P.T., Miedema, F., et al., 1992. Viral phenotype and immune response in primary human immunodeficiency virus type 1 infection. *J. Infect. Dis.* 165 (3), 427–432.
- Rouzine, I.M., Coffin, J.M., 1999. Linkage disequilibrium test implies a large effective population number for HIV in vivo. *Proc. Natl. Acad. Sci. USA* 96 (19), 10758–10763.
- Sarnagadharan, M.G., Popovic, M., Bruch, L., Schupbach, J., Gallo, R.C., 1984. Antibodies reactive with human T-lymphotropic retroviruses (HTLV-III) in the serum of patients with AIDS. *Science* 224 (4648), 506–508.
- Sato, H., Shiino, T., Kodaka, N., Taniguchi, K., Tomita, Y., Kato, K., et al., 1999. Evolution and biological characterization of human immunodeficiency virus type 1 subtype E gp120 V3 sequences following horizontal and vertical virus transmission in a single family. *J. Virol.* 73 (5), 3551–3559.
- Schmitz, J.E., Kuroda, M.J., Santra, S., Sasseville, V.G., Simon, M.A., Lifton, M.A., et al., 1999. Control of viremia in simian immunodeficiency virus infection by CD8 + lymphocytes. *Science* 283 (5403), 857–860.
- Schock, H.B., Garsky, V.M., Kuo, L.C., 1996. Mutational anatomy of an HIV-1 protease variant conferring cross-resistance to protease inhibitors in clinical trials. Compensatory modulations of binding and activity. *J. Biol. Chem.* 271 (50), 31957–31963.
- Schuitemaker, H., Koot, M., Kootstra, N.A., Dercksen, M.W., de Goede, R.E.Y., van Steenwijk, R.P., et al., 1992. Biological phenotype of human immunodeficiency virus type 1 clones at different stages of infection: progression of disease is associated with a shift from monocytotropic to T-cell-tropic virus population. *J. Virol.* 66 (3), 1354–1360.
- Schupbach, J., Popovic, M., Gilden, R.V., Gonda, M.A., Sarnagadharan, M.G., Gallo, R.C., 1984. Serological analysis of a subgroup of human T-lymphotropic retroviruses (HTLV-III) associated with AIDS. *Science* 224 (4648), 503–505.
- Seibert, S.A., Howell, C.Y., Hughes, M.K., Hughes, A.L., 1995. Natural selection on the *gag*, *pol*, and *env* genes of human immunodeficiency virus 1 (HIV-1). *Mol. Biol. Evol.* 12 (5), 803–813.
- Shankarappa, R., Gupta, P., Learn, G.H.J., Rodrigo, A.G., Rinaldo, C.R.Jr., Gorro, M.C., et al., 1998. Evolution of human immunodeficiency virus type 1 envelope sequences in infected individuals with differing disease progression profiles. *Virology* 241 (2), 251–259.
- Shankarappa, R., Margolick, J.B., Gange, S.J., Rodrigo, A.G., Upchurch, D., Farzadegan, H., et al., 1999. Consistent viral evolutionary changes associated with the progression of human immunodeficiency virus type 1 infection. *J. Virol.* 73 (12), 10489–10502.
- Sharp, P.M., Robertson, D.L., Hahn, B.H., 1995. Cross-species transmission and recombination of 'AIDS' viruses. *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* 349 (1327), 41–47.
- Shugars, D.C., Slade, G.D., Patton, L.L., Fiscus, S.A., 2000. Oral and systemic factors associated with increased levels of human immunodeficiency virus type 1 RNA in saliva. *Oral Surg. Oral Med. Oral Pathol. Oral Radiol. Endod.* 89, 432–440.
- Simmonds, P., Zhang, L.Q., McOmish, F., Balfe, P., Ludlam, C.A., Leigh Brown, A.J., 1991. Discontinuous sequence change of human immunodeficiency virus (HIV) type 1 *env* sequences in plasma viral and lymphocyte-associated proviral populations in vivo: implications for models of HIV pathogenesis. *J. Virol.* 65 (11), 6266–6276.
- Simon, F., Maucelere, P., Roques, P., Lousert-Ajaka, I., Muller-Trutwin, M.C., Saragosti, S., et al., 1998. Identification of a new human immunodeficiency virus type 1

- distinct from group M and group O. *Nat. Med.* 4 (9), 1032–1037.
- Smith, M.S., Koerber, K.L., Pagano, J.S., 1993. Zidovudine-resistant human immunodeficiency virus type 1 genomes detected in plasma distinct from viral genomes in peripheral blood mononuclear cells. *J. Infect. Dis.* 167 (2), 445–448.
- Starcich, B.R., Hahn, B.H., Shaw, G.M., McNeely, P.D., Modrow, S., Wolf, H., et al., 1986. Identification and characterization of conserved and variable regions in the envelope gene of HTLV-III/LAV, the retrovirus of AIDS. *Cell* 45 (5), 637–648.
- Steuler, H., Storch-Hagenlocher, B., Wildemann, B., 1992. Distinct populations of human immunodeficiency virus type 1 in blood and cerebrospinal fluid. *AIDS Res. Hum. Retroviruses* 8 (1), 53–59.
- Temin, H.M., 1992. Origin and general nature of retroviruses. In: Levy, J.A. (Ed.), *The Retroviridae*, vol. 1. Plenum Press, New York, pp. 1–18.
- Tersmette, M., de Goede, R.E.Y., Al, B.J.M., Winkel, I.N., Gruters, R.A., Cuypers, H.T., et al., 1988. Differential syncytium-inducing capacity of human immunodeficiency virus isolates: frequent detection of syncytium-inducing isolates in patients with acquired immunodeficiency syndrome (AIDS) and AIDS-related complex. *J. Virol.* 62 (6), 2026–2032.
- Tersmette, M., Gruters, R.A., de Wolf, F., de Goede, R.E.Y., Lange, J.M., Schellekens, P.T., et al., 1989. Evidence for a role of virulent human immunodeficiency virus (HIV) variants in the pathogenesis of acquired immunodeficiency syndrome: studies on sequential HIV isolates. *J. Virol.* 63 (5), 2118–2125.
- Trkola, A., Dragic, T., Arthos, J., Binley, J.M., Olson, W.C., Allaway, G.P., et al., 1996. CD4-dependent, antibody-sensitive interactions between HIV-1 and its co-receptor CCR-5. *Nature* 384 (6605), 184–187.
- Uccini, S., Riva, E., Antonelli, G., D'Offizi, G., Prozzo, A., Angelici, A., et al., 1999. The benign cystic lymphoepithelial lesion of the parotid gland is a viral reservoir in HIV type 1-infected patients. *AIDS Res. Hum. Retroviruses* 15 (15), 1339–1344.
- Vanden Haesevelde, M.M., Peeters, M., Jannes, G., Janssens, W., van der Groen, G., Sharp, P.M., et al., 1996. Sequence analysis of a highly divergent HIV-1-related lentivirus isolated from a wild captured chimpanzee. *Virology* 221 (2), 346–350.
- Varela-Echavarria, A., Garvey, N., Preston, B.D., Dougherty, J.P., 1992. Comparison of Moloney murine leukemia virus mutation rate with the fidelity of its reverse transcriptase in vitro. *J. Biol. Chem.* 267 (34), 24681–24688.
- Vartanian, J.P., Meyerhans, A., Asjo, B., Wain-Hobson, S., 1991. Selection, recombination, and G→A hypermutation of human immunodeficiency virus type 1 genomes. *J. Virol.* 65 (4), 1779–1788.
- Vernazza, P.L., Eron, J.J., Cohen, M.S., van der Horst, C.M., Troiani, L., Fiscus, S.A., 1994. Detection and biologic characterization of infectious HIV-1 in semen of seropositive men. *AIDS* 8 (9), 1325–1329.
- Vernazza, P.L., Eron, J.J., Fiscus, S.A., Cohen, M.S., 1999. Sexual transmission of HIV: infectiousness and prevention. *AIDS* 13 (2), 155–166.
- Vicenzi, E., Bordignon, P.P., Biswas, P., Brambilla, A., Bovolenta, C., Cota, M., et al., 1999. Envelope-dependent restriction of human immunodeficiency virus type 1 spreading in CD4(+) T lymphocytes: R5 but not X4 viruses replicate in the absence of T-cell receptor restimulation. *J. Virol.* 73 (9), 7515–7523.
- Vidal, N., Peeters, M., Mulanga-Kabeya, C., Nzilambi, N., Robertson, D., Ilunga, W., et al., 2000. Unprecedented degree of human immunodeficiency virus type 1 (HIV-1) group M genetic diversity in the Democratic Republic of Congo suggests that the HIV-1 pandemic originated in Central Africa. *J. Virol.* 74 (22), 10498–10507.
- Wei, X., Ghosh, S.K., Taylor, M.E., Johnson, V.A., Emini, E.A., Deutsch, P., et al., 1995. Viral dynamics in human immunodeficiency virus type 1 infection. *Nature* 373 (6510), 117–122.
- Wodarz, D., Nowak, M.A., 1999. Evolutionary dynamics of HIV-induced subversion of the immune response. *Immunol. Rev.* 168, 75–89.
- Wolfs, T.F., Zwart, G., Bakker, M., Goudsmit, J., 1992. HIV-1 genomic RNA diversification following sexual and parenteral virus transmission. *Virology* 189 (1), 103–110.
- Wolinsky, S.M., Korber, B.T., Neumann, A.U., Daniels, M., Kunstman, K.J., Whetsell, A.J., et al., 1996. Adaptive evolution of human immunodeficiency virus-type 1 during the natural course of infection. *Science* 272 (5261), 537–542.
- Wolinsky, S.M., Wike, C.M., Korber, B.T., Hutto, C., Parks, W.P., Rosenblum, L.L., et al., 1992. Selective transmission of human immunodeficiency virus type-1 variants from mothers to infants. *Science* 255 (5048), 1134–1137.
- Wong, J.K., Ignacio, C.C., Torriani, F., Havlir, D., Fitch, N.J., Richman, D.D., 1997. In vivo compartmentalization of human immunodeficiency virus: evidence from the examination of pol sequences from autopsy tissues. *J. Virol.* 71 (3), 2059–2071.
- Wong-Staal, F., Shaw, G.M., Hahn, B.H., Salahuddin, S.Z., Popovic, M., Markham, P., et al., 1985. Genomic diversity of human T-lymphotropic virus type III (HTLV-III). *Science* 229 (4715), 759–762.
- Wooley, D.P., Smith, R.A., Czajak, S., Desrosiers, R.C., 1997. Direct demonstration of retroviral recombination in a rhesus monkey. *J. Virol.* 71 (12), 9650–9653.
- Xiao, L., Owen, S.M., Goldman, I., Lal, A.A., de Jong, J.J., Goudsmit, J., et al., 1998. CCR5 coreceptor usage of non-syncytium-inducing primary HIV-1 is independent of phylogenetically distinct global HIV-1 isolates: delineation of consensus motif in the V3 domain that predicts CCR-5 usage. *Virology* 240 (1), 83–92.
- Yerly, S., Kaiser, L., Race, E., Bru, J.P., Clavel, F., Perrin, L., 1999. Transmission of antiretroviral-drug-resistant HIV-1 variants. *Lancet* 354 (9180), 729–733.
- Yi, Y., Isaacs, S.N., Williams, D.A., Frank, I., Schols, D., De Clercq, E., et al., 1999. Role of CXCR4 in cell-cell fusion

- and infection of monocyte-derived macrophages by primary human immunodeficiency virus type 1 (HIV-1) strains: two distinct mechanisms of HIV-1 dual tropism. *J. Virol.* 73 (9), 7117–7125.
- Zhang, J., Tang, L., Li, T., Ma, Y., Sapp, C.M., 2000. Most retroviral recombinations occur during minus-strand DNA synthesis. *J. Virol.* 74 (5), 2313–2322.
- Zhang, L., Diaz, R.S., Ho, D.D., Mosley, J.W., Busch, M.P., Mayer, A., 1997a. Host-specific driving force in human immunodeficiency virus type 1 evolution in vivo. *J. Virol.* 71 (3), 2555–2561.
- Zhang, L.Q., MacKenzie, P., Cleland, A., Holmes, E.C., Leigh Brown, A.J., Simmonds, P., 1993. Selection for specific sequences in the external envelope protein of human immunodeficiency virus type 1 upon primary infection. *J. Virol.* 67 (6), 3345–3356.
- Zhang, Y.M., Imamichi, H., Imamichi, T., Lane, H.C., Falloon, J., Vasudevachari, M.B., et al., 1997b. Drug resistance during indinavir therapy is caused by mutations in the protease gene and in its Gag substrate cleavage sites. *J. Virol.* 71 (9), 6662–6670.
- Zhao, L.J., Mukherjee, S., Narayan, O., 1994. Biochemical mechanism of HIV-I Vpr function. Specific interaction with a cellular protein. *J. Biol. Chem.* 269 (22), 15577–15582.
- Zhu, T., Korber, B.T., Nahmias, A.J., Hooper, E., Sharp, P.M., Ho, D.D., 1998. An African HIV-1 sequence from 1959 and implications for the origin of the epidemic. *Nature* 391 (6667), 594–597.
- Zhu, T., Mo, H., Wang, N., Nam, D.S., Cao, Y., Koup, R.A., et al., 1993. Genotypic and phenotypic characterization of HIV-1 patients with primary infection. *Science* 261 (5125), 1179–1181.