

Slopes and local change

Project: Image-pattern Network Investigation

1. Prepare the purpose, prerequisites and materials

Teach the learning cycle

Primary target: Calculate an average rate of change over a stated interval

Session · Sequence and minutes · Resources

Session: Session 1 · Understand · 45 min

Sequence and minutes: 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

Resources: Sections 1–6; record the brief recall in the workbook.

Session: Session 2 · Practise · 45 min

Sequence and minutes: 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.

Resources: Sections 7–10; use the matching workbook W2 and misconception check.

Session: Session 3 · Investigate and transfer · 45 min

Sequence and minutes: 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

Resources: Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

This is a starting allocation of 135 minutes, not a validated duration. Preserve all named topics. Schedule extra time for connected examples, the full investigation, independent cases and project building. Repeated copies of a case share one recorded attempt; they are not new evidence.

Retrieval answer guidance

Loss functions

Without opening the old lesson, explain one key idea from Loss functions. Give an example and a mistake that the idea helps you avoid.

Look for: Calculate per-example squared error and mean squared error

Reference example, not the only acceptable wording: Teacher explanation: Residuals are -1 and 2, squares 1 and 4, MSE 2.5.

Forward passes by hand

Without opening the old lesson, explain one key idea from Forward passes by hand. Give an example and a mistake that the idea helps you avoid.

Look for: Record scores and activations in dependency order

Reference example, not the only acceptable wording: Teacher explanation: $z = [-3, 3]$, $h = [0, 3]$, $y = -5$.

Bias terms and decision boundaries

Without opening the old lesson, explain one key idea from Bias terms and decision boundaries. Give an example and a mistake that the idea helps you avoid.

Look for: Calculate a weighted score with a bias term

Reference example, not the only acceptable wording: Teacher explanation: Weighted sum $10 - 4 = 6$; bias -3; score 3; prediction 1. Add the bias once after summing.

Introduce rate of change through finite differences, with no assumed calculus fluency. Use two 45-minute sessions, a calculator and a labelled parameter-versus-loss sketch. Reserve the word local for a stated centre and nearby values.

Outcomes

Calculate an average rate of change over a stated interval

Estimate a local slope with nearby loss values

Explain the limits of a local direction clue

Preparation

Printed lesson and workbook; paper and pencil

Calculator and optional local Python 3 editor; no packages or AI accounts needed

Use the fictional examples supplied; prepare the fresh retry before class

Prerequisite answer

The residual is 2 and the squared error is 4.

End goal for the pictorial case

By the end, I can explain slopes and local change using a traced example and a stated limitation.

This added worked case illustrates the week's main idea. Retain all original and connected topics; schedule its practice within the teaching cycle.

2. Explain and model the mechanism

Let a one-example model predict w and let its target be 2. Its squared loss is $L(w) = (w - 2)^2$. Here w is the parameter we vary, not a new data example. L is a nonnegative function of that parameter.

From $w = 0$ to $w = 1$, the loss changes by -3 while w changes by 1. The average slope is -3. From $w = 1$ to $w = 2$ it is -1. The curve does not have one fixed slope everywhere.

Near $w = 1$, use two symmetric nearby values: 0.9 and 1.1. Their losses are 1.21 and 0.81. Change in loss divided by change in w gives $(0.81 - 1.21) / (1.1 - 0.9) = -2$. This symmetric estimate describes change near the centre $w = 1$.

For this quadratic, the symmetric estimate equals the exact slope $2(w - 2)$ in exact arithmetic. Numerical estimates for other functions are generally approximate. Smaller steps can reduce some approximation errors, but extremely small steps can suffer from computer rounding.

Term · Meaning

Term: Slope

Meaning: Change in output divided by change in input over a stated interval

Term: Local change

Meaning: Change considered near a particular input or parameter value

Term: Step size

Meaning: The amount by which an input or parameter is changed

Term: Stationary point

Meaning: A point where a differentiable function has zero local slope

Compare slope clues on the same loss curve

Current w · Local slope · Nearby downhill direction

Current w: 1

Local slope: -2

Nearby downhill direction: Increase w a little

Current w: 2

Local slope: 0

Nearby downhill direction: No first order direction

Current w: 3

Local slope: 2

Nearby downhill direction: Decrease w a little

These are local clues for $L(w) = (w - 2)^2$. At $w = 2$ the displayed function has its minimum; zero slope does not establish a minimum for every possible function.

Guided: $L(2.9) = 0.81$ and $L(3.1) = 1.21$, so slope is $0.4 / 0.2 = 2$. A small decrease in w lowers this loss.
Exit: at $w = 1$ slope is negative, but increasing w to 4 raises loss from 1 to 4.

Model the reasoning aloud

Use the complete worked example below before asking learners to complete W2. Reveal a small part, invite a prediction, then explain the deciding step. These teacher notes contain solutions; do not reveal the independent case answer during modelling.

Compare slope clues on the same loss curve

Current w · Local slope · Nearby downhill direction

Current w: 1

Local slope: -2

Nearby downhill direction: Increase w a little

Current w: 2

Local slope: 0

Nearby downhill direction: No first order direction

Current w: 3

Local slope: 2

Nearby downhill direction: Decrease w a little

These are local clues for $L(w) = (w - 2)^2$. At $w = 2$ the displayed function has its minimum; zero slope does not establish a minimum for every possible function.

At $w = 3$, use 2.9 and 3.1. Calculate both losses and the symmetric slope. Which small change should lower loss here?

Thinking move · Teacher prompt

Thinking move: Plan

Teacher prompt: What is given? What is the question? Which procedure fits these facts?

Thinking move: Do

Teacher prompt: Point to each input and intermediate step in the worked trace. Name the rule that allows the next step.

Thinking move: Check

Teacher prompt: Does the output answer the question? Which assumption or limitation prevents a larger claim?

Use the diagram and verbal explanation together. Ask learners to match a sentence to a specific visual step; do not assign learners fixed visual or auditory learning styles.

A pictorial worked case: Slopes and local change

Use a local slope to propose a small move rather than claiming every move helps.

The facts we will use

Toy loss $J(a)=(a-2)^2$. At $a = 0$, use symmetric step 0.5.

All inputs and outcomes in this worked example are invented classroom data; no real model run is claimed.

AI ACADEMY / AI-5/9

Slopes and local change

Start with the supplied facts

Follow each arrow. Ask which fact allows the next step.



1. Evaluate nearby points

$J(0.5)=2.25$ and $J(-0.5)=6.25$.



2. Estimate slope

$\text{Slope}=(2.25-6.25)/(0.5-(-0.5))=-4$.



3. Check small move

Move to $a = 0.25$: loss 3.0625, below starting loss 4.

CHANGE ONE THING

Move from 0 all the way to 5.

Explain the changed result using the matching lesson.

Read the picture one step at a time

Step · What happens and why

Step: 1. Evaluate nearby points

What happens and why: $J(0.5)=2.25$ and $J(-0.5)=6.25$.

Step: 2. Estimate slope

What happens and why: $\text{Slope} = (2.25 - 6.25) / (0.5 - (-0.5)) = -4$.

Step: 3. Check small move

What happens and why: Move to $a = 0.25$: loss 3.0625, below starting loss 4.

Match each pictured step to the supplied facts. This is a worked teaching case, not an independent assessment.

Why this result follows

A slope describes how the loss changes near the current parameter. The symmetric calculation compares equal small moves on either side. Its negative result suggests a small increase may help at zero. It does not promise every increase helps: jumping to five passes the optimum at two and produces a larger loss. Check the new loss after each proposed step.

What this example does not establish

A finite difference approximates local change; numerical precision and nonsmooth points need care in more complex objectives.

Change one thing in the pictured case

Change: Move from 0 all the way to 5.

Prediction: Loss becomes 9, exceeding the starting 4.

Reason: A negative local slope suggests increasing a locally, not that every positive jump decreases loss.

Ask while showing the picture

Cover the result. Ask learners to name the inputs and predict the next step. Uncover one step, connect the icon to the actual procedure, and explain its evidence. Then change the supplied condition and compare. The picture is a representation; it does not document execution of a real AI model.

3. Demonstrate and guide practice

Time or stage · Teacher action

Time or stage: Session 1 start to 7 minutes

Teacher action: Retrieve squared loss and distinguish parameter w from an example input.

Time or stage: 7 to 18 minutes

Teacher action: Build the five-row table and sketch the curve. Calculate slopes across two different intervals.

Time or stage: 18 to 32 minutes

Teacher action: Complete W1 and the worked symmetric estimate near 1. Emphasise denominator 2 times step.

Time or stage: 32 to 45 minutes

Teacher action: Run or trace the optional code, complete W2 and the guided case.

Time or stage: Session 2 start to 8 minutes

Teacher action: Compare small changes with jumps across the minimum using W3.

Time or stage: 8 to 20 minutes

Teacher action: Use $F(w)$ to challenge zero-slope claims and complete W5.

Time or stage: 20 to 35 minutes

Teacher action: Collect W4 independently, including interval arithmetic and the oversized-step counterexample.

Time or stage: 35 to 45 minutes

Teacher action: Use the exit response and a fresh retry to check direction, scale and stated limits.

Use the worked example in Sections 3 and 6. Collect a prediction before revealing its result; ask the learner to point to each intermediate fact.

Provide an axis label and the fraction change in loss over change in parameter. Extension: algebraically expand the symmetric difference for $(w - 2)$ squared to recover $2(w - 2)$, noting exact arithmetic versus floating-point computation.

Reduce help deliberately

Completion task: At centre $w = 0$, use step 0.5. Calculate $L(-0.5)$, $L(0.5)$ and the symmetric slope.

Teacher check: Teacher explanation: Losses are 6.25 and 2.25. Denominator is 1, giving slope -4.

Start by identifying the given inputs together. Leave the intermediate steps blank. If needed, model one

step, then withdraw that help on the next step. Move to a full trace only when the partial trace is understood. For partners, alternate explainer and checker; collect a response from every learner before discussion.

Individual reasoning check

Use the worked case for Slopes and local change. Proposed explanation: “Divide by the step only”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Expected correction: A symmetric comparison spans from centre minus step to centre plus step, a width of twice the step.

Collect explanations, not only yes/no votes. If a misunderstanding is common, re-model the relevant step for the class. If it affects a few learners, give a short targeted group explanation while others solve a more demanding variation.

Pictorial practice question

Which calculation disproves “any increase helps”? Point to the relevant step in the picture, then write the changed result and the rule or evidence that supports it.

Reasoned teacher answer

$J(5)=(5-2)^2=9 > J(0)=4$. The local direction is useful only with an appropriately sized step and verification. Changed-case check: Loss becomes 9, exceeding the starting 4. A negative local slope suggests increasing a locally, not that every positive jump decreases loss.

Changed-case reasoning: Loss becomes 9, exceeding the starting 4. A negative local slope suggests increasing a locally, not that every positive jump decreases loss.

Collect each learner’s explanation before discussion. Use the first missing connection between input, deciding step and result to choose support. This question is worked-case practice, not fresh mastery evidence.

4. Run the investigation and diagnose misconceptions

Divide by the step only: A symmetric comparison spans from centre minus step to centre plus step, a width of twice the step.

Negative loss slope means negative loss: The slope describes change; squared loss itself remains nonnegative.

Zero slope always means finished: Stationary points need additional evidence; some are not minima.

Use Section 9 for the evidence ledger. Add a boundary or missing-input case and record the expected response before checking. More successful repeats of one case do not establish wider coverage.

Match the support to the first error

What the response shows · Next teaching action

What the response shows: Missing or misread facts

Next teaching action: Read the case aloud, mark supplied versus unknown facts, and let the learner restate it.

What the response shows: Wrong procedure

Next teaching action: Return to the deciding step in the worked case. Contrast it with the proposed incorrect explanation.

What the response shows: Arithmetic or implementation error

Next teaching action: Trace intermediate values or lines; preserve the first mismatch and retry that step.

What the response shows: Correct result without a reason

Next teaching action: Ask for a trace and a changed case before marking independent understanding.

What the response shows: An unsupported wider claim

Next teaching action: Ask which supplied evidence supports the claim and which further observation would be needed.

Offer an oral explanation, drawing, enlarged visual or fewer examples when that removes a reading/writing barrier. Keep the same reasoning goal. Stretch secure learners by changing a boundary or assumption and asking them to defend the effect, not simply by assigning more of the same questions.

5. Assess and explain the answers

Guided and exit questions

Guided: $L(2.9) = 0.81$ and $L(3.1) = 1.21$, so slope is $0.4 / 0.2 = 2$. A small decrease in w lowers this loss.
Exit: at $w = 1$ slope is negative, but increasing w to 4 raises loss from 1 to 4.

W1 Measure an interval slope

For $L(w) = (w - 2)^2$, calculate the average slope from $w = 0$ to $w = 2$. Compare it with the local slope at $w = 2$.

Teacher explanation: Average slope is $(0 - 4) / 2 = -2$. The local slope at 2 is zero. They describe different

locations or intervals.

W2 Check a symmetric estimate

At centre $w = 0$, use step 0.5. Calculate $L(-0.5)$, $L(0.5)$ and the symmetric slope.

Teacher explanation: Losses are 6.25 and 2.25. Denominator is 1, giving slope -4.

W3 Challenge a large step

Starting at $w = 3$, compare loss after moving to 2.9 and after moving to -1. Both moves decrease w .

Teacher explanation: Initial loss is 1; at 2.9 it is 0.81; at -1 it is 9. The small move helps, the large move overshoots.

W4 Estimate a new curve independently

Let $J(a) = (a + 1)$ squared. At $a = 1$ use step 0.25 to find the symmetric slope. Compare $J(1)$ with $J(0.9)$, then explain why moving to -4 does not follow safely from the slope clue.

Teacher explanation: $J(0.75) = 3.0625$; $J(1.25) = 5.0625$; slope is $2 / 0.5 = 4$. $J(1) = 4$ and $J(0.9) = 3.61$, so a small decrease helps. $J(-4) = 9$: a large decrease overshoots.

W5 Limit zero slope evidence

For $F(w) = (w \text{ squared} - 1) \text{ squared}$, compare $F(0)$ and $F(0.1)$. Does zero local slope at zero prove a minimum?

Teacher explanation: $F(0) = 1$ and $F(0.1) = 0.9801$. Nearby loss is lower, so zero slope alone does not prove a minimum.

Independent assessment and fresh retry

W4 requires losses 3.0625 and 5.0625, slope 4, and the finite-step limitation. Fresh retry: $K(c) = (c - 3)$ squared at $c = 2$, step 0.5, uses losses 2.25 and 0.25 and slope -2. Moving to 2.1 gives loss 0.81 rather than 1.

Assess independent transfer

Selected case: Let $J(a) = (a + 1)$ squared. At $a = 1$ use step 0.25 to find the symmetric slope. Compare $J(1)$ with $J(0.9)$, then explain why moving to -4 does not follow safely from the slope clue.

Selected case answer

Teacher explanation: $J(0.75) = 3.0625$; $J(1.25) = 5.0625$; slope is $2 / 0.5 = 4$. $J(1) = 4$ and $J(0.9) = 3.61$, so a small decrease helps. $J(-4) = 9$: a large decrease overshoots.

Dimension · Evidence required

Dimension: Explanation

Evidence required: Names the applicable idea or procedure and ties it to the supplied facts.

Dimension: Trace and result

Evidence required: Preserves the given inputs and shows the relevant intermediate steps; distinguishes a prediction from an observed execution.

Dimension: Limits

Evidence required: States the assumptions, missing evidence or conditions under which the result would change.

Record each dimension as not yet, with support, or independent. Accept a valid alternative procedure with a defensible trace. Do not penalise an honestly recorded model failure if the learner correctly explains its evidence and limits.

Feedback that the learner uses

Name the earliest unsupported step and give one action to repair it. Have the learner make that repair now and explain why. Then assess the relevant skill on a fresh, fully specified case. Confirm it was not previously practised with feedback; solve and check the new case before assigning it. A follow-up design prompt is a starting brief, not automatically an equivalent test.

6. Connect the project, support and next lesson

Add a labelled slopes and local change evidence sheet to Image-pattern Network Investigation. Show how you can calculate an average rate of change over a stated interval. Use the supplied fictional case and keep its inputs, intermediate steps, calculation or prediction, and limitation. Connect this record to last week's record; a matching answer without a trace is insufficient.

Provide an axis label and the fraction change in loss over change in parameter. Extension: algebraically expand the symmetric difference for $(w - 2)$ squared to recover $2(w - 2)$, noting exact arithmetic versus floating-point computation.

Week 10 extends local slopes to several parameters.

Retain the project ledger and assess a changed case after support. Check the relevant prerequisite before carrying this component forward.

Plan the delayed return

Suggested return points in this level: ai-5/10, ai-5/12, ai-5/15. Use actual classroom dates, adjusting for holidays and gaps. One, three and six teaching weeks are scheduling choices, not a claimed optimal

spacing.

At a delayed check, ask for the idea from memory and a changed application. Compare with the earlier independent response and record support used. For a new level, revisit the previous capstone before checking the new prerequisites.

Record · Teacher evidence

Record: Learner code and date

Teacher evidence: _____

Record: Concept and case used; prior exposure

Teacher evidence: _____

Record: Explanation / trace / limits

Teacher evidence: _____

Record: Support and first repair

Teacher evidence: _____

Record: Changed case and delayed outcome

Teacher evidence: _____

Record: Next teaching action

Teacher evidence: _____

Use the evidence to choose reteaching or the next step. A completed page, confidence rating or attractive project does not by itself demonstrate understanding.

Project evidence from this case

For your Image-pattern Network Investigation, save the input, procedure, observed result and limitation of this check. Label this worked example as practice; use a different, previously unreviewed case when checking independent understanding.

7. Prepare, teach and assess

Teaching target: Calculate an average rate of change over a stated interval

Prepare the labelled worked-case picture, the matching workbook, fictional input cards and a place for individual responses. Check a local Python 3 environment before class; have a paper-trace version ready.

Session · Plan · Resources

Session: Session 1 · Understand · 45 min

Plan: 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

Resources: Sections 1–6; record the brief recall in the workbook.

Session: Session 2 · Practise · 45 min

Plan: 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.

Resources: Sections 7–10; use the matching workbook W2 and misconception check.

Session: Session 3 · Investigate and transfer · 45 min

Plan: 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

Resources: Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

Before class, solve the supported practice: At centre $w = 0$, use step 0.5. Calculate $L(-0.5)$, $L(0.5)$ and the symmetric slope.

Reasoned practice answer: Teacher explanation: Losses are 6.25 and 2.25. Denominator is 1, giving slope -4.

Check every learner before discussion: Use the worked case for Slopes and local change. Proposed explanation: “Divide by the step only”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Misconception repair: A symmetric comparison spans from centre minus step to centre plus step, a width of twice the step.

Support: read the exact case aloud, enlarge the labelled picture, supply input cards and model only the first deciding step. Accept a spoken explanation or drawing; retain the reasoning goal.

Extension: Move from 0 all the way to 5. Require a prediction and a reason before comparing. Expected result: Loss becomes 9, exceeding the starting 4. Reason: A negative local slope suggests increasing a

locally, not that every positive jump decreases loss.

Independent assessment: withhold the reserved answer until the learner has attempted the new case.
Record support separately from correctness.

Criterion · Not yet · With support · Independent

Criterion: Explanation

Not yet: Names an answer without a deciding fact

With support: Uses a hint to link fact and decision

Independent: Explains the deciding fact in own words

Criterion: Trace or test

Not yet: Cannot connect input and result

With support: Completes steps with prompts

Independent: Traces or tests a changed case accurately

Criterion: Limits and revision

Not yet: Claims more than evidence supports

With support: Names a limit after a prompt

Independent: States a defensible limit and useful next test

Project checkpoint: Make a transparent first version. Keep a trace of how one input becomes an output; explain the rule or learned component.

Use the weekly teacher answer for topic-specific reasoning. A saved progress tick is an activity record; judge mastery from the learner's explanation and evidence.

8. Use feedback, labs and repair paths

Use the two formative questions as supported practice, not as independent mastery scores. Discuss the deciding evidence rather than accepting a guessed selection.

The feedback comes from the reviewed public worked case. Do not reveal the separately reserved independent teacher answer until the learner has attempted that case.

Ask students to predict before running code. Preserve errors and actual outputs; collect a changed test and an explanation. The branch-boundary and indexing lessons deliberately include failure cases.

Use the earlier worked-explanation links to address a specific missing building block, then return to the current question. Adapt this suggestion after observing the learner; it is not an automatic diagnosis.

Collect the completed-work portfolio as evidence. Record support separately from correctness, and assess explanation, trace/test and limitations. Progress ticks and quiz selections alone do not establish mastery.