

Gradients with a numerical check

Project: Image-pattern Network Investigation

1. Prepare the purpose, prerequisites and materials

Teach the learning cycle

Primary target: Estimate one partial derivative while other parameters stay fixed

Session · Sequence and minutes · Resources

Session: Session 1 · Understand · 45 min

Sequence and minutes: 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

Resources: Sections 1–6; record the brief recall in the workbook.

Session: Session 2 · Practise · 45 min

Sequence and minutes: 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.

Resources: Sections 7–10; use the matching workbook W2 and misconception check.

Session: Session 3 · Investigate and transfer · 45 min

Sequence and minutes: 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

Resources: Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

This is a starting allocation of 135 minutes, not a validated duration. Preserve all named topics. Schedule extra time for connected examples, the full investigation, independent cases and project building. Repeated copies of a case share one recorded attempt; they are not new evidence.

Retrieval answer guidance

Slopes and local change

Without opening the old lesson, explain one key idea from Slopes and local change. Give an example and a mistake that the idea helps you avoid.

Look for: Calculate an average rate of change over a stated interval

Reference example, not the only acceptable wording: Teacher explanation: Average slope is $(0 - 4) / 2 = -2$. The local slope at 2 is zero. They describe different locations or intervals.

Parameters and architecture

Without opening the old lesson, explain one key idea from Parameters and architecture. Give an example and a mistake that the idea helps you avoid.

Look for: Distinguish architectural choices from parameter values

Reference example, not the only acceptable wording: Teacher explanation: There are 8 weights and 2 biases, total 10.

Activation functions

Without opening the old lesson, explain one key idea from Activation functions. Give an example and a mistake that the idea helps you avoid.

Look for: Trace identity ReLU and sigmoid on a numerical score

Reference example, not the only acceptable wording: Teacher explanation: Identity gives -2, 0, 2; ReLU gives 0, 0, 2. ReLU retains the positive score itself; it is not a step classifier.

Use two 45-minute sessions to separate coordinate probes from combined changes. The first gradient check is deliberately an affine model with squared loss, avoiding hidden-layer calculus. Introduce the chain-rule reasoning in words before showing the compact formula.

Outcomes

Estimate one partial derivative while other parameters stay fixed

Assemble a gradient in a declared parameter order

Compare numerical and formula based gradients with a tolerance

Preparation

Printed lesson and workbook; paper and pencil

Calculator and optional local Python 3 editor; no packages or AI accounts needed

Use the fictional examples supplied; prepare the fresh retry before class

Prerequisite answer

The two evaluation points lie one step on either side of the centre, so their separation is two steps.

End goal for the pictorial case

By the end, I can explain gradients with a numerical check using a traced example and a stated limitation.

This added worked case illustrates the week's main idea. Retain all original and connected topics; schedule its practice within the teaching cycle.

2. Explain and model the mechanism

Use one example with input $x = 2$ and target $t = 3$. The model predicts $y = wx + b$, and loss $L(w, b) = (2w + b - 3)^2$. At $w = 1$, $b = 0$, prediction is 2, residual is -1 and loss is 1. There are two parameters in order $[w, b]$.

To estimate the partial slope for w , change w only and keep $b = 0$. To estimate the partial slope for b , change b only and keep $w = 1$. Moving both at once measures change along a combined direction; it does not isolate either partial derivative.

The gradient is $[-4, -2]$ in order $[w, b]$. Both entries refer to the same original parameter vector, not to a partially updated model.

The helper assumes finite numerical inputs and a positive nonzero step. The formula follows the squared-loss rule: local loss change is twice the residual times the prediction sensitivity. Changing w by a small amount changes prediction twice as much here; changing b changes prediction by the same amount.

Exact floating-point equality is not required. Record the step and a tolerance, then try more than one parameter setting. A passing numerical check supports agreement for those tested points; it does not prove all possible inputs.

Term · Meaning

Term: Partial derivative

Meaning: A local slope with respect to one parameter while others are held fixed

Term: Gradient

Meaning: The ordered vector of partial derivatives of a scalar function

Term: Finite difference

Meaning: An estimate of change computed from nearby function values

Term: Tolerance

Meaning: A stated allowance for numerical disagreement

Two separate probes at one parameter vector

Parameter being probed · Held fixed · Gradient entry

Parameter being probed: w near 1

Held fixed: b is 0 and x is 2

Gradient entry: First entry -4

Parameter being probed: b near 0

Held fixed: w is 1 and x is 2

Gradient entry: Second entry -2

Parameter being probed: Assemble in order [w, b]

Held fixed: Same original model

Gradient entry: Gradient [-4, -2]

The first two rows are independent local probes, not successive parameter updates. The final row records both results in the declared order.

Guided: prediction 1, residual -2, loss 4, gradient [-8, -4]. Exit: changing both parameters measures a directional change; each partial must vary just its own coordinate while holding the others fixed.

Model the reasoning aloud

Use the complete worked example below before asking learners to complete W2. Reveal a small part, invite a prediction, then explain the deciding step. These teacher notes contain solutions; do not reveal the independent case answer during modelling.

Two separate probes at one parameter vector

Parameter being probed · Held fixed · Gradient entry

Parameter being probed: w near 1

Held fixed: b is 0 and x is 2

Gradient entry: First entry -4

Parameter being probed: b near 0

Held fixed: w is 1 and x is 2

Gradient entry: Second entry -2

Parameter being probed: Assemble in order [w, b]

Held fixed: Same original model

Gradient entry: Gradient [-4, -2]

The first two rows are independent local probes, not successive parameter updates. The final row records both results in the declared order.

At $w = 0$, $b = 1$, find prediction, residual, loss and the gradient using [4 times residual, 2 times residual].

Thinking move · Teacher prompt

Thinking move: Plan

Teacher prompt: What is given? What is the question? Which procedure fits these facts?

Thinking move: Do

Teacher prompt: Point to each input and intermediate step in the worked trace. Name the rule that allows the next step.

Thinking move: Check

Teacher prompt: Does the output answer the question? Which assumption or limitation prevents a larger claim?

Use the diagram and verbal explanation together. Ask learners to match a sentence to a specific visual step; do not assign learners fixed visual or auditory learning styles.

A pictorial worked case: Gradients with a numerical check

Check each coordinate of a gradient while holding the other parameter fixed.

The facts we will use

Model $y=2w+b$, target 3. At $[w, b]=[1, 0]$, $y = 2$, residual $y-\text{target}=-1$; loss is residual².

All inputs and outcomes in this worked example are invented classroom data; no real model run is claimed.

AI ACADEMY / AI-5/10

Gradients with a numerical check

Start with the supplied facts

Follow each arrow. Ask which fact allows the next step.



1. Check w direction

Analytic derivative $2 \times \text{residual} \times 2 = -4$.



2. Check b direction

Analytic derivative $2 \times \text{residual} = -2$.



3. Numerical check

With step 0.1: w losses 0.64, 1.44 give -4; b losses 0.81, 1.21 give -2.

CHANGE ONE THING

Swap the two gradient components in the program output.

Explain the changed result using the matching lesson.

Read the picture one step at a time

Step · What happens and why

Step: 1. Check w direction

What happens and why: Analytic derivative $2 \times \text{residual} \times 2 = -4$.

Step: 2. Check b direction

What happens and why: Analytic derivative $2 \times \text{residual} = -2$.

Step: 3. Numerical check

What happens and why: With step 0.1: w losses 0.64, 1.44 give -4; b losses 0.81, 1.21 give -2.

Match each pictured step to the supplied facts. This is a worked teaching case, not an independent assessment.

Why this result follows

A gradient lists the slope for each parameter, in a declared order. Changing w by a small amount changes the prediction twice as much as changing b by the same amount. That explains the different derivative magnitudes. Perturb one coordinate while holding the other fixed, then compare the numerical slope with its analytic counterpart.

What this example does not establish

The exact agreement occurs for this quadratic toy case; finite differences in larger models involve rounding and step-size limits.

Change one thing in the pictured case

Change: Swap the two gradient components in the program output.

Prediction: The coordinate-wise numerical check exposes the mismatch.

Reason: The w perturbation changes y twice as strongly as the same b perturbation.

Ask while showing the picture

Cover the result. Ask learners to name the inputs and predict the next step. Uncover one step, connect the icon to the actual procedure, and explain its evidence. Then change the supplied condition and compare. The picture is a representation; it does not document execution of a real AI model.

3. Demonstrate and guide practice

Time or stage · Teacher action

Time or stage: Session 1 start to 7 minutes

Teacher action: Retrieve the symmetric denominator and compute the initial prediction and residual.

Time or stage: 7 to 18 minutes

Teacher action: Probe w alone with b fixed; record the two losses and slope.

Time or stage: 18 to 32 minutes

Teacher action: Probe b alone and assemble $[-4, -2]$. Complete $W1$ and $W2$.

Time or stage: 32 to 45 minutes

Teacher action: Trace the optional numerical checker and the guided $[0, 1]$ case.

Time or stage: Session 2 start to 8 minutes

Teacher action: Use $W3$ to distinguish a combined direction from two partial derivatives.

Time or stage: 8 to 20 minutes

Teacher action: Discuss the ReLU corner in $W5$ and a stated numerical tolerance.

Time or stage: 20 to 35 minutes

Teacher action: Collect $W4$ independently, requiring both pairs of losses and coordinate order.

Time or stage: 35 to 45 minutes

Teacher action: Use the exit response and fresh retry, then connect local gradient evidence to next week.

Use the worked example in Sections 3 and 6. Collect a prediction before revealing its result; ask the learner to point to each intermediate fact.

Provide separate w -probe and b -probe tables. Extension: at the lesson model $[1, 0]$, test steps 0.1, 0.01 and 0.001 and report absolute disagreements from $[-4, -2]$, without claiming universal tolerance guarantees.

Reduce help deliberately

Completion task: A w probe has losses 1.44 and 0.64 at 0.9 and 1.1 with b fixed. A learner divides their difference by 0.1. Correct the result.

Teacher check: Teacher explanation: The interval is 0.2. $(0.64 - 1.44) / 0.2 = -4$, not -8.

Start by identifying the given inputs together. Leave the intermediate steps blank. If needed, model one step, then withdraw that help on the next step. Move to a full trace only when the partial trace is understood. For partners, alternate explainer and checker; collect a response from every learner before discussion.

Individual reasoning check

Use the worked case for Gradients with a numerical check. Proposed explanation: “The gradient is one scalar”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Expected correction: It has one entry per parameter for a scalar loss, in a stated order.

Collect explanations, not only yes/no votes. If a misunderstanding is common, re-model the relevant step for the class. If it affects a few learners, give a short targeted group explanation while others solve a more demanding variation.

Pictorial practice question

Why should the numerical checks perturb one coordinate at a time? Point to the relevant step in the picture, then write the changed result and the rule or evidence that supports it.

Reasoned teacher answer

They estimate a partial derivative. Changing both coordinates would mix their effects and would not independently verify $[-4, -2]$. Changed-case check: The coordinate-wise numerical check exposes the mismatch. The w perturbation changes y twice as strongly as the same b perturbation.

Changed-case reasoning: The coordinate-wise numerical check exposes the mismatch. The w perturbation changes y twice as strongly as the same b perturbation.

Collect each learner’s explanation before discussion. Use the first missing connection between input, deciding step and result to choose support. This question is worked-case practice, not fresh mastery evidence.

4. Run the investigation and diagnose misconceptions

The gradient is one scalar: It has one entry per parameter for a scalar loss, in a stated order.

Use the updated weight for the bias probe: Both entries must refer to the same current parameter vector.

A symmetric value defines the derivative at any corner: Left and right slopes can differ; the ReLU example exposes this.

Use Section 9 for the evidence ledger. Add a boundary or missing-input case and record the expected response before checking. More successful repeats of one case do not establish wider coverage.

Match the support to the first error

What the response shows · Next teaching action

What the response shows: Missing or misread facts

Next teaching action: Read the case aloud, mark supplied versus unknown facts, and let the learner restate it.

What the response shows: Wrong procedure

Next teaching action: Return to the deciding step in the worked case. Contrast it with the proposed incorrect explanation.

What the response shows: Arithmetic or implementation error

Next teaching action: Trace intermediate values or lines; preserve the first mismatch and retry that step.

What the response shows: Correct result without a reason

Next teaching action: Ask for a trace and a changed case before marking independent understanding.

What the response shows: An unsupported wider claim

Next teaching action: Ask which supplied evidence supports the claim and which further observation would be needed.

Offer an oral explanation, drawing, enlarged visual or fewer examples when that removes a reading/writing barrier. Keep the same reasoning goal. Stretch secure learners by changing a boundary or assumption and asking them to defend the effect, not simply by assigning more of the same questions.

5. Assess and explain the answers

Guided and exit questions

Guided: prediction 1, residual -2, loss 4, gradient [-8, -4]. Exit: changing both parameters measures a directional change; each partial must vary just its own coordinate while holding the others fixed.

W1 Recover a gradient

For the lesson model at $w = 1$ and $b = 1$, calculate prediction, residual, loss and gradient.

Teacher explanation: Prediction 3, residual 0, loss 0, gradient $[0, 0]$.

W2 Catch a denominator error

A w probe has losses 1.44 and 0.64 at 0.9 and 1.1 with b fixed. A learner divides their difference by 0.1. Correct the result.

Teacher explanation: The interval is 0.2. $(0.64 - 1.44) / 0.2 = -4$, not -8.

W3 Separate a direction from a coordinate

Starting at $[w, b] = [1, 0]$, both coordinates move by plus or minus 0.1 together. Compute the symmetric slope and explain why it is not either partial derivative.

Teacher explanation: Residuals are -1.3 and -0.7; losses 1.69 and 0.49. Slope is -6. It measures the combined direction $[1, 1]$, equal here to the sum of partials -4 and -2.

W4 Check a new example independently

Use $x = 3$, target 2 and $y = wx + b$ at $w = 0$, $b = 1$. With step 0.1, calculate both numerical partials. Compare with $[2 \text{ times residual times } x, 2 \text{ times residual}]$.

Teacher explanation: Residual -1 and loss 1. w probes give predictions 0.7 and 1.3, losses 1.69 and 0.49, partial -6. b probes give predictions 0.9 and 1.1, losses 1.21 and 0.81, partial -2. Formula gradient is $[-6, -2]$.

W5 Recognise a corner

Compute the symmetric difference of ReLU at zero with step 0.1. Explain why this does not prove a derivative exists there.

Teacher explanation: The value is 0.5. The left slope is 0 and right slope is 1, so the slopes disagree and the derivative at zero does not exist.

Independent assessment and fresh retry

W4 requires numerical and formula gradients $[-6, -2]$ with losses shown. Fresh retry: $x = 1$, target 4, $w = 2$, $b = 1$ gives residual -1 and gradient $[-2, -2]$. Step 0.1 in either coordinate gives losses 1.21 and 0.81. Require the fixed coordinate to be named.

Assess independent transfer

Selected case: Use $x = 3$, target 2 and $y = wx + b$ at $w = 0$, $b = 1$. With step 0.1, calculate both numerical partials. Compare with $[2 \text{ times residual times } x, 2 \text{ times residual}]$.

Selected case answer

Teacher explanation: Residual -1 and loss 1. w probes give predictions 0.7 and 1.3, losses 1.69 and 0.49,

partial -6. b probes give predictions 0.9 and 1.1, losses 1.21 and 0.81, partial -2. Formula gradient is [-6, -2].

Dimension · Evidence required

Dimension: Explanation

Evidence required: Names the applicable idea or procedure and ties it to the supplied facts.

Dimension: Trace and result

Evidence required: Preserves the given inputs and shows the relevant intermediate steps; distinguishes a prediction from an observed execution.

Dimension: Limits

Evidence required: States the assumptions, missing evidence or conditions under which the result would change.

Record each dimension as not yet, with support, or independent. Accept a valid alternative procedure with a defensible trace. Do not penalise an honestly recorded model failure if the learner correctly explains its evidence and limits.

Feedback that the learner uses

Name the earliest unsupported step and give one action to repair it. Have the learner make that repair now and explain why. Then assess the relevant skill on a fresh, fully specified case. Confirm it was not previously practised with feedback; solve and check the new case before assigning it. A follow-up design prompt is a starting brief, not automatically an equivalent test.

6. Connect the project, support and next lesson

Add a labelled gradients with a numerical check evidence sheet to Image-pattern Network Investigation. Show how you can estimate one partial derivative while other parameters stay fixed. Use the supplied fictional case and keep its inputs, intermediate steps, calculation or prediction, and limitation. Connect this record to last week's record; a matching answer without a trace is insufficient.

Provide separate w-probe and b-probe tables. Extension: at the lesson model [1, 0], test steps 0.1, 0.01 and 0.001 and report absolute disagreements from [-4, -2], without claiming universal tolerance guarantees.

Week 11 applies gradients to simultaneous parameter updates.

Retain the project ledger and assess a changed case after support. Check the relevant prerequisite before

carrying this component forward.

Plan the delayed return

Suggested return points in this level: ai-5/11, ai-5/13, ai-5/16. Use actual classroom dates, adjusting for holidays and gaps. One, three and six teaching weeks are scheduling choices, not a claimed optimal spacing.

At a delayed check, ask for the idea from memory and a changed application. Compare with the earlier independent response and record support used. For a new level, revisit the previous capstone before checking the new prerequisites.

Record · Teacher evidence

Record: Learner code and date

Teacher evidence: _____

Record: Concept and case used; prior exposure

Teacher evidence: _____

Record: Explanation / trace / limits

Teacher evidence: _____

Record: Support and first repair

Teacher evidence: _____

Record: Changed case and delayed outcome

Teacher evidence: _____

Record: Next teaching action

Teacher evidence: _____

Use the evidence to choose reteaching or the next step. A completed page, confidence rating or attractive project does not by itself demonstrate understanding.

Project evidence from this case

For your Image-pattern Network Investigation, save the input, procedure, observed result and limitation of this check. Label this worked example as practice; use a different, previously unreviewed case when checking independent understanding.

7. Prepare, teach and assess

Teaching target: Estimate one partial derivative while other parameters stay fixed

Prepare the labelled worked-case picture, the matching workbook, fictional input cards and a place for individual responses. Check a local Python 3 environment before class; have a paper-trace version ready.

Session · Plan · Resources

Session: Session 1 · Understand · 45 min

Plan: 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

Resources: Sections 1–6; record the brief recall in the workbook.

Session: Session 2 · Practise · 45 min

Plan: 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.

Resources: Sections 7–10; use the matching workbook W2 and misconception check.

Session: Session 3 · Investigate and transfer · 45 min

Plan: 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

Resources: Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

Before class, solve the supported practice: A w probe has losses 1.44 and 0.64 at 0.9 and 1.1 with b fixed. A learner divides their difference by 0.1. Correct the result.

Reasoned practice answer: Teacher explanation: The interval is 0.2. $(0.64 - 1.44) / 0.2 = -4$, not -8.

Check every learner before discussion: Use the worked case for Gradients with a numerical check.

Proposed explanation: “The gradient is one scalar”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Misconception repair: It has one entry per parameter for a scalar loss, in a stated order.

Support: read the exact case aloud, enlarge the labelled picture, supply input cards and model only the first deciding step. Accept a spoken explanation or drawing; retain the reasoning goal.

Extension: Swap the two gradient components in the program output. Require a prediction and a reason before comparing. Expected result: The coordinate-wise numerical check exposes the mismatch. Reason: The w perturbation changes y twice as strongly as the same b perturbation.

Independent assessment: withhold the reserved answer until the learner has attempted the new case. Record support separately from correctness.

Criterion · Not yet · With support · Independent

Criterion: Explanation

Not yet: Names an answer without a deciding fact

With support: Uses a hint to link fact and decision

Independent: Explains the deciding fact in own words

Criterion: Trace or test

Not yet: Cannot connect input and result

With support: Completes steps with prompts

Independent: Traces or tests a changed case accurately

Criterion: Limits and revision

Not yet: Claims more than evidence supports

With support: Names a limit after a prompt

Independent: States a defensible limit and useful next test

Project checkpoint: Make a transparent first version. Keep a trace of how one input becomes an output; explain the rule or learned component.

Use the weekly teacher answer for topic-specific reasoning. A saved progress tick is an activity record; judge mastery from the learner's explanation and evidence.

8. Use feedback, labs and repair paths

Use the two formative questions as supported practice, not as independent mastery scores. Discuss the deciding evidence rather than accepting a guessed selection.

The feedback comes from the reviewed public worked case. Do not reveal the separately reserved

independent teacher answer until the learner has attempted that case.

Ask students to predict before running code. Preserve errors and actual outputs; collect a changed test and an explanation. The branch-boundary and indexing lessons deliberately include failure cases.

Use the earlier worked-explanation links to address a specific missing building block, then return to the current question. Adapt this suggestion after observing the learner; it is not an automatic diagnosis.

Collect the completed-work portfolio as evidence. Record support separately from correctness, and assess explanation, trace/test and limitations. Progress ticks and quiz selections alone do not establish mastery.