

Accuracy as a Fraction

Project: Transparent Book Recommender

1. Your mission

Your learning target

I can explain why overall accuracy can hide failure on a less common class.

I can compare a training-majority baseline with class-specific results.

I can calculate an equal-class average and report its limits alongside the confusion matrix.

Remember before reading

Close your notes first. A rough first answer helps us choose the right starting point; it is not a score for your ability.

Without opening the old lesson, explain one key idea from Training and Testing. Give an example and a mistake that the idea helps you avoid.


Without opening the old lesson, explain one key idea from Features and Distractors. Give an example and a mistake that the idea helps you avoid.

Without opening the old lesson, explain one key idea from Debugging Strategies. Give an example and a mistake that the idea helps you avoid.

After trying, check the relevant lesson or ask your teacher. Change the part of your explanation that the evidence shows was wrong.

Week 12: Accuracy as a Fraction

Accuracy is correct predictions divided by all predictions.



THINKER MISSION

Calculate and compare 7/10 and 18/20 models.

Thinker name and date:

Write or draw your reasoning and evidence.

Readiness for the connected explanation

With reference rows and prediction columns ordered clear then blocked, where are false negatives when blocked is positive?

Readiness for Summaries with counts and proportions

If 3 of 12 responses are Yes, what fraction are Yes?

My readiness explanation

This week's end goal


By the end, I can compare accuracy with class-specific performance in an imbalanced evaluation.

2. Notice the evidence

PICTURE STORY

Notice → Predict → Explain

Begin with visible evidence. Do not explain more than the picture and information support.



MYSTERY

The learners must use accuracy as a fraction to solve a problem. What information do they have? What decision or calculation do they make? What would count as evidence that it worked?

Three observations (what you can point to):

Write or draw your reasoning and evidence.

My prediction and the clue supporting it:

Write or draw your reasoning and evidence.

A closer explanation

Accuracy counts all correct predictions and divides by all evaluated examples. Each example has equal weight, so a common class contributes more to the score. A high result can hide failure on a rare class. This week keeps the confusion matrix visible and checks both classes separately.

New fictional training data has 16 clear and 4 blocked labels. Its majority baseline predicts clear for every query; choose that rule from training, not evaluation labels. A separate evaluation set also contains 16 clear and 4 blocked cases. A frozen candidate and the baseline are evaluated on the same 20 records.

Method and reference row · Predicted clear · Predicted blocked

Method and reference row: Baseline reference clear

Predicted clear: 16

Predicted blocked: 0

Method and reference row: Baseline reference blocked

Predicted clear: 4

Predicted blocked: 0

Method and reference row: Candidate reference clear

Predicted clear: 12

Predicted blocked: 4

Method and reference row: Candidate reference blocked

Predicted clear: 0

Predicted blocked: 4

Rows are references and columns are predictions, ordered clear then blocked. Blocked is positive. Both methods get $16/20=80$ percent accuracy. The baseline misses all four blocked cases. The candidate finds all four but falsely flags four clear cases. The same score conceals a different tradeoff.

Apply the connected idea: Summaries with counts and proportions


A fictional poll asks whether a short activity was useful. Responses are Yes, No or Unknown. We will report Yes among known responses and also disclose unknown responses. Choosing a denominator changes the question being answered.

3. See how it works

VISUAL MECHANISM

Input → Process → Output

Trace the information. At each stage, ask what changed and how you could test it.



↓



PROCESS

Accuracy is correct predictions divided by all predictions.





OUTPUT

Calculate and compare 7/10 and 18/20 models.

Which stage could fail even when the other two are correct? Explain:

Write or draw your reasoning and evidence.

A closer explanation

```
def scores(counts):  
    clear_n = sum(counts[0])  
    blocked_n = sum(counts[1])  
    if clear_n == 0 or blocked_n == 0:  
        raise ValueError("Need evaluation examples of both classes")  
    correct = counts[0][0] + counts[1][1]  
    accuracy = correct / (clear_n + blocked_n)  
    clear_fraction = counts[0][0] / clear_n  
    blocked_fraction = counts[1][1] / blocked_n  
    balanced = (clear_fraction + blocked_fraction) / 2  
    return accuracy, balanced  
  
baseline = [[16, 0], [4, 0]]  
candidate = [[12, 4], [0, 4]]  
  
for name, counts in [("baseline", baseline), ("candidate", candidate)]:  
    accuracy, balanced = scores(counts)  
    print(name, round(100 * accuracy, 1), round(100 * balanced, 1))
```

The output is baseline 80.0 50.0, then candidate 80.0 87.5. In each output line the first percentage is ordinary accuracy and the second is balanced accuracy. For the candidate, the clear-class correct fraction is $12/16=75$ percent and the blocked-class fraction is $4/4=100$ percent. Their equal average is 87.5 percent.

For the baseline, those fractions are $16/16=100$ percent and $0/4=0$ percent, averaging 50 percent. The code assumes a two-by-two table of nonnegative integer counts in the stated order. It stops when either reference class is absent. Then report the observed counts and the unavailable class result; do not invent a denominator or silently substitute zero.

Apply the connected idea: Summaries with counts and proportions

```
responses = ["Yes", "No", "Yes", "Unknown", "Yes"]  
  
yes = 0  
  
known = 0  
  
for response in responses:  
    if response == "Yes" or response == "No":
```

```
known = known + 1

if response == "Yes":
    yes = yes + 1

if known > 0:
    print(yes / known)
else:
    print("No known responses")
```

Follow the evidence

Step or case · Evidence · Meaning

Step or case: Yes

Evidence: 3

Meaning: Numerator

Step or case: Yes or No

Evidence: 4

Meaning: Known denominator

Step or case: All supplied

Evidence: 5

Meaning: Includes one unknown

Read each row from left to right and connect the evidence to the stated rule.

The printed proportion is 0.75, or 75 percent, among four known responses. Yes among all five supplied responses is 0.6, or 60 percent. Both calculations are correct when labelled; they describe different denominators. Unknown must not silently become No.

Name the group behind a percentage

Count the responses before dividing. In the example, three say Yes, one says No and one is Unknown. There are four known responses, so Yes among known responses is 3 divided by 4, or 75 percent. Yes among all five supplied responses is 3 divided by 5, or 60 percent. Neither number needs to be hidden. Label the question each answers, and keep the unknown count visible.

Connect the picture and the explanation



Point to the input, the deciding step and the result in this week's visual. Explain the same step in ordinary words. A picture helps when you can say what each part represents.

4. Compare the cases

CASE GALLERY

Four Tests, Four Kinds of Evidence

A system is not understood until it survives more than one comfortable example.




NORMAL CASE: predict and name evidence. BOUNDARY CASE: predict and name evidence.



MISSING

Needed information absent



TRANSFER

Fresh situation or user

MISSING INFORMATION: predict and name evidence. TRANSFER CASE: predict and name evidence.

Which case is most likely to reveal a hidden assumption? Why?

Write or draw your reasoning and evidence.

Change one thing in the pictured case

Change: A candidate instead marks every book repair needed.

Prediction: Recall is $4/4=100\%$, but accuracy is only $4/20=20\%$.

Reason: All 16 not-needed books become false alarms.

5. Words that unlock the idea

VOCABULARY LAB

Words You Can Use to Reason

A definition matters when you can recognize the idea, use it, and contrast it with a near-miss.

1

ACCURACY

correct predictions divided
by all predictions

2

NUMERATOR

top number: correct
predictions

3

DENOMINATOR

bottom number: all
predictions

4

PERCENTAGE

a fraction out of 100

Word · My own example · Not this / boundary

Word: accuracy

My own example: The number of correct predictions divided by the total number of scored predictions.

Not this / boundary: _____

Word: numerator

My own example: The top number of a fraction, identifying the counted portion.

Not this / boundary: _____

Word: denominator

My own example: The bottom number of a fraction, identifying the number of equal parts or total cases considered.

Not this / boundary: _____

Word: percentage

My own example: A proportion expressed per hundred; multiply a fraction by 100 to express it as a percent.

Not this / boundary: _____

Words used in the closer explanation

Word · Meaning

Word: Class imbalance

Meaning: Unequal numbers of examples in different classes

Word: Majority baseline

Meaning: A rule that always predicts the most common training label

Word: Class-specific correct fraction

Meaning: Correct predictions for one reference class divided by the number of examples of that class

Word: Balanced accuracy

Meaning: The average of the class-specific correct fractions with equal weight for each class

Terms for Summaries with counts and proportions

Term · Meaning

Term: Count

Meaning: The number of matching records

Term: Denominator

Meaning: The total number used in a fraction

Term: Proportion

Meaning: A part divided by its relevant whole

Term: Eligible

Meaning: Included under a stated rule

6. Follow a worked example

WORKED EVIDENCE

Reason Step by Step

Predict first. Then check each step. A correct answer without visible reasoning is incomplete evidence.

WEEK 12 • WORKED EVIDENCE

- 1 Model A: 7 correct out of 10 $\rightarrow 7/10 = 70\%$.
- 2 Model B: 18 correct out of 20 $\rightarrow 18/20 = 90\%$.
- 3 Model B has higher measured accuracy on these tests.
- 4 Still ask: were the test cases fair and were some errors more costly?

1. Model A: 7 correct out of 10 $\rightarrow 7/10 = 70\%$.
2. Model B: 18 correct out of 20 $\rightarrow 18/20 = 90\%$.
3. Model B has higher measured accuracy on these tests.
4. Still ask: were the test cases fair and were some errors more costly?

Recalculate, retrace, or restate the reasoning in your own representation:

Write or draw your reasoning and evidence.

CHECK Did the evidence answer the exact question? Did the total, path, or source record stay consistent?

A closer explanation

Same overall score different class results

Summary · Always clear baseline · Frozen candidate

Summary: Overall correct

Always clear baseline: 16 of 20 is 80 percent

Frozen candidate: 16 of 20 is 80 percent

Summary: Reference clear correct

Always clear baseline: 16 of 16 is 100 percent

Frozen candidate: 12 of 16 is 75 percent

Summary: Reference blocked correct

Always clear baseline: 0 of 4 is 0 percent

Frozen candidate: 4 of 4 is 100 percent

Overall accuracy weights classes by their number of evaluation examples. Balanced accuracy gives the two class-specific fractions equal weight.

Both methods score 80 percent. Explain which cases the baseline fails, how the candidate changes the errors, and why their balanced accuracies differ.

Balanced accuracy is a metric choice, not a repair to the classifier or a guarantee of safety. It gives classes equal influence in this summary; it does not encode the actual consequences of false alarms and missed cases. Four blocked evaluation examples are also too few for a strong general claim.

Apply the connected idea: Summaries with counts and proportions

For Yes, No, Unknown, Unknown, calculate Yes among known and Yes among all supplied.

Learn from the worked case

Cover the result before checking it. Say what is given, choose the procedure, and follow one step at a time. Then uncover the result and explain your first mismatch. Ask why a step is allowed, not only what number it produces.

A pictorial worked case: Accuracy as a Fraction

A fictional shelf classifier predicts repair needed or not needed.

The facts we will use

Evaluation reference labels: 16 not-needed and 4 needed. Positive class is needed.

Always-not-needed baseline is correct on 16. Candidate has TP=3,FN=1,FP=2,TN=14.

AI ACADEMY / AI-2/12

Accuracy can hide missed repair cases

percent



Bar length shows the supplied value; the marked baseline is zero.

Accuracy as a Fraction

Start with the supplied facts

Follow each arrow. Ask which fact allows the next step.



1. Reference population

20 labelled cases, including four repair-needed books.



2. Baseline

$16/20=80\%$ accuracy; catches 0/4 needed cases.



3. Candidate

$(3+14)/20=85\%$ accuracy; catches $3/4=75\%$ needed cases.



4. Compare costs

Candidate also raises two false alarms; discuss both missed repairs and extra checks.

CHANGE ONE THING

A candidate instead marks every book repair needed.

Explain the changed result using the matching lesson.

Read the picture one step at a time

Step · What happens and why

Step: 1. Reference population

What happens and why: 20 labelled cases, including four repair-needed books.

Step: 2. Baseline

What happens and why: $16/20=80\%$ accuracy; catches 0/4 needed cases.

Step: 3. Candidate

What happens and why: $(3+14)/20=85\%$ accuracy; catches $3/4=75\%$ needed cases.

Step: 4. Compare costs

What happens and why: Candidate also raises two false alarms; discuss both missed repairs and extra checks.

Match each pictured step to the supplied facts. This is a worked teaching case, not an independent assessment.

Why this result follows

A majority-class baseline can look accurate while missing every positive case. The candidate improves overall agreement and repair detection in this table, but false alarms still matter. Use the same cases, denominators and positive-class definition for both. Do not infer that a single metric identifies the best policy for every library.

What this example does not establish

These selected cases do not establish future rates or specify the real cost of each error.

7. Find and repair a mistake

ERROR DETECTIVE

Compare → Diagnose → Repair

Do not patch the answer. Find the earliest unsupported assumption, wrong step, or weak evidence.



COMPARE • DIAGNOSE

Weak idea: The model with more correct answers always has higher accuracy, even if test sizes differ. Strong move: locate evidence, change one cause, and retest.

Evidence record · Expected · Actual · Likely cause / next test

Evidence record: normal case

Expected: works

Actual: _____

Likely cause / next test: _____

Evidence record: boundary case

Expected: stated rule

Actual: _____

Likely cause / next test: _____

Evidence record: missing input

Expected: safe response

Actual: _____

Likely cause / next test: _____

Evidence record: fresh transfer

Expected: generalizes

Actual: _____

Likely cause / next test: _____

Repair one cause. Why is this repair better than changing several things?

Write or draw your reasoning and evidence.

A closer explanation

Compare methods on the same appropriate evaluation examples and report the training-derived baseline. Include ordinary accuracy, each class denominator, false positives and false negatives. An equal-class

summary adds information but cannot choose the acceptable tradeoff for the task. Keep development decisions separate from a final test.

A result of 4/4 correct blocked cases is 100 percent on those four examples, not proof of perfect future detection. If the evaluated set contains no blocked cases, it cannot measure blocked-case performance at all. Next week separates precision and recall by asking which denominator answers each question.

A team advertises 80 percent accuracy but its classifier predicts clear on every record. What additional evidence would you request, and what can you already conclude about the four blocked cases in our worked evaluation?

Apply the connected idea: Summaries with counts and proportions

A percentage from a few volunteers is not automatically representative of the whole class. Report counts, the question, who responded and exclusions alongside the percentage.

Three of five known volunteers said Yes. Can you say 60 percent of all students agree?

8. Practise together

LOGIC STUDIO

Trace Before You Run

Block code, pseudocode, tables, and diagrams are different ways to expose the same reasoning.



TRACE THE LOGIC

Predict the result. Trace every condition or calculation. Compare expected with actual. Explain the first difference.

SET evidence_record PREDICT expected_result RUN one normal case and one boundary case
COMPARE actual_result with expected_result IF different: locate first cause; change one factor; RETEST
EXPLAIN result, limitation, and next safe step

Trace one input. Which line changes the state or chooses a path?

Write or draw your reasoning and evidence.

A closer explanation

Consider a separate illustrative set with 8 clear and 8 blocked cases. Suppose the same within-class fractions hold: the baseline matrix is $[[8,0],[8,0]]$ and the candidate matrix is $[[6,2],[0,8]]$. Ordinary accuracy is now 50 percent versus 87.5 percent. Balanced accuracy remains 50 percent versus 87.5 percent. The class mix changed the overall score even though the within-class fractions stayed fixed.

This is an arithmetic comparison, not proof that a real classifier keeps the same behavior after a population change. Report sample size, class counts, full matrix and how the evaluation data was chosen. Do not remove inconvenient cases or duplicate rare evaluation cases just to improve a headline number. If training is resampled, keep that separate from evaluation and document it.

Apply the connected idea: Summaries with counts and proportions

Dividing by known when it equals 0 causes ZeroDivisionError. Test $\text{known} > 0$ before division and report no known responses if none exist. Reporting 0 percent would wrongly suggest known responses with no Yes answers.

Practise with less help: Compare both class fractions

Use workbook W2 for the connected example “Accuracy and class imbalance”. First fill the input and rule boxes. Try the middle steps before asking for a hint. After a hint, try the next step yourself.

Plan · Do · Check

Plan: What is given? Which rule or procedure applies?

Do: Record each intermediate step.

Check: Does the result follow? What remains unknown?

Explain your trace to a partner. Your partner points to one step to check. Swap roles on the next case. Each person writes their own explanation.

Check your reasoning

Use the worked case for Accuracy and class imbalance. Proposed explanation: “High accuracy means both classes work well”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Answer on your own before discussion. If you are unsure, say which step you need help with.

Practise the connected topic: Summaries with counts and proportions

Use its named workbook W2. Identify the given inputs, try the middle steps with less help, then explain your trace. Check this claim before group discussion: A successful run proves the conclusion

Explain the picture

Why is 100% repair detection not sufficient to declare the changed classifier best?

My explanation using a pictured step and the given facts

This is supported practice. Keep the separately named independent case for an attempt before feedback.

9. Run the investigation

EVIDENCE LAB

Predict → Test → Record → Explain

Keep expected and actual results separate. Surprises are useful data when recorded honestly.



PREDICT → TEST → RECORD → EXPLAIN

Score two prediction sheets, simplify or convert each fraction, compare percentages, then inspect which cases were wrong.

Score two prediction sheets, simplify or convert each fraction, compare percentages, then inspect which cases were wrong.

Test / input · Prediction · Actual · Evidence / calculation · Change or keep?

Test / input: normal

Prediction: _____

Actual: _____

Evidence / calculation: _____

Change or keep?: _____

Test / input: boundary

Prediction: _____

Actual: _____

Evidence / calculation: _____

Change or keep?: _____

Test / input: missing

Prediction: _____

Actual: _____

Evidence / calculation: _____

Change or keep?: _____

Test / input: fresh transfer

Prediction: _____

Actual: _____

Evidence / calculation: _____

Change or keep?: _____

What did the hardest test teach you that the normal test did not?

Write or draw your reasoning and evidence.

Plan → try → check


Before trying: what do I expect, and why? While trying: which step could first go wrong? Afterwards: what did I actually observe, and what should change? Keep “not run” if you only traced the procedure.

10. Build your understanding

MASTERY

From Knowing the Word to Owning the Idea

Move upward only when you can produce evidence without copying the worked example.



MASTERY LADDER
1 Name it • 2 Trace it • 3 Apply it • 4 Challenge it • 5 Explain limits

1 • NAME: Define the concept without repeating the textbook.

2 • TRACE: Show the information, state, branch, calculation, or evidence path.

3 • APPLY: Solve a different example independently.

4 • CHALLENGE: Find a boundary case, counterexample, or unfair outcome.

5 • EXPLAIN LIMITS: Say when the result should not be trusted or used alone.

My strongest evidence of independent understanding:

Write or draw your reasoning and evidence.

11. Connect your project

CUMULATIVE PROJECT

Transparent Book Recommender

Every week adds one inspectable piece. The system must show how it reached a result.



TRANSPARENT BOOK RECOMMENDER

Report correct recommendations / all test users, alongside the raw test count.

THIS WEEK'S CHECKPOINT Report correct recommendations / all test users, alongside the raw test count.

SYSTEM MAP Reader need → allowed inputs → transparent rule or score → ranked candidates → explanation → test record → human decision

What will you add, change, or verify in the project this week?

Write or draw your reasoning and evidence.

What should remain under human control?

Write or draw your reasoning and evidence.

RESPONSIBLE DESIGN Collect only necessary information. Show uncertainty and limits. Never confuse a ranking with a fact about a person.

Evidence for your project


Compare recommender choices with a simple baseline and report relevant rare-category failures, not only aggregate accuracy.

12. Show what you know

INDEPENDENT MASTERY PROOF

Explain • Transfer • Test • Defend

Complete this page without copying the worked example. Your reasoning matters more than decoration.



INDEPENDENT PROOF

Calculate and compare $7/10$ and $18/20$ models.

CHALLENGE Calculate and compare $7/10$ and $18/20$ models.

1. My fresh example, trace, calculation, or design:

Write or draw your reasoning and evidence.

2. My evidence that it works (include a boundary or missing-data case):

Write or draw your reasoning and evidence.

3. One limitation, fairness concern, or situation needing human review:

Write or draw your reasoning and evidence.

SELF-CHECK ☐ I explained the mechanism ☐ I used evidence ☐ I tested a fresh case ☐ I named a limitation ☐ I can answer 'why?'

13. Study a complete case

Week 12 Worked case

Use the supplied details to practise the weekly concept before attempting the independent question. This case extends the illustrated lesson.

The situation

Nine of twelve held-out classifications are correct.

The question

Compute accuracy and state the denominator.

Worked explanation

$9/12=3/4=75\%$. Count all scored cases, including errors.

Explain it in your own words

Which detail supports the answer, and why does it matter?

Write or draw your reasoning and evidence.

Trace the supplied case visually

All scored cases belong in the denominator

Evidence or stage · Value or result

Evidence or stage: Correct classifications

Value or result: 9/12

Nine of twelve held-out classifications are correct.

Compute accuracy and state the denominator.

$9/12=3/4=75\%$. Count all scored cases, including errors.

14. Try a new case independently

Independent case: Assess a new comparison independently

New training labels have a clear majority. A separate evaluation has 18 clear and 2 blocked records. The always-clear baseline is $[[18,0],[2,0]]$ and candidate is $[[15,3],[1,1]]$, with reference rows and clear then blocked on both axes. Calculate ordinary accuracy, both class fractions and balanced accuracy for each. Give candidate FP and FN, then explain why the metrics do not choose an automatic winner.

My independent explanation, trace and limits

Attempt this case before feedback. Record any help. Earlier examples below are further practice; a case already practised with feedback cannot establish fresh transfer.

Week 12 Independent application

Close the worked explanation. Apply the idea to this different question without copying.

A classifier gets 9 of 12 cases right. Another gets 8 of 10 right. Calculate both accuracies and identify the larger fraction.

My answer, with a trace, calculation or supporting evidence

Write or draw your reasoning and evidence.

Create and test

Change one relevant detail. Predict the result before testing. Include a boundary or missing-information case when it helps reveal a limit.

My changed input and expected result

Write or draw your reasoning and evidence.

Connect to your project

Report correct recommendations / all test users, alongside the raw test count.

The evidence I will keep

Write or draw your reasoning and evidence.

Your teacher has the independent answer and scoring criteria. Complete the matching workbook week to keep your practice evidence together.

Transfer practice from the original programme

Class A gets 8/10 cases right and class B gets 18/30 right. Compute each rate and the combined rate.

Why is averaging the two percentages misleading here?

Explain the mechanism, evidence and one limit.

Use feedback to improve

Attempt the independent case before hints. Record whether you worked alone, used a hint or followed a model. After feedback, fix the first unsupported step and explain why it changed. A later check uses a changed case, not a copied answer.