

# Mean, Median, and Baselines

Project: Responsible Plant Classifier

## 1. Mean, Median, and Baselines

### Teach the learning cycle

Primary target: Fit mean and median constant baselines on training targets

#### Session · Sequence and minutes · Resources

**Session:** Session 1 · Understand · 45 min

**Sequence and minutes:** 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

**Resources:** Sections 1–6; record the brief recall in the workbook.

---

**Session:** Session 2 · Practise · 45 min

**Sequence and minutes:** 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.

**Resources:** Sections 7–10; use the matching workbook W2 and misconception check.

---

**Session:** Session 3 · Investigate and transfer · 45 min

**Sequence and minutes:** 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

**Resources:** Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

---

This is a starting allocation of 135 minutes, not a validated duration. Preserve all named topics. Schedule extra time for connected examples, the full investigation, independent cases and project building. Repeated copies of a case share one recorded attempt; they are not new evidence.

## Retrieval answer guidance

### Regression and Residuals

Without opening the old lesson, explain one key idea from Regression and Residuals. Give an example and a mistake that the idea helps you avoid.

Look for: Distinguish numeric prediction from classification

Reference example, not the only acceptable wording: Teacher explanation: Residuals -1, 1, 0 cm; the first record is overpredicted.

### Precision and Recall

Without opening the old lesson, explain one key idea from Precision and Recall. Give an example and a mistake that the idea helps you avoid.

Look for: Choose the denominator that answers the question

Reference example, not the only acceptable wording: Teacher explanation: Predicted positives  $TP+FP=7$ ; reference positives  $TP+FN=4$ .

### Scores and Calibration Intuition

Without opening the old lesson, explain one key idea from Scores and Calibration Intuition. Give an example and a mistake that the idea helps you avoid.

Look for: Interpret calibration using groups of forecasts

Reference example, not the only acceptable wording: Teacher explanation:  $2/10=.2$  and  $6/10=.6$ .

Use the student module and workbook section with this exact week title. The worked case, independent question and answers below form one aligned lesson sequence.

### What the learner must demonstrate

For training targets [1,3,11], compute mean and median constant baselines. Compare total absolute training error for each. Where must final performance be measured?

### Concept to explain

Mean and median are transparent regression baselines; outliers affect them differently.

#### Term · Meaning

**Term:** mean

**Meaning:** sum divided by count

---

**Term:** median

**Meaning:** middle ordered value

---

**Term:** outlier

**Meaning:** value far from most observations

---

**Term:** location baseline

**Meaning:** single typical value predicted for all cases

---

## Prepare the class

Use the supplied scenario, student illustrations and blank paper. For numeric work, offer counters or a calculator after learners show the setup. Fictional records are supplied; personal data and paid accounts are unnecessary.

Allow 80 minutes: recall 8; explanation 12; worked demonstration 15; guided practice 20; independent assessment 15; project and reflection 10. Extend the lab into a second session when needed.

## Preparation for the connected explanation

Teach two 45-minute sessions. This week focuses on constant prediction baselines. Model the worked example first, collect the new W 4 case independently before feedback, and use the fresh retry to check transfer. Award credit for traceable reasoning and bounded claims, including justified uncertainty.

## Readiness answer

Opposite signs can cancel; a magnitude-based loss avoids that cancellation.

Printed student lesson and workbook

Paper, pencils and a calculator for fractions

The fictional examples and a blank evidence table

Optional Python 3 editor; no external packages or AI accounts required

## End goal for the pictorial case

By the end, I can explain mean, median, and baselines using a traced example and a stated limitation.

This added worked case illustrates the week's main idea. Retain all original and connected topics; schedule its practice within the teaching cycle.

# 2. Explain the mechanism

## Explain the mechanism

Trace the information, fitted behavior, and evaluation proof separately.

QUESTION Choose and justify a baseline under an extreme value.

MODEL IDEA Mean and median are transparent regression baselines; outliers affect them differently.

PROOF NEEDED Use an unseen case, visible calculation, error analysis, and a stated limitation.

UNDER THE HOOD Heights are 10, 11, 12, 13, and 54 cm. Mean =  $100 \div 5 = 20$  cm.

Draw a fresh data → model → prediction → evaluation trace:

Predict first. Then check each step. A correct answer without visible evidence cannot be audited.

STEP 1 Heights are 10, 11, 12, 13, and 54 cm.

STEP 2 Mean =  $100 \div 5 = 20$  cm.

STEP 3 Median is 12 cm, the middle ordered value.

STEP 4 For typical-height absolute error, median is more robust; still report why 54 is present.

CHECK Verify the denominator, unit, split, class definition, and data version before interpreting a score.

Recalculate, retrace, or restate the evidence independently:

## Concrete teaching case

Training heights are 2,4,12 cm. Compare constant mean and median baselines.

Compute both and training absolute errors.

## Exact explanation to model

Mean=6, median=4. Mean errors 4,2,6 total12; median errors 2,0,8 total10. Median minimises absolute error here.

## Explain the added worked example

Before a complex regression model, fit a simple constant predictor. A mean baseline uses the training average. A median baseline uses the middle training target. They ignore features but provide a comparison that is easy to reproduce. Evaluation targets must not determine the constants.

Use fictional classroom data throughout this investigation. Keep the task, units, reference labels and data split visible as you reason. Earlier weeks supplied the model-building process; this week examines constant prediction baselines.

Training targets [2, 4, 6, 12] have mean  $24/4=6$  and median  $(4+6)/2=5$ . On separate targets [4, 8], the mean baseline predicts [6, 6] with absolute errors [2, 2], averaging 2. The median baseline predicts [5, 5] with errors [1, 3], also averaging 2.

Trace the calculation on paper before running this optional Python check. Use the stated fictional inputs

and compare intermediate values. The code checks this example; your independent workbook case uses different evidence.

## Key terms

### Term · Meaning

**Term:** Constant baseline

**Meaning:** One fixed prediction for every query

---

**Term:** Training mean

**Meaning:** Sum of training targets divided by their count

---

**Term:** Training median

**Meaning:** Middle value of sorted training targets

---

**Term:** Absolute error

**Meaning:** Magnitude of observed minus predicted

---

## Model the reasoning aloud

Use the complete worked example below before asking learners to complete W2. Reveal a small part, invite a prediction, then explain the deciding step. These teacher notes contain solutions; do not reveal the independent case answer during modelling.

## Constant prediction baselines at a glance

### Part · Evidence or action · Interpretation

**Part:** Training targets

**Evidence or action:** Fit mean 6

**Interpretation:** Freeze constant

---

**Part:** Training targets

**Evidence or action:** Fit median 5

**Interpretation:** Freeze constant

---

**Part:** Evaluation targets

**Evidence or action:** Compare fixed predictions

**Interpretation:** Both MAE 2 here

---

Mean and median answer different training objectives. A mean minimizes total squared error among constants; a median minimizes total absolute error. This does not guarantee either wins on a new evaluation set. With extreme values, the mean can move much more than the median.

**Calculate MAE for both baselines on [4, 8]. Explain the steps or evidence behind your response.**

---

---

---

---

**Thinking move · Teacher prompt**

**Thinking move:** Plan

**Teacher prompt:** What is given? What is the question? Which procedure fits these facts?

---

**Thinking move:** Do

**Teacher prompt:** Point to each input and intermediate step in the worked trace. Name the rule that allows the next step.

---

**Thinking move:** Check

**Teacher prompt:** Does the output answer the question? Which assumption or limitation prevents a larger claim?

---

Use the diagram and verbal explanation together. Ask learners to match a sentence to a specific visual step; do not assign learners fixed visual or auditory learning styles.

**A pictorial worked case: Mean, Median, and Baselines**

Fit two constant height baselines and compare them on held-out examples.

## The facts we will use

Training heights [2, 4, 6, 20] cm. Evaluation heights [4, 10] cm. Fit constants only from training.

All inputs and outcomes in this worked example are invented classroom data; no real model run is claimed.

AI ACADEMY / AI-4/20

## Mean, Median, and Baselines

### Start with the supplied facts

Follow each arrow. Ask which fact allows the next step.



### 1. Fit mean

Training mean= $(2+4+6+20)/4=8$  cm.



### 2. Fit median

Training median= $(4+6)/2=5$  cm.



### 3. Evaluate constants

Mean MAE= $(4+2)/2=3$  cm; median MAE= $(1+5)/2=3$  cm.

### CHANGE ONE THING

Replace training 20 with 8 while keeping evaluation unchanged.

Explain the changed result using the matching lesson.

## Read the picture one step at a time

### Step · What happens and why

**Step:** 1. Fit mean

**What happens and why:** Training mean= $(2+4+6+20)/4=8$  cm.

---

**Step: 2. Fit median**

**What happens and why:** Training median= $(4+6)/2=5$  cm.

---

**Step: 3. Evaluate constants**

**What happens and why:** Mean MAE= $(4+2)/2=3$  cm; median MAE= $(1+5)/2=3$  cm.

---

Match each pictured step to the supplied facts. This is a worked teaching case, not an independent assessment.

### Why this result follows

Read the picture in order. Mean MAE= $(4+2)/2=3$  cm; median MAE= $(1+5)/2=3$  cm. The arrows show which recorded input or intermediate result each decision uses. Explain the deciding step aloud before checking the changed case; pointing to the final answer alone does not show how it follows.

### What this example does not establish

Mean and median have different loss properties, but their held-out performance depends on the particular evaluation cases.

### Change one thing in the pictured case

Change: Replace training 20 with 8 while keeping evaluation unchanged.

Prediction: Both fitted constants become 5 cm and both evaluation MAEs remain 3 cm.

Reason: Mean is now  $20/4=5$ ; median stays 5. The earlier training outlier affected the mean.

### Ask while showing the picture

Cover the result. Ask learners to name the inputs and predict the next step. Uncover one step, connect the icon to the actual procedure, and explain its evidence. Then change the supplied condition and compare. The picture is a representation; it does not document execution of a real AI model.

## 3. Demonstrate and guide practice

### Demonstrate and guide practice

#### Board sequence

Write the given facts: Training heights are 2, 4, 12 cm. Compare constant mean and median baselines.

Write the question to explain: Compute both and training absolute errors.

Reveal after predictions: Mean=6, median=4. Mean errors 4,2,6 total12; median errors 2,0,8 total10.

Median minimises absolute error here.

Keep for an independent attempt: For training heights 2, 3 and 10 cm, calculate mean and median. Which constant is less influenced by the high value?

## Classroom dialogue

Ask: Which facts are given, and which would we have to assume? Learners mark relevant details before discussing answers.

Say: Predict first. Show the rule, information path or calculation that would produce the result. A correct guess is not yet an explanation.

Reveal reasoning one step at a time. Explain why a tempting alternative fails. Write units and denominators for arithmetic, trace conditions for decisions, and identify supporting sources for claims.

In pairs, one learner solves while the other checks evidence; exchange roles. Remove the worked answer before independent assessment.

## Model these guided questions

Guided response: Both score 2; their per-record absolute errors differ. Exit response: It fits using evaluation answers and makes the baseline unavailable for genuinely unseen cases.

## Case-specific teaching sequence

### Stage · Teacher action

**Stage:** Session 1 start to 7 minutes

**Teacher action:** Use the prerequisite question: Why can signed residuals hide error size? Check the explanation before proceeding.

---

**Stage:** 7 to 18 minutes

**Teacher action:** Read the opening and model the worked evidence. Ask students to identify constant baseline in context.

---

**Stage:** 18 to 32 minutes

**Teacher action:** Complete W 1 and W 2 with support: fit two constants; score separately. Require the evidence, not only the final response.

---

**Stage:** 32 to 45 minutes

**Teacher action:** Read the visual row by row. Collect the guided explanation and compare reasoning in pairs. Trace the Python check after the paper calculation.

---

**Stage:** Session 2 start to 8 minutes

**Teacher action:** Retrieve the distinction between constant baseline and training mean. Invite a counterexample to an overconfident claim.

---

**Stage:** 8 to 20 minutes

**Teacher action:** Discuss the application and complete W 3: Replace 12 by 40. What changes?

---

**Stage:** 20 to 35 minutes

**Teacher action:** Collect W4 independently before giving hints. Assess these outcomes: Fit mean and median constant baselines on training targets; Compare constant predictions on separate records. Record any support used.

---

**Stage:** 35 to 45 minutes

**Teacher action:** Use W 5 and the exit response to assess limits. Give targeted feedback, then the fresh retry. Save evidence of independent understanding.

---

## Reduce help deliberately

Completion task: Calculate MAE for both baselines on [4, 8].

Teacher check: Teacher explanation: Both score 2; their per-record absolute errors differ.

Start by identifying the given inputs together. Leave the intermediate steps blank. If needed, model one step, then withdraw that help on the next step. Move to a full trace only when the partial trace is understood. For partners, alternate explainer and checker; collect a response from every learner before discussion.

## Individual reasoning check

Use the worked case for Constant prediction baselines. Proposed explanation: “A final answer is enough”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Expected correction: Ask for the relevant counts, units, procedure or saved record. For this week: Mean 6; median 5.

Collect explanations, not only yes/no votes. If a misunderstanding is common, re-model the relevant step

for the class. If it affects a few learners, give a short targeted group explanation while others solve a more demanding variation.

### Pictorial practice question

Does either original baseline win under evaluation MAE? Point to the relevant step in the picture, then write the changed result and the rule or evidence that supports it.

### Reasoned teacher answer

No, both have 3 cm MAE on the supplied evaluation set; do not claim a strict winner from the training statistic alone. Changed-case check: Both fitted constants become 5 cm and both evaluation MAEs remain 3 cm. Mean is now  $20/4=5$ ; median stays 5. The earlier training outlier affected the mean.

Changed-case reasoning: Both fitted constants become 5 cm and both evaluation MAEs remain 3 cm. Mean is now  $20/4=5$ ; median stays 5. The earlier training outlier affected the mean.

Collect each learner's explanation before discussion. Use the first missing connection between input, deciding step and result to choose support. This question is worked-case practice, not fresh mastery evidence.

## 4. Run the weekly evidence lab

### Run the weekly evidence lab

Keep the test conditions fixed, record every result, and change only one planned factor.

1. Write the hypothesis and expected evidence before running.
2. Baseline Tournament: compute mean and median, predict every case, compare MAE, and explain the outlier's effect.
3. Run one normal, one boundary, one missing, and one shifted case.
4. Calculate the metric from raw counts or row-level errors.
5. Interpret what changed, what did not, and what remains unsafe or unknown.

### Make the lab runnable

Use the concrete case as the shared normal case and the independent question as a second case. Each pair creates a boundary or missing-information case and predicts its result before testing. Keep the expected result separate from an observed mistake.

#### Record · What to capture

**Record:** Input or evidence

**What to capture:** Exact data, instruction or claim

---

**Record:** Expected result

**What to capture:** Answer or safe fallback declared before the run

---

**Record:** Observed result

**What to capture:** Actual result, including errors

---

**Record:** Explanation

**What to capture:** Rule, calculation or evidence explaining the outcome

---

**Record:** Next test

**What to capture:** One justified change and a new prediction

---

## Diagnose and reteach

MISCONCEPTION TO REPAIR The mean is always the fairest and most scientific summary.

For guessing, return to given facts and request a trace. For vocabulary without application, use one example and one non-example. For arithmetic errors, check the setup before the calculation. Finish with a different uncoached case before recording mastery.

## Address these additional misconceptions

A final answer is enough: Ask for the relevant counts, units, procedure or saved record. For this week: Mean 6; median 5.

A stronger claim makes a stronger project: It fits using evaluation answers and makes the baseline unavailable for genuinely unseen cases.

The new case can be copied from the example: The independent task changes the evidence. Require a fresh explanation: Mean 6 and median 4. Mean errors [3, 3] MAE 3; median errors [1, 5] MAE 3.

## Match the support to the first error

### What the response shows · Next teaching action

**What the response shows:** Missing or misread facts

**Next teaching action:** Read the case aloud, mark supplied versus unknown facts, and let the learner restate it.

---

**What the response shows:** Wrong procedure

**Next teaching action:** Return to the deciding step in the worked case. Contrast it with the proposed incorrect explanation.

---

**What the response shows:** Arithmetic or implementation error

**Next teaching action:** Trace intermediate values or lines; preserve the first mismatch and retry that step.

---

**What the response shows:** Correct result without a reason

**Next teaching action:** Ask for a trace and a changed case before marking independent understanding.

---

**What the response shows:** An unsupported wider claim

**Next teaching action:** Ask which supplied evidence supports the claim and which further observation would be needed.

---

Offer an oral explanation, drawing, enlarged visual or fewer examples when that removes a reading/writing barrier. Keep the same reasoning goal. Stretch secure learners by changing a boundary or assumption and asking them to defend the effect, not simply by assigning more of the same questions.

## 5. Assess and explain the answers

### Assess and explain the answers

#### Worked case answer

Mean=6, median=4. Mean errors 4,2,6 total12; median errors 2,0,8 total10. Median minimises absolute error here.

#### Independent question

For training heights 2, 3 and 10 cm, calculate mean and median. Which constant is less influenced by the high value?

#### Expected independent answer

Mean is  $15/3 = 5$  cm; median is 3 cm. The median is less affected by the high value. Fit a baseline from training data only.

#### What counts as evidence

## Criterion · 0 points · 1 point · 2 points

### Criterion: Result

**0 points:** Incorrect or absent

**1 point:** Partly correct

**2 points:** Correct with necessary conditions

---

### Criterion: Reasoning

**0 points:** No supporting trace

**1 point:** Some steps

**2 points:** Clear mechanism, calculation or source support

---

### Criterion: Limit

**0 points:** Overclaims certainty

**1 point:** Vague limitation

**2 points:** Relevant uncertainty or missing fact

---

### Criterion: Transfer

**0 points:** Copies without adapting

**1 point:** Adapts with prompts

**2 points:** Solves a changed case independently

---

Use 6/8 as a classroom checkpoint, with 2 required for both Result and Reasoning. This is a proposed teaching rule, not a validated certification threshold. Do not count a coached answer as independent mastery; correct unsafe or unsupported claims regardless of total.

## Feedback

Ask the learner to find the evidence that challenges this claim: “Mean and median must always be equal.” Keep the first attempt and an uncoached retry separate.

## Original transfer answer

Mean=5, median=3. Mean absolute-error total=4+2+6=12; median total=2+0+8=10. These are training results. Fit the constant using training targets and measure final performance on untouched test targets.

## Explanations for the additional workbook examples

### W1 Fit two constants

Find the mean and median of the worked training targets.

Teacher explanation: Mean 6; median 5.

For the mean baseline, sum the worked training targets and divide by their count to obtain 6. For the median baseline, sort the training targets and take the middle value, 5. Freeze these choices before examining evaluation targets; choosing a new constant from evaluation outcomes would contaminate the comparison.

### W2 Score separately

Calculate MAE for both baselines on [4, 8].

Teacher explanation: Both score 2; their per-record absolute errors differ.

For constant 6, absolute errors on targets 4 and 8 are 2 and 2, so MAE is  $(2+2)/2 = 2$ . For constant 5, the errors are 1 and 3, so MAE is  $(1+3)/2 = 2$ . Equal averages can hide different per-record errors; retain the error table as well as the summary.

### W3 Inspect an extreme value

Replace 12 by 40. What changes?

Teacher explanation: Mean becomes 13; median stays 5. Investigate validity before deciding treatment.

### W4 Apply independently

Training [1, 3, 5, 15], evaluation [3, 9]. Fit both constants and calculate MAE.

Teacher explanation: Mean 6 and median 4. Mean errors [3, 3] MAE 3; median errors [1, 5] MAE 3.

### W5 Prevent leakage

Why not predict the evaluation mean as the baseline?

Teacher explanation: It fits using evaluation answers and makes the baseline unavailable for genuinely unseen cases.

### Assess independent transfer

Selected case: Training [1, 3, 5, 15], evaluation [3, 9]. Fit both constants and calculate MAE.

### Selected case answer

Teacher explanation: Mean 6 and median 4. Mean errors [3, 3] MAE 3; median errors [1, 5] MAE 3.

## Dimension · Evidence required

**Dimension:** Explanation

**Evidence required:** Names the applicable idea or procedure and ties it to the supplied facts.

---

**Dimension:** Trace and result

**Evidence required:** Preserves the given inputs and shows the relevant intermediate steps; distinguishes a prediction from an observed execution.

---

**Dimension:** Limits

**Evidence required:** States the assumptions, missing evidence or conditions under which the result would change.

---

Record each dimension as not yet, with support, or independent. Accept a valid alternative procedure with a defensible trace. Do not penalise an honestly recorded model failure if the learner correctly explains its evidence and limits.

## Feedback that the learner uses

Name the earliest unsupported step and give one action to repair it. Have the learner make that repair now and explain why. Then assess the relevant skill on a fresh, fully specified case. Confirm it was not previously practised with feedback; solve and check the new case before assigning it. A follow-up design prompt is a starting brief, not automatically an equivalent test.

# 6. Connect and extend learning

## Connect and extend learning

### Responsible Plant Classifier

Document any numeric baseline used in Plant Classifier feature checks, including unit, distribution, and outlier decision.

Keep the decision, input, result, explanation and remaining limitation in the project evidence record. Retrieve this record next week rather than restarting the project.

### Support and stretch

Use a tiny labelled dataset, explicit units and a named positive class. Stretch: require a baseline, held-out evaluation and subgroup or shifted-condition analysis with denominators.

## Exit reflection

Ask for one decision the learner can now justify and one situation requiring more evidence or human review. Confidence ratings are separate from correct independent performance.

## Bridge to the next week

Next: Week 21: Absolute Error and MAE. Retrieve today's evidence and identify what carries forward and what is new.

## Support and fresh reassessment

Provide a blank evidence table and allow a verbal explanation before writing. Highlight the given conditions without supplying the W 4 conclusion. Use constant baseline and training mean in full sentences.

Extension: alter one condition in the worked example and explain which conclusion changes and which remains supported.

For independent W 4, require: Mean 6 and median 4. Mean errors [3, 3] MAE 3; median errors [1, 5] MAE 3. Fresh retry training [2, 2, 8], evaluation [1, 7]: mean 4 median 2; MAE 3 for each. Require training-only fitting. Mark each criterion as independent, supported or not yet demonstrated. Reassess the reasoning after feedback; do not replace the first attempt silently.

Next week: Absolute and squared loss. Carry forward the saved evidence and the distinction between development choices and justified claims.

## Plan the delayed return

Suggested return points in this level: ai-4/21, ai-4/23, ai-4/26. Use actual classroom dates, adjusting for holidays and gaps. One, three and six teaching weeks are scheduling choices, not a claimed optimal spacing.

At a delayed check, ask for the idea from memory and a changed application. Compare with the earlier independent response and record support used. For a new level, revisit the previous capstone before checking the new prerequisites.

### Record · Teacher evidence

**Record:** Learner code and date

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

**Record:** Concept and case used; prior exposure

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

**Record:** Explanation / trace / limits

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

**Record:** Support and first repair

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

**Record:** Changed case and delayed outcome

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

**Record:** Next teaching action

**Teacher evidence:** \_\_\_\_\_

\_\_\_\_\_

Use the evidence to choose reteaching or the next step. A completed page, confidence rating or attractive project does not by itself demonstrate understanding.

### Project evidence from this case

For your Responsible Plant Classifier, save the input, procedure, observed result and limitation of this check. Label this worked example as practice; use a different, previously unreviewed case when checking independent understanding.

## 7. Prepare, teach and assess

Teaching target: Fit mean and median constant baselines on training targets

Prepare the labelled worked-case picture, the matching workbook, fictional input cards and a place for individual responses. Check a local Python 3 environment before class; have a paper-trace version ready.

### Session · Plan · Resources

**Session:** Session 1 · Understand · 45 min

**Plan:** 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.

**Resources:** Sections 1–6; record the brief recall in the workbook.

\_\_\_\_\_

**Session:** Session 2 · Practise · 45 min

**Plan:** 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation;

40–45: prepare the investigation.

**Resources:** Sections 7–10; use the matching workbook W2 and misconception check.

---

**Session:** Session 3 · Investigate and transfer · 45 min

**Plan:** 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.

**Resources:** Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

---

Before class, solve the supported practice: Calculate MAE for both baselines on [4, 8].

Reasoned practice answer: Teacher explanation: Both score 2; their per-record absolute errors differ.

Check every learner before discussion: Use the worked case for Constant prediction baselines. Proposed explanation: “A final answer is enough”. Decide whether this explanation is supported. Point to the step or supplied fact that decides; explain your repair if needed.

Misconception repair: Ask for the relevant counts, units, procedure or saved record. For this week: Mean 6; median 5.

Support: read the exact case aloud, enlarge the labelled picture, supply input cards and model only the first deciding step. Accept a spoken explanation or drawing; retain the reasoning goal.

Extension: Replace training 20 with 8 while keeping evaluation unchanged. Require a prediction and a reason before comparing. Expected result: Both fitted constants become 5 cm and both evaluation MAEs remain 3 cm. Reason: Mean is now  $20/4=5$ ; median stays 5. The earlier training outlier affected the mean.

Independent assessment: withhold the reserved answer until the learner has attempted the new case. Record support separately from correctness.

**Criterion · Not yet · With support · Independent**

**Criterion:** Explanation

**Not yet:** Names an answer without a deciding fact

**With support:** Uses a hint to link fact and decision

**Independent:** Explains the deciding fact in own words

---

**Criterion:** Trace or test

**Not yet:** Cannot connect input and result

**With support:** Completes steps with prompts

**Independent:** Traces or tests a changed case accurately

---

**Criterion:** Limits and revision

**Not yet:** Claims more than evidence supports

**With support:** Names a limit after a prompt

**Independent:** States a defensible limit and useful next test

---

Project checkpoint: Evaluate and revise. Compare with your starting method, explain errors and record a justified revision.

Use the weekly teacher answer for topic-specific reasoning. A saved progress tick is an activity record; judge mastery from the learner's explanation and evidence.