Sampling variability

Learn to understand, design, build, test and explain AI systems responsibly.

Project: Reproducible AI Research Study

TEACHER GUIDE · 1 OF 9

Prepare the purpose, prerequisites and materials

> Prepare Sampling variability: check the target, prerequisites, resources and adaptable teaching schedule.

Teach the learning cycle

Primary target: Explain the **mechanism** and terminology of **sampling variability**

Session · Sequence and minutes · Resources

> **Session:** Session 1 · Understand · 45 min
>
> **Sequence and minutes:** 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.
>
> **Resources:** Sections 1–6; record the brief recall in the workbook.

> **Session:** Session 2 · Practise · 45 min
>
> **Sequence and minutes:** 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.
>
> **Resources:** Sections 7–10; use the matching workbook W2 and misconception check.

> **Session:** Session 3 · Investigate and transfer · 45 min

> **Sequence and minutes:** 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.
>
> **Resources:** Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

This is a starting allocation of 135 minutes, not a validated duration. Preserve all named topics. Schedule extra time for connected examples, the full investigation, independent cases and project building.

Repeated copies of a case share one recorded attempt; they are not new **evidence**.

Retrieval answer guidance

Confounders and controls

Without opening the old lesson, explain one key idea from Confounders and controls. Give an example and a mistake that the idea helps you avoid.

Look for: Explain the **mechanism** and terminology of confounders and controls

Reference example, not the only acceptable wording: Teacher explanation: A confounder changes alongside the intended factor and can offer another explanation. Controls aim to isolate the proposed **mechanism**.

For a toy comparison vary only retrieval method while holding corpus, queries, cutoff and relevance rubric fixed. Randomise or counterbalance order when order effects matter.

Baselines and fair comparisons

Without opening the old lesson, explain one key idea from Baselines and fair comparisons. Give an example and a mistake that the idea helps you avoid.

Look for: Explain the **mechanism** and terminology of baselines and fair comparisons

Reference example, not the only acceptable wording: Teacher explanation: Fair baselines use credible procedures on matched **evidence** and resource conditions. A weak arbitrary reference can exaggerate improvement.

For a room-reporting study use manual authoritative lookup or a deterministic same-room rule, not random room names. Match cases, available sources and scoring.

Replication and reproduction

Without opening the old lesson, explain one key idea from Replication and reproduction. Give an example and a mistake that the idea helps you avoid.

Look for: Explain the **mechanism** and terminology of replication and reproduction

Reference example, not the only acceptable wording: Teacher explanation: Reproducing a computation uses recorded artifacts; replication collects new **evidence** under a comparable question.

Naming conventions vary, so define the terms in the project. Here replay means same artifacts/procedure; new-data replication means newly collected eligible cases under the same plan.

Preserve environment, data versions and raw results.

Two 45-minute sessions. **Sampling variability** means different eligible samples can give different estimates even under the same population. Separate this from procedure changes.

Use the supplied constructed scenarios to make the **mechanism** inspectable. Collect W4 before sharing solutions.

Students should name intermediate **evidence** and limitations; classroom pacing and learning effectiveness remain unvalidated.

Outcomes

Explain the **mechanism** and terminology of **sampling variability**

Apply the stated procedure to a new case with a traceable result

Identify an assumption or limitation and design a revealing follow-up

Preparation

Lesson and workbook; paper and pencil

Calculator and optional local Python 3 for arithmetic; no live API or private data

Prepare the fresh retry separately and withhold its solution until after collection

Prerequisite answer

A result depends on the **input**s, procedure and conditions that produced it. Preserve data roles and distinguish observed outcomes from **prediction**s or unmeasured conditions.

If the prior concept cannot be explained, use a short retrieval activity before this chapter.

End goal for the pictorial case

Explain **sampling variability** using the records and boundaries of this fictional investigation.

This added worked case illustrates the week's main idea. Retain all original and connected topics; schedule its practice within the teaching cycle.

Explain and model the mechanism

> Model Sampling variability using the supplied case, labelled mechanism and reasoning. Make the scope of each conclusion explicit.

Sampling variability means different eligible samples can give different estimates even under the same population. Separate this from procedure changes.

Explain Sampling variability · 1

Begin by writing the question the procedure is intended to answer. Identify the information already supplied, the information it will produce and the information still missing.

These are constructed classroom examples; they do not require live accounts, private records or an external service. Do not treat a plausible explanation as a measured outcome.

Consider finite population values1, 1, 3, 3 with distinct IDs. Enumerate the six unordered samples of size2 **without replacement**. Their means are1, 2, 2, 2, 2, 3.


Explain Sampling variability · 2

Before applying the procedure, write its units, denominator or decision boundary where relevant. A partner should be able to repeat each step using the same **input**s.

If a required **input** is absent or violates the stated contract, record that condition explicitly instead of quietly replacing it with a convenient value.

Term	Meaning
Sampling variability	Changes in estimates across eligible samples
Sampling distribution	The distribution of an estimate under a sampling procedure
Without replacement	Selecting distinct units within a sample

Sampling variability — evidence trace



The figure connects this chapter's worked input, declared procedure and result. Read the complete worked reasoning for qualifiers and intermediate steps; the abbreviated labels do not replace that reasoning.

Explain Sampling variability · 3

Guided: Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ. Qualification: This finite example is not a general normality guarantee or a model of a biased convenience sample.

Exit: accept a concrete variation with its predicted outcome and a stated condition that would challenge the **mechanism**; reject a claim resting on unmeasured **evidence**.

Model the reasoning aloud

Use the complete worked example below before asking learners to complete W2. Reveal a small part, invite a **prediction**, then explain the deciding step.

These teacher notes contain solutions; do not reveal the independent case answer during modelling.

Find the mean of sample means and the chance of mean2 under uniform selection of those six samples.

Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ.

Model the reasoning aloud · 4

Compare the intermediate **evidence** with the final statement. Explain why each number or decision follows from the stated **input**s.

The conclusion belongs to this scenario and procedure; changing the **input**s, evaluation population or authority can change what it establishes. Retain the raw record so a later revision can be compared honestly.

Model the reasoning aloud · 5

Optional local lab: predict the printed result before running this standalone Python 3 example. Change one relevant boundary **input**, record the outcome and state the contract or statistical assumptions.

The program uses fictional values and the Python standard library; it performs no external operations.

```
from itertools import combinations

def sample_means(population, size):
    if type(size) is not int or not 1 <= size <= len(population):
        raise ValueError('sample size')
    return [sum((population[i] for i in indices)) / size for indices in
combinations(range(len(population)), size)]

print(sample_means([1, 1, 3, 3], 2))
```

Thinking move	Teacher prompt
Plan	What is given? What is the question? Which procedure fits these facts?
Do	Point to each input and intermediate step in the worked trace. Name the rule that allows the next step.
Check	Does the output answer the question? Which assumption or limitation prevents a larger claim?

Use the diagram and verbal explanation together. Ask learners to match a sentence to a specific visual step; do not assign learners fixed visual or auditory learning styles.

A pictorial worked case: Sampling variability

Two small samples can differ. A fixed classifier is scored on two distinct four-case samples from a declared collection.


The facts we will use

> **Supplied facts**
>
> Sample A has 3 correct of 4; sample B has 2 correct of 4. The model and rubric are unchanged.

AI ACADEMY / AI-7/11

Two distinct samples, fixed model

correct cases out of 4 in each sample



Bar length shows the supplied value; the marked baseline is zero.

Follow the labelled picture with the matching explanation. Open the vector diagram

AI ACADEMY / AI-7/11

Sampling variability

Start with the supplied facts

Follow each arrow. Ask which fact allows the next step.



1. Keep model fixed

No training or threshold change occurs between samples.



2. Compute observed rates

A is 75%; B is 50%, differing by 25 percentage points.



3. Consider sampling

Different included cases can change a small-sample rate.

CHANGE ONE THING

Evaluate the exact same fixed four cases twice with deterministic predictions.

Explain the changed result using the matching lesson.

Follow the labelled picture with the matching explanation. Open the vector diagram

Read the picture one step at a time

STEP	WHAT HAPPENS AND WHY
1. Keep model fixed	No training or threshold change occurs between samples.
2. Compute observed rates	A is 75%; B is 50%, differing by 25 percentage points.
3. Consider sampling	Different included cases can change a small-sample rate.

Match each pictured step to the supplied facts. This is a worked teaching case, not an independent assessment.

Why this result follows

Why the result follows

Sampling variability means a statistic can vary because different units were observed. Do not automatically attribute every rate change to a model improvement or deterioration.

What this example does not establish

Keep the stated limit

These two samples do not supply a probability model or confidence interval; none is invented.

Change one thing in the pictured case

Change one thing

Change: Evaluate the exact same fixed four cases twice with deterministic **prediction**s.

Predicted result

Prediction: The counts should agree unless some condition, data or scoring changed.

Why the result follows

Reason: The sampling explanation no longer applies when the records are identical.

Ask while showing the picture

Ask while showing the picture · 6

Cover the result. Ask learners to name the **input**s and predict the next step. Uncover one step, connect the icon to the actual procedure, and explain its **evidence**. Then change the supplied condition and compare.

The picture is a representation; it does not document execution of a real AI model.

Demonstrate and guide practice

> Demonstrate Sampling variability, reduce support deliberately and check individual reasoning before discussing answers.

Time or stage	Teacher action
Session 1: 0–7 minutes	Retrieve the prerequisite; identify the input, procedure and evidence source.
7–18 minutes	Explain section 1 and vocabulary. Complete W1 with a mechanism-based explanation.
18–32 minutes	Trace the declared procedure and worked case. Complete W2, recording intermediate steps.
32–45 minutes	Discuss the diagram with the worked input; use W3 to distinguish measured claims and limitations.
Session 2: 0–8 minutes	Retrieve the worked reasoning and collect the guided response before discussing it.
8–20 minutes	Compare a proposed boundary variant with the original case; identify what remains fixed.
20–35 minutes	Collect W4 independently. Require the procedure trace and the bounded conclusion.
35–45 minutes	Use W5 and exit responses for feedback. Administer the fresh retry after feedback and preserve both attempts.

Use the worked example in Sections 3 and 6. Collect a **prediction** before revealing its result; ask the learner to point to each intermediate fact.

Provide an **input**/procedure/result/limit frame and a calculator when needed. Ask students to identify the first step before completing the whole trace.

Extension: compare two concrete variants of this **mechanism** while holding unrelated conditions fixed; report both successful and unsuccessful outcomes.

Reduce help deliberately

Completion practice

Completion task: Find the mean of sample means and the chance of mean2 under uniform selection of those six samples. Show the intermediate steps and give the conclusion with its scope.

Reasoning to listen for

Teacher check: Teacher explanation: Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ.

Start by identifying the given **input**s together. Leave the intermediate steps blank. If needed, model one step, then withdraw that help on the next step. Move to a full trace only when the partial trace is understood.

For partners, alternate explainer and checker; collect a response from every learner before discussion.

Individual reasoning check

Use the worked case for **Sampling variability**. Proposed explanation: “The worked result establishes every future case”. Decide whether this explanation is supported.

Point to the step or supplied fact that decides; explain your repair if needed.

Expected repair

Expected correction: This finite example is not a general normality guarantee or a model of a biased convenience sample.

Collect explanations, not only yes/no votes. If a misunderstanding is common, re-model the relevant step for the class.

If it affects a few learners, give a short targeted group explanation while others solve a more demanding variation.

Pictorial practice question

Calculate the two four-case rates and their difference. Explain how changed included cases can affect a fixed model, and why identical deterministic records should not differ on a rerun.

Reasoned teacher answer

No **training** or threshold change occurs between samples. A is 75%; B is 50%, differing by 25 percentage points. Different included cases can change a small-sample rate.

Changed case: The counts should agree unless some condition, data or scoring changed. The sampling explanation no longer applies when the records are identical.

Changed-case reasoning: The counts should agree unless some condition, data or scoring changed. The sampling explanation no longer applies when the records are identical.

Collect each learner's explanation before discussion. Use the first missing connection between **input**, deciding step and result to choose support. This question is worked-case practice, not fresh mastery **evidence**.

TEACHER GUIDE · 4 OF 9

Run the investigation and diagnose misconceptions

> Run the Sampling variability investigation and use the first unsupported step to choose a teaching response.

The worked result establishes every future case: This finite example is not a general normality guarantee or a model of a biased convenience sample.

A missing **input** may be replaced without recording the change: A replacement changes the **evidence** or contract. Record the change and its justification; do not silently turn unknown into a measured value.

Repeating an assessed case is new independent **evidence**: A replay checks consistency. An example used for feedback or revision cannot later be described as untouched final evaluation.

Use Section 9 for the **evidence** ledger. Add a boundary or missing-**input** case and record the expected response before checking. More successful repeats of one case do not establish wider coverage.

Match the support to the first error

What the response shows	Next teaching action
Missing or misread facts	Read the case aloud, mark supplied versus unknown facts, and let the learner restate it.
Wrong procedure	Return to the deciding step in the worked case. Contrast it with the proposed incorrect explanation.
Arithmetic or implementation error	Trace intermediate values or lines; preserve the first mismatch and retry that step.
Correct result without a reason	Ask for a trace and a changed case before marking independent understanding.
An unsupported wider claim	Ask which supplied evidence supports the claim and which further observation would be needed.

Offer an oral explanation, drawing, enlarged visual or fewer examples when that removes a reading/writing barrier. Keep the same reasoning goal.

Stretch secure learners by changing a boundary or assumption and asking them to defend the effect, not simply by assigning more of the same questions.

Assess and explain the answers

> Use the Sampling variability questions, teacher keys and criteria after the relevant attempts. Previously discussed cases are practice; protect the separately selected independent task.

Guided and exit questions

Guided: Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ. Qualification: This finite example is not a general normality guarantee or a model of a biased convenience sample.

Exit: accept a concrete variation with its predicted outcome and a stated condition that would challenge the **mechanism**; reject a claim resting on unmeasured **evidence**.

W1 Explain the mechanism

Workbook question · W1

Explain the weekly **mechanism** in your own words. Use the worked scenario to name its **input**, decision rule and **output**.

Teacher reasoning · W1

Teacher explanation: **Sampling variability** means different eligible samples can give different estimates even under the same population. Separate this from procedure changes.

Consider finite population values1, 1, 3, 3 with distinct IDs. Enumerate the six unordered samples of size2 **without replacement**. Their means are1, 2, 2, 2, 2, 3.

W2 Trace the worked case

Workbook question · W2

Find the mean of sample means and the chance of mean2 under uniform selection of those six samples. Show the intermediate steps and give the conclusion with its scope.

Teacher reasoning · W2

Teacher explanation: Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ.

W3 Audit the limit

Workbook question · W3

A report omits this qualification: This finite example is not a general normality guarantee or a model of a biased convenience sample.

Explain how omitting it could change a reader's interpretation, and write an accurate replacement.

Teacher reasoning · W3

Teacher explanation: This finite example is not a general normality guarantee or a model of a biased convenience sample.

State the observed result separately from the unverified or out-of-scope claim; preserve the original **evidence** rather than inventing a missing fact.

W4 Apply a new case independently

Workbook question · W4

Population0, 0, 4, 4 with distinct IDs; enumerate size2 sample means and their average.

Teacher reasoning · W4

Teacher explanation: Means0, 2, 2, 2, 2, 4; average2; mean2 occurs4/6. Equal values have distinct IDs and count as separate eligible units.

W5 Design a revealing follow-up

Workbook question · W5

For the worked scenario, propose one additional case that could reveal a failure, describe its expected outcome before running it, and explain what would need to remain fixed for a fair comparison.

Do not reuse W4 as a new final test after receiving feedback.

Teacher reasoning · W5

Teacher explanation: Accept a concrete variant of the worked **input** or a missing/conflicting/boundary condition, with an expectation derived from the declared procedure.

Require preserved settings, original **output**s and data roles. This finite example is not a general normality guarantee or a model of a biased convenience sample.

The new case should test this particular **mechanism** rather than merely restate the original result.

Independent assessment and fresh retry

Teacher key · Independent assessment and fresh retry

W4: Means0, 2, 2, 2, 2, 4; average2; mean2 occurs4/6. Equal values have distinct IDs and count as separate eligible units. Require intermediate reasoning and the stated scope.

Fresh retry: Population2, 2, 6, 6 under same sampling. Expected reasoning: Means2, 4, 4, 4, 4, 6; average4; mean4 probability2/3.

If students need support, record it and reassess with another new case rather than awarding independence to a copied answer.

Assess independent transfer

Independent question · withhold the key

Selected case: Population0, 0, 4, 4 with distinct IDs; enumerate size2 sample means and their average.

Selected case answer

Teacher key · Selected case answer

Teacher explanation: Means0, 2, 2, 2, 2, 4; average2; mean2 occurs4/6. Equal values have distinct IDs and count as separate eligible units.

Dimension	Evidence required
Explanation	Names the applicable idea or procedure and ties it to the supplied facts.
Trace and result	Preserves the given inputs and shows the relevant intermediate steps; distinguishes a prediction from an observed execution.
Limits	States the assumptions, missing evidence or conditions under which the result would change.

Teacher key · Selected case answer

Record each dimension as not yet, with support, or independent. Accept a valid alternative procedure with a defensible trace.

Do not penalise an honestly recorded model failure if the learner correctly explains its **evidence** and limits.

Feedback that the learner uses

Name the earliest unsupported step and give one action to repair it. Have the learner make that repair now and explain why. Then assess the relevant skill on a fresh, fully specified case.

Confirm it was not previously practised with feedback; solve and check the new case before assigning it. A follow-up design prompt is a starting brief, not automatically an equivalent test.

TEACHER GUIDE · 6 OF 9

Connect the project, support and next lesson

> Connect Sampling variability to Reproducible AI Research Study, adapt support and plan a later changed-case check.

Add a labelled **sampling variability evidence** sheet to Reproducible AI Research Study. Show how you can explain the **mechanism** and terminology of **sampling variability**.

Use the supplied fictional case and keep its **input**s, intermediate steps, calculation or **prediction**, and limitation. Connect this record to last week's record; a matching answer without a trace is insufficient.

Provide an **input**/procedure/result/limit frame and a calculator when needed. Ask students to identify the first step before completing the whole trace.

Extension: compare two concrete variants of this **mechanism** while holding unrelated conditions fixed; report both successful and unsuccessful outcomes.

Next connection: Uncertainty intervals. Preserve the current case record and retrieve its **evidence** boundary before moving on.

Retain the project ledger and assess a changed case after support. Check the relevant prerequisite before carrying this component forward.

Plan the delayed return

Suggested return points in this level: ai-7/12, ai-7/14, ai-7/17. Use actual classroom dates, adjusting for holidays and gaps. One, three and six teaching weeks are scheduling choices, not a claimed optimal spacing.

At a delayed check, ask for the idea from memory and a changed application. Compare with the earlier independent response and record support used.

For a new level, revisit the previous capstone before checking the new prerequisites.

Record	Teacher evidence
Learner code and date	
Concept and case used; prior exposure	
Explanation / trace / limits	
Support and first repair	
Changed case and delayed outcome	
Next teaching action	

Use the **evidence** to choose reteaching or the next step. A completed page, confidence rating or attractive project does not by itself demonstrate understanding.

Project evidence from this case

For the Reproducible AI Research Study, retain an annotated worked trace and the changed-case reasoning. Connect the claim to its records, units and assumptions so another investigator can challenge or reproduce it.

Prepare, teach and assess

> Use this preparation summary alongside the detailed Sampling variability lesson; it is a checklist rather than a replacement for the full sequence.

Teaching target

Teaching target: Explain the **mechanism** and terminology of **sampling variability**

Prepare the labelled worked-case picture, the matching workbook, fictional **input** cards and a place for individual responses. Check a local Python 3 environment before class; have a paper-trace version ready.

Session · Plan · Resources

> **Session:** Session 1 · Understand · 45 min
>
> **Plan:** 0–7: recall and prerequisite; 7–15: inspect the concrete case and predict; 15–30: teacher models the worked trace; 30–40: learners explain a step using the text and visual; 40–45: check the core idea.
>
> **Resources:** Sections 1–6; record the brief recall in the workbook.

> **Session:** Session 2 · Practise · 45 min
>
> **Plan:** 0–5: retrieve the procedure; 5–15: W2 completion practice; 15–23: explain and check with a partner; 23–30: each learner answers the misconception check; 30–40: targeted reteaching or a harder variation; 40–45: prepare the investigation.
>
> **Resources:** Sections 7–10; use the matching workbook W2 and misconception check.

> **Session:** Session 3 · Investigate and transfer · 45 min
>
> **Plan:** 0–5: retrieve the decision rule; 5–20: investigate or trace; 20–30: start the independent case; 30–38: connect the project; 38–43: act on feedback; 43–45: plan the return check.
>
> **Resources:** Sections 9 and 11–14. Give extra time to finish the full task set; use Section 15 when the week includes a coding lab.

Before class, solve the supported practice: Find the mean of sample means and the chance of mean2 under uniform selection of those six samples. Show the intermediate steps and give the conclusion with its scope.

Supported practice answer

Reasoned practice answer: Teacher explanation: Mean=(1+2+2+2+2+3)/6=2; mean2 occurs4/6=2/3. The population mean is2, while individual samples differ.

Check every learner before discussion: Use the worked case for **Sampling variability**. Proposed explanation: “The worked result establishes every future case”. Decide whether this explanation is supported.

Point to the step or supplied fact that decides; explain your repair if needed.

Repair the misconception

Misconception repair: This finite example is not a general normality guarantee or a model of a biased convenience sample.

Support without giving away the answer

Support: read the exact case aloud, enlarge the labelled picture, supply **input** cards and model only the first deciding step. Accept a spoken explanation or drawing; retain the reasoning goal.

Extend the reasoning

Extension: Evaluate the exact same fixed four cases twice with deterministic **prediction**s. Require a **prediction** and a reason before comparing.

Expected result: The counts should agree unless some condition, data or scoring changed. Reason: The sampling explanation no longer applies when the records are identical.

Protect the independent attempt

Independent assessment: withhold the reserved answer until the learner has attempted the new case. Record support separately from correctness.

Criterion	Not yet	With support	Independent
Explanation	Names an answer without a deciding fact	Uses a hint to link fact and decision	Explains the deciding fact in own words

Criterion	Not yet	With support	Independent
Trace or test	Cannot connect input and result	Completes steps with prompts	Traces or tests a changed case accurately
Limits and revision	Claims more than evidence supports	Names a limit after a prompt	States a defensible limit and useful next test

Project checkpoint

Project checkpoint: Make a transparent first version. Keep a trace of how one **input** becomes an **output**; explain the rule or learned component.

Use the weekly teacher answer for topic-specific reasoning. A saved progress tick is an activity record; judge mastery from the learner's explanation and **evidence**.

TEACHER GUIDE · 8 OF 9

Use feedback, labs and repair paths

> Use feedback, earlier review and portfolios purposefully. Apply the coding routine only when this lesson has an executable lab.

Use the two formative questions as supported practice, not as independent mastery scores. Discuss the deciding **evidence** rather than accepting a guessed selection.

The feedback comes from the reviewed public worked case. Do not reveal the separately reserved independent teacher answer until the learner has attempted that case.

Ask students to predict before running code. Preserve errors and actual **output**s; collect a changed test and an explanation. The branch-boundary and indexing lessons deliberately include failure cases.

Use the earlier worked-explanation links to address a specific missing building block, then return to the current question. Adapt this suggestion after observing the learner; it is not an automatic diagnosis.


Collect the completed-work portfolio as **evidence**. Record support separately from correctness, and assess explanation, trace/test and limitations. Progress ticks and quiz selections alone do not establish mastery.

Visual concept journey

> Show one Sampling variability graphic at a time, request a prediction and discuss its trace before the independent assessment.

Use one picture at a time. Hide a worked result before asking for a **prediction**, then reveal it for feedback. These are supported teaching cases; retain the separate independent assessment for a fresh first attempt.

01. Sampling variability: the case



Follow the labelled picture with the matching explanation. Open the vector diagram

Two small samples can differ. A fixed classifier is scored on two distinct four-case samples from a declared collection. Supplied facts: Sample A has 3 correct of 4; sample B has 2 correct of 4.


The model and rubric are unchanged.

Point to the given fact that matters most. Explain why it matters.

Discuss while showing the picture

Teacher prompt: Point to the given fact that matters most. Explain why it matters. Ask learners to describe one label or arrow in ordinary words. Use the adjacent transcript if the image is difficult to see.

02. Follow the deciding steps



Follow the labelled picture with the matching explanation. Open the vector diagram


Keep model fixed: No **training** or threshold change occurs between samples. → Compute observed rates: A is 75%; B is 50%, differing by 25 percentage points. → Consider sampling: Different included cases can change a small-sample rate.

Point to each numbered stage and explain which recorded **input** it uses.

Discuss while showing the picture

Teacher prompt: Point to each numbered stage and explain which recorded **input** it uses. Ask learners to describe one label or arrow in ordinary words. Use the adjacent transcript if the image is difficult to see.

03. Two distinct samples, fixed model



Follow the labelled picture with the matching explanation. Open the vector diagram

Two distinct samples, fixed model. Units: correct cases out of 4 in each sample. Sample A: 3. Sample B: 2.


Read the labels and values. Explain the relationship the model shows.

Discuss while showing the picture

Teacher prompt: Read the labels and values. Explain the relationship the model shows. Ask learners to describe one label or arrow in ordinary words. Use the adjacent transcript if the image is difficult to see.

Check row/column meanings, units and denominators before discussing a difference. Supplied toy values are not real class or model results.

04. Change one thing. Trace what follows.



Follow the labelled picture with the matching explanation. Open the vector diagram

Original case: Different included cases can change a small-sample rate. Changed condition: Evaluate the exact same fixed four cases twice with deterministic **prediction**s.

Expected result: The counts should agree unless some condition, data or scoring changed. Reason: The sampling explanation no longer applies when the records are identical.


Predict before checking the worked changed-case result. Name what changed and what the supplied facts allow you to conclude.

Discuss while showing the picture

Teacher prompt: Predict before checking the worked changed-case result. Name what changed and what the supplied facts allow you to conclude. Ask learners to describe one label or arrow in ordinary words.

Use the adjacent transcript if the image is difficult to see. Changed-case reasoning: The sampling explanation no longer applies when the records are identical.

05. Connect the explanation to its limit



Follow the labelled picture with the matching explanation. Open the vector diagram

Sampling variability means a statistic can vary because different units were observed. Do not automatically attribute every rate change to a model improvement or deterioration.

Limit: These two samples do not supply a probability model or confidence interval; none is invented.


Explain the case, then point to a limit on the claim.

Discuss while showing the picture

Teacher prompt: Explain the case, then point to a limit on the claim. Ask learners to describe one label or arrow in ordinary words. Use the adjacent transcript if the image is difficult to see.

Keep the stated scope: These two samples do not supply a probability model or confidence interval; none is invented.

06. Make the idea visible in your project



Follow the labelled picture with the matching explanation. Open the vector diagram

For the Reproducible AI Research Study, retain an annotated worked trace and the changed-case reasoning. Connect the claim to its records, units and assumptions so another investigator can challenge or reproduce it.

Draw your **input**, deciding step, result and limit. Supported question: Calculate the two four-case rates and their difference.

Explain how changed included cases can affect a fixed model, and why identical deterministic records should not differ on a rerun.

Draw your project trace and justify the deciding step using **evidence**.

Discuss while showing the picture

Teacher prompt: Draw your project trace and justify the deciding step using **evidence**. Ask learners to describe one label or arrow in ordinary words. Use the adjacent transcript if the image is difficult to see.

Supported practice key: No **training** or threshold change occurs between samples. A is 75%; B is 50%, differing by 25 percentage points. Different included cases can change a small-sample rate.

Changed case: The counts should agree unless some condition, data or scoring changed. The sampling explanation no longer applies when the records are identical.

Save or print your responses for your portfolio. Progress and drafts on the hosted website stay in the current browser. Use only fictional or teacher-approved data.