

A probability-free proof of an explicit CrooŁ-Laba-Sisask Lemma

Olivier Ramaré

December 14, 2022

Abstract

File CrooŁ-Laba-Sisask-Lemma-02.tex This note proposes a proof free of probabilistic language of the famous CrooŁ-Laba-Sisask Lemma. In between, we do the same for the Khintchine and Marcinkiewicz-Zygmund inequalities and explicitate the implied constants.

1 Introduction

We propose a proof free of probabilistic language of (a variant of) [2, Lemma 3.2] by E. CrooŁ, I. Łaba and O. Sisask, which we now state. We follow their notation and in particular, when z is a complex number, z° is defined to be $z/|z|$ when $z \neq 0$, and to be 0 when $z = 0$.

Theorem 1.1. *Let (X, μ) be a probability space and $p \geq 2$. Given a function f in the form*

{CLS}

$$f = \sum_{k \leq K} \lambda_k g_k$$

where $(g_k)_k$ is a collection of measurable functions on X of $L^p(\mu)$ -norm at most 1. Let $\varepsilon > 0$. There exists an L -tuple $(k_1, \dots, k_L) \in \{1, \dots, K\}^L$ of length $L \leq 20p/\varepsilon^2$ such that

$$\int_X \left| \frac{f(x)}{\|\lambda\|_1} - \frac{1}{L} \sum_{\ell \leq L} \lambda_{k_\ell}^\circ g_{k_\ell}(x) \right|^p d\mu \leq \varepsilon^p,$$

where $\|\lambda\|_1 = \sum_{k \leq K} |\lambda_k|$.

This theorem has its origin in the paper [3] by E. CrooŁ and O. Sisask. We refer to the paper [6] by L. Pierce for much deeper background on the Khintchine and Marcinkiewicz-Zygmund inequalities.

[2] E. CrooŁ, I. Łaba, and O. Sisask, 2013, ‘‘Arithmetic progressions in sumsets and L^p -almost-periodicity’’.

[3] E. CrooŁ and O. Sisask, 2010, ‘‘A probabilistic technique for finding almost-periods of convolutions’’.

[6] L. B. Pierce, 2021, ‘‘On superorthogonality’’.

2 An upper explicit Khintchine Inequality

{KI}

Theorem 2.1. *We have, when $p \geq 1$,*

$$(1/2^N) \sum_{(\varepsilon_n) \in \{\pm 1\}^N} \left| \sum_{n \leq N} c_n \varepsilon_n \right|^p \leq p^{p/2} \left(\sum_{n \leq N} |c_n|^2 \right)^{p/2}.$$

This is only half of the Khintchine Inequality and in a special context, but this will be enough for us. We followed [1, Chapter 10, Theorem 1, page 354] by Y.S. Chow and H. Teicher.

Proof. Let us start with $p = 2k \geq 2$, so that we may open the inner sum and get

$$\begin{aligned} 2^N S(2k) &= \sum_{(\varepsilon_n) \in \{\pm 1\}^N} \left| \sum_{n \leq N} c_n \varepsilon_n \right|^p \\ &= \sum_{\substack{s_1 + s_2 + \dots + s_N = 2k, \\ s_n \geq 0}} \binom{2k}{s_1, s_2, \dots, s_N} \prod_{1 \leq n \leq N} c_n^{s_n} \sum_{(\varepsilon_n) \in \{\pm 1\}^N} \prod_{n \leq N} \varepsilon_n^{s_n} \end{aligned}$$

by the multinomial theorem. The inner summand vanishes as soon as some s_n is odd, whence, by letting $2t_n = s_n$, we get

$$\begin{aligned} 2^N S(2k) &= \sum_{\substack{t_1 + t_2 + \dots + t_N = k, \\ t_n \geq 0}} \binom{2k}{2t_1, 2t_2, \dots, 2t_N} \prod_{n \leq N} (c_n^2)^{t_n} \\ &\leq C \sum_{\substack{t_1 + t_2 + \dots + t_N = k, \\ t_n \geq 0}} \binom{k}{t_1, t_2, \dots, t_N} \prod_{n \leq N} (c_n^2)^{t_n} = C \left(\sum_{n \leq N} c_n^2 \right)^k \end{aligned}$$

where

$$\begin{aligned} C &= \max \binom{2k}{2t_1, 2t_2, \dots, 2t_N} \binom{k}{t_1, t_2, \dots, t_N}^{-1} \\ &\leq \max \frac{2k(2k-1) \dots (k+1)}{\prod_j 2t_j(2t_j-1) \dots (t_j+1)} \\ &\leq \max \frac{k^k}{2^{t_1 + \dots + t_k}} \leq (k/2)^k. \end{aligned}$$

As $S(p)$ is increasing, we simply choose $k = \lceil p/2 \rceil$. This gives us $2k \geq p + 2$ and thus

$$(k/2)^k \leq (p+2)^{1+p/2} \leq (30p)^{p/2}.$$

This concludes the main part of the proof, except for the constant 30. We will not continue the proof but simply refer to the paper [5] by U. Haagerup who shows that best constant is (be careful: the abstract of this paper misses a closing parenthesis for the value of B_p , but the value of B_p displayed in the

[1] Y. S. Chow and H. Teicher, 1997, *Probability theory*.

[5] U. Haagerup, 1981, "The best constants in the Khintchine inequality".

middle of page 232 misses a squareroot-sign around the π , as an inspection of the proof at the end the paper rapidly reveals)

$$\begin{cases} 1 & \text{when } 0 < p \leq 2, \\ \sqrt{2} \left(\frac{\Gamma((p+1)/2)}{\sqrt{\pi}} \right)^{1/p} & \text{when } 2 < p. \end{cases}$$

We readily check that this implies that the constant 1 rather than 30 is admissible. \square

3 An upper explicit Marcinkiewicz-Zygmund Inequality

Theorem 3.1. *Let (X, μ) be a probability space. When $p \geq 1$, let $(f_n)_{n \leq N}$ be a system of functions such that $\int_X f_n(x) d\mu = 0$. We have*

{MZI}

$$\begin{aligned} \int_{(x_n) \in X^N} \left| \sum_{1 \leq n \leq N} f_n(x_n) \right|^p d(x_n) \\ \leq (4p)^{p/2} \int_{(x_n) \in X^N} \left(\sum_{1 \leq n \leq N} |f_n(x_n)|^2 \right)^{p/2} d(x_n). \end{aligned}$$

The power of this inequality is that the implied constant do not depend on N , the effect of some orthogonality. Again, this is only half of the Marcinkiewicz-Zygmund Inequality and in a special context, but this will be enough for us. We followed [1, Chapter 10, Theorem 2, page 356] by Y.S. Show and H. Teicher. The relevant constant is the subject of [7] by Y.-F. Ren and H.-Y. Liang (their value is slightly worse than ours) and [4] by D. Ferger, where the best constant is determined provided the f_n 's are "symmetric".

Proof. We first notice that, since $\int_0^1 f_n(x) dx = 0$, we may introduce a symmetrization through

$$\sum_{n \leq N} f_n(x_{2n-1}) = - \int_{(x_{2n}) \in X^N} \sum_{n \leq 2N} (-1)^n f_{[n/2]}(x_n) d(x_{2n}).$$

Jensen's inequality gives us that

$$\begin{aligned} \int_{(x_{2n-1}) \in X^N} \left| \int_{(x_{2n}) \in X^N} \sum_{n \leq 2N} (-1)^n f_{[n/2]}(x_n) d(x_{2n}) \right|^p d(x_{2n-1}) \\ \leq \int_{(x_{2n-1}) \in X^N} \int_{(x_{2n}) \in X^N} \left| \sum_{n \leq 2N} (-1)^n f_{[n/2]}(x_n) \right|^p d(x_{2n}) d(x_{2n-1}) \end{aligned}$$

from which we deduce that the L^p -norm of the symmetrization controls the one of the initial sum:

$$\int_{(x_{2n-1}) \in X^N} \left| \sum_{n \leq N} f_n(x_{2n-1}) \right|^p d(x_{2n-1}) \leq \int_{(x_n) \in X^{2N}} \left| \sum_{n \leq 2N} (-1)^n f_{[n/2]}(x_n) \right|^p d(x_n).$$

[7] Y.-F. Ren and H.-Y. Liang, 2001, "On the best constant in Marcinkiewicz-Zygmund inequality".

[4] D. Ferger, 2014, "Optimal constants in the Marcinkiewicz-Zygmund inequalities".

We next introduce Rademacher's system by noticing that, by successively exchanging x_{2n-1} and x_{2n} , have

$$\begin{aligned} (1/2^N) \sum_{(\varepsilon_n) \in \{\pm 1\}^N} \int_{(x_n) \in X^{2N}} \left| \sum_{n \leq 2N} \varepsilon_{[n/2]} (-1)^n f_{[n/2]}(x_n) \right|^p d(x_n) \\ = \int_{(x_n) \in X^{2N}} \left| \sum_{n \leq 2N} (-1)^n f_{[n/2]}(x_n) \right|^p d(x_n). \end{aligned}$$

We may now remove the symmetrization since:

$$\begin{aligned} \int_{(x_n) \in X^{2N}} \left| \sum_{n \leq 2N} \varepsilon_{[n/2]} (-1)^n f_{[n/2]}(x_n) \right|^p d(x_n) \\ \leq \int_{(x_n) \in X^{2N}} 2^{p-1} \left(\left| \sum_{n \leq N} \varepsilon_n f_n(x_{2n}) \right|^p + \left| \sum_{n \leq N} \varepsilon_n f_n(x_{2n-1}) \right|^p \right) d(x_n) \\ \leq 2^p \int_{(x_{2n}) \in X^N} \left| \sum_{n \leq N} \varepsilon_n f_n(x_{2n}) \right|^p d(x_{2n}). \end{aligned}$$

The Khintchine Inequality from Theorem 2.1 now gives us that

$$(1/2^N) \sum_{(\varepsilon_n) \in \{\pm 1\}^N} \left| \sum_{n \leq N} \varepsilon_n f_n(x_{2n}) \right|^p \leq p^{p/2} \left(\sum_{n \leq N} |f_n(x_{2n})|^2 \right)^{p/2}.$$

The proof is then complete. Concerning the constant 4 in the Theorem, the paper [7] by Y.-F. Ren and H.-Y. Liang gives the upper bound 9/2, which is worse than the above one. \square

4 Proof of Theorem 1.1

Proof. We define $\Omega = \{1, \dots, K\}$ which we equip with the probability measure defined by $\nu(\{k\}) = |\lambda_k|/\|\lambda\|_1$. Given a positive integer L , we consider the family of functions φ_ℓ , for $\ell \leq L$ given by

$$\begin{aligned} \varphi_\ell : \Omega^L \times X &\rightarrow \mathbb{C} \\ ((u_h)_{h \leq L}, x) &\mapsto \lambda_{u_\ell}^\circ g_{u_\ell}(x) \end{aligned}$$

so that

$$\int_{\Omega^L} \varphi_\ell((u_h)_{h \leq L}, x) d\nu = \sum_{k \in \Omega} \frac{|\lambda_k|}{\|\lambda\|_1} \lambda_k^\circ g_k(x) = \frac{f}{\|\lambda\|_1} = f_0$$

say. We aim at showing that $(1/L) \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x)$ closely approximates f_0 for most values of $(u_h)_{h \leq L}$. Selecting one such value gives qualitatively our result. To do so, we write

$$\begin{aligned} \int_{\Omega^L} \int_X \left| \frac{1}{L} \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x) - f_0 \right|^p d((u_h)_{h \leq L}) dx \\ = \frac{1}{L^p} \int_{\Omega^L} \int_X \left| \sum_{\ell \leq L} (\varphi_\ell((u_h)_{h \leq L}, x) - f_0) \right|^p d((u_h)_{h \leq L}) dx. \end{aligned}$$

[7] Y.-F. Ren and H.-Y. Liang, 2001, "On the best constant in Marcinkiewicz-Zygmund inequality".

We apply the Marcinkiewicz-Zygmung Inequality, i.e. Theorem 3.1, to this latter expression, getting

$$\begin{aligned}
& \int_{\Omega^L} \int_X \left| \frac{1}{L} \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x) - f_0 \right|^p d((u_h)_{h \leq L}) dx \\
& \leq \frac{(4p)^{p/2}}{L^{p/2}} \int_{\Omega^L} \int_X \left| \frac{1}{L} \sum_{\ell \leq L} |\varphi_\ell((u_h)_{h \leq L}, x) - f_0|^2 \right|^{p/2} d((u_h)_{h \leq L}) dx \\
& \leq \frac{(4p)^{p/2}}{L^{p/2}} \int_X \int_{\Omega^L} \left| \frac{1}{L} \sum_{\ell \leq L} |\varphi_1((u_h)_{h \leq L}, x) - f_0|^2 \right|^{p/2} d((u_h)_{h \leq L}) dx.
\end{aligned}$$

The end is straightforward:

$$\begin{aligned}
& \int_{\Omega^L} \int_X \left| \frac{1}{L} \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x) - f_0 \right|^p d((u_h)_{h \leq L}) dx \\
& \leq \frac{(4p)^{p/2}}{L^{p/2}} \int_X \int_{\Omega^L} |\varphi_1((u_h)_{h \leq L}, x) - f_0|^p d((u_h)_{h \leq L}) dx.
\end{aligned}$$

Concerning the relevant p -norms, we make the following observations:

$$\begin{aligned}
& \int_X \int_{\Omega^L} |\varphi_1((u_h)_{h \leq L}, x)|^p d((u_h)_{h \leq L}) dx \\
& = \int_{\Omega^L} \left(\int_X |\varphi_1((u_h)_{h \leq L}, x)|^p dx \right) d((u_h)_{h \leq L}) \leq 1,
\end{aligned}$$

on the one side while on the other side, by the triangle inequality, we have

$$\|f_0\|_p \leq \sum_{k \leq K} \frac{|\lambda_k|}{\|\lambda\|_1} \|g_k\|_p \leq 1.$$

Therefore

$$\begin{aligned}
& \left(\int_{\Omega^L} \int_X \left| \frac{1}{L} \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x) - f_0 \right|^p d((u_h)_{h \leq L}) dx \right)^{1/p} \\
& \leq \frac{(4p)^{1/2}}{L^{1/2}} (1 + 1) = \sqrt{16p/L}.
\end{aligned}$$

We deduce from this inequality that the set of $(u_h)_{h \leq L}$ for which

$$\int_X \left| \frac{1}{L} \sum_{\ell \leq L} \varphi_\ell((u_h)_{h \leq L}, x) - f_0 \right|^p dx > \varepsilon^p$$

has measure at most $\sqrt{16p/(\varepsilon^2 L)}$ which is strictly less than 1 by our assumption on L . The theorem follows readily. \square

References

- [1] Y. S. Chow and H. Teicher. *Probability theory*. Third. Springer Texts in Statistics. Independence, interchangeability, martingales. Springer-Verlag, New York, 1997, pp. xxii+488. ISBN: 0-387-98228-0. DOI: [10.1007/978-1-4612-1950-7](https://doi.org/10.1007/978-1-4612-1950-7) (cit. on pp. 2, 3).

- [2] E. Croot, I. Laba, and O. Sisask. “Arithmetic progressions in sumsets and L^p -almost-periodicity”. In: *Combin. Probab. Comput.* 22.3 (2013), pp. 351–365. ISSN: 0963-5483. DOI: [10.1017/S0963548313000060](https://doi.org/10.1017/S0963548313000060) (cit. on p. 1).
- [3] E. Croot and O. Sisask. “A probabilistic technique for finding almost-periods of convolutions”. In: *Geom. Funct. Anal.* 20.6 (2010), pp. 1367–1396. ISSN: 1016-443X. DOI: [10.1007/s00039-010-0101-8](https://doi.org/10.1007/s00039-010-0101-8) (cit. on p. 1).
- [4] D. Ferger. “Optimal constants in the Marcinkiewicz-Zygmund inequalities”. In: *Statist. Probab. Lett.* 84 (2014), pp. 96–101. ISSN: 0167-7152. DOI: [10.1016/j.spl.2013.09.029](https://doi.org/10.1016/j.spl.2013.09.029) (cit. on p. 3).
- [5] U. Haagerup. “The best constants in the Khintchine inequality”. In: *Studia Math.* 70.3 (1981), 231–283 (1982). ISSN: 0039-3223. DOI: [10.4064/sm-70-3-231-283](https://doi.org/10.4064/sm-70-3-231-283) (cit. on p. 2).
- [6] L. B. Pierce. “On superorthogonality”. In: *J. Geom. Anal.* 31.7 (2021), pp. 7096–7183. ISSN: 1050-6926. DOI: [10.1007/s12220-021-00606-3](https://doi.org/10.1007/s12220-021-00606-3) (cit. on p. 1).
- [7] Y.-F. Ren and H.-Y. Liang. “On the best constant in Marcinkiewicz-Zygmund inequality”. In: *Statist. Probab. Lett.* 53.3 (2001), pp. 227–233. ISSN: 0167-7152. DOI: [10.1016/S0167-7152\(01\)00015-3](https://doi.org/10.1016/S0167-7152(01)00015-3) (cit. on pp. 3, 4).