

DEFINABILITY OF HENSELIAN VALUATIONS IN POSITIVE (RESIDUE) CHARACTERISTIC

Margarete Ketelsen and Simone Ramello

joint with Piotr Szewczyk (Dresden)

Institut für Mathematische Logik und Grundlagenforschung
University of Münster, Germany

PhDs in Logic, Granada, 04.10.2023



THE ROAD AHEAD

$\underbrace{\text{model theory}}_{\text{MATHEMATICAL LOGIC}}$ of $\underbrace{\text{valued fields}}_{\text{ALGEBRA}}.$

THE ROAD AHEAD

$\underbrace{\text{model theory}}_{\text{MATHEMATICAL LOGIC}}$ of $\underbrace{\text{valued fields}}_{\text{ALGEBRA}}.$

Today, we will try to:

- Tell you what valued fields are.

THE ROAD AHEAD

$\underbrace{\text{model theory}}_{\text{MATHEMATICAL LOGIC}}$ of $\underbrace{\text{valued fields}}_{\text{ALGEBRA}}.$

Today, we will try to:

- ▶ Tell you what valued fields are.
- ▶ Give you an idea of what our results look like.

THE ROAD AHEAD

$\underbrace{\text{model theory}}_{\text{MATHEMATICAL LOGIC}}$ of $\underbrace{\text{valued fields}}_{\text{ALGEBRA}}.$

Today, we will try to:

- ▶ Tell you what valued fields are.
- ▶ Give you an idea of what our results look like.
- ▶ Tell you about an obstacle in this area and how we turned it into a tool.

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

► $v(xy) = v(x) + v(y),$

multiplying two elements sums their valuations

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

► $v(xy) = v(x) + v(y),$

multiplying two elements sums their valuations

► $v(x + y) \geq \min\{v(x), v(y)\}.$

all triangles are isosceles

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

► $v(xy) = v(x) + v(y),$

multiplying two elements sums their valuations

► $v(x + y) \geq \min\{v(x), v(y)\}.$

all triangles are isosceles

(Counter)intuition: an element $r \in K^\times$ is *large* if its valuation $v(r) \in \Gamma$ is *small*, i.e. close to 0.

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

► $v(xy) = v(x) + v(y),$

multiplying two elements sums their valuations

► $v(x + y) \geq \min\{v(x), v(y)\}.$

all triangles are isosceles

(Counter)intuition: an element $r \in K^\times$ is *large* if its valuation $v(r) \in \Gamma$ is *small*, i.e. close to 0. Along this intuition, we usually set $v(0) := \infty$.

VALUATIONS AND WHERE TO FIND THEM

Definition

A valuation on a field K is a surjective map $v: K^\times \rightarrow \Gamma$, where $(\Gamma, +, \leq, 0)$ is an ordered abelian group, such that:

- ▶ $v(xy) = v(x) + v(y)$, *multiplying two elements sums their valuations*
- ▶ $v(x + y) \geq \min\{v(x), v(y)\}$. *all triangles are isosceles*

(Counter)intuition: an element $r \in K^\times$ is *large* if its valuation $v(r) \in \Gamma$ is *small*, i.e. close to 0. Along this intuition, we usually set $v(0) := \infty$.

The ordered abelian group Γ is called the *value group*. We also denote it by vK .

OUR FAVOURITE EXAMPLE

Fix a prime number p .

► If $a \in \mathbb{Z} \setminus \{0\}$, then

$$v_p(a) := \max\{n \in \mathbb{N} : p^n \text{ divides } a\}.$$

OUR FAVOURITE EXAMPLE

Fix a prime number p .

► If $a \in \mathbb{Z} \setminus \{0\}$, then

$$v_p(a) := \max\{n \in \mathbb{N} : p^n \text{ divides } a\}.$$

For example, $v_3(6560) = 0$. According to v_3 , then, 6560 is “big”. But $v_3(6561) = 8$, which is then “smaller” than 6560.

OUR FAVOURITE EXAMPLE

Fix a prime number p .

► If $a \in \mathbb{Z} \setminus \{0\}$, then

$$v_p(a) := \max\{n \in \mathbb{N} : p^n \text{ divides } a\}.$$

For example, $v_3(6560) = 0$. According to v_3 , then, 6560 is “big”. But $v_3(6561) = 8$, which is then “smaller” than 6560. If $a, b \in \mathbb{Z} \setminus \{0\}$ are coprime, then

$$v_p\left(\frac{a}{b}\right) := v_p(a) - v_p(b).$$

OUR FAVOURITE EXAMPLE

Fix a prime number p .

- If $a \in \mathbb{Z} \setminus \{0\}$, then

$$v_p(a) := \max\{n \in \mathbb{N} : p^n \text{ divides } a\}.$$

For example, $v_3(6560) = 0$. According to v_3 , then, 6560 is “big”. But $v_3(6561) = 8$, which is then “smaller” than 6560. If $a, b \in \mathbb{Z} \setminus \{0\}$ are coprime, then

$$v_p\left(\frac{a}{b}\right) := v_p(a) - v_p(b).$$

- This defines a valuation $v_p: \mathbb{Q} \setminus \{0\} \rightarrow \mathbb{Z}$, called *the p -adic valuation*. With it, we can define a distance on \mathbb{Q} by setting $d_p(a, b) := p^{-v_p(a-b)}$.

OUR FAVOURITE EXAMPLE

Fix a prime number p .

- If $a \in \mathbb{Z} \setminus \{0\}$, then

$$v_p(a) := \max\{n \in \mathbb{N} : p^n \text{ divides } a\}.$$

For example, $v_3(6560) = 0$. According to v_3 , then, 6560 is “big”. But $v_3(6561) = 8$, which is then “smaller” than 6560. If $a, b \in \mathbb{Z} \setminus \{0\}$ are coprime, then

$$v_p\left(\frac{a}{b}\right) := v_p(a) - v_p(b).$$

- This defines a valuation $v_p: \mathbb{Q} \setminus \{0\} \rightarrow \mathbb{Z}$, called *the p -adic valuation*. With it, we can define a distance on \mathbb{Q} by setting $d_p(a, b) := p^{-v_p(a-b)}$.
- If we *complete* the corresponding metric space, we obtain a (new) valued field called \mathbb{Q}_p , with its own valuation v_p . These are the *p -adic numbers*.

WHY YOU SHOULD LIKE THE p -ADICS

- ▶ (\mathbb{Q}_p, v_p) is crucial for algebraic purposes. But we are logicians (allegedly)!

WHY YOU SHOULD LIKE THE p -ADICS

- ▶ (\mathbb{Q}_p, v_p) is crucial for algebraic purposes. But we are logicians (allegedly)!
- ▶ A valuation is “the same” as its valuation ring, i.e. the subring

$$\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}.$$

This is the part where we should tell you that ∞ is larger than all elements of Γ , and thus $0 \in \mathcal{O}_v$.

WHY YOU SHOULD LIKE THE p -ADICS

- ▶ (\mathbb{Q}_p, v_p) is crucial for algebraic purposes. But we are logicians (allegedly)!
- ▶ A valuation is “the same” as its valuation ring, i.e. the subring

$$\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}.$$

This is the part where we should tell you that ∞ is larger than all elements of Γ , and thus $0 \in \mathcal{O}_v$.

- ▶ In the case of \mathbb{Q}_p , this subring is called \mathbb{Z}_p (guess why!).

WHY YOU SHOULD LIKE THE p -ADICS

- ▶ (\mathbb{Q}_p, v_p) is crucial for algebraic purposes. But we are logicians (allegedly)!
- ▶ A valuation is “the same” as its valuation ring, i.e. the subring

$$\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}.$$

This is the part where we should tell you that ∞ is larger than all elements of Γ , and thus $0 \in \mathcal{O}_v$.

- ▶ In the case of \mathbb{Q}_p , this subring is called \mathbb{Z}_p (guess why!). Julia Robinson pointed out something remarkable about \mathbb{Z}_p (for $p \neq 2$):

$$\mathbb{Z}_p = \{x \in \mathbb{Q}_p \mid \exists Y(Y^2 = 1 + px^2)\}.$$

There is a similar formula for $p = 2$.

- ▶ \mathbb{Z}_p is given, as a subset of \mathbb{Q}_p , by a polynomial equation together with some quantifiers.

WHY YOU SHOULD LIKE THE p -ADICS

- ▶ (\mathbb{Q}_p, v_p) is crucial for algebraic purposes. But we are logicians (allegedly)!
- ▶ A valuation is “the same” as its valuation ring, i.e. the subring

$$\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}.$$

This is the part where we should tell you that ∞ is larger than all elements of Γ , and thus $0 \in \mathcal{O}_v$.

- ▶ In the case of \mathbb{Q}_p , this subring is called \mathbb{Z}_p (guess why!). Julia Robinson pointed out something remarkable about \mathbb{Z}_p (for $p \neq 2$):

$$\mathbb{Z}_p = \{x \in \mathbb{Q}_p \mid \exists Y(Y^2 = 1 + px^2)\}.$$

There is a similar formula for $p = 2$.

- ▶ \mathbb{Z}_p is given, as a subset of \mathbb{Q}_p , by a polynomial equation together with some quantifiers. We say that it is a *definable* set in the language of rings.

LOGICIANS, ASSEMBLE! CONT'D

Big question: Is this common? When is some valuation ring definable in the language of rings?

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?

Yes, but often in several different ways.

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- ▶ Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?
Yes, but often in several different ways.
- ▶ v is *henselian* if there is a **unique** way to extend v to any algebraic extension of K .

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- ▶ Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?
Yes, but often in several different ways.
- ▶ v is *henselian* if there is a **unique** way to extend v to any algebraic extension of K . A henselian valuation is a bit like a *fill the gaps* exercise in a textbook.

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- ▶ Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?
Yes, but often in several different ways.
- ▶ v is *henselian* if there is a **unique** way to extend v to any algebraic extension of K . A henselian valuation is a bit like a *fill the gaps* exercise in a textbook.
- ▶ v_p is henselian.

THE PROBLEM OF HENSELIANITY

Not all valuations are created equal.

- ▶ Take a field K with a valuation v . We give you an algebraic extension L of K , e.g. $L = K(\alpha)$ where α is the root of some polynomial over K . Can you extend v to L ?
Yes, but often in several different ways.
- ▶ v is *henselian* if there is a **unique** way to extend v to any algebraic extension of K . A henselian valuation is a bit like a *fill the gaps* exercise in a textbook.
- ▶ v_p is henselian. *We will only care about henselian valuations.*

THE BIG QUESTION, TAKE 2

Big question: when is
a **henselian** valuation ring definable in the language of rings?

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}$.

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K \mid v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p)

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K : v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p) both have residue field \mathbb{F}_p , the finite field with p elements

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K : v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p) both have residue field \mathbb{F}_p , the finite field with p elements
In fact, $\mathbb{Q} \subseteq \mathbb{Q}_p$ is an *immediate extension*:

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K : v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p) both have residue field \mathbb{F}_p , the finite field with p elements
In fact, $\mathbb{Q} \subseteq \mathbb{Q}_p$ is an *immediate extension*: They have the same value groups and residue fields.

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K : v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p) both have residue field \mathbb{F}_p , the finite field with p elements
In fact, $\mathbb{Q} \subseteq \mathbb{Q}_p$ is an *immediate extension*: They have the same value groups and residue fields.
- So a valued field consists of *two fields*: the “big” valued field and the “smaller” residue field.

TWO FIELDS IN DISGUISE

- To any valued field (K, v) we can associate another “smaller” field, called the *residue field*,

$$Kv := \{x \in K : v(x) \geq 0\} / \{x \in K : v(x) > 0\}.$$

Indeed, $\mathfrak{m}_v := \{x \in K : v(x) > 0\}$ is the unique maximal ideal of $\mathcal{O}_v = \{x \in K : v(x) \geq 0\}$.

- **Example:** (\mathbb{Q}, v_p) and (\mathbb{Q}_p, v_p) both have residue field \mathbb{F}_p , the finite field with p elements
In fact, $\mathbb{Q} \subseteq \mathbb{Q}_p$ is an *immediate extension*: They have the same value groups and residue fields.
- So a valued field consists of *two fields*: the “big” valued field and the “smaller” residue field. If we talk about the characteristic of a valued field, we talk about the characteristics of the two fields
 - **equicharacteristic zero:** $\text{char}(K) = \text{char}(Kv) = 0$
 - **mixed characteristic:** $\text{char}(K) = 0 < p = \text{char}(Kv)$, where p is prime
 - **positive characteristic:** $\text{char}(K) = \text{char}(Kv) = p$, where p is prime

A CANONICAL FRIEND

- ▶ Henselian valuations on a given field K arrange themselves nicely according to whether their residue field is separably closed or not,

$$H_1(K) := \{v: Kv \text{ is not separably closed}\} \text{ vs. } H_2(K) := \{v: Kv \text{ is separably closed}\}.$$


A CANONICAL FRIEND

- Henselian valuations on a given field K arrange themselves nicely according to whether their residue field is separably closed or not,

$$H_1(K) := \{v: Kv \text{ is not separably closed}\} \text{ vs. } H_2(K) := \{v: Kv \text{ is separably closed}\}.$$

- $H_1(K)$ is linearly ordered by inclusion.

The “middle point” between $H_1(K)$ and $H_2(K)$ is the *canonical henselian valuation* v_K .



THE GIST OF IT

$$\underbrace{\exists \text{ definable (non-trivial) henselian valuation}}_{\text{Logic question}} \iff \underbrace{\text{Conditions on the canonical henselian valuation}}_{\text{(Almost) algebra answer}}$$

WHAT WE PROVED

Theorem (Jahnke, Koenigsmann, 2017; Ketelsen, R., Szewczyk, 2023)

Let K be a non-separably closed henselian field.

If $\text{char}(K) = p > 0$, then assume that K is perfect.

If $\text{char}(K) = 0 < p = \text{char}(Kv_K)$, then assume that \mathcal{O}_{v_K}/p is semi-perfect.

Then,

$$K \text{ admits a definable non-trivial henselian valuation} \iff \begin{cases} Kv_K = Kv_K^{\text{sep}}, & \text{or} \\ Kv_K \text{ is not } t\text{-henselian}, & \text{or} \\ \exists L \succeq Kv_K \text{ with } v_L L \text{ divisible}, & \text{or} \\ v_K K \text{ is not divisible}, & \text{or} \\ (K, v_K) \text{ is not defectless}, & \text{or} \\ \exists L \succeq Kv_K \text{ with } (L, v_L) \text{ not defectless.} \end{cases}$$

WHAT WE HAD BEFORE

Theorem (Jahnke, Koenigsmann, 2017; Ketelsen, R., Szewczyk, 2023)

Let K be a non-separably closed henselian field, $\text{char}(Kv) = 0$.

If $\text{char}(K) = p > 0$, then assume that K is perfect.

If, further, $\text{char}(K) = 0 < p = \text{char}(Kv_K)$, then further assume that \mathcal{O}_{v_K}/p is semi-perfect.

Then,

$$K \text{ admits a definable non-trivial henselian valuation} \iff \begin{cases} Kv_K = Kv_K^{\text{sep}}, & \text{or} \\ Kv_K \text{ is not } t\text{-henselian}, & \text{or} \\ \exists L \succeq Kv_K \text{ with } v_L L \text{ divisible}, & \text{or} \\ v_K K \text{ is not divisible}, & \text{or} \\ (K, v_K) \text{ is not defectless}, & \text{or} \\ \exists L \succeq Kv_K \text{ with } (L, v_L) \text{ not defectless.} \end{cases}$$

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again.

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

- ▶ We say that $(K, v) \subseteq (L, v)$ is *defectless* if

$$[L : K] = [Lv : Kv](vL : vK).$$

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

- ▶ We say that $(K, v) \subseteq (L, v)$ is *defectless* if

$$[L : K] = [Lv : Kv](vL : vK).$$

In particular, then, if $\text{char}(Kv) = 0$, then $p = 1$ and so equality holds.

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

- ▶ We say that $(K, v) \subseteq (L, v)$ is *defectless* if

$$[L : K] = [Lv : Kv](vL : vK).$$

In particular, then, if $\text{char}(Kv) = 0$, then $p = 1$ and so equality holds. Otherwise, not being defectless (= *having defect*) is a problem.

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

- ▶ We say that $(K, v) \subseteq (L, v)$ is *defectless* if

$$[L : K] = [Lv : Kv](vL : vK).$$

In particular, then, if $\text{char}(Kv) = 0$, then $p = 1$ and so equality holds. Otherwise, not being defectless (= *having defect*) is a problem.

- ▶ For us, however, defect is a **source of information!**

WE DON'T TALK ABOUT DEFECT

Actually, we do now.

- ▶ Given a henselian valuation v and a finite field extension $K \subseteq L$, then there is a unique extension of v to L , which we denote by v again. Then, we have

$$[L : K] \geq [Lv : Kv](vL : vK).$$

More precisely,

$$[L : K] = p^d [Lv : Kv](vL : vK),$$

where $p = \text{char}(Kv)$, if the latter is positive, and $p = 1$ if $\text{char}(Kv) = 0$.

- ▶ We say that $(K, v) \subseteq (L, v)$ is *defectless* if

$$[L : K] = [Lv : Kv](vL : vK).$$

In particular, then, if $\text{char}(Kv) = 0$, then $p = 1$ and so equality holds. Otherwise, not being defectless (= *having defect*) is a problem.

- ▶ For us, however, defect is a **source of information!** (At least when it is “of independent type”).