1. The probability of a leap year selected at random contain 53							
Sunday is:							
(a) 53/366	(b) 1/7	(c) 2/7	(d) 53/365				
2. A bag contains	2. A bag contains 3 red and 2 blue marbles. A marble is drawn at						
random. The prol	bability of drav	wing a black ball	is:				
• •	` '	(c) $0/5$					
3. The probabilit	y that it will ra	in tomorrow is 0	.85. What is the				
probability that it							
* *	• •		(d) none of these				
•	•		ed from the numbers				
(1, 2, 3,,15	•		4.0				
	, ,	(c) 2/15	• •				
5. What are the							
		(c) 8					
-	-	number selecte	d at random from the				
numbers (1,2,3, .		() 10 10 7	(1)				
• •	1	` '	(d) none of these				
7. The sum of th							
		0 (d) none					
	_	are given; choos	e the correct answer				
for that which is							
			(d) none of these.				
		nultaneously, tha	n the probability of				
getting at least to			(1) 1 (0)				
		(c) ½					
10. A letter is ch							
ASSASSINATIO	JNQ. The pro	bability that the	letter chosen has:				
(a) 6/13	(b) //13	(c) 1	(d) none of these.				
44 A dia : :	5 :		!				
	-		ing an even number.				
(A) 2/3	(B) I	(C) 5/6	D) 1/2				
12. Two coins are thrown at the same time. Find the probability of getting both heads.							
(A) 3/4 (B) 1/4		(D) 0					
13. Two dice are thrown simultaneously. The probability of getting a							

sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4	/9			
14. 100 cards are numbered from 1 to 100. Find the probability of getting a prime number.							
(A) 3/4		(C) 1/4	(D)	29/100			
_	a blue ball is do	ouble that of a		If the probability en the number of			
16. A box of	f 600 bulbs con trandom from t ve bulb is:	itains 12 defe	ctive bulbs.				
mixed thoro	narked with nur oughly. One car lity that the nur (B) 1/10	d is drawn fro	m this box r s a perfect	andomly, then square.			
18. What is (A) 1/7	the probability (B) 53/366	•					
19. A card is drawn from a well shuffled deck of 52 cards. Find the probability of getting a king of red suit. (A) 1/26 (B) 3/26 (C) 7/52 (D) 1/13							
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is: (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12							
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game. (A) 3/4 (B) 1/2 (C) 1 (D) 1/4							

same birth	day is the sa	me birthd	ay is:	at both will ha		
(A) 364/36	5 (B)	31/365	(C) 1/365	(D)	1/133225	
2. Then the	ber x is chose probability (B) 2/5	that $x^2 < 2$	is?	ne numbers -2	2, -1, 0 , 1,	
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is 2/3, then the number of white marbles in the jar is: (A) 10 (B) 6 (C) 8 (D) 7						
Then the pr	per is select robability th (B) 4/25	at it is a m	ultiple of 3	st 50 natural ı and 4 is:	numbers.	
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?						
a) $\frac{1}{7}$	b) $\frac{5}{42}$		c) $\frac{1}{21}$	d) $\frac{4}{21}$		
	cored by ba The standa		_	atches are 50	, 70, 82,	
	b) 25.49				59	
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.						
a) 13, 15	b) 13,	18	c) 18, 15		d) 13, 16	
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is						
a) $^{1}/_{2}$	b) $^{1}/_{\mathrm{p}}$		3. The value	of E(X²) is _ d) 9		
31.The ran	31. The random variables X and Y have variances 0.2 and 0.5					

respectively. Let Z= 5X-2Y. The variance of Z is?

<pre>probability? a) P(x) = 1</pre>	•	P(x) = 3	one is not possi	ble in			
• •	2 and E(z) = 4 b) 6	c) 0	•	ufficient data			
34.The cov	ariance of two	independent	random variabl	e is			
a) 1	b) 0	c) - 1	d) Und	defined			
35.If Σ P(x) a) 0	b) = k ² – 8 then, b) 1	the value of I		ufficient data			
36.If P(x) = a) 1	0.5 and x = 4, b) 0.5	then E(x) = ? c) 4	d) 2				
is always?	r ete probabilit b) Infinite	y distribution c) 1	, the sum of all d) Und				
38.If the probability of hitting the target is 0.4, find mean and variance. a) 0.4, 0.24							
39.If the probability that a bomb dropped from a place will strike the target is 60% and if 10 bombs are dropped, find mean and variance? a) 0.6, 0.24 b) 6, 2.4 c) 0.4, 0.16 d) 4, 1.6 40. Find the mean of tossing 8 coins. a) 2 b) 4 c) 8 d) 1							
,		,	standard norm	al distribution?			

c) 5

d) 7

a) 3

b) 4

a) Mean is 0 c) Mean is 0								
42.Varianc (a) E(X)						d) (E(X))2		
43.Mean of a random variable X is given by a) E(X)								
44.Mean of a constant 'a' is a) 0 b) a c) a/2 d) 1								
45.Variance of a constant 'a' is . a) 0								
46.Find the mean and variance of X?								
×	0	1	2	3	4			
f(x)	1/9	2/9	3/9	2/9	1/9			
a) 2, 4/3 47.Find the	·) 3, 4/3 tion of a ı		c) 2, 2/3 variable :		d) 3, 2/3		

	Х	0	1	2	3	
	f(x)	1/6	2/6	2/6	1/6	
a) ().5		b) 1.5		c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by



c) np2q

d) npq2

- 49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.**
- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx
- 50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?**

- a) \sqrt{np} b) \sqrt{pq} c) (np)2