

Right-orders on Free Groups

George Metcalfe and Simon Santschi

Mathematical Institute, University of Bern, Switzerland
{george.metcalfe,simon.santschi}@unibe.ch

Let \mathbf{G} be a group (or monoid). A *right-order* on \mathbf{G} is a total order \leq on G such that for all $a, b, c \in G$, $a \leq b$ implies $ac \leq bc$. For a subset $S \subseteq G$ of a group \mathbf{G} we say that S *extends to a right-order* on \mathbf{G} if there exists a right-order \leq on \mathbf{G} such that for all $g \in S$, $e < g$, where e is the neutral element of \mathbf{G} . Free groups are well-known to be right-orderable and in [1] it is shown that given a free group $\mathbf{F}_g(X)$ over a set X the problem of checking for a finite subset $S \subseteq \mathbf{F}_g(X)$ if it extends to a right-order on $\mathbf{F}_g(X)$ is decidable.

In this talk, which is based on results from [5], we show that this problem is NP-complete for $|X| \geq 2$. For the proof we exploit the following connection between right-orders on free groups and the validity of certain equations in the class \mathbf{LG} of ℓ -groups¹ obtained in [3]:

Proposition. *Let $\mathbf{F}_g(X)$ be the free group over a set X . Then a finite subset $\{s_1, \dots, s_n\} \subseteq \mathbf{F}_g(X)$ extends to a right-order on $\mathbf{F}_g(X)$ if and only if $\mathbf{LG} \not\models e \leq s_1 \vee \dots \vee s_n$.*

Since the equational theory of \mathbf{LG} is co-NP-complete (see [4]), it follows that the problem of checking if a finite set of elements of a free group extends to a right-order is in NP. To show that the problem is NP-hard we use a method of eliminating meets from ℓ -group equations. This method allows us to reduce the validity of certain equations, the set of which is known to be co-NP-hard, to equations of the form $e \leq s_1 \vee \dots \vee s_n$ with group terms s_1, \dots, s_n , yielding:

Theorem. *Let $\mathbf{F}_g(X)$ be the free group over a set X with $|X| \geq 2$. Then the problem of checking for $\{s_1, \dots, s_n\} \subseteq \mathbf{F}_g(X)$ if it extends to a right-order is NP-complete.*

Using a method of eliminating inverses in ℓ -group equations first considered in [2] we also obtain an analogue of the result for free monoids in the case where the generating set is infinite:

Corollary. *Let $\mathbf{F}_m(X)$ be the free monoid over an infinite set X . Then the problem of checking for $s_1, t_1, \dots, s_n, t_n \in \mathbf{F}_m(X)$ if there exists a right-order \leq on $\mathbf{F}_m(X)$ with $s_1 < t_1, \dots, s_n < t_n$ is NP-complete.*

References

- [1] A. Clay, L.H. Smith. Corrigendum to: “On ordering free groups”. *J. Symb. Comput.*, 44(10) 1529–1532, 2009.
- [2] A. Colacito, N. Galatos, G. Metcalfe, and S. Santschi. From distributive ℓ -monoids to ℓ -groups, and back again. *J. Algebra*, 601:129–148, 2022.
- [3] A. Colacito and G. Metcalfe. Ordering groups and validity in lattice-ordered groups. *J. Pure Appl. Algebra*, 223(12):5163–5175, 2019.
- [4] N. Galatos and G. Metcalfe. Proof theory for lattice-ordered groups. *Ann. Pure Appl. Logic*, 8(167):707–724, 2016.
- [5] G. Metcalfe and S. Santschi. Equational theories of idempotent semifields, Manuscript. Available at <https://arxiv.org/abs/2402.09876>, 2024.

¹An ℓ -group is an algebra $\mathbf{L} = \langle L, \wedge, \vee, \cdot, {}^{-1}, e \rangle$ such that $\langle L, \wedge, \vee \rangle$ is a lattice, $\langle L, \cdot, {}^{-1}, e \rangle$ is a group, and for all $a, b, c, d \in L$, $a \leq b \implies cad \leq cbd$.