

# AFFINE PAVINGS OF QUIVER FLAG VARIETIES

XIAOXIANG ZHOU

ABSTRACT. In this article, we construct affine pavings for quiver partial flag varieties when the quiver is of Dynkin type. To achieve our results, we extend methods from Cerulli-Irelli–Esposito–Franzen–Reineke and Maksimau as well as techniques from Auslander–Reiten theory.

## CONTENTS

1. Introduction	1
Acknowledgement	3
2. Preliminaries	3
2.1. Extended quiver	3
2.2. Canonical functor $\Phi$	4
2.3. Dimension vector	5
2.4. Ext-vanishing properties	7
3. Main Theorem	9
4. Application: Dynkin Case	13
5. affine paving for big order representations	14
6. Application: Affine Case	21
References	22

## 1. INTRODUCTION

Affine pavings are an important concept in algebraic geometry similar to cellular decompositions in topology. A complex algebraic variety  $X$  has an affine paving if  $X$  has a filtration

$$\emptyset = X_0 \subset X_1 \subset \cdots \subset X_d = X$$

with  $X_i$  closed and  $X_{i+1} \setminus X_i$  isomorphic to some affine space  $\mathbb{A}_{\mathbb{C}}^k$ .

Affine pavings imply nice properties about the cohomology of varieties, for example the vanishing of cohomology in odd degrees. For other properties see [3, 1.7].

Affine pavings have been constructed in many cases, as for Grassmannians, flag varieties, as well as certain Springer fibers, quiver Grassmannians, and quiver flag varieties. This article focuses on the case of (strict) partial flag varieties which parameterize subrepresentations of a fixed indecomposable representation of a quiver. In particular, we consider quivers of Dynkin type or affine type. In this case, affine pavings have been constructed in [6] for quiver Grassmannians in all types and in [7] for partial flag varieties of type  $A$  and  $D$  (see Table 1). Besides, affine pavings have been constructed in [5, Theorem 6.3] for

strict partial flag varieties in type  $\tilde{A}$  with cyclic orientation, which generalized the result in [8] for complete quiver flag varieties in nilpotent representations of an oriented cycle. Recently, De Concini and Maffei [4] introduced an independent method to construct affine pavings for Springer fibers of type  $G_2$ ,  $F_4$ , and  $E_7$ , which differs from the technique we describe. In this paper, we will tackle the remaining cases.

**Theorem 1.1.** *Denote  $Q$  a quiver and  $M$  a representation of  $Q$ .*

- (1) *If  $Q$  is Dynkin, then any (strict) partial flag variety  $\text{Flag}(M)$  has an affine paving;*
- (2) *If  $Q$  is of type  $\tilde{A}$  or  $\tilde{D}$ , then for any indecomposable representation  $M$ , the (strict) partial flag variety  $\text{Flag}(M)$  has an affine paving;*
- (3) *If  $Q$  is of type  $\tilde{E}$ , assume that  $\text{Flag}(N)$  has an affine paving for any regular quasi-simple representation  $N \in \text{rep}(Q)$ , then  $\text{Flag}(M)$  has an affine paving for any indecomposable representation  $M$ .*

	$\text{Gr}^{KQ}(X)$	$\text{Flag}_{\text{d}}(X)$	$\text{Flag}_{\text{d, str}}(X)$
$A$	[6, Section 5]	[7, Theorem 2.20]	Theorem 4.1
$D$			
$E$		Theorem 4.1	
$\tilde{A}$	[6, Section 6]	Theorem 6.3	
$\tilde{D}$			
$\tilde{E}$		reduced to the regular quasi-finite case.	

TABLE 1

We proceed as follows. In Section 2, we discuss basic definitions and properties of partial flags. In Section 3 we will prove key Theorems 3.2 and 3.3, which allow us to construct affine pavings for quiver partial flag varieties inductively. We apply these theorems to partial flag varieties of Dynkin type, see Section 4, and to partial flag varieties of affine type, see Section 6. We will combine and extend results from [6] and [7]. Following the arguments of [7] would require studying millions of cases when we consider the Dynkin quivers of type  $E$ . To avoid this, we extend the methods of [6] from quiver Grassmannian to quiver partial flag variety. This will reduce the case by case analysis to a feasible computation of (mostly) 8 critical cases, which we carry out in Section 4 and Section 5.

**Conventions and Notations.** *Throughout this article,  $K = \mathbb{C}$ ,  $R$  is a  $K$ -algebra with unit, and  $\text{mod}(R)$  denotes the category of left  $R$ -modules of finite dimension. Let  $Q$  be a quiver equipped with the set of finite vertices  $v(Q)$  and the set of finite edges  $a(Q)$ . For an arrow  $b$ , we call  $s(b)$  the starting vertex and  $t(b)$  the terminal vertex of  $b$ . We denote by  $KQ$  the path algebra and  $\text{rep}(Q) = \text{mod}(KQ)$  the category of quiver representations of finite dimension. For a representation  $X \in \text{rep}(Q)$ , we denote by  $X_i := e_i X$  the  $K$ -linear space at the vertex  $i \in v(Q)$ . We denote by  $P(i)$ ,  $I(i)$  and  $S(i)$  the indecomposable projective, injective, simple modules corresponding to the vertex  $i$ , respectively.*

## ACKNOWLEDGEMENT

First, I would like to thank my supervisor, Jens Niklas Eberhardt, for introducing me this specific problem, discussing earlier drafts, as well as providing help with the write-up. I would also like to thank Hans Franzen for answering some questions regarding [6], and thank Ruslan Maksimau, Francesco Esposito for some comments and suggestions. This article is part of my master thesis project.

## 2. PRELIMINARIES

**2.1. Extended quiver.** In this subsection, we introduce the notion of extended quiver which allows to view partial flag varieties as quiver Grassmannians. Intuitively, a flag of quiver representations can be encoded as a subspace of a representation of the extended quiver.

**Definition 2.1** (Extended quiver). *For a quiver  $Q$  and an integer  $d \geq 1$ , the extended quiver  $Q_d$  is defined as follows:*

- The vertex set of  $Q_d$  is defined as the Cartesian product of the vertex set of  $Q$  and  $\{1, \dots, d\}$ , i.e.,  

$$v(Q_d) = v(Q) \times \{1, \dots, d\}.$$
- There are two types of arrows: for each  $(i, r) \in v(Q) \times \{1, \dots, d-1\}$ , there is one arrow from  $(i, r)$  to  $(i, r+1)$ ; for each arrow  $i \rightarrow j$  in  $Q$  and  $r \in \{1, \dots, d\}$ , there is one arrow from  $(i, r)$  to  $(j, r)$ .


The extended quiver  $Q_d$  is exactly the same quiver as  $\hat{\Gamma}_d$  in [7, Definition 2.2]. The next definition is a small variation:

**Definition 2.2** (Strict extended quiver). *For a quiver  $Q$  and an integer  $d \geq 2$ , the strict extended quiver  $Q_{d,\text{str}}$  is defined as follows:*

- The vertex set of  $Q_d$  is defined as the Cartesian product of the vertex set of  $Q$  and  $\{1, \dots, d\}$ , i.e.,  

$$v(Q_{d,\text{str}}) = v(Q) \times \{1, \dots, d\}.$$
- We have two types of arrows: for each  $(i, r) \in v(Q) \times \{1, \dots, d-1\}$ , there is one arrow from  $(i, r)$  to  $(i, r+1)$ ; for each arrow  $i \rightarrow j$  in  $Q$  and  $r \in \{2, \dots, d\}$ , there is one arrow from  $(i, r)$  to  $(j, r-1)$ .

**Example 2.3.** The (strict) extended quiver for a Dynkin quiver  $Q$  of type  $A_4$  looks as follows.



Next, we define the quiver algebras for later use.

**Definition 2.4** (Algebra of an extended quiver). *For an extended quiver  $Q_d$ , let  $KQ_d$  be the corresponding path algebra, and  $I$  be the ideal of  $KQ_d$  identifying all the paths with the same sources and targets. The algebra of the extended quiver  $Q_d$  is defined as*

$$R_d := KQ_d/I.$$

*Similarly, we define the algebra  $R_{d,\text{str}} := KQ_{d,\text{str}}/I$  for the strict extended quiver.*

By abuse of notation, we often abbreviate  $R_d$  and  $R_{d,\text{str}}$  by  $R$ .

**2.2. Canonical functor  $\Phi$ .** We follow [7, 2.3] in this subsection with a few variations.

**Definition 2.5** (Partial flag). *For a quiver representation  $X \in \text{rep}(Q)$ , a partial flag of  $X$  is defined as an increasing sequence of subrepresentations of  $X$ . For an integer  $d \geq 1$ , we denote*

$$\text{Flag}_d(X) := \{0 \subseteq M_1 \subseteq \cdots \subseteq M_d \subseteq X\}$$

*as the collection of all partial flags of length  $d$ , and call it the partial flag variety.*

**Definition 2.6** (Strict partial flag). *For a quiver representation  $X \in \text{rep}(Q)$ , a strict partial flag of  $X$  is defined as an increasing sequence of subrepresentations  $(M_k)_k$  of  $X$  such that for any arrow  $x \in v(Q)$  and any  $k$ , we have  $x.M_{k+1} \subseteq M_k$ . For an integer  $d \geq 2$ , we denote*

$$\text{Flag}_{d,\text{str}}(X) := \{0 \subseteq M_1 \subseteq \cdots \subseteq M_d \subseteq X \mid x.M_{k+1} \subseteq M_k\}$$

*as the collection of all strict partial flags of length  $d$ , and call it the strict partial flag variety.*

**Definition 2.7** (Grassmannian). *Let  $R$  be the bounded quiver algebra defined in Definition 2.5 or 2.6. For a module  $T \in \text{mod}(R)$ , the Grassmannian  $\text{Gr}^R(T)$  is defined as the set of all submodules of  $T$ , i.e.,*

$$\text{Gr}^R(T) := \{T' \subseteq T \text{ as the submodule}\}.$$

**Definition 2.8** (Canonical functor  $\Phi$ ). *The canonical functor  $\Phi : \text{rep}(Q) \longrightarrow \text{mod}(R)$  is defined as follows:*

- $(\Phi(X))_{(i,r)} := X_i$ ;
- $(\Phi(X))_{(i,r) \rightarrow (i,r+1)} := \text{Id}_{X_i}$ ;
- Either  $(\Phi(X))_{(i,r) \rightarrow (j,r)} := X_{i \rightarrow j}$  for  $R = R_d$ ,  
or  $(\Phi(X))_{(i,r) \rightarrow (j,r-1)} := X_{i \rightarrow j}$  for  $R = R_{d,\text{str}}$ .

The functor  $\Phi$  helps to realize a partial flag as a quiver subrepresentation.

**Proposition 2.9.** *For a representation  $X \in \text{rep}(Q)$ , the canonical functor  $\Phi$  induces isomorphisms*

$$\text{Flag}_d(X) \cong \text{Gr}^{R_d}(\Phi(X)) \quad \text{Flag}_{d,\text{str}}(X) \cong \text{Gr}^{R_{d,\text{str}}}(\Phi(X)).$$

*Proof.* The isomorphism maps a flag  $M : M_1 \subseteq \cdots \subseteq M_d$  to a representation  $\Phi'(M)$  with  $\Phi'(M)_{(i,r)} = M_{i,r}$  and obvious morphisms for arrows. The non-strict case is mentioned in [7, page 4] and the strict case works similarly.  $\square$



FIGURE 1

**Example 2.10.** Consider the quiver  $Q: x \longrightarrow y \longleftarrow z \longrightarrow w$ , and let  $X: X_x \longrightarrow X_y \longleftarrow X_z \longrightarrow X_w$  be a representation. The varieties  $\text{Flag}_3(X), \text{Flag}_{3,\text{str}}(X)$  then arise as quiver Grassmannian as shown in Figure 1.

In many cases, the proof of the strict case and the non-strict case is the same, so we often treat them in the same way. For example, we may abbreviate the formula in Proposition 2.9 as

$$\text{Flag}(X) \cong \text{Gr}(\Phi(X)).$$

**2.3. Dimension vector.** In this subsection we recall some notations of dimension vectors.

**Definition 2.11** (Dimension vector). *For a quiver  $Q$  and a representation  $M \in \text{rep}(Q)$ , the set of dimension vectors of  $Q$  is defined as  $\prod_{i \in v(Q)} \mathbb{Z}$ , and the dimension vector of  $M$  is defined as*

$$\underline{\dim} M := (\dim_K M_i)_{i \in v(Q)}.$$

Moreover, if  $R = KQ/I$  is a bounded quiver algebra, then every module  $T \in \text{mod}(R)$  can be viewed as a representation of  $Q$ , so we automatically have a notion of dimension vector for  $R$  and  $T$ .

Now we can write the (strict) partial flag variety and Grassmannian as disjoint union of several pieces. Since  $v(Q_{d,(\text{str})}) = v(Q) \times \{1, \dots, d\}$ , any dimension vector  $\mathbf{f}$  of  $R$  can be viewed as  $d$  dimension vectors  $(\mathbf{f}_1, \dots, \mathbf{f}_d)$ . Define

$$\begin{aligned} \text{Flag}_{\mathbf{f}}(X) &:= \{0 \subseteq M_1 \subseteq \dots \subseteq M_d \subseteq X \mid \underline{\dim} M_k = \mathbf{f}_k\} && \subseteq \text{Flag}_d(X), \\ \text{Flag}_{\mathbf{f}, \text{str}}(X) &:= \{0 \subseteq M_1 \subseteq \dots \subseteq M_d \subseteq X \mid x.M_{k+1} \subseteq M_k, \underline{\dim} M_k = \mathbf{f}_k\} && \subseteq \text{Flag}_{d, \text{str}}(X), \\ \text{Gr}_{\mathbf{f}}^R(T) &:= \{T' \subseteq T \text{ with } \underline{\dim} T' = \mathbf{f}\} && \subseteq \text{Gr}^R(T). \end{aligned}$$

Then from the Proposition 2.9 we get

$$\text{Flag}_{\mathbf{f}}(X) \cong \text{Gr}_{\mathbf{f}}^{R_d}(\Phi(X)) \quad \text{Flag}_{\mathbf{f}, \text{str}}(X) \cong \text{Gr}_{\mathbf{f}}^{R_{d, \text{str}}}(\Phi(X)).$$

*Remark 2.12.* All the spaces we defined here have natural topologies and variety structures. For example, by the standard embedding

$$\text{Gr}_{\mathbf{f}}^R(T) \hookrightarrow \prod_{(i,r) \in v(Q_{d,(\text{str})})} \text{Gr}_{\mathbf{f}_{i,r}}(T_{(i,r)}),$$

$\text{Gr}_{\mathbf{f}}^R(T)$  is then endowed with the subspace topology and subvariety structure.

Finally, we need to define the Euler form of two dimension vectors. For this we need to define the set of virtual arrows of the quivers  $Q_d$  and  $Q_{d, \text{str}}$ . Following Example 2.15, the virtual arrows of the quivers  $Q_3$  and  $Q_{3, \text{str}}$  are depicted in red.

**Definition 2.13** (Virtual arrows of the quiver  $Q_d$ ). *For  $d \geq 1$ , the virtual arrows of the quiver  $Q_d$  is defined as a triple  $(va(Q_d), s, t)$ , where*

$$va(Q_d) := a(Q) \times \{1, \dots, d-1\}$$

*is a finite set, and  $s, t : va(Q_d) \rightarrow v(Q_d)$  are maps defined by*

$$s((i \rightarrow j, r)) = (i, r) \quad t((i \rightarrow j, r)) = (j, r+1).$$

**Definition 2.14** (Virtual arrows of the quiver  $Q_{d, \text{str}}$ ). *For  $d \geq 2$ , the virtual arrows of the quiver  $Q_{d, \text{str}}$  is defined as a triple  $(va(Q_{d, \text{str}}), s, t)$ , where*

$$va(Q_{d, \text{str}}) := a(Q) \times \{2, \dots, d-1\}$$

*is a finite set, and  $s, t : va(Q_{d, \text{str}}) \rightarrow v(Q_{d, \text{str}})$  are maps defined by*

$$s((i \rightarrow j, r)) = (i, r) \quad t((i \rightarrow j, r)) = (j, r).$$

**Example 2.15.**



**Definition 2.16** (Euler form of  $R$ ). *Let  $R$  be a bounded quiver algebra defined in Definition 2.4. We denote*

$$\begin{aligned} v(R) &:= \{\text{vertices in } Q_d \text{ or } Q_{d,\text{str}}\}, \\ a(R) &:= \{\text{arrows in } Q_d \text{ or } Q_{d,\text{str}}\}, \\ va(R) &:= \{\text{virtual arrows in } Q_d \text{ or } Q_{d,\text{str}}\}. \end{aligned}$$

For two dimension vectors  $\mathbf{f}, \mathbf{g}$  of  $R$ , the Euler form  $\langle \mathbf{f}, \mathbf{g} \rangle_R$  is defined by

$$\langle \mathbf{f}, \mathbf{g} \rangle_R := \sum_{i \in v(R)} f_i g_i - \sum_{b \in a(R)} f_{s(b)} g_{t(b)} + \sum_{c \in va(R)} f_{s(c)} g_{t(c)}.$$

**2.4. Ext-vanishing properties.** We will show that some higher rank extension group are zero, which will be a key ingredient in the proofs of the next section.

For a bounded quiver algebra  $R$  defined in Definition 2.4, we have a standard resolution for every  $R$ -module  $T$ :

$$\begin{aligned} 0 \rightarrow \bigoplus_{c \in va(Q)} Re_{t(c)} \otimes_K e_{s(c)} T \rightarrow \bigoplus_{b \in a(Q)} Re_{t(b)} \otimes_K e_{s(b)} T \rightarrow \bigoplus_{i \in v(Q)} Re_i \otimes_K e_i T \rightarrow T \rightarrow 0 \\ r \otimes x \longmapsto \begin{pmatrix} rc_1 \otimes x + r \otimes b_1 x \\ -rc_2 \otimes x - r \otimes b_2 x \end{pmatrix} \qquad r \otimes x \longmapsto rx \\ r \otimes x \longmapsto rb \otimes x - r \otimes bx \end{aligned}$$

There are exactly two paths of length two from  $s(c)$  to  $t(c)$  for any virtual arrow  $c$ , which we denoted by  $b_1 c_1$  and  $b_2 c_2$  in the above. By definition, these paths are identified in  $R$ .

**Lemma 2.17.** *Let  $M, N \in \text{rep}(Q)$ .*

- (1)  $\text{gl. dim } R \leq 2$ ;
- (2) *The functor  $\Phi : \text{rep}(Q) \rightarrow \text{mod}(R)$  is exact and fully faithful;*
- (3)  *$\Phi$  maps projective module to projective module, and maps injective module to injective module;*
- (4)  $\text{Ext}_{KQ}^i(M, N) \cong \text{Ext}_R^i(\Phi(M), \Phi(N))$ ;
- (5)  $\text{proj. dim } \Phi(M) \leq 1, \text{inj. dim } \Phi(M) \leq 1$ ;

*Proof.*

For (1), this follows from the standard resolution.

For (2), it follows by direct inspection, see [7, Lemma 2.3].

For (3), we reduce to the case of indecomposable projective modules, and observe that

$$\Phi(P(i)) = P((i, 1)), \quad \Phi(I(i)) = I((i, d)).$$

For (4), it comes from the fact that  $\Phi$  is fully faithful and maps projective module to projective module.

For (5), notice that the minimal projective resolution of  $M$  is of length 1, and  $\Phi(-)$  sends the projective resolution of  $M$  to the projective resolution of  $\Phi(M)$  by (3), thus we get  $\text{proj. dim } \Phi(M) \leq 1$ . The injective dimension of  $\Phi(M)$  is computed in a similar way.  $\square$

The following key lemma will be crucial later.

**Lemma 2.18.** *Let  $X, S \in \text{rep}(Q)$  and  $V \subseteq \Phi(X), W \subseteq \Phi(S), T \in \text{mod}(R)$ . Then  $\text{Ext}_R^2(W, T) = 0$  and  $\text{Ext}_R^2(T, \Phi(X)/V) = 0$ .*

*Proof.* The short exact sequence

$$0 \longrightarrow W \longrightarrow \Phi(S) \longrightarrow \Phi(S)/W \longrightarrow 0$$

induces the long exact sequence

$$\cdots \longrightarrow \text{Ext}_R^2(\Phi(S), T) \longrightarrow \text{Ext}_R^2(W, T) \longrightarrow \text{Ext}_R^3(\Phi(S)/W, T) \longrightarrow \cdots.$$

By Lemma 2.17 (1) and (5),  $\text{Ext}_R^3(\Phi(S)/W, T)$  and  $\text{Ext}_R^2(\Phi(S), T)$  are both 0, so  $\text{Ext}_R^2(W, T) = 0$ .

Similarly, from the short exact sequence

$$0 \longrightarrow V \longrightarrow \Phi(X) \longrightarrow \Phi(X)/V \longrightarrow 0$$

we get the induced long exact sequence

$$\cdots \longrightarrow \text{Ext}_R^2(T, \Phi(X)) \longrightarrow \text{Ext}_R^2(T, \Phi(X)/V) \longrightarrow \text{Ext}_R^3(T, V) \longrightarrow \cdots,$$

so  $\text{Ext}_R^2(T, \Phi(X)/V) = 0$ .  $\square$

We will frequently use extension groups as well as long exact sequences, so we introduce some abbreviations. For  $Q$ -representations  $M, N$  and  $R$ -modules  $T, T'$ , we denote

$$\begin{aligned} [M, N]^i &:= \dim_K \text{Ext}_{KQ}^i(M, N) & [M, N] &:= \dim_K \text{Hom}_{KQ}(M, N) \\ [T, T']^i &:= \dim_K \text{Ext}_R^i(T, T') & [T, T'] &:= \dim_K \text{Hom}_R(T, T') \end{aligned}$$

and write the Euler form as

$$\langle T, T' \rangle_R := \sum_{i=0}^{\infty} (-1)^i [T, T']^i = [T, T'] - [T, T']^1 + [T, T']^2.$$

**Lemma 2.19** (Homological interpretation of the Euler form). *For two  $R$ -modules  $T, T'$ , we have*

$$\langle T, T' \rangle_R = \langle \underline{\dim} T, \underline{\dim} T' \rangle_R.$$

*Proof.* Compute  $\langle T, T' \rangle_R$  by applying the functor  $\text{Hom}_R(-, T')$  to the standard resolution of the  $R$ -module  $T$ .  $\square$



### 3. MAIN THEOREM

In this section we state and prove the main theorems, which are essential in Section 4 and 6.

Let  $\eta : 0 \longrightarrow X \xrightarrow{\iota} Y \xrightarrow{\pi} S \longrightarrow 0$  be a short exact sequence in  $\text{rep}(Q)$ . Consider the canonical **non-continuous** map

$$\Psi : \text{Gr}(\Phi(Y)) \longrightarrow \text{Gr}(\Phi(X)) \times \text{Gr}(\Phi(S)) \quad U \longmapsto ([\Phi(\iota)]^{-1}(U), [\Phi(\pi)](U)).$$

Denote the set

$$\text{Gr}(\Phi(Y))_{\mathbf{f}, \mathbf{g}} := \Psi^{-1} \left( \text{Gr}_{\mathbf{f}}(\Phi(X)) \times \text{Gr}_{\mathbf{g}}(\Phi(S)) \right)$$

and let  $\Psi_{\mathbf{f}, \mathbf{g}}$  be the map  $\Psi$  restricted to  $\text{Gr}(\Phi(Y))_{\mathbf{f}, \mathbf{g}}$ , i.e.,

$$\Psi_{\mathbf{f}, \mathbf{g}} : \text{Gr}(\Phi(Y))_{\mathbf{f}, \mathbf{g}} \longrightarrow \text{Gr}_{\mathbf{f}}(\Phi(X)) \times \text{Gr}_{\mathbf{g}}(\Phi(S)).$$

*Remark 3.1.* Even though  $\Psi$  is not continuous,  $\Psi_{\mathbf{f}, \mathbf{g}}$  is continuous. Moreover, for any dimension vectors  $\mathbf{f}, \mathbf{g}$ , the set

$$\text{Gr}(\Phi(Y))_{\geq \mathbf{f}, \leq \mathbf{g}} := \left\{ U \in \text{Gr}(\Phi(Y)) \mid \begin{array}{l} \underline{\dim}[\Phi(\iota)]^{-1}(U) \geq \mathbf{f} \\ \underline{\dim}[\Phi(\pi)](U) \leq \mathbf{g} \end{array} \right\}$$

is closed in  $\text{Gr}(\Phi(Y))$ . This gives us a filtration

$$0 = Z_0 \subset Z_1 \subset \cdots \subset Z_d = \text{Gr}_h(\Phi(Y))$$

with  $Z_i$  closed and  $Z_{i+1} \setminus Z_i$  isomorphic to  $\text{Gr}(\Phi(Y))_{\mathbf{f}, \mathbf{g}}$  for some  $\mathbf{f}, \mathbf{g}$ . Therefore, from the affine pavings of  $\text{Gr}(\Phi(Y))_{\mathbf{f}, \mathbf{g}}$  (for every  $\mathbf{f}, \mathbf{g}$ ) one can construct one affine paving of  $\text{Gr}_h(\Phi(Y))$ .

**Theorem 3.2.** *If  $\eta$  splits, then  $\Psi$  is surjective. Moreover, if  $[S, X]^1 = 0$ , then  $\Psi_{\mathbf{f}, \mathbf{g}}$  is a Zariski-locally trivial affine bundle of rank  $\langle \mathbf{g}, \underline{\dim} \Phi(X) - \mathbf{f} \rangle_R$ .*

**Theorem 3.3** (Generalizes [6, Theorem 32]). *When  $\eta$  does not split and  $[S, X]^1 = 1$ ,*

$$\text{Im } \Psi_{\mathbf{f}, \mathbf{g}} = \left( \text{Gr}_{\mathbf{f}}(\Phi(X)) \times \text{Gr}_{\mathbf{g}}(\Phi(S)) \right) \setminus \left( \text{Gr}_{\mathbf{f}}(\Phi(X_S)) \times \text{Gr}_{\mathbf{g} - \underline{\dim} \Phi(S^X)}(\Phi(S/S^X)) \right)$$

where

$$X_S := \max \{ M \subseteq X \mid [S, X/M]^1 = 1 \} \subseteq X,$$

$$S^X := \max \{ M \subseteq S \mid [M, X]^1 = 1 \} \subseteq S.$$

Moreover,  $\Psi_{\mathbf{f}, \mathbf{g}}$  is a Zarisky-locally trivial affine bundle of rank  $\langle \mathbf{g}, \underline{\dim} \Phi(X) - \mathbf{f} \rangle_R$  over  $\text{Im } \Psi_{\mathbf{f}, \mathbf{g}}$ .

We will spend the rest of the section proving these theorems. We investigate the image as well as the fiber of  $\Psi$  respectively.

**Lemma 3.4** (Follows [6, Lemma 21]). *The element  $(V, W) \in \text{Gr}(\Phi(X)) \times \text{Gr}(\Phi(S))$  lies in the image of  $\Psi$  if and only if the canonical map  $\text{Ext}^1(\Phi(S), \Phi(X)) \longrightarrow \text{Ext}^1(W, \Phi(X)/V)$  maps  $\eta$  to 0.*

*Proof.* The canonical map is defined as follows:

$$\begin{array}{ccccccc}
\eta \in \text{Ext}^1(\Phi(S), \Phi(X)) & 0 \longrightarrow & \Phi(X) & \longrightarrow & \Phi(Y) & \xrightarrow{\Phi(\pi)} & \Phi(S) \longrightarrow 0 \\
\downarrow & & \parallel & & \uparrow & & \uparrow \\
& \text{Ext}^1(W, \Phi(X)) & 0 \longrightarrow & \Phi(X) & \longrightarrow & \pi^{-1}(W) & \longrightarrow W \longrightarrow 0 \\
\downarrow & & \downarrow & & \downarrow & & \parallel \\
\bar{\eta} \in \text{Ext}^1(W, \Phi(X)/V) & 0 \longrightarrow & \Phi(X)/V & \longrightarrow & \pi^{-1}(W)/V & \longrightarrow & W \longrightarrow 0
\end{array}$$

so  $\bar{\eta} = 0$  if and only if the last short exact sequence splits, that means, there exists a submodule  $U \subseteq \Phi(Y)$ , such that  $\Phi(\pi)(U) = W$  and  $U \cap \Phi(X) = V$ .  $\square$

**Corollary 3.5.** *Resume the notations of Lemma 3.4 When  $\eta$  splits, then  $\Psi$  is surjective.*

**Lemma 3.6.** *The canonical map  $\text{Ext}^1(\Phi(S), \Phi(X)) \rightarrow \text{Ext}^1(W, \Phi(X)/V)$  is surjective.*

*Proof.* By using the long exact sequence of extension groups and the fact that  $\text{Ext}^2(\Phi(S)/W, \Phi(X)) = 0$  and  $\text{Ext}^2(W, V) = 0$  by Lemma 2.18, the maps

$$\text{Ext}^1(\Phi(S), \Phi(X)) \rightarrow \text{Ext}^1(W, \Phi(X)) \quad \text{Ext}^1(W, \Phi(X)) \rightarrow \text{Ext}^1(W, \Phi(X)/V)$$

are both surjective. Thus the composition is also surjective.  $\square$

**Corollary 3.7.** *Let  $W \subseteq \Phi(S), V \subseteq \Phi(X)$  be  $R$ -submodules, then*

$$[W, \Phi(X)/V]^1 \leq [\Phi(S), \Phi(X)]^1 = [S, X]^1.$$

*In particular, when  $[S, X]^1 = 1$ , we get  $[W, \Phi(X)/V]^1 = 0$  or  $1$ ; when  $\eta$  generates  $\text{Ext}^1(S, X)$ , we get*

$$(V, W) \in \text{Im } \Psi \iff [W, \Phi(X)/V]^1 = 0.$$

In the case where  $\eta$  generates  $\text{Ext}^1(S, X)$ , we want to describe  $\text{Im } \Psi$  more precisely. For this reason we need to introduce two new  $R$ -modules:

$$\widetilde{X}_S := \max \{V \subseteq \Phi(X) \mid [\Phi(S), \Phi(X)/V]^1 = 1\} \subseteq \Phi(X),$$

$$\widetilde{S}^X := \max \{W \subseteq \Phi(S) \mid [W, \Phi(X)]^1 = 1\} \subseteq \Phi(S).$$

$\widetilde{X}_S$  and  $\widetilde{S}^X$  are well-defined because of the following lemma:

**Lemma 3.8** (Follows [6, Lemma 27]).

- (i) *Let  $V, V' \subset \Phi(X)$  such that  $[\Phi(S), \Phi(X)/V]^1 = [\Phi(S), \Phi(X)/V']^1 = 1$ . Then  $[\Phi(S), \Phi(X)/(V + V')]^1 = 1$ .*
- (ii) *Let  $W, W' \subset \Phi(S)$  such that  $[W, \Phi(X)]^1 = [W', \Phi(X)]^1 = 1$ . Then  $[W \cap W', \Phi(X)]^1 = 1$ .*

*Proof.* We only prove (i). (ii) is similar.

From the short exact sequence

$$0 \longrightarrow \Phi(X)/(V \cap V') \longrightarrow \Phi(X)/V \oplus \Phi(X)/V' \longrightarrow \Phi(X)/(V + V') \longrightarrow 0,$$

we get the long exact sequence

$$\cdots \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V \cap V'}\right) \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V}\right) \oplus \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V'}\right) \rightarrow \text{Ext}^1\left(\Phi(S), \frac{\Phi(X)}{V + V'}\right) \rightarrow \cdots.$$

By Corollary 3.7,  $[\Phi(S), \Phi(X)/(V \cap V')]^1 \leq 1$ ,  $[\Phi(S), \Phi(X)/(V + V')]^1 \leq 1$ , and this forces  $[\Phi(S), \Phi(X)/(V + V')]^1 = 1$ .  $\square$

**Lemma 3.9** (Follows [6, Lemma 31(1)(2)], with the same proof). *Let  $\tau$  be the Auslander–Reiten translation.*

*Let  $f : X \rightarrow \tau S$  be a non-zero morphism,<sup>1</sup> then  $X_S = \ker(f)$ ;*

*also,  $\Phi(f) : \Phi(X) \rightarrow \Phi(\tau S)$  is a non-zero morphism,  $\widetilde{X}_S = \ker(\Phi(f))$ .*

*Proof.* For any  $M \subseteq X$ , we have

$$\begin{aligned} \text{Ext}^1(S, X/M)^\vee &\cong \overline{\text{Hom}}(X/M, \tau S) \\ &\cong \{g \in \text{Hom}(X, \tau S) \mid g|_M = 0\} \\ &\cong \begin{cases} \mathbb{C}, & M \subseteq \ker f \\ 0, & M \not\subseteq \ker f. \end{cases} \end{aligned}$$

so  $[S, X/M]^1 = 1$  exactly when  $M \subseteq \ker f$ . Thus  $X_S = \ker f$ .

For  $\Phi(f)$  it is similar. For any  $V \subseteq \Phi(X)$ , we have

$$\begin{aligned} \text{Ext}^1(\Phi(S), \Phi(X)/V)^\vee &\cong \overline{\text{Hom}}(\Phi(X)/V, \tau\Phi(S)) \\ &\cong \overline{\text{Hom}}(\Phi(X)/V, \Phi(\tau S)) \\ &\cong \{g \in \text{Hom}(\Phi(X), \Phi(\tau S)) \mid g|_V = 0\} \\ &\cong \begin{cases} \mathbb{C}, & V \subseteq \ker \Phi(f) \\ 0, & V \not\subseteq \ker \Phi(f). \end{cases} \end{aligned}$$

so  $[\Phi(S), \Phi(X)/V]^1 = 1$  exactly when  $V \subseteq \ker \Phi(f)$ . Thus  $\widetilde{X}_S = \ker(\Phi(f))$ .  $\square$

**Corollary 3.10.**  $\widetilde{X}_S = \Phi(X_S)$ . (since  $\widetilde{X}_S = \ker(\Phi(f)) = \Phi(\ker(f)) = \Phi(X_S)$ )

By a dual argument, one can show that  $\widetilde{S}^X = \Phi(S^X)$ .

**Lemma 3.11** (Follows [6, Lemma 31(6)]). *For  $V \subseteq \Phi(X)$  and  $W \subseteq \Phi(S)$ , we have*

$$[W, \Phi(X)/V]^1 = 0 \iff V \not\subseteq \Phi(X_S) \text{ or } W \not\supseteq \Phi(S^X).$$

*Proof.*  $\Leftarrow$ : Without loss of generality suppose  $V \not\subseteq \Phi(X_S)$ , then

$$V \not\subseteq \Phi(X_S) \iff [\Phi(S), \Phi(X)/V]^1 = 0 \Rightarrow [W, \Phi(X)/V]^1 = 0.$$

$\Rightarrow$ : If not, then  $V \subseteq \Phi(X_S)$  and  $W \supsetneq \Phi(S^X)$ , and<sup>2</sup>

$$[W, \Phi(X)/V]^1 \geq [\Phi(S^X), \Phi(X)/\Phi(X_S)]^1 = [S^X, X/X_S]^1 = 1. \quad \square$$

**Corollary 3.12.** *When  $\eta$  generates  $\text{Ext}^1(S, X)$ , we have*

$$\text{Im } \Psi_{f,g} = \left( \text{Gr}_f(\Phi(X)) \times \text{Gr}_g(\Phi(S)) \right) \setminus \left( \text{Gr}_f(\Phi(X_S)) \times \text{Gr}_{g-\underline{\dim} \Phi(S^X)}(\Phi(S/S^X)) \right).$$

<sup>1</sup>Since  $X$  is not injective,  $[X, \tau S] = [S, X]^1 = 1$ ,  $f$  is uniquely determined up to a constant.


<sup>2</sup> $[S^X, X/X_S]^1 = 1$  follows from [6, Lemma 31(5)].

**Lemma 3.13.** *For  $(V, W) \in \text{Im } \Psi$ , the preimage of  $(V, W)$  is a torsor of  $\text{Hom}_R(W, \Phi(X)/V)$ . Hence, there is a non-canonical isomorphism*

$$\Psi^{-1}((V, W)) \cong \text{Hom}_R(W, \Phi(X)/V).$$

*Proof.* Recall the commutative diagram

$$\begin{array}{ccccccc}
 \eta \in \text{Ext}^1(\Phi(S), \Phi(X)) & 0 & \longrightarrow & \Phi(X) & \longrightarrow & \Phi(Y) & \xrightarrow{\Phi(\pi)} \Phi(S) \longrightarrow 0 \\
 \downarrow & & & \parallel & & \uparrow & \uparrow \\
 & \text{Ext}^1(W, \Phi(X)) & 0 & \longrightarrow & \Phi(X) & \longrightarrow & \pi^{-1}(W) \longrightarrow W \longrightarrow 0 \\
 \downarrow & \downarrow & & \downarrow & & \downarrow & \parallel \\
 \bar{\eta} \in \text{Ext}^1(W, \Phi(X)/V) & 0 & \longrightarrow & \Phi(X)/V & \xrightarrow{\iota} & \pi^{-1}(W)/V & \xrightarrow{\pi'} W \longrightarrow 0
 \end{array}$$



When  $(V, W) \in \text{Im } \Psi$ ,  $\bar{\eta}$  is split, and each split morphism  $\theta$  give us an element in  $\Psi^{-1}((V, W))$ . If we fix one split morphism  $\theta_0$ , then the other split morphisms are all of the form  $\theta_0 + \iota \circ f$  where  $f \in \text{Hom}_R(W, \Phi(X)/V)$  (and this form is unique). So

$$\Psi^{-1}((V, W)) \cong \{\theta : \text{split morphism}\} \cong \text{Hom}_R(W, \Phi(X)/V). \quad \square$$

*Remark 3.14.* Any point  $(V, W) \in \text{Im } \Psi_{f,g}$  can be also viewed as a morphism

$$f : \text{Spec } K \longrightarrow \text{Im } \Psi_{f,g} \subseteq \text{Gr}_f(\Phi(X)) \times \text{Gr}_g(\Phi(S))$$

where Grassmannian are viewed as moduli spaces over  $K$ . Essentially by replacing  $\text{Spec } K$  by any locally closed reduced subscheme  $\text{Spec } A$  of  $\text{Im } \Psi_{f,g}$  in Lemma 3.13, we can run the machinery of algebraic geometry, and mimic the proof of [6, Theorem 24] to show that  $\Psi_{f,g}$  is a Zariski-locally trivial affine bundle over  $\text{Im } \Psi_{f,g}$  when  $\eta$  generates  $\text{Ext}^1(S, X)$ . Roughly, there are 4 steps:

1. Realise Grassmannians as representable functors, and replace  $K$ -modules by  $A$ -modules;
2. Verify that  $\Psi_{f,g}^{-1}(\text{Spec } A)$  is a  $\text{Hom}_A(\mathcal{W}, \Phi(X)_A/\mathcal{V})$ -torsor, where

$$(\mathcal{V}, \mathcal{W}) \in \text{Gr}_f(\Phi(X))(A) \times \text{Gr}_g(\Phi(S))(A)$$

corresponds to the immersion  $\text{Spec } A \hookrightarrow \text{Im } \Psi_{f,g}$ ;

3. Verify that  $\text{Hom}_A(\mathcal{W}, \Phi(X)_A/\mathcal{V})$  is a vector bundle over  $\text{Spec } A$  of constant dimension  $\langle f, \underline{\dim} \Phi(X) - g \rangle_R$ ;
4. Find a section of  $\Psi_{f,g}^{-1}(\text{Spec } A) \longrightarrow \text{Spec } A$ , which is essentially the splitting  $\theta$  in [6, Lemma 22].

*Proof of Theorem 3.2 and 3.3.* We have already computed  $\text{Im } \Psi$  in Corollary 3.5 and 3.12. In both cases  $\eta$  generates  $\text{Ext}^1(S, X)$ , so by Corollary 3.7 we get

$$\begin{aligned}
 (V, W) \in \text{Im } \Psi_{f,g} &\iff [W, \Phi(X)/V]^1 = 0 \\
 &\implies [W, \Phi(X)/V] = \langle W, \Phi(X)/V \rangle_R = \langle f, \underline{\dim} \Phi(X) - g \rangle_R.
 \end{aligned}$$

From Remark 3.14,  $\Psi_{f,g}$  is a Zariski-locally trivial affine bundle.  $\square$

**Lemma 4.2** (Follows [7, Lemma 2.22]). *For an indecomposable representation  $M \in \text{rep}(Q)$  with  $\text{ord}(M) \leq 2$ , the variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  is either empty or a direct product of some copies of  $\mathbb{P}^1$ . Especially, the partial flag variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  has an affine paving.*

*Proof.* For every  $i \in v(Q)$ ,  $\dim_K M_i \leq 2$ . Since  $Q$  is a tree and  $M$  is indecomposable, for every  $b \in a(Q)$  satisfying  $\dim_K M_{s(b)} = \dim_K M_{t(b)} = 2$ , the map  $M_{s(b)} \rightarrow M_{t(b)}$  is an isomorphism. Therefore, when  $\text{Gr}_{\mathbf{f}}(\Phi(M)) \neq \emptyset$ ,<sup>3</sup> we get the natural embedding

$$\text{Gr}_{\mathbf{f}}(\Phi(M)) \longrightarrow \prod_{\substack{i \in v(Q) \text{ s.t.} \\ \dim_K M_i = 2 \\ \mathbf{f}_{(i,r)} = 1 \text{ for some } r}} \mathbb{P}^1,$$

and the information of non-vertical arrows in the extended quiver (see Example 2.3) just reduce the number of  $\mathbb{P}^1$ . Precisely, one need to carefully discuss three cases of  $M_i \rightarrow M_j$ :

$$K \hookrightarrow K^2 \quad K^2 \twoheadrightarrow K \quad \text{and} \quad K^2 \xrightarrow{\cong} K^2. \quad \square$$

*Proof of Theorem 4.1, assuming Theorem 5.1.* First of all, for any indecomposable representation  $M \in \text{rep}(Q)$  we obtain an affine paving. This follows from Theorem 5.1 when  $\text{ord}(M) > 2$ , and follows from Lemma 4.2 when  $\text{ord}(M) \leq 2$ .

The general case follows by induction on the dimension vector. The indecomposable representations  $\{N_i\}_{i \in Q_0}$  of quiver  $Q$  can be ordered such that  $[N_i, N_j] = 0$  for all  $i > j$ . Therefore, every non-indecomposable representation  $M$  can be decomposed as the direct sum of two nonzero representations  $M_1, M_2$  satisfying  $[M_2, M_1]^1 = 0$ . By applying Theorem 3.2 to the short exact sequence

$$0 \rightarrow M_1 \rightarrow M \rightarrow M_2 \rightarrow 0,$$

we get an affine paving from the affine pavings of  $M_1$  and  $M_2$ , see Remark 3.1.  $\square$

*Remark 4.3.* By the same technique one can show that, for Dynkin quiver  $Q$  and any representation  $M$  with  $\max_{i \in v(Q)} \dim_K M_i \leq 2$ , the variety  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  has an affine paving.

This result does not depend on Theorem 5.1.

## 5. AFFINE PAVING FOR BIG ORDER REPRESENTATIONS

This section aims to establish the following theorem:

**Theorem 5.1.** *Suppose  $Q$  is of Dynkin type. For any indecomposable representation  $M \in \text{rep}(Q)$  with  $\text{ord}(M) > 2$ , the (strict) partial flag variety  $\text{Gr}(\Phi(M))$  has an affine paving.*


When the quiver  $Q$  is of type  $A$  or  $D$ , Theorem 5.1 is trivially true since no indecomposable representation can have order bigger than two. So we only concentrate on type  $E$ .

The idea of the proof is as follows. For any indecomposable representation  $Y$  with  $\text{ord}(Y) > 2$ , we put  $Y$  into a short exact sequence

$$\eta : 0 \rightarrow X \rightarrow Y \rightarrow S \rightarrow 0$$

fulfilling the assumptions of Theorem 3.3, and then  $\text{Gr}(\Phi(Y))$  has an affine paving if  $\text{Im } \Psi$  has. If additionally the map  $X \hookrightarrow Y$  is a minimal sectional mono, then  $\text{Im } \Psi_{\mathbf{f}, \mathbf{g}}$  can be written as the product space, which makes  $\text{Im } \Psi$  easier to understand.

<sup>3</sup>This condition imposes very strong restrictions on  $\mathbf{f}$ .



Here, we recall two results from [6]. ???

**Definition 5.2** (Sectional morphism). *Let  $Q$  be a quiver of Dynkin type, and  $M, N \in \text{rep}(Q)$  be two indecomposable representations of  $Q$ . A morphism  $f \in \text{Hom}_{KQ}(M, N)$  is called sectional if  $f$  can be written as the composition*

$$f : M = X_0 \xrightarrow{f_1} X_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{t-1}} X_{t-1} \xrightarrow{f_t} X_t = N$$

where  $f_i \in \text{Hom}_{KQ}(X_{i-1}, X_i)$  are irreducible morphisms between indecomposable representations, and  $\tau X_{i+2} \not\cong X_i$  for any suitable  $i$ .

A sectional morphism  $f \in \text{Hom}_{KQ}(M, N)$  is called as a sectional mono if  $f$  is injective; a sectional mono is called minimal if  $f_t \circ \cdots \circ f_{i+1} : X_i \rightarrow N$  are surjective for any  $i \in \{1, 2, \dots, t\}$ .

**Lemma 5.3** (Happel–Ringel). *Let  $M$  and  $N$  be two indecomposable  $Q$ -representations. Any sectional morphism  $f \in \text{Hom}_{KQ}(M, N)$  is either surjective or injective.*

*Proof.* When  $Q$  is a quiver without oriented cycles, then  $[N, M]^1 \leq [M, \tau N] = 0$ , thus by [6, Lemma 7] we get the result; when  $Q$  is of type  $\tilde{A}$ , the result comes from [6, Lemma 51].  $\square$

The next two lemmas tell us the existence of the desired short exact sequence.

**Lemma 5.4.** *For every indecomposable representation  $Y$  of type  $E$  with  $\text{ord}(Y) > 2$ , there is a minimal sectional mono  $f : X \rightarrow Y$  such that  $\underline{\dim}(X) < \underline{\dim}(Y)$ .*

*Proof.* Suppose that  $Y$  is an indecomposable representation of type  $E$  such that  $\text{ord}(Y) > 2$ , then  $s_{P(e)}(Y) = \text{ord}(Y) > 2$ . By direct inspection on every table on Figure 2, one sees that there is a sectional path (which can hence be chosen to be minimal) ending in  $[Y]$  and starting at some  $[M]$  with  $s_{P(e)}(M) < s_{P(e)}(Y)$ , so  $\underline{\dim}(M) < \underline{\dim}(Y)$ . Write this sectional path as  $M = X_0 \rightarrow \cdots \rightarrow X_t = Y$ , and take the maximal  $i$  such that  $\underline{\dim}(X_i) < \underline{\dim}(Y)$ . Take  $X = X_i$ . By Lemma 5.4, the sectional mono  $f : X \rightarrow Y$  is injective. It follows that there exists a minimal sectional mono from  $X$  to  $Y$ .  $\square$

**Remark 5.5.** The chosen minimal sectional monos are presented in Figure 3, where the arrows ends in  $[Y]$  and starts at  $[X]$ . From those tables it is clear that in type  $E_6$  the only indecomposable representations with order strictly bigger than two are those lying in the  $\tau^-$ -orbit of  $P(e)$ . In particular, they are the middle term of an almost split sequence,




FIGURE 3. Minimal sectional monos

and hence in type  $E_6$  Theorem 4.1 is clear. Additionally, the tables demonstrate that the condition  $\text{ord}(Y) > 2$  in the lemma is essential and cannot be omitted.

**Lemma 5.6.** *Let  $\iota : X \hookrightarrow Y$  be a minimal sectional mono, and  $S := Y/X$  be the quotient. Then  $S$  is indecomposable and rigid, the short exact sequence*

$$\eta : 0 \longrightarrow X \longrightarrow Y \longrightarrow S \longrightarrow 0$$

*generates  $\text{Ext}^1(S, X)$ , and  $S^X = S$ . Moreover, one gets*

$$\begin{cases} [X, X] = [Y, Y] = [S, S] = 1 \\ [X, Y] = [Y, S] = [S, X]^1 = 1 \\ [X, S] = [Y, X] = [S, Y] = [S, X] = 0 \\ [X, X]^1 = [Y, Y]^1 = [S, S]^1 = 0 \\ [X, Y]^1 = [Y, S]^1 = [X, S]^1 = [Y, X]^1 = [S, Y]^1 = 0 \end{cases}$$

*Proof.* Since every indecomposable representation of Dynkin quiver is a brick and rigid, we get  $[X, X] = [Y, Y] = 1$  and  $[X, X]^1 = [Y, Y]^1 = 0$ . Since  $Q$  is a of type  $E$ , By the



definition of minimal sectional mono, we get  $[Y, X]^1 = 0$ . Then by Unger's lemma [6] one gets that  $S$  is indecomposable and  $[S, X]^1 = 1$ . In particular  $\eta$  is generating. Since  $\iota$  is minimal, the pullback  $j^*(\eta)$  by any (proper) monomorphism  $j : N \hookrightarrow S$  splits; it follows that  $S^X = S$ . By applying the functors  $[Y, -], [-, S], [X, -], [-, X], [-, Y]$  to the short exact sequence  $\eta$  we get all the dimensions.  $\square$

**Lemma 5.7** (Follows [6, Lemma 36], with the same proof). *The notation  $X, Y$  and  $S$  remains as in the previous lemma. Let  $X_1$  be the indecomposable module in the sectional path  $X = X_0 \rightarrow \cdots \rightarrow X_t = Y$ , and let  $E \rightarrow X$  be the minimal right almost split morphism ending in  $X$ . We have the decomposition  $E = E' \oplus \tau X_1$  for some  $Q$ -representation  $E'$ . When  $Y$  is not projective,  $X_S$  is isomorphic to  $\ker(E \rightarrow \tau Y) \cong E' \oplus \ker(\tau X_1 \rightarrow \tau Y)$ ; when  $Y$  is projective,  $X_S \cong E$ .*

**Corollary 5.8.** *When  $\iota : X \rightarrow Y$  is irreducible monomorphism, the representation  $X_S$  is either 0 or an indecomposable representation with property that  $X_S \rightarrow X$  is also an irreducible monomorphism.*

*Proof.* If  $\iota$  is an irreducible monomorphism, it follows that  $X_1 = Y$ , and  $X$  does not belong to the  $\tau^-$ -orbit of  $P(e)$ . It follows that the representation  $E$  has at most two indecomposable components. By Lemma 5.7, we obtain  $X_S \cong E'$ :

- If  $Y$  is not projective, then  $\tau X_1 = \tau Y$ , so  $X_S \cong E' \oplus \ker(\tau X_1 \rightarrow \tau Y) \cong E'$ ;
- If  $Y$  is projective, then  $\tau X_1 = 0$ , so  $X_S \cong E \cong E'$ .

In both cases,  $E' \subset E$  is either 0 or indecomposable. Since  $E \rightarrow X$  is the minimal right almost split morphism,  $E' \rightarrow X$  is an irreducible monomorphism.  $\square$

For convenience, we simplify the notations: write  $\text{Gr}_{\mathbf{f}}(\Phi(M))$  as  $\text{Gr}(M)$ ,  $\text{Gr}_{\mathbf{f}}(\Phi(M)) \setminus \text{Gr}_{\mathbf{f}}(\Phi(N))$  as  $U(M, N)$ , where we omit subscripts which indicate the dimension vectors.

**Lemma 5.9** (Follows [6, Theorem 59]). *Let  $f : X \hookrightarrow Y$  be a minimal sectional mono and  $S := Y/X$  be the quotient. When  $X_S = F \oplus T$  with  $F$  and  $T$  nonzero indecomposable,  $F \hookrightarrow X$  irreducible and  $T \hookrightarrow X/F$  sectional mono, we have*

$$U(X, X_S) \longrightarrow \text{Gr}(F) \times U(X/F, T) \quad \text{or} \quad U(F, F_{X/F})$$

as a (Zariski) locally-trivial affine bundle.

*Proof.* We have two short exact sequence satisfying the conditions in 3.3:

$$\begin{array}{ccccccc} \eta : & 0 & \longrightarrow & F & \longrightarrow & X & \xrightarrow{\pi} X/F \longrightarrow 0 \\ \xi : & 0 & \longrightarrow & T & \longrightarrow & X/F & \xrightarrow{\pi'} X/X_S \longrightarrow 0. \end{array}$$

Let  $N \in \text{Gr}(X)$  be a subrepresentation, it is obvious that  $N \in \text{Gr}(X_S) \iff \pi' \circ \pi(N) = 0$ , so

$$\begin{aligned} N \in U(X, X_S) &\iff \pi' \circ \pi(N) \neq 0 \\ &\iff \pi(N) \notin \text{Gr}(T) \\ &\iff \pi(N) \in U(X/F, T) \\ &\iff \Psi_{\eta}(N) \in \text{Gr}(F) \times U(X/F, T). \end{aligned}$$

Thus the (Zariski) locally-trivial affine bundle map

$$U(X, F) \longrightarrow \text{Gr}(F) \times \text{Gr}(X/F)$$

restricted to the (Zariski) locally-trivial affine bundle map

$$U(X, X_S) \longrightarrow \text{Gr}(F) \times U(X/F, T). \quad \square$$

We conclude Theorem 3.3, Lemma 3.9, 5.6, 5.7, 5.9 in the following proposition.

**Proposition 5.10.** *Let  $f : X \hookrightarrow Y$  be a minimal sectional mono and  $S := Y/X$  be the quotient. One get (Zariski) locally-trivial affine maps*

$$\begin{aligned} \text{Gr}(Y) &\longrightarrow \text{Gr}(X) \times \text{Gr}(S) \quad \text{or} \quad U(X, X_S), \\ U(Y, X) &\longrightarrow \text{Gr}(X) \times \text{Gr}(S) \quad \text{or} \quad U(X, X_S), \end{aligned}$$

where

$$\begin{aligned} X_S &:= \max \{M \subseteq X \mid [S, X/M]^1 = 1\} \\ &= \ker(X \longrightarrow \tau S) \\ &= \begin{cases} E' \oplus \ker(\tau X_1 \longrightarrow \tau Y) & Y \text{ not projective,} \\ E & Y \text{ projective.} \end{cases} \end{aligned}$$

Moreover, when  $X_S = F \oplus T$  with  $F$  and  $T$  nonzero indecomposable,  $F \hookrightarrow X$  irreducible and  $T \hookrightarrow X/F$  sectional mono, one get a (Zariski) locally-trivial affine map

$$U(X, X_S) \longrightarrow \text{Gr}(F) \times U(X/F, T) \quad \text{or} \quad U(F, F_{X/F}).$$

With Proposition 5.10 in hand, we prove Theorem 5.1 by cases, which turns out to be a pure combinatorial problem.

*Proof of Theorem 5.1.* When the minimal sectional mono  $X \longrightarrow Y$  is irreducible, we use Theorem 3.3 to get morphism

$$\text{Gr}(Y) \longrightarrow \text{Gr}(X) \times \text{Gr}(S) \quad \text{or} \quad U(X, X_S).$$

By observation of Figure 3,  $s_{P(e)}(S) = s_{P(e)}(Y) - s_{P(e)}(X)$  is smaller or equal to 2, so by Lemma 4.2  $\text{Gr}(S)$  has the affine paving property. Let  $Y' := X$ ,  $X' := X_S$ ,  $S' := Y'/X'$ , we again use Theorem 3.3 to get (Zariski) locally-trivial affine maps

$$\begin{aligned} \text{Gr}(X) &\longrightarrow \text{Gr}(X') \times \text{Gr}(S') \quad \text{or} \quad U(X', X'_{S'}) \\ U(X, X_S) &\longrightarrow \text{Gr}(X') \times \text{Gr}(S') \quad \text{or} \quad U(X', X'_{S'}). \end{aligned}$$

Luckily  $s_{P(e)}(S')$  is still smaller or equal to 2. We can continue this process until the order of representations are small enough. This solved all the cases for type  $E_6$ ,  $E_7$ , and the rests are exception cases in type  $E_8$ , as shown in Figure 3. These exceptional cases are addressed in a similar manner, although the argument becomes more intricate.

*Remark 5.11.* We can not copy everything in [6, Lemma 56], sometimes it would happen that  $X_S = F \oplus T$  with  $F$  and  $T$  indecomposable,  $F \hookrightarrow X$  is irreducible but  $T \longrightarrow X/F$  is not a sectional mono. Even when  $X_S$  is indecomposable, the map  $X_S \longrightarrow X$  can be not a sectional mono.

For example, take the quiver of type  $E_7$ :

$$\begin{array}{ccccccc} & & & & \bullet & & \\ & & & & \downarrow & & \\ \bullet & \longrightarrow & \bullet & \longrightarrow & \bullet & \longrightarrow & \bullet \longleftarrow \bullet \longleftarrow \bullet \end{array}$$

take  $Y = {}_{111111}^0$ ,  $X = {}_{111110}^0$ , then  $X_S = {}_{111100}^0$ , the map  $X_S \longrightarrow X$  is not a sectional mono.

Luckily, we can avoid this bad situation by carefully choosing the minimal sectional mono  $X \longrightarrow Y$ . The minimal sectional monos we chose are presented in Figure 3. We will write down the induction process in detail for some examples.



FIGURE 4

**Example 5.12.** In the case of Figure 8(a), if  $X_1 \longrightarrow Y$  is injective, then we obtain some (Zariski) locally-trivial affine maps

$$\begin{aligned} \mathrm{Gr}(Y) &\longrightarrow \mathrm{Gr}(X_1) \times \mathrm{Gr}(Y/X_1) && \text{or } U(X_1, X) \\ \mathrm{Gr}(X_1) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(X_1/X) && \text{or } U(X, A) \\ U(X_1, X) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(X_1/X) && \text{or } U(X, A) \\ U(X, A) &\longrightarrow \mathrm{Gr}(A) \times \mathrm{Gr}(X/A) && \text{or } U(A, -).^4 \end{aligned}$$

When  $X_1 \longrightarrow Y$  is not injective, we get

$$\begin{aligned} \mathrm{Gr}(Y) &\longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(X/Y) && \text{or } U(X, A) \\ U(X, A) &\longrightarrow \mathrm{Gr}(A) \times \mathrm{Gr}(X/A) && \text{or } U(A, -). \end{aligned}$$

These maps give the variety  $\mathrm{Gr}(Y)$  an affine paving from bottom to top.




FIGURE 5

**Example 5.13.** In Figure 8(b), we would like to prove that  $\mathrm{Gr}(Y)$  has the affine paving property. We have

$$\mathrm{Gr}(Y) \longrightarrow \mathrm{Gr}(X) \times \mathrm{Gr}(Y/X) \quad \text{or } U(X, A).$$

<sup>4</sup> $\mathrm{Gr}(A)$  is empty or a singleton, so is  $U(A, -)$ , no matter what representation is in the dash mark.

When the map  $M \rightarrow X$  is not monomorphism, we get

$$U(X, A) \rightarrow \text{Gr}(A) \times \text{Gr}(X/A) \quad \text{or} \quad U(A, -);$$

when the map  $M \rightarrow X$  is monomorphism, we get

$$\begin{aligned} U(X, A) &= U(X, M) \bigsqcup U(M, A) \\ U(X, M) &\rightarrow \text{Gr}(M) \times \text{Gr}(X/M) \quad \text{or} \quad U(M, A \oplus B) \\ U(M, A) &\rightarrow \text{Gr}(A) \times \text{Gr}(M/A) \quad \text{or} \quad U(A, -) \\ U(M, A \oplus B) &\rightarrow \begin{cases} \text{Gr}(A) \times \text{Gr}(M/A) & \text{or} \quad U(A, -), & B = 0, \\ \text{Gr}(A) \times U(M/A, B) & \text{or} \quad U(A, -), & B \neq 0. \end{cases} \end{aligned}$$

Since the order of  $X, Y/X, A, X/A, M, X/M, M/A$  are smaller or equal to 2 as well as  $\text{ord}(A) = \text{ord}(M/A) = 1$ , the induction process stops, we get an affine paving of  $\text{Gr}(Y)$ .




FIGURE 6

**Example 5.14.** In the case of Figure 8(c), we have

$$\text{Gr}(Y) \rightarrow \text{Gr}(X) \times \text{Gr}(Y/X) \quad \text{or} \quad U(X, A).$$

If  $A = 0$ , we're done; if not, we decompose  $A \rightarrow Y$  as compositions of minimal sectional monos:

Case 1:  $M \rightarrow X$  is not injective, then

$$\begin{aligned} U(X, A) &= U(X, N) \bigsqcup U(N, A) \\ U(X, N) &\rightarrow \text{Gr}(N) \times \text{Gr}(X/N) \quad \text{or} \quad U(N, -) \end{aligned}$$

Case 2:  $M \rightarrow X$  is injective, then

$$\begin{aligned} U(X, A) &= U(X, M) \bigsqcup U(M, N) \bigsqcup U(N, A) \\ U(X, M) &\rightarrow \text{Gr}(M) \times \text{Gr}(X/M) \quad \text{or} \quad U(M, N \oplus T) \\ U(M, N) &\rightarrow \text{Gr}(N) \times \text{Gr}(M/N) \quad \text{or} \quad U(N, -), \end{aligned}$$

Since  $M \rightarrow X$  is injective, we get  $B = 0$ , thus  $C = 0$  also, and then the sectional map  $T \rightarrow M/N$  is injective. We get

$$U(M, N \oplus T) \rightarrow \text{Gr}(N) \times U(M/N, T) \quad \text{or} \quad U(N, -).$$

Since  $\text{Gr}(X)$ ,  $\text{Gr}(Y/X)$ ,  $\text{Gr}(N)$ , ... have the property of affine paving, we conclude that  $\text{Gr}(Y)$  also has the property of affine paving.



**Lemma 6.2.** *Assume that  $Q$  is a quiver of type  $\tilde{A}$  or  $\tilde{D}$ ,  $M \in \text{rep}(Q)$  is the **regular quasi-simple** representation, then the Grassmannian  $\text{Gr}(\Phi(M))$  has an affine paving.*

*Proof.* The concept “quasi-simple” is defined in [6, Definition 15]; the concepts “preprojective”, “preinjective” and “regular” are defined in [6, 2.1.1]. It’s shown in [2, Section 9, Lemma 3] that the regular quasi-simple representation  $M$  have dimension vector smaller or equal to the minimal positive imaginary root, thus  $s_{P(e)}(M) \leq 2$  for the quiver of type  $\tilde{D}$  and  $s_{P(e)}(M) \leq 1$  for the quiver of type  $\tilde{A}$ .  $\square$

**Theorem 6.3.**

- (1) Assume that  $Q$  is a quiver of type  $\tilde{A}$  or  $\tilde{D}$ , then for any indecomposable representation  $M$ , the Grassmannian  $\text{Gr}(\Phi(M))$  has an affine paving;
- (2) Assume that  $Q$  is an affine quiver of type  $\tilde{E}$ , and  $\text{Gr}(\Phi(N))$  has an affine paving for any regular quasi-simple representation  $N \in \text{rep}(Q)$ . The Grassmannian  $\text{Gr}(\Phi(M))$  then has an affine paving for any indecomposable representation  $M$ .

For a regular quasi-simple representation  $Y$  of type  $\tilde{E}$ , it’s possible that there’s no short exact sequence

$$\eta : 0 \longrightarrow X \longrightarrow Y \longrightarrow S \longrightarrow 0$$

such that  $[S, X]^1 \leq 1$ . Then we can no longer use Theorem 3.2 or 3.3. Hence, the new methods are needed for this case.

REFERENCES

- [1] Klaus Bongartz. Critical simply connected algebras. *manuscripta mathematica*, 46(1):117–136, 1984.
- [2] William Crawley-Boevey. Lectures on representations of quivers. *unpublished notes*, 1992.
- [3] C. de Concini, George Lusztig, and C. Procesi. Homology of the zero-set of a nilpotent vector field on a flag manifold. *J. Am. Math. Soc.*, 1(1):15–34, 1988.
- [4] Corrado De Concini and Andrea Maffei. Paving Springer fibers for  $E7$ . *Proc. Indian Acad. Sci., Math. Sci.*, 132(2):21, 2022. Id/No 69.
- [5] Jens Niklas Eberhardt and Catharina Stroppel. Motivic Springer theory. *Indag. Math., New Ser.*, 33(1):190–217, 2022.
- [6] Giovanni Cerulli Irelli, Francesco Esposito, Hans Franzen, and Markus Reineke. Cell decompositions and algebraicity of cohomology for quiver Grassmannians. *Adv. Math.*, 379:48, 2021. Id/No 107544.
- [7] Ruslan Maksimau. Flag versions of quiver Grassmannians for Dynkin quivers have no odd cohomology over  $\mathbb{Z}$ , 2019.
- [8] Julia Sauter. Cell decompositions of quiver flag varieties for nilpotent representations of the oriented cycle. *arXiv preprint arXiv:1509.08026*, 2015.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF BONN, BONN, 53115, GERMANY,  
Email address: email:xx352229@mail.ustc.edu.cn




FIGURE 8. Special cases