

# AFFINE PAVING OF PARTIAL FLAG QUIVER VARIETY

XIAOXIANG ZHOU

ABSTRACT. In this article, we establish an affine paving for partial flag quiver varieties when the quiver is of Dynkin type. By copying results in [1, section 6] word by word, the same problem for affine quiver reduced to the case where the representation is regular quasi-simple. The idea of the proof mainly comes from [1], and the result is a natural continuation of [2].

## CONTENTS

|                               |   |
|-------------------------------|---|
| 1. Introduction               | 1 |
| 2. Preliminary Facts          | 1 |
| 2.1. Ext-vanishing properties | 1 |
| 3. Main Theorem               | 2 |
| 4. Application: Dynkin Case   | 2 |
| 5. Application: Affine Case   | 2 |
| References                    | 2 |

## 1. INTRODUCTION

Let  $Q$  be a quiver of Dynkin or affine type (without loops),  $X \in \text{Rep}(Q)$  be an quiver representation.<sup>1</sup> We are interested in three objects related to  $X \in \text{Rep}(Q)$ :

quiver Grassmannian  $\text{Gr}^{\text{KQ}}(X) := \{M_1 \mid M_1 \subseteq X\}$   
 partial flag variety  $d \geq 1$   $\text{Flag}_d(X) := \{0 \subseteq M_1 \subseteq \cdots M_d \subseteq X\}$   
 strict partial flag variety  $d \geq 2$   $\text{Flag}_{d,\text{str}}(X) := \{0 \subseteq M_1 \subseteq \cdots M_d \subseteq X \mid x.M_{i+1} \subseteq M_i\}$ <sup>2</sup>

It's easy to see that  $\text{Flag}_1(X) = \text{Gr}^{\text{KQ}}(X)$ . These geometrical objects can be divided into different pieces according to the dimension vectors of  $M_1, \dots, M_d$ , and each piece have its own natural (complex/Zarisky) topology. It was proved in [1] that  $\text{Gr}^{\text{KQ}}(X)$  have an affine paving, and in [2] that  $\text{Flag}_d(X)$  have the same property when  $Q$  is Dynkin quiver of type  $A/E$ . Here we go one step further, the results are concluded in the ???.

???Here is one table

The idea of proof is very simple: first, we view the partial flag quiver variety as the quiver Grassmannian of the more complicated quiver; then we establish the decomposition so that one may solve the problem by induction; finally we set a special way of decomposition for

<sup>1</sup>We fix the base field  $K = \mathbb{C}$  for convinience.

<sup>2</sup>for any  $x \in Q_1, i \in \{2, \dots, d\}$ .

each indecomposable module so that we can avoid meeting the bad decomposition. These contents are in Section ??, accordingly.

## 2. PRELIMINARY FACTS

### 2.1. Ext-vanishing properties.

## 3. MAIN THEOREM

### 4. APPLICATION: DYNKIN CASE

### 5. APPLICATION: AFFINE CASE

## REFERENCES

- [1] Giovanni Cerulli Irelli, Francesco Esposito, Hans Franzen, and Markus Reineke. Cell decompositions and algebraicity of cohomology for quiver grassmannians, 2019.
- [2] Ruslan Maksimau. Flag versions of quiver grassmannians for dynkin quivers have no odd cohomology over  $\mathbb{Z}$ , 2019.

SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF SCIENCE AND TECHNOLOGY OF CHINA, HEFEI, 230026, P.R. CHINA,

*Email address:* `email:xx352229@mail.ustc.edu.cn`